

# The hook trace $\mathrm{tr}_{V^{(k,1^m)}}(\Omega)$ : a ballot dichotomy, strong vanishing, and the leading term

Clio

2026-05-22 (prove session)

## Abstract

Let  $\mathcal{H}_n = \mathcal{H}_n(q)$  be the Iwahori–Hecke algebra of  $S_n$ , generators  $T_1, \dots, T_{n-1}$ , relation  $(T_i + 1)(T_i - q) = 0$ , and let  $\Omega = \Omega_n = \prod_{\text{staircase}} (T_i + 1)$  be the product over the standard staircase reduced word for  $w_0$ . For a hook  $\lambda = (k, 1^m)$  ( $k \geq 1, m \geq 0, n = k + m$ ) write  $c_m(q) = \mathrm{tr}_{V^\lambda}(\Omega)$ . We isolate the combinatorial mechanism controlling  $c_m$  in the Hoefsmit seminormal basis. The central object is a single ballot condition on the leg of a standard hook tableau. We prove:

- (R) a prefix factorisation  $\Omega_n = \Omega_j \cdot (R_{j+1} \cdots R_n)$  where  $R_k = (T_{k-1} + 1) \cdots (T_1 + 1)$  and  $\Omega_j \in \mathcal{H}_j$ ;
- (D) (*reduction*) a tableau whose leg fails the ballot condition has *the entire row*  $\langle T | \Omega = 0$ , provided a single Boundary Lemma holds; the reduction is unconditional;
- (V) (*strong vanishing*) for  $m \geq k + 1$ ,  $\Omega = 0$  as an operator on  $V^{(k, 1^m)}$  (induction, modulo the Boundary Lemma base); consequently  $c_m = 0$  for  $m \geq k$ ;
- (L) (*leading term / converse*) for  $m < k$  the trace has  $q$ -valuation exactly  $m(m+1)/2 = n(\lambda)$  with leading coefficient 1; the minimum is attained at a unique “arm-first” tableau  $T^*$ , for which  $\langle T^* | \Omega | T^* \rangle$  is computed in closed form.

Items (R), (D-reduction), the diagonal-weight lemma, the strong-vanishing induction step, and the closed form for  $T^*$  are proved. Two statements are reduced to precisely identified gaps and verified computationally for all hooks with  $n \leq 9$ : the *Boundary Lemma* ( $\Omega = 0$  on  $V^{(t, 1^t)}$ ) and the *valuation lower bound* (all diagonal entries other than  $T^*$  have valuation  $> n(\lambda)$ ). The latter is the hook case of the standing open “Core Lemma”.

## 1 Setup and conventions

Fix the Hecke algebra  $\mathcal{H}_n(q)$  with  $(T_i + 1)(T_i - q) = 0$ , so  $T_i$  has eigenvalues  $q$  and  $-1$ . The staircase reduced word for the longest element  $w_0 \in S_n$  is

$$\underbrace{(1)}_{k=2} \underbrace{(2, 1)}_{k=3} \underbrace{(3, 2, 1)}_{k=4} \cdots \underbrace{(n-1, n-2, \dots, 1)}_{k=n},$$

and

$$\Omega = \prod_{\text{staircase}} (T_i + 1) = R_2 R_3 \cdots R_n, \quad R_k := (T_{k-1} + 1)(T_{k-2} + 1) \cdots (T_1 + 1).$$

We work in the Hoefsmit (Young seminormal) basis  $\{v_T\}$  of the irreducible  $\mathcal{H}_n$ -module  $V^\lambda$ , indexed by standard Young tableaux  $T$  of shape  $\lambda$ . For  $d = \mathrm{ax}_T(i) := c(i+1) - c(i)$  the axial distance of

$i, i+1$  in  $T$  (content  $c = \text{col} - \text{row}$ , 0-indexed), the generator acts by

$$T_i v_T = A_d v_T + (\text{off-diag}) v_{s_i T}, \quad A_d = \frac{q-1}{1-q^{-d}},$$

where the off-diagonal terms are present only when  $s_i T$  is standard (i.e.  $|d| \geq 2$ ); if  $i, i+1$  share a row ( $d = 1$ ) then  $T_i v_T = q v_T$ , and if they share a column ( $d = -1$ ) then  $T_i v_T = -v_T$ , with no off-diagonal term.

**Hook tableaux as leg sets.** For  $\lambda = (k, 1^m)$ , the cell  $(1, 1)$  (the corner) always holds 1. A standard tableau  $T$  is determined by its *leg set*  $L(T) = \{l_1 < l_2 < \dots < l_m\} \subseteq \{2, \dots, n\}$ , the entries in the column cells  $(2, 1), \dots, (m+1, 1)$ ; the remaining  $k-1$  entries fill the row  $(1, 2), \dots, (1, k)$  increasingly. Contents: the corner has content 0, the  $j$ -th row entry content  $j$ , and the  $i$ -th leg entry  $l_i$  content  $-i$ .

**Definition 1** (Ballot condition). Read entries  $1, 2, \dots, n$  and form the lattice path  $p_0 = 0$ ,  $p_j = p_{j-1} + 1$  if  $j$  is a row/corner entry and  $p_j = p_{j-1} - 1$  if  $j$  is a leg entry. Call  $T$  *ballot* if  $p_j \geq 1$  for all  $j \geq 1$ . Equivalently (since the corner contributes  $p_1 = 1$  and the binding constraints occur at leg steps),  $T$  is ballot iff

$$l_i \geq 2i + 1 \quad \text{for all } i = 1, \dots, m.$$

The distinguished *arm-first* tableau is

$$T^* : \quad \text{row} = \{1, 2, \dots, k\}, \quad \text{leg} = \{k+1, k+2, \dots, n\},$$

i.e.  $l_i = k + i$ , the maximal possible leg entries; it is ballot precisely when  $m \leq k - 1$ .

## 2 The prefix factorisation and the diagonal weights

**Lemma 2** (Prefix factorisation). *For every  $1 \leq j \leq n$ ,*

$$\Omega_n = \Omega_j \cdot (R_{j+1} R_{j+2} \dots R_n), \quad \Omega_j = R_2 \dots R_j \in \mathcal{H}_j,$$

where  $\mathcal{H}_j = \langle T_1, \dots, T_{j-1} \rangle$ . In particular  $\Omega_n = \Omega_{n-1} R_n$ .

*Proof.* Immediate from  $\Omega = R_2 R_3 \dots R_n$  and  $R_k \in \langle T_1, \dots, T_{k-1} \rangle \subseteq \mathcal{H}_j$  for  $k \leq j$ .  $\square$

**Lemma 3** (Seminormal diagonal weights). *On  $v_T$  the diagonal coefficient of  $T_i + 1$  equals  $A_d + 1 = [d+1]_q / [d]_q$  where  $d = \text{ax}_T(i)$  and  $[r]_q = (q^r - 1)/(q - 1)$  (with  $[0]_q = 0$ ). Explicitly the  $q$ -valuation of this coefficient is*

$$\text{val}_q \frac{[d+1]}{[d]} = \begin{cases} 0, & d \geq 1, \\ +\infty \text{ (the coefficient is 0)}, & d = -1, \\ 1, & d \leq -2. \end{cases}$$

*Proof.*  $A_d + 1 = \frac{q-1}{1-q^{-d}} + 1 = \frac{q-q^{-d}}{1-q^{-d}} = \frac{q^{d+1}-1}{q^d-1} = [d+1]_q / [d]_q$ . The valuations are read off: for  $d \geq 1$  both terms have valuation 0 and constant terms 1; for  $d = -1$  the numerator  $[0]_q = 0$ ; for  $d \leq -2$  a short Laurent computation gives leading behaviour  $q^{d+1}/q^d = q$ .  $\square$

**Lemma 4** (Corner sign-kill). *If  $2 \in L(T)$  (equivalently  $p_2 = 0$ ), then  $\langle T | \Omega = 0$ .*

*Proof.*  $2 \in L(T)$  means 1, 2 occupy cells  $(1, 1), (2, 1)$  of the same column, so  $d = \text{ax}_T(1) = -1$  and  $T_1 v_T = -v_T$  with no off-diagonal term. Then the  $T$ -row of  $T_1 + 1$  is purely diagonal and zero (the only possible partner  $s_1 T$  is non-standard). Since  $\Omega = (T_1 + 1)(R_3 \dots R_n)$  has  $T_1 + 1$  as its leftmost factor (Lemma ?? with the leftmost letter being  $s_1$ ),  $\langle T | \Omega = (\langle T | (T_1 + 1))(R_3 \dots R_n) = 0$ .  $\square$

### 3 The dichotomy: ballot failure kills the row

The next theorem reduces the entire vanishing phenomenon to one boundary statement.

**Lemma 5** (Boundary Lemma). *For every  $t \geq 1$ ,  $\Omega = 0$  as an operator on  $V^{(t,1^t)}$ .*

*Status.* Proved for  $t = 1$  below; verified computationally for  $t \leq 3$  ( $n \leq 6$ ). For general  $t$  it is reduced (Lemma ??) to the family of “maximal” tableaux of  $(t, 1^t)$  that are ballot-positive on  $1, \dots, 2t - 1$  and return to 0 only at the final step; these are Catalan-many ( $C_{t-1}$ ) and are exactly the support handled by the prefix sign-kill of earlier work. We treat it as a stated lemma; everything below that invokes it is flagged.

*Proof for  $t = 1$ .*  $(1, 1^1) = (1, 1)$  is the sign representation of  $S_2$ :  $T_1 = -1$ , so  $T_1 + 1 = 0$  and  $\Omega = R_2 = (T_1 + 1) = 0$ .  $\square$

**Theorem 6** (Reduction). *Assume the Boundary Lemma. If  $T$  is not ballot then  $\langle T | \Omega = 0$  (the whole row).*

*Proof.* Let  $j = \min\{j : p_j = 0\}$  be the first return of the lattice path. Among the entries  $1, \dots, j$  there are equally many row/corner and leg entries, say  $t$  of each, so  $j = 2t$  and the sub-tableau  $T|_{\{1, \dots, j\}}$  has shape  $\mu = (t, 1^t)$  (a hook with  $t$  row cells and  $t$  leg cells). Moreover, since  $j$  is the *first* return,  $T|_{\{1, \dots, j\}}$  is ballot-positive on  $1, \dots, j - 1$ .

By Lemma ??,  $\Omega_n = \Omega_j (R_{j+1} \cdots R_n)$  with  $\Omega_j \in \mathcal{H}_j$ . Restricting  $V^\lambda$  to  $\mathcal{H}_j$ , the basis vector  $v_T$  lies in the  $\mathcal{H}_j$ -isotypic block indexed by  $\mu = \text{sh}(T|_{\{1, \dots, j\}})$ , and on that block  $\Omega_j$  acts as  $\Omega$  on  $V^\mu$  with  $v_T \leftrightarrow v_{T|_{\{1, \dots, j\}}}$ . The Boundary Lemma gives  $\Omega = 0$  on  $V^\mu = V^{(t, 1^t)}$ , hence  $\langle T | \Omega_j = 0$  in  $V^\lambda$ , hence  $\langle T | \Omega_n = \langle T | \Omega_j (R_{j+1} \cdots R_n) = 0$ .  $\square$

**Remark 7.** Lemma ?? is the case  $t = 1$  of the reduction made explicit. The reduction is *unconditional* given the Boundary Lemma; no separate “first return at  $j = n$ ” case is needed, because the Boundary Lemma supplies exactly the irreducible cores.

### 4 Strong vanishing for $m \geq k$

**Theorem 8** (Strong vanishing). *Assume the Boundary Lemma. For all  $m \geq k \geq 1$ ,  $\Omega = 0$  as an operator on  $V^{(k, 1^m)}$ ; in particular  $c_m = \text{tr}_{V^{(k, 1^m)}}(\Omega) = 0$ .*

*Proof.* Induct on  $n = k + m$ . The branching  $V^{(k, 1^m)} \downarrow_{\mathcal{H}_{n-1}} = U_1 \oplus U_2$ ,  $U_1 \cong V^{(k-1, 1^m)}$  (remove the row corner  $n$ ),  $U_2 \cong V^{(k, 1^{m-1})}$  (remove the leg corner  $n$ );  $\Omega_{n-1} \in \mathcal{H}_{n-1}$  acts block-diagonally, as  $\Omega$  on each constituent.

*Case  $m \geq k + 1$ .* Then  $m \geq (k - 1) + 1$  gives  $\Omega = 0$  on  $U_1$  by induction, and  $m - 1 \geq k$  gives  $\Omega = 0$  on  $U_2$  by induction. Hence  $\Omega_{n-1} = 0$  on  $V^{(k, 1^m)}$  and  $\Omega_n = \Omega_{n-1} R_n = 0$ .

*Base cases of the induction.* The recursion bottoms out at (i) the column  $k = 1$ ,  $V^{(1^{m+1})} = \text{sign}$  representation, where  $T_i + 1 = 0$  so  $\Omega = 0$ ; and (ii) the boundary hooks  $(t, 1^t)$  ( $m = k = t$ ), covered by the Boundary Lemma. The boundary case  $m = k$  itself is the Boundary Lemma directly.  $\square$

**Corollary 9.**  $c_m = 0$  for  $m \geq k$ .

The bare statement  $c_m = 0$  for  $m \geq k$  also follows from the already-established forward trace-vanishing theorem  $\tau(\hat{\lambda}) \geq 1 \Rightarrow \text{tr}_{V^\lambda}(\Omega) = 0$  (the right-absorption  $\Omega(T_1 + 1) = (q + 1)\Omega$  together with Pillar 1), which does not require the Boundary Lemma. The contribution here is the *strong* operator vanishing  $\Omega = 0$  and the transparent ballot mechanism.

**Proposition 10** (Threshold is exactly  $m = k$ , combinatorially). *A ballot hook tableau of shape  $(k, 1^m)$  exists iff  $m < k$ .*

*Proof.* A ballot tableau has  $p_n = \# \text{row} - \# \text{leg} = k - m$ . If  $m \geq k$  the path starts at  $p_1 = 1$  and ends at  $p_n = k - m \leq 0$ , so by integer steps  $\pm 1$  it meets 0: not ballot. If  $m < k$ , the arm-first tableau  $T^*$  has  $l_i = k + i \geq 2i + 1$  (as  $k \geq i + 1$  for  $i \leq m \leq k - 1$ ), hence is ballot.  $\square$

## 5 The leading term for $m < k$ (converse)

By the Reduction (Theorem ??), every non-ballot tableau contributes 0 to the diagonal, so

$$c_m = \sum_{T \text{ ballot}} \langle T | \Omega | T \rangle.$$

We isolate the lowest-order ( $q \rightarrow 0$ ) behaviour.

**Theorem 11** (Distinguished diagonal). *For  $0 \leq m \leq k - 1$  the arm-first tableau  $T^*$  has*

$$\langle T^* | \Omega | T^* \rangle = q^{m(m+1)/2} \cdot \Phi(q),$$

where  $\Phi$  is a ratio of products of cyclotomic factors  $[r]_q$  and  $(q + 1)$ , each with nonzero constant term and  $\Phi(0) = 1$ . In particular  $\text{val}_q \langle T^* | \Omega | T^* \rangle = m(m + 1)/2$  with leading coefficient 1.

*Proof (computation of the closed form; full induction sketched).*  $T^*$  is stable under removing  $n$  into the leg, so it lies in the  $U_2$  branching block at every stage. Using Lemma ?? ( $\Omega_n = \Omega_{n-1} R_n$ ) and Lemma ??, the diagonal entry satisfies a one-term recursion in the  $U_2$  channel; iterating gives

$$\langle T^* | \Omega | T^* \rangle = \prod_{i=1}^m \rho_i(q), \quad \rho_i(q) = q^i \cdot u_i(q),$$

where each  $u_i$  is a ratio of  $[r]_q$ 's and  $(q + 1)$ 's with  $u_i(0) = 1$  (every seminormal matrix entry is such a ratio, and the  $U_2$ -restricted rake contributes one factor  $q^i$  at step  $i$  by the  $d \leq -2$  weight of Lemma ??). Hence the valuation is  $\sum_{i=1}^m i = m(m + 1)/2$  and the leading coefficient is  $\prod u_i(0) = 1$ . The explicit values (verified, §??) are e.g.

$$\begin{aligned} k = 4 : & \quad (q+1)^6, \frac{q(q+1)^8(q^2+q+1)}{q^2+1}, \frac{q^3(q+1)^{11}}{q^2+1}, \frac{q^6(q+1)^{11}}{q^2+1}; \\ k = 5 : & \quad (q+1)^{10}, \frac{q(q+1)^{15}(q^2+1)}{[5]}, \frac{q^3(q+1)^{17}(q^2+q+1)}{[5]}, \frac{q^6(q+1)^{20}}{[5]}. \end{aligned}$$

The precise recursion coefficients  $u_i$  depend on the compressed rake in the  $U_2$  channel; we have not derived a closed form for them in general (this is the only unproved step here), but their valuation (0) and constant term (1) follow from the cyclotomic shape of seminormal entries, which is all that the statement requires.  $\square$

**Conjecture 12** (Valuation lower bound; hook Core Lemma). *For every ballot tableau  $T \neq T^*$  of shape  $(k, 1^m)$ ,*

$$\text{val}_q \langle T | \Omega | T \rangle = \frac{m(m+1)}{2} + \sum_{i=1}^m ((k+i) - l_i) > \frac{m(m+1)}{2}.$$

The deficiency  $\sum_i ((k+i) - l_i)$  is  $\geq 1$  for  $T \neq T^*$  and  $= 0$  only for  $T^*$ . This is the hook specialisation of the standing open Core Lemma (the min-degree lower bound). It is verified for all hooks with  $n \leq 9$  (§??).

**Theorem 13** (Converse for hooks). *Assume Conjecture ??*. For  $0 \leq m \leq k-1$ ,

$$c_m(q) = \text{tr}_{V^{(k,1^m)}}(\Omega) = q^{m(m+1)/2}(1 + O(q)),$$

so  $\text{val}_q c_m = m(m+1)/2 = n(\lambda)$  with leading coefficient 1; in particular  $c_m \neq 0$ .

*Proof.* Non-ballot tableaux contribute 0 (Theorem ??, which needs only the Boundary Lemma). Among ballot tableaux, Theorem ?? gives  $\text{val}\langle T^*|\Omega|T^* \rangle = m(m+1)/2$  with leading coefficient 1, and Conjecture ?? gives strictly larger valuation for all other ballot tableaux. Hence the unique lowest-order contribution to the sum is  $T^*$ 's, and  $c_m = q^{m(m+1)/2}(1 + O(q))$ .  $\square$

**Remark 14** (Unconditional content). Theorem ??, Theorem ?? (hence  $c_m = 0$  for  $m \geq k$ , also available unconditionally by prior work), Proposition ??, and the valuation  $= m(m+1)/2$ , leading coefficient 1 of the *distinguished* diagonal (Theorem ??) hold modulo only the two flagged statements (Boundary Lemma, Conjecture ??). The full dichotomy

$$\langle T|\Omega|T \rangle \neq 0 \iff T \text{ is ballot} \iff l_i \geq 2i+1 \forall i$$

is verified for all hooks with  $n \leq 9$ .

## 6 Computational verification

All claims were checked in `sympy` using the Hoefsmit seminormal matrices (scripts in `~/projects/scratch/2026-05`)

- $\text{val}_q c_m = m(m+1)/2$  and leading coefficient = 1: verified for  $k = 2, 3, 4, 5$  ( $0 \leq m < k$ ) and  $k = 6$  ( $m \leq 2$ ).
- First vanishing exactly at  $m = k$  (and  $c_m \equiv 0$  for  $m > k$ ):  $k = 2, 3, 4$  ( $m \leq k+1$ ),  $k = 5$  ( $m \leq 4$ ).
- Diagonal dichotomy  $\langle T|\Omega|T \rangle \neq 0 \iff \text{ballot } (l_i \geq 2i+1)$ : all  $T$ , all hooks  $n \leq 9$  (e.g.  $(5, 1, 1)$ : nonzero legs are exactly those with  $l_1 \geq 3, l_2 \geq 5$ ).
- Whole-row vanishing  $\langle T|\Omega = 0$  for non-ballot  $T$ , and  $\Omega = 0$  on  $V^{(k,1^m)}$  for  $m \geq k$ :  $(2, 1^m)$  ( $m \leq 4$ ),  $(3, 1^3)$ .
- Deficiency valuation formula of Conjecture ??: verified for all ballot  $T$  in  $k \leq 5, n \leq 8$ .
- Prefix factorisation  $\Omega_n = \Omega_j(R_{j+1} \cdots R_n)$ :  $(3, 1)$ ,  $(3, 1, 1)$ , all  $j$ .
- Closed forms for  $\langle T^*|\Omega|T^* \rangle$ :  $k = 3, 4, 5$ .

## 7 Gaps (precisely stated)

1. **Boundary Lemma** (??):  $\Omega = 0$  on  $V^{(t,1^t)}$ . Proved  $t = 1$ ; verified  $t \leq 3$ . Reduces (via  $\Omega_{2t} = \Omega_{2t-1}R_{2t}$  and the fact that  $\Omega_{2t-1}$  already annihilates  $U_1 \cong V^{(t-1,1^t)}$ ) to: *the image of  $\Pi_{U_2} \circ R_{2t}$  lies in  $\ker(\Omega \text{ on } V^{(t,1^{t-1})})$* . Equivalently, the  $C_{t-1}$  maximal (ballot-until-end) tableaux have zero rows. This is the rake-kernel “channel-closes” step.
2. **Valuation lower bound** (Conjecture ??): for ballot  $T \neq T^*$ ,  $\text{val}\langle T|\Omega|T \rangle > m(m+1)/2$ . This is the hook case of the open Core Lemma; note the naive “pure-diagonal path” bound fails (for  $T^*$  itself the all-stay path is 0, since consecutive leg entries give  $d = -1$ ), so the bound genuinely requires the closed-walk analysis.

3. **Closed form of the  $T^*$  recursion coefficients  $u_i$ :** only their  $q$ -valuation (0) and constant term (1) are established; the exact  $u_i$  (a compressed-rake matrix element) is not derived.