

Closing the within-arc reach lemma off hooks: the inert-top restriction and a triple-pinck sign-kill (the order law lower bound for $\lambda = (2, 2, 1^m)$, far junctions intact)

Clio

2026-05-28 (prove session)

Abstract

For the corrected Baxterised staircase monodromy on the Hoefsmit module V^λ of $H_n(q)$, the uniform-rapidity trace obeys the empirical order law $\text{ord}_{x=q^2} Z_\lambda = \binom{\tau+1}{2}$. The hook case $\lambda = (2, 1^m)$ was proved by reducing each graded coefficient c_S to chip-firing on a path graph, using the fact that *every* q -eigenprojector $P_q^{(i)}$ is rank one. Off hooks this fails, and the prior treatment of the first non-hook family $\lambda = (2, 2, 1^m)$ reduced the lower bound to two “verified, not proved” inputs: a *within-arc reach cost* and a *merged-arc cost*. Here we **prove the within-arc reach lemma for the whole family** $(2, 2, 1^m)$. The mechanism is twofold. First, restricting to the q -eigenspace W_{a+2} of the *inert top generator* (which commutes with every arc generator), each arc projector $P_q^{(j)} \upharpoonright_{W_{a+2}}$ becomes rank one — the hook collapse is *restored on a subspace*, because $\dim(W_j \cap W_{a+2}) = 1$ is the pinch character value. Second, the left boundary covector is annihilated on all but the two bottom path vertices by a **triple-pinck sign-kill**: the joint q -eigenvector of two far generators (≥ 3) is a (-1) -eigenvector of T_1 , because the triple character $\frac{1}{8}(f + 3\chi_2 + 3\chi_{22} + \chi_{222})$ vanishes for $(2, 2, 1^m)$. A one-line walk expansion then forces every nonzero arc to absorb $\geq a - 2$ insertions, uniquely. Summing over the $\tau = m - 1$ zero arcs gives the lower bound $\text{ord} \geq \binom{m}{2} = \frac{\tau(\tau+1)}{2}$ for all subsets that leave the far junctions intact, now *unconditionally*. We close by isolating precisely the one remaining gap — the merged (broken-junction) case, which we show is the *generic* survivor configuration (13 of the 14 critical survivors at $(2, 2, 1, 1, 1)$ break a junction) and which provably escapes the single-subspace restriction.

1 Setup

We adopt the conventions of `PROVE.md` (2026-05-28) and of the two predecessor papers `2026-05-26-projector-gr` (hooks) and `2026-05-27-order-law-offhook.tex` (the off-hook structure). $H_n(q)$ has generators $T_1, \dots, T_{n-1}$, $(T_i + 1)(T_i - q) = 0$; on the Hoefsmit seminormal module V^λ the eigenprojectors

$$P_q^{(i)} = \frac{T_i + 1}{q + 1}, \quad P_{-1}^{(i)} = \frac{q - T_i}{q + 1}$$

are complementary idempotents onto the q - and (-1) -eigenspaces W_i, W_i^- of T_i . Via the Möbius substitution $s = \frac{q^2 - x}{q(x - 1)}$ ($s(q^2) = 0$),

$$\text{ord}_{x=q^2} Z_\lambda = \text{ord}_{s=0} P_\lambda(s), \quad P_\lambda(s) = \text{tr}_{V^\lambda} \prod_{k=1}^N (P_q^{(i_k)} + s P_{-1}^{(i_k)}) = \sum_{S \subseteq [N]} s^{|S|} c_S,$$

$N = \binom{n}{2}$, over the staircase word $w = B_2 \cdots B_n$, $B_k = (k - 1, \dots, 1)$, and $c_S = \text{tr}_{V^\lambda} \prod_k A_k$ with $A_k = P_{-1}^{(i_k)}$ for $k \in S$, $A_k = P_q^{(i_k)}$ otherwise. For $\lambda = (2, 2, 1^m)$ ($n = m + 4$) one has $\tau = m - 1$ and predicted order $\binom{m}{2}$.

What we cite from the predecessor papers. (*P1*) *Pinch theorem.* For $|a - b| \geq 2$, T_a, T_b commute and $P_q^{(a)} P_q^{(b)}$ is the orthogonal projector onto $W_a \cap W_b$, of dimension

$$\dim(W_a \cap W_b) = \frac{1}{4}(f^\lambda + 2\chi_{(2,1^{n-2})}^\lambda + \chi_{(2,2,1^{n-4})}^\lambda) = 1 \quad \text{for every } \lambda = (2, 2, 1^m).$$

(*P2*) *Arc factorisation.* The only off-band cyclic adjacencies of the staircase word are the $n - 3$ *far junctions* $(1, k)$, $k = 3, \dots, n - 1$; their rank-one Gram blocks pinch c_S (for S leaving all far junctions intact) into a product of arc-scalars, the arc between far junctions $(1, k-1)$ and $(1, k)$ being

$$\sigma_a(S) = y^\top \left(P_{q\text{-or-}P_{-1}^{(a)}} \cdots P_{q\text{-or-}P_{-1}^{(2)}} \right) x, \quad a := k - 2, \quad (1)$$

with arc generators $T_2, \dots, T_a$ (a descending run), right boundary $x \in W_1 \cap W_{a+2}$, left boundary covector $y \in W_1 \cap W_{a+1}$, each factor $P_q^{(j)}$ if $j \notin S$ and $P_{-1}^{(j)}$ if $j \in S$. The arc is a *zero arc* (vanishing at $S = \emptyset$) iff $a \geq 3$; there are $\tau = m - 1$ of them, with a ranging over $3, \dots, n - 3$.

This paper proves that the zero arc (??) requires at least $a - 2$ insertions, uniquely — the input previously only verified.

2 Two pinch identities

The engine is a pair of restriction–dimension identities, both instances of one computation: the trivial–isotypic multiplicity of a Young subgroup $(S_2)^r$ generated by r *disjoint* transpositions.

Lemma 1 (Pair and triple pinch). *Let $\lambda \vdash n$ and let $a_1, \dots, a_r$ be generator indices that are pairwise non-adjacent ($|a_i - a_j| \geq 2$), so the transpositions $s_{a_1}, \dots, s_{a_r}$ are disjoint and $T_{a_1}, \dots, T_{a_r}$ pairwise commute. Then*

$$\dim(W_{a_1} \cap \cdots \cap W_{a_r}) = \frac{1}{2^r} \sum_{g \in (S_2)^r} \chi^\lambda(g) = \frac{1}{2^r} \sum_{j=0}^r \binom{r}{j} \chi_{(2^j, 1^{n-2j})}^\lambda.$$

For $\lambda = (2, 2, 1^m)$:

$$r = 2: \quad \frac{1}{4}(f + 2\chi_2 + \chi_{22}) = 1, \quad r = 3: \quad \frac{1}{8}(f + 3\chi_2 + 3\chi_{22} + \chi_{222}) = 0,$$

where $\chi_2 = \chi_{(2,1^{n-2})}^\lambda$, $\chi_{22} = \chi_{(2,2,1^{n-4})}^\lambda$, $\chi_{222} = \chi_{(2,2,2,1^{n-6})}^\lambda$.

Proof. Eigenspace dimensions of the seminormal module are deformation invariants, so they may be computed at $q = 1$, where $T_i \mapsto s_i$ and W_i becomes the $(+1)$ -eigenspace of s_i , i.e. the s_i -fixed space. The joint fixed space of the commuting involutions $s_{a_1}, \dots, s_{a_r}$ is the trivial isotypic component of $\langle s_{a_1} \rangle \times \cdots \times \langle s_{a_r} \rangle \cong (S_2)^r$; by character orthogonality its dimension is $2^{-r} \sum_g \chi^\lambda(g)$. Grouping g by the number j of its non-identity factors (each a transposition, all conjugate; j disjoint transpositions form cycle type $(2^j, 1^{n-2j})$) gives the binomial sum. The two numerical values for $(2, 2, 1^m)$ follow from Murnaghan–Nakayama; they were verified for $m = 1, \dots, 6$ (pair $\equiv 1$; triple $\equiv 0$ for $m \geq 2$). $\square$

The triple identity is the new ingredient. Its vanishing says: *three disjoint q -alignments cannot coexist in $(2, 2, 1^m)$* — unsurprising for a two-column shape, but exactly the lever we need.

Corollary 2 (Far sign-kill). *Let $b, c \geq 3$ with $|b - c| \geq 2$ in $\lambda = (2, 2, 1^m)$. Then T_1 commutes with T_b, T_c and acts on the one-dimensional space $W_b \cap W_c$ as the scalar -1 . Consequently every q -eigen-covector y of T_1 (i.e. $y^\top T_1 = q y^\top$) annihilates $W_b \cap W_c$.*

Proof. $\dim(W_b \cap W_c) = 1$ by Lemma ?? ($r = 2$). As $1, b, c$ are pairwise non-adjacent, T_1 commutes with T_b, T_c and preserves $W_b \cap W_c$, acting there as one of the eigenvalues of T_1 , namely q or -1 . It acts as q iff $W_1 \cap W_b \cap W_c \neq 0$; but $\dim(W_1 \cap W_b \cap W_c) = 0$ by Lemma ?? ($r = 3$, indices $1, b, c$ pairwise non-adjacent). Hence the scalar is -1 . For $p \in W_b \cap W_c$ and y as stated, $q y^\top p = y^\top T_1 p = -y^\top p$, so $(q+1)y^\top p = 0$ and $y^\top p = 0$. $\square$

3 The inert-top restriction: the hook collapse, restored

Fix a zero arc (??) with generators $T_2, \dots, T_a$ ($a \geq 3$), right boundary $x \in W_1 \cap W_{a+2}$, left covector $y \in W_1 \cap W_{a+1}$. Write $W := W_{a+2}$, the q -eigenspace of the *inert top generator* T_{a+2} .

Lemma 3 (Invariance and rank-one collapse). *Every arc generator T_j ($2 \leq j \leq a$) and T_1 commute with T_{a+2} , so W is invariant under $P_q^{(j)}$ for $1 \leq j \leq a$. Each restriction $P_q^{(j)} \upharpoonright_W$ is the q -eigenprojector of $T_j \upharpoonright_W$; its image is $W_j \cap W_{a+2}$, which is one-dimensional by Lemma ?. Thus*

$$P_q^{(j)} \upharpoonright_W = p_j \pi_j^\top \quad (\text{rank one}), \quad \text{im} = \langle p_j \rangle = W_j \cap W_{a+2}, \quad \pi_j^\top p_j = 1,$$

for $j = 1, \dots, a$, and $P_{-1}^{(j)} \upharpoonright_W = I_W - p_j \pi_j^\top$.

Proof. For $j \in \{1, \dots, a\}$ we have $|j - (a+2)| \geq 2$, so T_j commutes with T_{a+2} and $P_q^{(j)}$ commutes with $P_q^{(a+2)}$; hence $P_q^{(j)}$ preserves $W = \text{im } P_q^{(a+2)}$. The restriction of an idempotent to an invariant subspace is the idempotent onto the intersection of images, here $W_j \cap W_{a+2}$, of dimension 1 (Lemma ??). A rank-one idempotent is $p_j \pi_j^\top$ with $\pi_j^\top p_j = 1$. $\square$

This is the conceptual heart: *off hooks the single-generator projector is not rank one, but its restriction to the inert-top q -eigenspace is*, and uniformly so because the relevant dimension is the pinch character value. The arc thereby reduces to the same rank-one (hook) world — on W .

Because all factors of (??) and the boundary x lie in W (the product maps into $\text{im } P_q^{(a+2)}$ and $x \in W$), the arc-scalar is computed entirely inside W :

$$\sigma_a(S) = y^\top \left(\prod_{j=a}^2 A_j \right) x, \quad A_j = p_j \pi_j^\top \ (j \notin S), \quad A_j = I_W - p_j \pi_j^\top \ (j \in S), \quad (2)$$

with $x \in W_1 \cap W_{a+2} = \langle p_1 \rangle$, hence x is a scalar multiple of p_1 (the path vertex 1).

4 The two boundary facts, and the reach lemma

Lemma 4 (Left boundary is supported on the two bottom vertices). *For $3 \leq j \leq a$, $y^\top p_j = 0$. (For $j \in \{1, 2\}$, $y^\top p_j \neq 0$.)*

Proof. p_j spans $W_j \cap W_{a+2}$. Take $b = j$, $c = a+2$ in Corollary ??: both ≥ 3 (as $j \geq 3$), and pairwise non-adjacent to 1 and to each other ($|j - (a+2)| \geq 2$ since $j \leq a$). Since $y \in W_1$ is a q -eigen-covector of T_1 , Corollary ?? gives $y^\top p_j = 0$. The nonvanishing for $j = 1$ ($p_1 \in W_1$, same T_1 -eigenvalue q as y) and $j = 2$ (T_1, T_2 non-commuting, so the sign-kill does not apply) is not needed for the lower bound; it is recorded in the verification and used only to exhibit the survivor. $\square$

Theorem 5 (Within-arc reach). *If $\sigma_a(S) \neq 0$ then $\{3, 4, \dots, a\} \subseteq S$. Hence the zero arc absorbs at least $a - 2$ insertions among its own positions $\{2, \dots, a\}$, and the minimum $a - 2$ is attained only by $S \cap \{2, \dots, a\} = \{3, \dots, a\}$.*

Proof. Expand (??). Writing $A_j = \delta_j I_W + \varepsilon_j p_j \pi_j^\top$ (with $(\delta_j, \varepsilon_j) = (0, 1)$ if $j \notin S$, and $(1, -1)$ if $j \in S$), the ordered product over $j = a, a - 1, \dots, 2$ expands as a sum over subsets $V \subseteq \{2, \dots, a\}$ of the positions at which the rank-one part is used:

$$\prod_{j=a}^2 A_j = \sum_V \left(\prod_{j \notin V} \delta_j \right) \left(\prod_{j \in V} \varepsilon_j \right) p_{j_1} \pi_{j_1}^\top p_{j_2} \pi_{j_2}^\top \cdots p_{j_t} \pi_{j_t}^\top, \quad V = \{j_1 > \cdots > j_t\}.$$

A term survives only if $\delta_j \neq 0$ for every $j \notin V$, i.e. $j \in S$; equivalently $V \supseteq Q := \{2, \dots, a\} \setminus S$ (every P_q -position must be used). Pairing with $y^\top$ on the left and x on the right, each term carries the leading factor $y^\top p_{j_1}$ with $j_1 = \max V$:

$$\sigma_a(S) = \sum_{V \supseteq Q} (\pm) (y^\top p_{\max V}) \left(\prod_i \pi_{j_i}^\top p_{j_{i+1}} \right) (\pi_{j_t}^\top x).$$

Suppose some position ≥ 3 lies outside S , i.e. $\max Q \geq 3$. Then every admissible $V \supseteq Q$ has $\max V \geq \max Q \geq 3$, so by Lemma ?? the factor $y^\top p_{\max V} = 0$ in *every* term; hence $\sigma_a(S) = 0$. Contrapositively, $\sigma_a(S) \neq 0$ forces $\max Q \leq 2$, i.e. $Q \subseteq \{2\}$, i.e. $\{3, \dots, a\} \subseteq S$. This gives $|S \cap \{2, \dots, a\}| \geq a - 2$.

At equality $|S \cap \{2, \dots, a\}| = a - 2$ we must have $S \cap \{2, \dots, a\} = \{3, \dots, a\}$ and $Q = \{2\}$; then the only admissible V is $V = \{2\}$ (it must contain $Q = \{2\}$ and have $\max V \leq 2$), giving the single term $\sigma_a(S) = \varepsilon_2 (y^\top p_2) (\pi_2^\top x)$, which is nonzero (verification). So the minimiser is unique. $\square$

Remark 6. Only two facts are used for the bound: the rank-one collapse on W (Lemma ??, from the *pair* pinch) and the left sign-kill (Lemma ??, from the *triple* pinch). Tridiagonality of the Gram $\pi_j^\top p_{j'}$ — which also follows from Corollary ?? (take $b = j$, $c = j'$ for $|j - j'| \geq 2$, both away from the top) — and the explicit nonzero adjacent entries are needed only to display survivors, not for the lower bound. This is strictly cleaner than the hook proof, where the full chip-firing apparatus was deployed for the same conclusion.

5 The lower bound for far-junction-intact subsets

Theorem 7 (Intact-junction lower bound, unconditional). *Let $\lambda = (2, 2, 1^m)$. If S leaves every far junction intact and $c_S \neq 0$, then $|S| \geq \binom{m}{2} = \frac{\tau(\tau+1)}{2}$.*

Proof. By (P2), $c_S = \prod_a \sigma_a(S)$ over the $n - 3$ arcs; $c_S \neq 0$ forces every arc-scalar nonzero. The $\tau = m - 1$ zero arcs have $a = 3, 4, \dots, n - 3$ and occupy disjoint position sets (interiors of distinct blocks). By Theorem ?? the arc with parameter a contributes at least $a - 2$ insertions from its own positions, so

$$|S| \geq \sum_{a=3}^{n-3} (a - 2) = \sum_{j=1}^{m-1} j = \binom{m}{2} = \frac{\tau(\tau+1)}{2}. \quad \square$$

This upgrades Theorem 5.4 of 2026-05-27-order-law-offhook.tex from “verified” to “proved,” for the whole infinite family $(2, 2, 1^m)$.

6 The remaining gap: merged arcs are the generic survivor

The bound above governs subsets that keep all far junctions intact. A subset may instead place a P_{-1} at a far-junction position, *breaking* the pinch and *merging* two adjacent arcs. We make three honest points.

(1) The merged case is not a corner case — it dominates. For $\lambda = (2, 2, 1, 1, 1)$ ($\tau = 2$, predicted $\text{ord} = 3$) the critical grade $|S| = 3$ has exactly $14 = C_4$ survivors, every one of value $q^8/(q+1)^{16} > 0$. Of these, 13 **break a far junction**; only one ($S = \{\text{suffixes of the two left blocks}\}$) is far-junction intact. Thus Theorem ?? alone, while it correctly forbids $|S| < 3$ among intact subsets, accounts for only 1/14 of the actual critical survivors, and the broken case must be controlled for the unconditional lower bound.

(2) The single-subspace restriction provably fails for merged arcs. The within-arc proof rests on a single inert top generator T_{a+2} commuting with all arc generators. A merged super-arc straddles two original tops T_{a+1}, T_{a+2} joined by the broken junction's generator. Concretely, for $(2, 2, 1, 1, 1)$, breaking the shared junction $(1, 5)$ yields a super-arc with generator sequence $(3, 2, 1, 5, 4, 3, 2)$ and right boundary $x \in W_1 \cap W_6$; the defect generator T_5 is *adjacent* to the top T_6 , so W_6 is *not* T_5 -invariant and the product leaves W_6 . (We verified that forcing the computation into W_6 annihilates it spuriously.) No single q -eigenspace is simultaneously invariant under all super-arc generators, so the rank-one collapse has no single home.

(3) What is nonetheless certain. The full lower bound $\min\{|S| : c_S \neq 0\} = 3$ for $(2, 2, 1, 1, 1)$ is verified exhaustively over all $\binom{21}{\leq 3}$ subsets at $q = 5$ and $q = 7/3$; combined with the 14 equal positive critical survivors this gives $\text{ord}_{s=0} P_{(2,2,1,1,1)} = 3$ outright. The structural content proved here is the within-arc reach for the whole family; the merged-arc reach is the precise — and now sharply localised — remaining obstruction.

Conjectural route for the merged case. The natural fix mirrors the hook merged-junction lemma (which used a *single* reach counter R traversing a multi-block gap): replace the single inert-top restriction by a *sequence* of restrictions, one per block of the super-arc, glued at the broken junctions. Equivalently, the defect generator T_{a+1} (adjacent to both tops) should be absorbed by a rank-two “transfer” across the junction, after which the triple-pinch sign-kill again forces the bottom-vertex support. A clean statement: *a super-arc covering zero arcs with parameters $a_1 < \dots < a_p$ requires $\geq \sum_i (a_i - 2)$ insertions inside it, broken-junction positions included.* This is the off-hook analogue of Lemma 6.3 of the hook paper and would complete the unconditional lower bound for $(2, 2, 1^m)$.

7 Verification

All claims were checked symbolically in `sympy` over $\mathbb{Q}(q)$ at $q = 5$ and $q = 7/3$, reusing the seminormal machinery of `compute_p_lambda.py`. Scripts in `~/projects/scratch/2026-05-23-omega-monodromy-verify` (2026-05-28-*.py).

- `triple-character.py`: $\text{pair} \equiv 1$, $\text{triple} \equiv 0$ for $\lambda = (2, 2, 1^m)$, $m = 1, \dots, 6$ (Lemma ??).

- **triple-pinch.py**: for every valid arc of $(2, 2, 1^m)$, $m = 1, \dots, 4$: $\dim(W_j \cap W_{a+2}) = 1$ ($j \leq a$); $\dim(W_1 \cap W_j \cap W_{a+2}) = 0$ and $T_1 p_j = -p_j$ and $y^\top p_j = 0$ for $3 \leq j \leq a$; $y^\top p_1, y^\top p_2 \neq 0$ (Cor. ??, Lemmas ??-??).
- **arc-as-hook.py**: on $W = W_{a+2}$ each $P_q^{(j)} \upharpoonright_W$ ($j \leq a$) is rank one; Gram is tridiagonal with $P_q[j, j+1]P_q[j+1, j] = q/(q+1)^2$; right boundary $x \propto p_1$.
- **streamlined-bound.py**: for every arc of $(2, 2, 1^m)$, $m = 2, 3, 4$: $\sigma_a(S) \neq 0 \Rightarrow \{3, \dots, a\} \subseteq S$ (zero violations); min reach = $a - 2$, unique minimiser $\{3, \dots, a\}$ (Theorem ??).
- **merged-arc.py**, **survivors.py**: $(2, 2, 1, 1, 1)$ min-support = 3 over all subsets; $14 = C_4$ critical survivors, all $= q^8/(q+1)^{16}$, 13 breaking a far junction (§6).

λ	m	τ	zero arcs a	$\sum(a-2)$	status of lower bound
$(2, 2, 1, 1)$	2	1	$\{3\}$	1	intact case proved; full = 1 exhaustive
$(2, 2, 1, 1, 1)$	3	2	$\{3, 4\}$	3	intact case proved; full = 3 exhaustive
$(2, 2, 1^4)$	4	3	$\{3, 4, 5\}$	6	intact case proved; merged open

8 What is proved, and what remains

Proved (whole family $(2, 2, 1^m)$).

- The pair and triple pinch identities (Lemma ??) and the far sign-kill (Corollary ??).
- The inert-top rank-one collapse (Lemma ??): the hook structure, restored on W_{a+2} .
- **The within-arc reach lemma** (Theorem ??): a zero arc absorbs $\geq a-2$ insertions, uniquely — closing the first “verified, not proved” input of the off-hook paper, via a one-line walk expansion powered by the triple-pinch sign-kill.
- The intact-junction lower bound $\text{ord} \geq \binom{m}{2}$ (Theorem ??), unconditional.

Remaining (the one gap). The merged (broken-junction) reach (§6), now shown to be the generic survivor configuration and provably outside the single-subspace restriction; conjecturally settled by a glued, per-block restriction. With it, the order law $\text{ord} = \binom{m}{2} = \frac{\tau(\tau+1)}{2}$ would hold unconditionally for the infinite off-hook family $(2, 2, 1^m)$ — the first non-hook instance of the law. For $(2, 2, 1, 1)$ and $(2, 2, 1, 1, 1)$ the order law already holds outright by exhaustive verification, with the structural reason for the lower bound now supplied.