


Math Bootcamp

Day 1 — Foundations (Sections 1.1–1.5)

Math Camp 2026

1.1 Number systems — names and definitions

Symbol	English name	What it is
$\mathbb{N}$	natural numbers	Non-negative integers for counting
$\mathbb{Z}$	integers	$\dots, -2, -1, 0, 1, 2, \dots$
$\mathbb{Q}$	rational numbers	Fractions p/q ($q \neq 0$)
$\mathbb{R}$	real numbers	All points on the number line
$\mathbb{R}^n$	n -dimensional real space	Lists / vectors of length n



1.1 Symbols and vectors — names and definitions

Common symbols	$\in$	“is an element of”	$\forall$	“for all”
	$\exists$	“there exists”	$\Rightarrow$	“implies”
			$\Leftrightarrow$	“if and only if” (iff)

Scalar vs. vector

- **Scalar:** one number, e.g. income $w \in \mathbb{R}$
- **Vector:** a list of numbers $\mathbf{x} = (x_1, \dots, x_n) \in \mathbb{R}^n$
- **Dot product:** $\mathbf{p} \cdot \mathbf{x} = \sum_{i=1}^n p_i x_i$

1.1 — Which set? Together

Smallest set that fits?

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

→ $\mathbb{N}$

counting: $0, 1, 2, \dots$

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

→ $\mathbb{N}$

counting: $0, 1, 2, \dots$

2. Money you spend at a vending machine (e.g. HK\$8.50)

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

→ $\mathbb{N}$

counting: $0, 1, 2, \dots$

2. Money you spend at a vending machine (e.g. HK\$8.50)

→ $\mathbb{Q}$

e.g. $8.50 = 17/2$ dollars

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

→ $\mathbb{N}$

counting: $0, 1, 2, \dots$

2. Money you spend at a vending machine (e.g. HK\$8.50)

→ $\mathbb{Q}$

e.g. $8.50 = 17/2$ dollars

3. Order as a vector: (HK\$ spent on drink, # snacks) = (8.50, 2)

1.1 — Which set? Together

Smallest set that fits?

1. The number of girlfriends / boyfriends you can have

→ $\mathbb{N}$

counting: $0, 1, 2, \dots$

2. Money you spend at a vending machine (e.g. HK\$8.50)

→ $\mathbb{Q}$

e.g. $8.50 = 17/2$ dollars

3. Order as a vector: (HK\$ spent on drink, # snacks) = (8.50, 2)

→ $(8.50, 2) \in \mathbb{Q} \times \mathbb{N}$

1.2 Logic — names and definitions

English	Symbol	Meaning
P only if Q	$P \Rightarrow Q$	Q is necessary for P (without Q , no P)
P if Q	$Q \Rightarrow P$	Q is sufficient for P (Q alone is enough)
P if and only if Q	$P \Leftrightarrow Q$	Q is necessary and sufficient
Contrapositive	$\neg Q \Rightarrow \neg P$	same truth as $P \Rightarrow Q$: if Q fails, P must fail

1.1 — IKEA run Together

First week in HK: you go to IKEA and spot three plush toys.



Otter HK\$179.9



Hedgehog HK\$79.9



BLÅHAJ HK\$179.9

You have **HK\$500** pocket money ($B = 500$).

Basket $\mathbf{x} \in \mathbb{N}^3$: # otters, # hedgehogs, # sharks. E.g. $(1, 1, 0)$ = one otter, one hedgehog, no shark. 6/13
Price vector $\mathbf{p} = (179.9, 79.9, 179.9) \in \mathbb{Q}^3$. Budget set: $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$.

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in] (0, 0, 3) \in \mathcal{B}$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in] (0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)

2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. **True or false?**

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. **True or false?**

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ **if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. **True or false?**

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ **if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$

6. $[\Rightarrow]$ $(0, 0, 3) \in \mathcal{B}$ **if** $(1, 1, 1) \in \mathcal{B}$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ only if $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ if $\mathbf{p} \cdot (1, 1, 1) \leq 500$

6. $[\Rightarrow]$ $(0, 0, 3) \in \mathcal{B}$ if $(1, 1, 1) \in \mathcal{B}$ **False** ($(1, 1, 1) \in \mathcal{B}$ but $539.7 > 500$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. **True or false?**

1. $[\in]$ $(0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset]$ $\mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall]$ $\forall \mathbf{x} \in \mathcal{B}$, $\mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists]$ $\exists \mathbf{x} \in \mathcal{B}$, $\mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow]$ $(1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ **if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$

6. $[\Rightarrow]$ $(0, 0, 3) \in \mathcal{B}$ **if** $(1, 1, 1) \in \mathcal{B}$ **False** ($(1, 1, 1) \in \mathcal{B}$ but $539.7 > 500$)
7. [contrapositive] **If** $\mathbf{p} \cdot (1, 1, 1) + \mathbf{p} \cdot (0, 1, 0) > 500$, **then** $(1, 1, 1) \notin \mathcal{B}$ **or** $(0, 1, 0) \notin \mathcal{B}$

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. **True or false?**

1. $[\in] (0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset] \mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall] \forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists] \exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow] (1, 1, 1) \in \mathcal{B}$ **only if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ **if** $\mathbf{p} \cdot (1, 1, 1) \leq 500$

6. $[\Rightarrow] (0, 0, 3) \in \mathcal{B}$ **if** $(1, 1, 1) \in \mathcal{B}$ **False** ($(1, 1, 1) \in \mathcal{B}$ but $539.7 > 500$)
7. [contrapositive] If $\mathbf{p} \cdot (1, 1, 1) + \mathbf{p} \cdot (0, 1, 0) > 500$, then $(1, 1, 1) \notin \mathcal{B}$ **or** $(0, 1, 0) \notin \mathcal{B}$ **False** ($519.6 > 500$, yet each basket alone is in $\mathcal{B}$)

1.2 — Plush symbols practice Together

$\mathbf{p} = (179.9, 79.9, 179.9)$, $\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$. True or false?

1. $[\in] (0, 0, 3) \in \mathcal{B}$ **False** ($539.7 > 500$)
2. $[\subset] \mathbb{N}^3 \subset \mathcal{B}$ **False** (e.g. $(3, 0, 0) \notin \mathcal{B}$)
3. $[\forall] \forall \mathbf{x} \in \mathcal{B}, \mathbf{x} \neq (0, 1, 0)$ **False** ($(0, 1, 0) \in \mathcal{B}$)
4. $[\exists] \exists \mathbf{x} \in \mathcal{B}, \mathbf{x} = (3, 0, 0)$ **False** ($539.7 > 500$)
5. $[\Rightarrow] (1, 1, 1) \in \mathcal{B}$ only if $\mathbf{p} \cdot (1, 1, 1) \leq 500$ **True** ($439.7 \leq 500$)

also True: $(1, 1, 1) \in \mathcal{B}$ if $\mathbf{p} \cdot (1, 1, 1) \leq 500$

6. $[\Rightarrow] (0, 0, 3) \in \mathcal{B}$ if $(1, 1, 1) \in \mathcal{B}$ **False** ($(1, 1, 1) \in \mathcal{B}$ but $539.7 > 500$)
7. [contrapositive] If $\mathbf{p} \cdot (1, 1, 1) + \mathbf{p} \cdot (0, 1, 0) > 500$, then $(1, 1, 1) \notin \mathcal{B}$ or $(0, 1, 0) \notin \mathcal{B}$ **False** ($519.6 > 500$, yet each basket alone is in $\mathcal{B}$)

each affordable $\nRightarrow$ jointly affordable

1.3 Sets and operations — names and definitions

Symbol	Name	Meaning
$\in, \notin$	element / not in	x belongs (or not) to A
$\subset, =$	subset, equality	$A \subset B$; $A = B$ iff mutual $\subset$
$\emptyset, \{ \}, \{x \in S : \dots\}$	sets	empty, roster, set-builder
$A \cup B, A \cap B$	union, intersection	in A or B ; in both
$A^c, A \setminus B$	complement, difference	not in A ; in A not B
$A \times B$	Cartesian product	pairs (a, b) , $a \in A$, $b \in B$

In economics: budget set $\mathcal{B} \subset \mathbb{N}^3$; $(8.50, 2) \in \mathbb{Q} \times \mathbb{N}$.

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets or baskets with ≥ 1 hedgehog

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets or baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. (0, 1, 0))
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets **or** baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$ affordable baskets with **no** hedgehog (e.g. (1, 0, 0))

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets **or** baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$ affordable baskets with **no** hedgehog (e.g. $(1, 0, 0)$)
4. $(179.9, 79.9) \in \mathbb{Q} \times \mathbb{Q}$ but $(179.9, 79.9) \notin \mathcal{B}$

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets or baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$ affordable baskets with no hedgehog (e.g. $(1, 0, 0)$)
4. $(179.9, 79.9) \in \mathbb{Q} \times \mathbb{Q}$ but $(179.9, 79.9) \notin \mathcal{B}$ **True** Cartesian product $\neq$ budget set — pairs $\neq$ baskets

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets or baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$ affordable baskets with no hedgehog (e.g. $(1, 0, 0)$)
4. $(179.9, 79.9) \in \mathbb{Q} \times \mathbb{Q}$ but $(179.9, 79.9) \notin \mathcal{B}$ True Cartesian product $\neq$ budget set — pairs $\neq$ baskets
5. $(1, 0, 0) \in \mathcal{B}$

1.3 — Set operations (IKEA) Together

$\mathcal{B} = \{\mathbf{x} \in \mathbb{N}^3 : \mathbf{p} \cdot \mathbf{x} \leq 500\}$, $\mathcal{H} = \{\mathbf{x} \in \mathbb{N}^3 : \text{middle entry} \geq 1\}$ (at least one hedgehog).

Describe the set.

1. $\mathcal{B} \cap \mathcal{H}$ affordable baskets with ≥ 1 hedgehog (e.g. $(0, 1, 0)$)
2. $\mathcal{B} \cup \mathcal{H}$ affordable baskets or baskets with ≥ 1 hedgehog
3. $\mathcal{B} \setminus \mathcal{H}$ affordable baskets with no hedgehog (e.g. $(1, 0, 0)$)
4. $(179.9, 79.9) \in \mathbb{Q} \times \mathbb{Q}$ but $(179.9, 79.9) \notin \mathcal{B}$ **True** Cartesian product $\neq$ budget set — pairs $\neq$ baskets
5. $(1, 0, 0) \subset \mathcal{B}$ **False** $(1, 0, 0)$ is a basket: use $(1, 0, 0) \in \mathcal{B}$ (True); but $\{(1, 0, 0)\} \subset \mathcal{B}$ is True

1.4 Functions — names and definitions

Term	Notation	Meaning
Function	$f: A \rightarrow B$	each $x \in A$ maps to exactly one $f(x) \in B$
Domain	A	allowed inputs
Codomain	B	target set — every output must lie in B
Range	$f(A)$	outputs hit: $\{f(x) : x \in A\} \subseteq B$
Injective (one-to-one)		$f(x_1) = f(x_2) \Rightarrow x_1 = x_2$
Surjective (onto)		every $y \in B$ equals some $f(x)$
Operations	$(f \pm g)(x)$, $(cf)(x)$, $(fg)(x)$	$(f \pm g)(x) = f(x) \pm g(x)$; $(cf)(x) = c f(x)$; $(fg)(x) = f(x)g(x)$
Composition	$(f \circ g)(x)$	apply g first: $(f \circ g)(x) = f(g(x))$

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

2. Codomain C ?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

2. Codomain C ? $\{18, 19, \dots, 80\}$

plausible partner ages

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

2. Codomain C ? $\{18, 19, \dots, 80\}$

plausible partner ages

3. Is $80 \in f(D)$?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

2. Codomain C ? $\{18, 19, \dots, 80\}$

plausible partner ages

3. Is $80 \in f(D)$? **No, sugar mommy?**

$(80 \in C, \text{ but } f(D) \subseteq \{17, \dots, 35\})$

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$

typical ages in a master's program

2. Codomain C ? $\{18, 19, \dots, 80\}$

plausible partner ages

3. Is $80 \in f(D)$? **No, sugar mommy?**

($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)

4. Range $f(D)$?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)? **No** ($f(D) \neq C$; e.g. $80 \in C$ but $80 \notin f(D)$)

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)? **No** ($f(D) \neq C$; e.g. $80 \in C$ but $80 \notin f(D)$)
6. Is f injective (one-to-one)?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)? **No** ($f(D) \neq C$; e.g. $80 \in C$ but $80 \notin f(D)$)
6. Is f injective (one-to-one)? **Probably not** (e.g. $f(22) = f(30) = 22$)

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)? **No** ($f(D) \neq C$; e.g. $80 \in C$ but $80 \notin f(D)$)
6. Is f injective (one-to-one)? **Probably not** (e.g. $f(22) = f(30) = 22$)
7. If $f(25) = \{22, 80\}$, is f a function?

1.4 — Ages Together

Define $f: D \rightarrow C$ by: $f(x)$ = your girlfriend / boyfriend's age when **your age** is x .

1. Domain D ? $\{22, 23, \dots, 30\}$ typical ages in a master's program
2. Codomain C ? $\{18, 19, \dots, 80\}$ plausible partner ages
3. Is $80 \in f(D)$? **No, sugar mommy?** ($80 \in C$, but $f(D) \subseteq \{17, \dots, 35\}$)
4. Range $f(D)$? **your age ± 5 on D** $\{17, 18, \dots, 35\}$
5. Is f surjective (onto C)? **No** ($f(D) \neq C$; e.g. $80 \in C$ but $80 \notin f(D)$)
6. Is f injective (one-to-one)? **Probably not** (e.g. $f(22) = f(30) = 22$)
7. If $f(25) = \{22, 80\}$, is f a function? **No — a correspondence** (a function gives **exactly one** output per input; it's immoral to have a correspondence ☺)

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept?

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept? **No maximum in P** (179.89 works; so does 179.899; ...)

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept? **No maximum in P** (179.89 works; so does 179.899; ...)

Reservation price (WTP)?

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept? **No maximum in P** (179.89 works; so does 179.899; ...)

Reservation price (WTP)? **$\sup P = 179.9$** least upper bound — the choke price)

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept? **No maximum in P** (179.89 works; so does 179.899; ...)

Reservation price (WTP)? **$\sup P = 179.9$** least upper bound — the choke price)

Is $179.9 \in P$?

1.5 — Reservation price (example first) Together

BLÅHAJ sticker price: **HK\$179.9**. You buy **only on sale**: when $p < 179.9$.

Prices where you still buy: $P = \{p \in \mathbb{R}_+ : p < 179.9\}$.

Highest sale price you would accept? **No maximum in P** (179.89 works; so does 179.899; ...)

Reservation price (WTP)? $\sup P = 179.9$ least upper bound — the choke price)

Is $179.9 \in P$? **No** at sticker price you walk away $\Rightarrow$ no max P)

1.5 Supremum and infimum — names and definitions

Term	Notation	Meaning
Maximum	$\max S$	largest element in S
Minimum	$\min S$	smallest element in S
Supremum	$\sup S$	least upper bound of S
Infimum	$\inf S$	greatest lower bound of S

Example (least upper bound): $S = [0, 1)$. Upper bounds include $1, 2, 3, \dots$; $\sup S = 1$ — the **smallest** upper bound ($1 \notin S$).

Relations: if $\max S$ exists, $\max S = \sup S$; if $\min S$ exists, $\min S = \inf S$.

Always: $\inf S \leq x \leq \sup S$ for all $x \in S$.

In our example: $\sup P = 179.9$, but $\sup P \notin P$ — so no $\max P$.

Use $\sup$ when a bound exists but $\max$ may not — reservation prices, demand choke prices, limits in equilibrium.