

- FOKKENS, H. 1998. The Ussen Project: the first decade of excavations at Oss, *Analecta Praehistorica Leidensia* 30.
- GEBERS, W. 1985. Jungbronzezeitliche und eisenzeitliche Getreidevorratshaltung in Rullstorf, in K. Wilhelmi (ed.), *Ausgrabungen in Niedersachsen. Archäologische Denkmalpflege 1979–1984*: 146–50. Stuttgart: Berichte zu Denkmalpflege in Niedersachsen, Beiheft 1.
- GUTTMANN, E. & J. LAST. 2000. A Late Bronze Age landscape at South Hornchurch, Essex, *Proceedings of the Prehistoric Society* 66: 319–59.
- HORST, F. 1985. *Zedau. Eine jungbronze- und eisenzeitliche Siedlung in der Altmark*. Berlin: Akademie Verlag.
- JEUNESSE, C. & M. EHRETMANN. 1988. La jeune femme, le cheval et le silo, *Cahiers Alsaciens d'Archéologie et d'Art et d'Histoire* 31: 43–54.
- KALIFF, A. 1998. Grave structures and altars: archaeological traces of Bronze Age eschatological conceptions, *European Journal of Archaeology* 1.2: 177–98.
- KOSSACK, G. 1954. *Studien zur Symbolgut der Urnenfelder und Hallstattzeit Mitteleuropas*. Berlin: de Gruyter. Römisch-Germanische Forschungen 20.
- KRISTIANSEN, K. 1998. *Europe before history*. Cambridge: Cambridge University Press.
- LUCAS, A.T. 1958. An Fhóir: a straw rope granary, *Gwern* 1.1: 2–20.
- MONTELIUS, O. 1897. Hausurnen und Gesichturnen, *Korrespondenzblatt der Deutschen Gesellschaft für Anthropologie, Ethnologie und Vorgeschichte* 28: 16–25.
- MÜLLER, R. 1999. Hausurnen, in R. Müller (ed.), *Reallexikon der Germanischen Altertumskunde* 14: 85–9. Berlin: De Gruyter.
- OELMANN, F. 1959. Pfahlhausurnen, *Germania* 37: 205–23.
- OLAUSSEN, D. 1986. Piledal and Svarte. A comparison between two Late Bronze Age cremation cemeteries in Scania, *Acta Archaeologica* 57: 121–52.
- RASSMUSSEN, M. & C. ANDERSEN. 1993. Bronze Age settlement, in B. Storgaard (ed.), *Digging into the past: 25 years of Danish archaeology*: 136–44. Aarhus: Jutland Archaeological Society.
- RITTERSHOFER, K.-F. (ed.). 1997. *Sonderbestattungen in der Bronzezeit in östlichen Mitteleuropa*. Espelkamp: Verlag Marie Leidorf.
- ROYMANS, N. 1999. Urnfield symbolism: ancestors and the land in the lower Rhine region, in F. Theuvs & N. Roymans (ed.), *Land and ancestors*: 11–61. Amsterdam: Amsterdam University Press.
- STJERNQVIST, B. 1961. *Simris II. Bronze Age problems in the light of the Simris excavations*. Lund: Gleerup Forlag.
- TESCH, S. 1998. Tradition and change during the Bronze and Iron Ages. Houses as archaeological sources for the study of changes in the cultural landscape, in R. Berglund (ed.), *The cultural landscape during 6000 years in southern Sweden*: 326–36. Copenhagen: Munksgaard. Ecological Bulletin 41.
- VILLES, A. 1981. Les bâtiments domestiques Hallstattiennes de la Chaussée-sur-Marne and la problème de la maison de l'Âge du Fer en France septentrionale, in V. Kruta (ed.), *L'Âge du Fer en France septentrionale*: 49–97. Reims: Mémoires de la Société archéologique champenoise 2.
- WHIMSTER, R. 1981. *Burial practices in Iron Age Britain*. Oxford: British Archaeological Reports. British series 90.

## The age of the common bean (*Phaseolus vulgaris* L.) in the northern Eastern Woodlands of North America

JOHN P. HART, DAVID L. ASCH, C. MARGARET SCARRY  
& GARY W. CRAWFORD\*

*This study addresses the lack of chronometric research on the common bean (Phaseolus vulgaris L.) to establish precisely the timing of its adoption and spread across the northern Eastern Woodlands of North America. Bean and directly associated maize samples were subjected to accelerator mass spectrometry (AMS) dating. The results show that the common bean apparently spread rapidly upon its introduction to the region, becoming archaeologically visible from the Illinois River valley to southern New England in the calibrated late 13th century AD, some 200–300 years later than previously thought.*

**Key-words:** northeastern North America, maize-beans-squash agriculture, common bean, *Phaseolus vulgaris*, AMS dating

Research in the Eastern Woodlands of North America during the last quarter of the 20th century established that prehistoric agricultural systems centring on a suite of indigenous oily and starchy seeded annual plants began to evolve

by 4000 BP (c. 2550 cal BC),<sup>1</sup> becoming wide-

1 In this article, years BP refers to radiocarbon age without adjustment for past atmospheric <sup>14</sup>C fluctuations; the BC/AD calendrical scale is reserved for dendrocalibrated dates. All calibrations were done with CALIB 4.3 (Stuiver *et al.* 1998).

\* Hart & Asch, Research & Collections Division, New York State Museum, Albany NY 12230, USA. jph\_nysm@mail.nysed.gov Scarry, Department of Anthropology, University of North Carolina, Chapel Hill NC 27599, USA. Crawford, Department of Anthropology, University of Toronto, Mississauga ONT L5L 1C6, Canada.

Received 15 June 2001, accepted 31 October 2001, revised 30 November 2001




FIGURE 1. General locations of archaeological sites listed in TABLES 1 & 2 (● sites with samples dated for the current study, ■ sites with previously dated beans).

| site, provenance <sup>1</sup>    | material | lab no. <sup>2</sup> | $\delta^{13}\text{C}$ (‰) | $^{14}\text{C}$ BP | AD cal 2 $\sigma$ (intercepts) |
|----------------------------------|----------|----------------------|---------------------------|--------------------|--------------------------------|
| (1) Burnham-Shepard <sup>3</sup> | bean     | B-29619              |                           | 550±60             | 1297 (1406) 1445               |
| (2) Skitchewaung, F 32           | bean     | AA-29119             | -25.9                     | 670±45             | 1274 (1297) 1399               |
| (2) Skitchewaung, F5a&b          | bean     | AA-29120             | -24.4                     | 765±50             | 1188 (1275) 1299               |
| (2) Skitchewaung, F46            | bean     | AA-29121             | -25.0                     | 600±50             | 1289 (1327, 1346, 1393) 1428   |
| (3) Broome Tech                  | bean     | AA-31007             | -27.3                     | 380±40             | 1437 (1481) 1638               |
| (4) Roundtop, F35                | bean     | AA-23106             | -27.2                     | 658±48             | 1276 (1299, 1374, 1376) 1404   |
| (4) Roundtop, F35                | maize    | AA-21979             | -8.7                      | 675±55             | 1260 (1297) 1403               |
| (4) Roundtop, F35                | wood     | AA-21980             | -27.6                     | 670±55             | 1262 (1297) 1403               |
| (4) Roundtop, post mold          | bean     | AA-26540             | -25.0                     | 315±45             | 1453 (1528, 1551, 1633) 1661   |
| (5) Thomas/Luckey, F78           | bean     | AA-29122             | -27.7                     | 695±45             | 1260 (1292) 1392               |
| (8) Gnagey, F30                  | bean     | AA-29117             | -23.3                     | 610±55             | 1283 (1323, 1350, 1390) 1428   |
| (8) Gnagey, F13D                 | bean     | AA-29118             | -25.0                     | 635±45             | 1282 (1303, 1368, 1383) 1409   |

1 Numbers correspond to site locations plotted on FIGURE 1.

2 AA = NSF Arizona AMS Facility, B = Beta Analytic, Inc.

3 Originally published in Bendremer *et al.* (1991). The Roundtop dates were originally published in Hart (1999b). All other dates were originally published in Hart & Scarry (1999).

TABLE 1. AMS dates of common beans and associated maize and wood in the northeastern US.



spread by 2000 BP (c. cal AD 15) (Asch 1994; Smith 1992). Except for the squash species *Cucurbita pepo* L., however, these were not the crops that dominated agricultural systems across the region at the time of European contact. Instead the 'three sisters' — squash, maize (*Zea mays* L.) and the common bean (*Phaseolus vulgaris* L.) — were the principal Native American crops of the early Historic period (Hurt 1987). *Cucurbita pepo* was domesticated both in eastern North America (ssp. *ovifera* (L.) Decker) and Mexico (ssp. *pepo*) (Decker 1988). Subspecies *ovifera* spread throughout the northern Eastern Woodlands as a result of human interactions, perhaps at first as a form with hard-shelled, bitter, inedible fruits (Asch 1994; Conard *et al.* 1984; Hart & Asch Sidell 1997; Smith 1992). Maize and a second squash species (*C. argyrosperma* Huber) were initially domesticated in Mesoamerica, and the common bean (hereafter 'beans') in Mesoamerica and northern South America (Fritz 1994; Kami *et al.* 1995; MacNeish & Eubanks 2000). These spread to the northern Eastern Woodlands through human interactions, perhaps *via* the American Southwest and Plains. The timing of the spread of these crops and the processes involved remain important issues for our understanding of prehistoric agricultural evolution (Asch 1994; Hart 1999a; Riley *et al.* 1990). In this article we provide new dates on beans that change our understanding of the history of this crop and maize-beans-squash agriculture in the northern Eastern Woodlands.

### Northern Eastern Woodlands beans

The history of squash and maize in the northern Eastern Woodlands has been clarified over the past two decades as the result of intensive efforts to recover and identify archaeological plant remains, together with developments in accelerator mass spectrometry (AMS) that made it possible to date small, critical crop specimens directly. Remains of presumably cultivated *C. pepo* have been directly dated to as early as 7100±300 BP (NSRL-298) (cal 2σ 6498 (5988, 5940, 5929) 5476 BC) in the Midwest (Asch & Asch 1985) and 5695±100 BP (AA-7491) (cal 2σ 4775 (4522, 4509, 4503) 4341 BC) in the northeastern US (Petersen & Asch Sidell 1996), and there are some indications of its biological domestication as early as c. 4250 BP (c. cal 2900 BC) (Asch 1995: 62–3). The cushaw squash,

*C. argyrosperma*, entered the region probably only a millennium ago (Fritz 1994). Maize has been directly dated to as early as 2077±70 BP (AA-8717) (cal 2σ 354 BC (90, 76, 59 BC) AD 72) in Illinois (Riley *et al.* 1994) and 1570±90 BP (TO-5307) (cal 2σ AD 258 (442, 448, 468, 482, 530) 657) southern Ontario (Crawford *et al.* 1997). The earliest direct date in the northeastern US is 1100±70 BP (B-53452) (cal. 2σ AD 776 (904, 910, 976) 1145) in eastern New York (Cassedy & Webb 1999). Stable carbon isotope analyses of human bone and measurements of maize quantities on archaeological sites suggest maize did not become an important crop throughout the region until after 1100 BP (Hart 1999a; Smith 1992).

For beans, in contrast to squash and maize, there has been a lack of direct AMS dating to establish firmly the first archaeological visibility of this crop in the northern Eastern Woodlands. Probably because beans arrived relatively late in time, the oldest apparent contexts in which they were found were not deemed controversial. It was long the conventional wisdom that beans were present in Mississippian and contemporaneous cultural contexts of the region, and palaeoethnobotanists estimated their probable time of arrival at 1150–950 BP (Ford 1985a; Griffin 1967; Yarnell 1976; 1986). Recent research in the northeastern US, however, has raised doubts about the validity of previous indirect dating of beans, which was based on apparent associations of the beans with radiocarbon-dated organic materials or temporally diagnostic pottery. In the northeastern US, direct AMS dating of beans from purportedly early contexts now suggests that they do not become archaeologically visible there until approximately cal AD 1300 (FIGURE 1, TABLE 1), some 200–300 years later than previously thought (Hart & Scarry 1999). Notably the Roundtop site in New York, one of the redated sites (Hart 1999b), was once cited by Yarnell (1976: 272) as having 'the earliest verified report of beans in the East' on the basis of a date of 880±60 BP (Y-1534) (cal 2σ AD 1005 (1163, 1173, 1180) 1282) on charcoal from a storage pit (not the pit from which beans were recovered).

Three areas remain in the northern Eastern Woodlands to the east of the Mississippi River valley where there are repeated reports of beans occurring in pre-AD 1300 contexts: the upper Ohio River basin, the central Ohio River basin, and the central and lower Illinois River




FIGURE 2. Bean cotyledon fragment from the Sunwatch site: a inner face of cotyledon, b plumule, c notch in cotyledon for radicle.

basin. In southern Ontario, beans have not been reported before AD 1200 and rarely before AD 1300 (Smith & Crawford 1997). Checking the potentially early Ontario records, Crawford found that only one bean identification can be confidently supported. An AMS date on this specimen from the Kelly site was obtained as part of the current study.

Many of the sites with beans supposedly in early contexts are multicomponent; have radio-

carbon dates (usually on charcoal) that often are widely dispersed, have large standard deviations, or were obtained from cultural features that did not contain the beans; or they lack radiocarbon dates altogether. In light of the revised chronology that resulted from direct dating of beans in the northeastern US, there was an evident need to investigate the age of beans from purportedly early contexts across a wider region. The present project was undertaken to clarify the history of beans across the northern Eastern Woodlands from the Illinois River valley to the east and north. Rather than reassessing the age of any particular site or site components in order to date beans indirectly, the goal was to obtain direct evidence for the age of the earliest beans in the various drainage basins.

#### Sample selection and identification

Since a radiocarbon date potentially can be compromised by unrecognized organic contamination or by large counting or non-counting errors in the radiocarbon lab, paired samples of maize recovered from the same contexts were sought for dating from some sites. A discordant maize date could serve as a signal that dating results for one of the samples is erroneous. Twenty-five bean and 12 maize samples from 20 sites (FIGURE 1) were obtained from various curatorial facilities.<sup>2</sup> The sites included one from southern Ontario, four from the upper Ohio River basin, eight from the central Ohio River basin<sup>3</sup> and seven from the central and lower Illinois River basin. Lawrence Kaplan provided previously unpublished bean and maize dates from the Blain Village site in the central Ohio River basin. Together with the previously published dates from the northeastern US, there are now 51 dates from 26 sites available to address the issue of the spread of beans across the northern Eastern Woodlands from the Illinois River valley to southern New England.

2 Carnegie Museum of Natural History, Center for American Archeology, Dayton Museum of Natural History, Grove Creek Mound Historic Site, Illinois State Museum, Ohio Historical Society, University of Indiana, University of Massachusetts-Boston, University of Toronto, Western Illinois University, William S. Webb Museum.

3 Original plans were to date matching bean and maize samples from the Muir Site in Kentucky, from which beans have been reported in c. AD 1000 contexts (Rossen 1988: 258). Neither the beans nor any maize from the same contexts as the reported beans could be located at the curatorial facility.



| site, provenance <sup>1</sup>                | material | lab no. <sup>2</sup> | $\delta^{13}\text{C}$ (‰) | $^{14}\text{C}$ BP        | AD cal 2 $\sigma$ (intercepts)     |
|--|----------|----------------------|---------------------------|---------------------------|------------------------------------|
| (6) Onoville Bridge, F103                    | bean     | AA-38454             | -25.9                     | 628 $\pm$ 33              | 1290 (1304, 1366, 1385) 1404       |
| (6) Onoville Bridge, F103                    | maize    | AA-38455             | -14.8                     | 624 $\pm$ 33              | 1291 (1305, 1355, 1365, 1386) 1405 |
| (7) Kelly, F7                                | bean     | TO-8963              |                           | 770 $\pm$ 50 <sup>3</sup> | 1164 (1271) 1298                   |
| (9) Campbell Farm, F352                      | bean     | AA-40132             | -27.1                     | 462 $\pm$ 38              | 1408 (1438) 1481                   |
| (9) Campbell Farm, F352                      | maize    | AA-40133             | -8.4                      | 794 $\pm$ 38              | 1164 (1258) 1286                   |
| (10) Portman, Trench 44                      | bean     | AA-38456             | -25.5                     | 682 $\pm$ 33              | 1277 (1295) 1390                   |
| (11) Saddle, F35                             | bean     | AA-38457             | -25.6                     | 675 $\pm$ 33              | 1278 (1297) 1392                   |
| (11) Saddle, F35                             | maize    | AA-38458             | -12.8                     | 605 $\pm$ 34              | 1296 (1326, 1348, 1391) 1412       |
| (12) Blennerhassett F23 N1/2                 | bean     | AA-38463             | -28.6                     | 277 $\pm$ 33              | 1518 (1643) 1789                   |
| (12) Blennerhassett F23 S1/2                 | bean     | AA-38464             | -27.9                     | 301 $\pm$ 33              | 1484 (1637) 1658                   |
| (13) Blain Village, House 1 <sup>4</sup>     | bean     | AA-16854             |                           | 510 $\pm$ 60              | 1304 (1421) 1478                   |
| (13) Blain Village, Pit 4 18-28 <sup>4</sup> | maize    | AA-16853             |                           | 420 $\pm$ 60              | 1408 (1448) 1637                   |
| (14) Baldwin                                 | bean     | AA-38459             | -26.7                     | 542 $\pm$ 33              | 1321 (1408) 1437                   |
| (14) Baldwin                                 | bean     | AA-38460             | -27.4                     | 494 $\pm$ 33              | 1401 (1428) 1447                   |
| (15) Gartner                                 | bean     | AA-38461             | -27.7                     | 579 $\pm$ 33              | 1301 (1332, 1339, 1398) 1423       |
| (15) Gartner                                 | bean     | AA-38462             | -28.1                     | 593 $\pm$ 33              | 1298 (1328, 1344, 1394) 1416       |
| (16) Sunwatch, F1/77                         | bean     | A-0175               | -26.5                     | 652 $\pm$ 42              | 1280 (1300, 1373, 1378) 1402       |
| (17) Fox Farm, FG09 Tu2                      | bean     | AA-38466             | -26.0                     | 683 $\pm$ 33              | 1277 (1295) 1389                   |
| (17) Fox Farm, FG09 Tu2                      | maize    | AA-38467             | -15.7                     | 592 $\pm$ 33              | 1299 (1329, 1344, 1395) 1418       |
| (18) Baker's Trails, F11                     | bean     | AA-40134             | -28.0                     | 539 $\pm$ 39              | 1314 (1409) 1440                   |
| (19) Murphy, 142.43.41                       | bean     | AA-38966             | -25.3                     | 603 $\pm$ 36              | 1297 (1326, 1347, 1392) 1412       |
| (20) Noble-Weiting, F8                       | bean     | AA-38964             | -27.1                     | 621 $\pm$ 36              | 1290 (1312, 1354, 1387) 1408       |
| (20) Noble-Weiting, F8                       | bean     | AA-38965             | -28.1                     | 634 $\pm$ 36              | 1286 (1303, 1368, 1383) 1404       |
| (21) Orendorf, F761                          | bean     | A-0177               | -27.0                     | 655 $\pm$ 43              | 1277 (1298) 1399                   |
| (21) Orendorf, F761                          | maize    | AA-38968             | -10.0                     | 698 $\pm$ 33              | 1269 (1291) 1385                   |
| (21) Orendorf, F782                          | bean     | AA-38967             | -27.6                     | 712 $\pm$ 33              | 1262 (1286) 1381                   |
| (22) Morton, F3                              | bean     | AA-38473             | -25.2                     | 675 $\pm$ 33              | 1278 (1297) 1392                   |
| (22) Morton, F3                              | maize    | AA-38474             | -9.7                      | 692 $\pm$ 33              | 1274 (1293) 1387                   |
| (23) Larson, F140                            | bean     | A-0176               | -25.4                     | 757 $\pm$ 44              | 1212 (1277) 1298                   |
| (23) Larson, F140                            | maize    | AA-38476             | -8.8                      | 704 $\pm$ 33              | 1265 (1289) 1383                   |
| (23) Larson, House 75                        | bean     | A-0174               | -27.1                     | 650 $\pm$ 43              | 1280 (1301, 1372, 1378) 1403       |
| (23) Larson, House 75                        | maize    | AA-38478             | -9.9                      | 719 $\pm$ 33              | 1259 (1284) 1379                   |
| (24) Hill Creek, F1-07C                      | bean     | AA-38471             | -25.9                     | 734 $\pm$ 33              | 1241 (1281) 1299                   |
| (24) Hill Creek, F1-07C                      | maize    | AA-38472             | -9.1                      | 772 $\pm$ 37              | 1211 (1268) 1293                   |
| (24) Hill Creek, F1-04PB                     | bean     | AA-40135             | -24.2                     | 641 $\pm$ 39              | 1283 (1302, 1370, 1381) 1403       |
| (24) Hill Creek, F1-04PB                     | maize    | AA-40136             | -10.0                     | 733 $\pm$ 55              | 1213 (1282) 1387                   |
| (25) Worthy-Merrigan, 1243B                  | bean     | AA-40138             | -25.9                     | 594 $\pm$ 49              | 1291 (1328, 1344, 1394) 1430       |
| (25) Worthy-Merrigan, 1243B                  | maize    | AA-38470             | -15.1                     | 615 $\pm$ 41              | 1289 (1319, 1352, 1389) 1413       |
| (26) Pere Marquette, F22-01P                 | maize    | AA-40137             | -9.2                      | 642 $\pm$ 36              | 1284 (1302, 1370, 1381) 1402       |

1 Numbers correspond to site locations plotted on FIGURE 1.

2 A = Illinois State Geological Survey, AA = NSF Arizona AMS Facility, TO = AMS Facility at the University of Toronto.

3 Date corrected to a nominal base of  $\delta^{13}\text{C} = -25\text{‰}$ .

4 Lawrence Kaplan pers. comm. 2000.

TABLE 2. AMS dates of common beans and associated maize from the current project.

All specimens were examined by the authors prior to submission for assay: Scarry and Asch examined all but three of the bean samples to confirm the original identifications. Lawrence Kaplan identified the two beans from the Noble-Weiting site, and Crawford identified the bean from the Kelly site. Two thirds of the beans were also examined by Richard Yarnell. All

specimens were carbonized; no evidence was found of preservatives or other contaminants that might affect sample carbon content. The bean samples mostly consisted of whole or partial cotyledons. From the position of the impression of the hypocotyl and plumule on the inner surface of the cotyledon, the notching of the cotyledon margin where the radicle

was present before carbonization, and from the reniform shape of the cotyledon (FIGURE 2), one may infer that the specimens represent species of the subfamily Faboideae (Fabaceae), in which typically the embryonic axis is curved and the hilum marginal according to seed length (Delorit & Gunn 1986). On the basis of cotyledon size and shape, together with geographic considerations and knowledge that only one species of domesticated bean was present in eastern North America at European contact, the specimens could be identified as *Phaseolus vulgaris* (Kirkbride *et al.* 2000).<sup>4</sup> Rare specimens preserved the hilar area and portions of the thin seed coat. If a specimen was large enough, it was divided and a portion was retained for future analyses or for additional dating should the need arise. To substantiate identifications, samples were digitally photographed under low magnification prior to division. Sample width, length, thickness and weight were recorded.

## Results

Results are presented in TABLE 2. No date is earlier than the late calibrated 13th century AD, consistent with the results obtained previously for the northeastern US. Of the 12 paired bean and maize samples from the current project and the northeastern US study, only 1 pair differed significantly at the .05 level of significance (Ward & Wilson 1978). The discordant pair, from the Campbell Farm site in southwestern Pennsylvania, was excavated from a cultural feature with active rodent burrowing, which may have mixed beans and maize from different temporal contexts. Multiple bean dates were obtained from 10 sites during the two projects. Only at Roundtop (TABLE 1) are the dates significantly different, representing temporally distinct occupations at this site (Hart 1999b). The paired bean and maize dates and paired bean dates indicate that the dates obtained for this project are reliable.

## Conclusions

The results of this project, coupled with the earlier findings for the northeastern US, suggest that the common bean spread rapidly

throughout the region, becoming a regularly grown crop from the Illinois River valley to southern New England by the late calibrated 13th century AD, two to three centuries later than previously thought. Thus, all crop elements of maize-beans-squash agricultural systems were in place in the northern Eastern Woodlands only for about 250 years before initial observation by Europeans (Biggar 1924: 158, 183). One implication of the revised chronology is that what once seemed an anomalous absence of beans in the American Bottom during the Mississippian florescence at Cahokia (Lopinot 1992) actually is consistent with the crop's history elsewhere in the region. More generally, it is clear that the growing of beans in the northern Eastern Woodlands was *not* associated with the major intensification and range extension of maize agriculture that occurred around 1100 BP. This expansion has sometimes been attributed in part to the arrival of beans, whose high protein content and amino acid composition compensated for deficiencies of maize in the critical amino acids lysine and tryptophane (Kaplan 1965; Stoltman & Baerreis 1983). But the nutritional complementation model, and others that attempt to account for the c. 1100 BP agricultural expansion by pointing to benefits gained by growing and consuming maize and beans together, now can be rejected.

In some respects the history of beans in the northern Eastern Woodlands is similar to what Kaplan & Lynch (1999) have documented for Mexico and Andean South America: beans entered agricultural systems relatively late and once adopted, their spread was very quick relative to maize and squash. This in turn indicates that the processes involved in the spread of the three crops differed dramatically, affected by each of the crop's genetics and reproduction systems, human population structure, intra- and inter-human population interactions and the agro-ecological systems into which the crops were adopted (Asch 1994; Hart 1999a; 2001). To conclude, rather than basing an agricultural chronology solely on the apparent associations of crop remains with other dated materials or with temporally diagnostic artefacts, this study further emphasizes the importance of directly dating early crop remains to determine their earliest date of archaeological visibility (Conard *et al.* 1984; Crawford *et al.* 1997; Hart 1999a).

4 A poorly preserved specimen from the Pere Marquette site in Illinois could only be identified as a probable bean cotyledon by these criteria, making its sacrifice for an AMS date inadvisable. Consequently, only a sample of maize from the same context was submitted for assay (TABLE 2).



**Acknowledgements.** This study was funded by National Science Foundation Grant BCS-9910438 and the New York State Museum. We thank Leslie Bush, Lawrence Conrad, Verna Cowin, Richard George, Alan Harn, James Heilman, Bennett Horter, William Johnson, Lawrence Kaplan, Andrea Keller, Jody O'Gorman, Nancy O'Malley, Terrance Martin, Robert McCullough, Cheryl Munson, Martha Otto, Marjorie Schroeder, Sissel Schroeder, Patrick Trader, Gail Wagner

and Michael Wiant for their help in obtaining samples for this project. We thank Lawrence Kaplan for allowing us to use the previously unpublished dates from the Blain Village site and Richard Yarnell for his assistance in identifying specimens. We thank Nancy Asch Sidell, Richard Mitchell, Charles Sheviak and two anonymous referees for their comments and suggestions on earlier versions of this paper.

## References

- ASCH, D.L. 1994. Aboriginal specialty-plant cultivation in eastern North America: Illinois prehistory and a post-contact perspective, in W. Green (ed.), *Agricultural origins and development in the Midcontinent*: 25–86. Iowa City (IA): University of Iowa. Report 19, Office of the State Archaeologist.
1995. Aboriginal specialty-plant propagation: Illinois prehistory and an eastern North American post-Contact perspective. Unpublished Ph.D. thesis, Department of Anthropology, University of Michigan.
- ASCH, D.L. & N.B. ASCH. 1985. Prehistoric plant cultivation in west-central Illinois, in Ford 1985b: 149–204.
- BENDREMER, J.C.M., E.A. KELLOGG & T.B. LARGY. 1991. A grass-lined maize storage pit and early maize horticulture in central Connecticut, *North American Archaeologist* 12: 325–49.
- BIGGAR, H.P. 1924. *The voyages of Jacques Cartier*. Ottawa: Public Archives of Canada. Publication 11.
- CASSEDY, D. & P. WEBB. 1999. New data on the chronology of maize horticulture in eastern New York and southern New England, in Hart 1999c: 85–100.
- CONARD, N., D.L. ASCH, N.B. ASCH, D. ELMORE, H. GOVE, M. RUBIN, J.A. BROWN, M.D. WIAIT, K.B. FARNSWORTH & T.G. COOK. 1984. Accelerator radiocarbon dating of evidence for prehistoric horticulture in Illinois, *Nature* 308: 443–6.
- CRAWFORD, G.W., D.G. SMITH & V.E. BOWYER. 1997. Dating the entry of corn (*Zea mays*) into the lower Great Lakes, *American Antiquity* 62: 112–19.
- DECKER, D.S. 1988. Origin(s), evolution, and systematics of *Cucurbita pepo* (Cucurbitaceae), *Economic Botany* 42: 4–13.
- DELORIT, R.J. & C.R. GUNN. 1986. *Seeds of continental United States legumes (Fabaceae)*. River Falls (WI): Agronomy Publications.
- FORD, R.I. 1985a. Patterns of prehistoric food production in North America, in Ford 1985b: 341–64.
- (Ed.). 1985b. *Prehistoric food production in North America*. Ann Arbor: University of Michigan, Museum of Anthropology. Anthropological Papers 75.
- FRITZ, G.J. 1994. Precolumbian *Cucurbita argyrosperma* ssp. *argyrosperma* (Cucurbitaceae) in the Eastern Woodlands of North America, *Economic Botany* 48: 280–92.
- GRIFFIN, J.B. 1967. Eastern North American archaeology: a summary, *Science* 156: 175–91.
- HART, J.P. 2001. Maize, matrilocality, migrations and northern Iroquoian evolution, *Journal of Archaeological Method and Theory* 8: 151–82.
- 1999a. Maize agriculture evolution in the Eastern Woodlands of North America: a Darwinian perspective, *Journal of Archaeological Method and Theory* 6: 137–80.
- 1999b. Dating Roundtop's domesticates: implications for northeastern late prehistory, in Hart 1999c: 47–68.
- (Ed.). 1999c. *Current Northeast paleoethnobotany*. Albany (NY): University of the State of New York. New York State Museum Bulletin 494.
- HART, J.P. & N. ASCH SIDELL. 1997. Additional evidence for early cucurbit use in the northern Eastern Woodlands east of the Allegheny Front, *American Antiquity* 62: 523–37.
- HART, J.P. & C.M. SCARRY. 1999. The age of common beans (*Phaseolus vulgaris*) in the northeastern United States, *American Antiquity* 64: 653–8.
- HURT, R.D. 1987. *Indian agriculture in America: prehistory to the present*. Lawrence (KS): University Press of Kansas.
- KAMI, J., V.B. VELÁSQUEZ, D.G. DEBOUCK & P. GEPTS. 1995. Identification of presumed ancestral DNA sequences of phaseolin in *Phaseolus vulgaris*, *Proceedings of the National Academy of Sciences USA* 92: 1101–4.
- KAPLAN, L. 1965. Archaeology and the domestication in American *Phaseolus* (beans), *Economic Botany* 19: 358–68.
- KAPLAN, L. & T.F. LYNCH. 1999. *Phaseolus* (Fabaceae) in archaeology: AMS radiocarbon dates and their significance for pre-Columbian agriculture, *Economic Botany* 53: 261–72.
- KIRKBRIDE, J.H., JR., C.R. GUNN, A.L. WEITZMANN & M.J. DALLWITZ. 2000. *Legume (Fabaceae) fruits and seeds*. Interactive CD-ROM. Boone (NC): Parkway Publishers.
- LOPINOT, N.H. 1992. Spatial and temporal variability in Mississippian subsistence: the archaeological record, in W.I. Woods (ed.), *Late prehistoric agriculture: observations from the Midwest*: 44–94. Springfield (IL): Illinois Historic Preservation Agency. Studies in Illinois Archaeology 8.
- MACNEISH, R.S. & M.W. EUBANKS. 2000. Comparative analysis of the Rio Balsas and Tehuacán models for the origin of maize, *Latin American Antiquity* 11: 1–18.
- PETERSEN, J.B. & N. ASCH SIDELL. 1996. Mid-Holocene evidence of *Cucurbita* sp. from central Maine, *American Antiquity* 61: 685–98.
- RILEY, T.J., R. EDGING & J. ROSSEN. 1990. Cultigens in prehistoric eastern North America: changing paradigms, *Current Anthropology* 31: 525–42.
- RILEY, T.J., G.R. WALTZ, C.J. BAREIS, A.C. FORTIER & K.E. PARKER. 1994. Accelerator mass spectrometry (AMS) dates confirm early *Zea mays* in the Mississippi River valley, *American Antiquity* 59: 490–97.
- ROSSEN, J. 1988. Botanical remains, in C.A. Turnbow & W.E. Sharp (ed.), *Muir: an early Fort Ancient site in the Inner Bluegrass*: 243–64. Lexington (KY): University of Kentucky. Archaeological Report 171. Program for Cultural Resource Assessment.
- SMITH, B.D. 1992. *Rivers of change: essays on early agriculture in eastern North America*. Washington (DC): Smithsonian Institution Press.
- SMITH, D.G. & G.W. CRAWFORD. 1997. Recent developments in the archaeology of the Princess Point complex in southern Ontario, *Journal Canadien d'Archéologie* 21: 9–32.
- STOLTMAN, J.B. & D.A. BAERREIS. 1983. The evolution of human ecosystems in the eastern United States, in H.E. Wright, Jr (ed.), *Late-Quaternary environments of the United States, Vol. 2. The Holocene*: 252–68. Minneapolis (MN): University of Minnesota Press.
- STUIVER, M., P.J. REIMER, E. BARD, J.W. BECK, G.S. BURR, K.A. HUGHEN, B. KROMER, F.G. MCCORMAC, J. VAN DER PLICHT & M. SPURK. 1998. INTCAL98 Radiocarbon age calibration 24,000–0 cal BP, *Radiocarbon* 40: 1041–83.
- WARD, G.K. & S.R. WILSON. 1978. Procedures for comparing and combining radiocarbon age determinations: a critical review, *Archaeometry* 20: 19–31.
- YARNELL, R.A. 1976. Early plant husbandry in eastern North America, in C. Cleland (ed.), *Cultural change and continuity: essays in honor of James Bennett Griffin*: 265–74. New York (NY): Academic Press.
1986. A survey of prehistoric crop plants in eastern North America, *Missouri Archaeologist* 47: 47–59.