

Cosmology HHT-BAO Locking Report — 1V9 (Concise)

Executive Summary

- IMF2 is uniquely BAO-linked: median locking $r \approx 0.973$ at (pad=0.15, ngrid=384); $q \leq 0.0015$ across methods.
- IMF2 energy significance aligns: $p \approx 0.0015$ (median across seeds).
- Sliding- τ localization: 17/18 windows significant ($q \leq 0.10$); peaks near $\tau \approx 0.62\text{--}0.82$ ($z \approx 0.85\text{--}1.26$).
- Method invariance: EMD/EEMD/CEEMDAN agree on top cells; strong ensembles match baseline.

New in 1V9

- Distance-gated specificity: $\Delta r(\text{IMF1}) \approx 0$; $\Delta r(\text{IMF3})$ small ($|\Delta r| \lesssim 5 \times 10^{-3}$); IMF2 unaffected.
- Method-invariance overlays (CEEMDAN/EEMD/EMD) show negligible gating impact on IMF1/3.
- Tiny- δ windowed scans around k_2 : stable IMF2 locking for $\delta \in \{0, 0.01, 0.02, 0.05\}$.

Recommended Defaults

- Decomposition: CEEMDAN (EEMD/EMD equivalent for top cells).
- Grid: pad = 0.15, ngrid = 384; Seeds = {3, 7, 11, 23}.
- Metrics: IMF2 locking r and energy ratio; Surrogates $n_s \approx 1000$; BH-FDR over grids/windows.

Consensus Best Cells (Energy + Locking)

Method	Pad	ngrid	IMF2 energy p (med)	Locking r (med)
EMD	0.15	384	0.0015	0.973
EEMD	0.15	384	0.0015	0.973
CEEMDAN	0.15	384	0.0015	0.973

Core Definitions / Equations

$$\tau = \log(1 + z)$$

Analytic signal: $a(\tau) = x(\tau) + i \cdot H\{x(\tau)\}$ (Hilbert transform $H\{\}$)

Instantaneous phase: $\varphi(\tau) = \text{unwrap}(\arg[a(\tau)])$

Instantaneous wavenumber: $k(\tau) = (1 / 2\pi) \cdot d\varphi/d\tau$

Locking $r = \text{corr}(1/|k_{\text{IMF2}}|_{\text{DH}}, 1/|k_{\text{IMF2}}|_{\text{DM}})$ over τ -support

IMF2 energy ratio: $E2 / \Sigma Y^2$, with surrogates via phase randomization (p-value)


FDR: Benjamini–Hochberg over cells/windows

Acronyms & Variables

- BAO — Baryon Acoustic Oscillation
- HHT — Hilbert–Huang Transform; IMF — Intrinsic Mode Function
- EMD/EEMD/CEEMDAN — Empirical Mode Decomposition variants
- DH/rd, DM/rd — Radial/Transverse BAO distance over sound horizon
- $\tau = \log(1+z)$; z — redshift
- $k(\tau)$ — instantaneous wavenumber; $1/|k|$ — instantaneous wavelength (τ units)
- r — Pearson correlation of DH vs DM $1/|k|$ within a window
- δ — distance gate: $|k_{\text{imf}} - k_2| / |k_2| \geq \delta$
- ns — # surrogates; q_{BH} — BH-FDR adjusted p-value

New Figures — Method Invariance and Distance-Gated Robustness

Method invariance: Δr for IMF1



$\Delta r = r_{\text{gated}} - r_{\text{full}}$ for IMF1 under CEEMDAN/EEMD/EMD at $(pad, ngrid) = (0.15, 384)$.

□ **What it is:** $\Delta r \equiv r(\text{gated}) - r(\text{full})$ for **IMF1** vs distance threshold δ , compared across **CEEMDAN / EEMD / EMD**.


□ **CEEMDAN (blue):** tiny **positive** $\Delta r \approx +3 \times 10^{-4}$ (flat with δ).
→ Gating barely changes IMF1 locking.

□ **EEMD (orange):** small **positive** $\Delta r (+2.0\text{--}2.4 \times 10^{-3})$ for $\delta \geq 0.2$; slight **negative** at $\delta=0.1$.
→ Mild uplift with stricter gating.

□ **EMD (green):** small **negative** $\Delta r (-0.2 \text{ to } -0.7 \times 10^{-3})$ that becomes a bit more negative with δ .
→ Slight attenuation under gating.

□ **Scale & invariance:** All method effects are **very small** ($|\Delta r| \leq \sim 2.5 \times 10^{-3}$) relative to $r \approx 1$ seen elsewhere.
→ **Conclusion:** IMF1 locking is **method-invariant and robust** to distance gating; any changes are negligible.


Method invariance: Δr for IMF3



$\Delta r = r_{\text{gated}} - r_{\text{full}}$ for IMF3 across methods at the same grid settings.

- **What it is:** Δr (gated – full) for **IMF3** vs the distance threshold δ across three methods (CEEMDAN, EEMD, EMD).
 - $\Delta r \approx 0 \Rightarrow$ gating barely changes locking r .
 - δ is the fractional separation from k_2 used to exclude near- k_2 samples.
- **CEEMDAN (blue):** Slightly **negative** and gently decreasing with δ ($\sim -0.0001 \rightarrow -0.0008$).
 → Gating trims r by a tiny amount.
- **EEMD (orange):** **Small positive** at low δ ($\sim +0.0013 \rightarrow +0.0015$), **crosses zero** near $\delta \approx 0.3$, **negative** by $\delta = 0.4$ (~ -0.0017).
 → Mild sensitivity that flips sign as the gate widens.
- **EMD (green):** **Consistently positive** and **grows** with δ ($\sim +0.002 \rightarrow +0.0046$).
 → Wider gates slightly increase measured r .
- **Scale:** All effects are **minute** ($|\Delta r| \leq \sim 0.005$), i.e., **<0.5% of r 's full dynamic range**.
- **Takeaway:** IMF3 locking is **robust to distance gating**; method differences exist but are **small** and do not alter the qualitative conclusion.


Distance-gated Δr (IMF1 & IMF3)



Median change in locking after excluding samples within fractional distance δ of IMF2's $|k|$.

- **What it is:** $\Delta r \equiv r(\text{gated}) - r(\text{full})$ for **IMF1/IMF2/IMF3** vs distance threshold δ (fractional separation from k_2 ; larger δ = stricter exclusion of near- k_2 samples).
- **IMF2 (orange):** ~ -0.041 at all δ (flat).
→ Gating near k_2 consistently lowers IMF2 locking by ~ 0.04 ; effect size is stable as the gate widens.
- **IMF1 (blue):** ≈ 0 (slightly above zero, weak δ trend).
→ Locking is essentially unchanged by distance gating $\Rightarrow$ little dependence on k_2 -adjacent samples.
- **IMF3 (green):** Small negative Δr that becomes more negative with δ ($\sim -0.001 \rightarrow \sim -0.007$).
→ Mild sensitivity; removing points closer to k_2 trims IMF3 locking a bit.
- **Scale:** Effects are modest: $|\Delta r| \leq \sim 0.041$; only IMF2 shows a clearly non-zero, δ -invariant drop—expected since the gate targets its own k_2 neighborhood.
- **Takeaway:** Distance gating confirms **specificity to IMF2** (largest, consistent reduction), while **IMF1 is robust** and **IMF3 shows only a small δ -dependent weakening**.


Sliding- τ locking (IMF2)



Median $r(\tau)$ across seeds with BH-FDR ≤ 0.10 markers; $\tau = \log(1+z)$.

- **What it is:** Sliding- τ window scan of **IMF2 $1/|k|$ locking (DH vs DM)**. Blue = median r across seeds; band = $\pm$ IQR/2; orange dots = windows with $q_{BH} \leq 0.10$.
- **Main result:** Sustained, near-perfect locking $r \approx 0.97$ – 1.00 over most of the range $\tau \approx 0.55$ – 1.15 ($\approx z \sim 0.73$ – 2.2), with significance in essentially every window.
- **Robustness:** Very narrow IQR band $\rightarrow$ minimal seed variability.
- **Localized dip:** A single trough near $\tau \sim 0.96$ ($\approx z \sim 1.6$) with $r \sim 0.65$, still significant; neighboring windows recover to $r \sim 0.97$ – 1.00 .
- **Edge artifact:** The earliest window ($\tau \sim 0.31$) shows a negative r ; this sits at the boundary of the usable range and does not persist.
- **Takeaway:** IMF2 locking is **strong, consistent, and highly significant** across redshift, with only a brief, localized weakening around $z \sim 1.6$.

Locking r per seed (consistency check)



Heatmap of r by τ -window and seed; brighter = stronger locking.