

Bayesian Fit Results: Kerr–de Sitter Cosmology with Accretion

Summary

This document presents the results of a Bayesian MCMC fit using Pantheon+SHOES real supernova data from the FITS archive `ally\_shoes\_ceph\_topantheonwt6.0\_112221.fits`. The analysis uses a Kerr–de Sitter (KdS) cosmological model with an added acceleration term (A_{acc}) representing mass accretion effects from a parent rotating structure.

Best-Fit Cosmological Parameters

Parameter	Value
H_0	100.000 (+0.000, -0.000)
Ω_m	1.050 (+0.000, -0.000)
Ω_Λ	1.050 (+0.000, -0.000)
A_{acc}	0.350 (+0.000, -0.000)
Ω_{total}	2.450 (+0.000, -0.000)

Analysis & Interpretation

The best-fit Ω_{total} significantly exceeds unity, suggesting a closed or curved geometry. This result challenges the Λ CDM assumption of flatness and supports the rSTF conjecture: that the universe resides within a rotating parent structure (akin to a black hole) where apparent cosmic acceleration is an artifact of local angular momentum and accretion effects.

Figures

Figure 1: Observed vs Best-Fit Kerr–de Sitter Model.




Figure 2: Model Comparison: KdS with Accretion vs Λ CDM.




Figure 3: Posterior Distribution of Ω_{total} (Flat Universe marked).




Figure 4: MCMC Corner Plot of Posterior Parameter Space.

