Insuffisance veineuse des membres inférieurs [uni & bilatéralité] Ages

[362 patients - 724 membres - 876 CEAP] [période 2019-11-18 à 2024-06-05] 1.0 Patients M&F MAD: 11.0 ans; 47.0(left) 69.0(right) 80 Smoothed distribution (362 patients) Smoothed distribution 156(M) 0.8 Smoothed distribution 206(F) Normal distribution Cumulative distribution (M) (F) Skeness: -0.34; -0.44(M); -0.28(F) 60 Mean: 56.7 ans; 56.7(M); 56.7(F) Median: 58.0 ans; 57.0(M); 58.0(F) -réquence Mode: 52.0 ans; 75.0(M); 52.0(F) MAD: 11.0 ans; 47.0(left) 69.0(right) Q1: 47.0 ans; 48.0(M); 46.0(F) 40 Q3: 69.0 ans; 68.2(M); 69.0(F) 0.4 Below Q1: 87 24.0% patients; 36 23.1%(M); 50 24.3%(F) Within IQR: 190 52.5% patients; 81 51.9%(M); 106 51,5%(F) Above Q3: 85 23.5% patients; 39 25.0%(M); 50 24.3%(F 20 0.2 0 0.0 30 80 10 20 40 50 60 70 90 Age du patient

pl13 pati agco sexe dia1 uti2(c3c6 full.abso)