
Table of Contents

.....	1
Problem 2.1	1
Problem 2.2	1
Problem 2.3	3

```
% SIO 229 Homework 2
clear all; close all; clc
```

Problem 2.1

Use MacCullagh's Formula to compare the difference in gravity at the pole and at the equator without Earth's spin and assume present shape and interior structure.

```
% g = -gradient of V
% V is MacCullagh's Formula
mEarth = 5.972e24;           % Mass of Earth (kg)
J = 0.001;                   % J_2 value for flattening roughly,
    unitless
a = 6378000;                 % Radius of Earth at equator m
G = 6.6743e-11;              % Gravitational constant N*m^2/kg^2

% radius and theta
rPole = 6357000;             % radius at the pole meters
thetaPole = 0;               % theta at the pole degrees

rEquator = a;                % radius at the equator meters
thetaEquator = 90;           % theta at the equator degrees

% component for g at the poles without phi
g_rPole = G*(mEarth/rPole^2-(3*a^2*J*(3*cosd(thetaPole)^2-1))/(2*rPole^4));
g_thetaPole = G*(-3*a^2*J*sind(thetaPole)*cosd(thetaPole))/rPole^4;

% component for g at the equator without phi
g_rEquator = G*(mEarth/rEquator^2-(3*a^2*J*(3*cosd(thetaEquator)^2-1))/(2*rEquator^4));
g_thetaEquator = G*(-3*a^2*J*sind(thetaEquator)*cosd(thetaEquator))/rEquator^4;
```

Problem 2.2

```
% Calculate the value of the geopotential U on the geoid

w = 7.2921150e-5;           % Earth's spin rate (rad/s)
theta = 0:180;               % degrees of latitude
```

```

f = (3/2)*J+(a^3*w^2)/(2*G*mEarth);      % flattening, unitless
r0 = a*(1-f*cosd(theta).^2);              % changing radius value in
    meters
P = (1/2)*(3*cosd(theta).^2-1);           % Legendre Polynomial,
    unitless

mono = -G*mEarth./r0;                     % Monopole N*m/kg
quad = (G*mEarth*a^2)*(J*P)./(r0.^3);     % Quadrupole N*m/kg
spin = -(1/2)*w^2*r0.^2.*sind(theta).^2;  % Rotation m^2/s^2

U0= mono+quad+spin;                       % Geopotential U on the geoid


U0mean = mean(U0)

figure(1)
subplot(2,1,1)
plot(theta, U0, 'LineWidth', 1)
hold on
plot(theta, mono, 'LineWidth', 1)
ylabel('Potential (m^2/s^2 or Nm/kg)', 'FontSize', 15)
legend('U_0', 'Monopole')
hold off
title('U_0 Summation of Monopole, Quadrupole, and
    Rotation', 'FontSize', 15)
subplot(2,1,2)
plot(theta, quad, 'r', 'LineWidth', 1)
hold on
plot(theta, spin, 'k', 'LineWidth', 1)
ylabel('Potential (m^2/s^2 or Nm/kg)', 'FontSize', 15)
legend('Quadrupole', 'Rotation')
xlabel('Colatitude \theta (degrees)', 'FontSize', 15)
set(gcf, 'color', 'w');
hold off

U0mean =

    -6.2634e+07

```



Problem 2.3

```
% Find the rotation rate, omega, when the Mississippi starts to flow
% backward from South to North. Assuming the Earth does not change
% shape.
% The source of the Mississippi is 47 degrees North and the delta is
% 29
% degrees North of the Equator.

MissLatS = 43;           % Mississippi source colatitude from the pole
degrees
MissLatD = 61;           % Mississippi delta colatitude from the pole
degrees

radiusSource = r0(44)+450; % radius of source in meters
Psource = (1/2)*(3*cosd(MissLatS).^2-1); % Legendre Polynomial
for Source

% Vary omega for when potential for the source and delta are equal
omega = (-1e-4:1e-7:1e-4); % Spin radian/sec

% Calculate the source gravitational potential, V source in N*m/kg or
% m^2/s^2
Vsource = -G*mEarth/radiusSource + G*mEarth*a^2*J*Psource/
(radiusSource^3);
```

```

% Calculate the source apparent centrifugal force, depends on omega
m^2/s^2
SpinSource = -(1/2)*omega.^2*radiusSource^2*sind(MissLatS)^2;

% Calculate the delta gravitational potential, geoid potential, V in
N*m/kg
% depends on omega
f = (3/2)*J+(a^3*omega.^2)/(2*G*mEarth); % flattening depends on
rot. rate
r0Delta = a*(1-f.*cosd(MissLatD).^2); % radius meters
depends on rotation rate
Pdelta = (1/2)*(3*cosd(MissLatD)^2-1); % Legendre Polynomial,
unitless
Vdelta = -G*mEarth./r0Delta + G*mEarth*a^2*J*Pdelta./(r0Delta.^3);

% Calculate the delta apparent centrifugal force m^2/s^2, depends on
omega
SpinDelta = -(1/2)*omega.^2.*r0Delta.^2*sind(MissLatD)^2;

% Source and Delta potential equalling zero
SandDequal0 = -Vsource - SpinSource + Vdelta + SpinDelta;

T = (2*pi./omega)/(60*60); % Period of rotation of Earth in hours
Topp = T(300) % Period in the opposite direction
Tsame = T(1702) % Period for the same direction today

figure(2)
plot(omega, SandDequal0,'LineWidth',1)
hold on
plot(omega, zeros(length(omega)), 'LineWidth',1)
plot([omega(1702) omega(300)], [SandDequal0(1702)
SandDequal0(300)], 'k*', 'LineWidth',1)
% plot(omega(300), SandDequal0(300), 'k*', 'LineWidth',1)
plot(w, 0, 'mx', 'LineWidth',1)
ylabel('Potential (m^2/s^2 or Nm/kg)', 'FontSize',15)
xlabel('\omega rad/sec', 'FontSize',15)
title('Potential of Varying \omega', 'FontSize',15)
set(gcf, 'color', 'w')
hold off


Topp =

-24.8977

Tsame =

24.8977

```



Published with MATLAB® R2020a