

Tylenol Cold + Flu + Cough Night

Drug Facts

<i>Active ingredients (in each 15 mL)</i>	<i>Purpose</i>
Acetaminophen 325 mg	Pain reliever/fever reducer
Dextromethorphan HBr 10 mg	Cough suppressant
Doxylamine Succinate 6.25 mg	Antihistamine
Phenylephrine HCl 5 mg	Nasal decongestant

Uses

- temporarily relieves these common cold/flu symptoms:
 - minor aches and pains
 - headache
 - sore throat
 - nasal congestion
 - runny nose and sneezing
 - cough
 - sinus congestion and pressure
- helps clear nasal passages
- relieves cough to help you sleep
- temporarily reduces fever

Warnings

Liver warning

This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: acetaminophen may cause severe skin reactions. Symptoms may include:

- skin reddening
- blisters
- rash

If a skin reaction occurs, stop use and seek medical help right away.

Sore throat warning

If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If

you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.

- if you have ever had an allergic reaction to this product or any of its ingredients

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- trouble urinating due to an enlarged prostate gland
- a breathing problem such as emphysema or chronic bronchitis
- glaucoma
- persistent or chronic cough such as occurs with smoking, asthma or emphysema
- cough that occurs with too much phlegm (mucus)

Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

When using this product

- **do not exceed recommended dose**
- excitability may occur, especially in children
- marked drowsiness may occur
- alcohol, sedatives and tranquilizers may increase drowsiness
- avoid alcoholic drinks
- be careful when driving a motor vehicle or operating machinery

Stop use and ask a doctor if

- nervousness, dizziness, or sleeplessness occur
- pain, nasal congestion or cough gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts

These could be signs of a serious condition.

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children.

Overdose warning

In case of overdose, get medical help or contact a Poison Control Center right away. (1-800-222-1222) Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- **do not take more than directed** (see **overdose warning**)
- mL = milliliter
- use only enclosed dosing cup designed for use with this product. Do not use any other dosing device.

adults and children 12 years and over	<ul style="list-style-type: none"> ▪ take 30 mL in the dosing cup provided every 4 hours while symptoms last ▪ do not take more than 150 mL in 24 hours, unless directed by a doctor
children under 12 years	ask a doctor

Other information

- each 15 mL contains: **sodium 5 mg**
- store between 20-25°C (68-77°F). Do not refrigerate.
- **do not use if neck band imprinted with "TYLENOL COLD+FLU+COUGH" or foil inner seal imprinted with "SAFETY SEAL®" is broken or missing**

Inactive ingredients

acetic acid, alcohol, anhydrous citric acid, FD&C blue no. 1, FD&C red no. 40, flavors, glycerin, propylene glycol, purified water, sodium benzoate, sorbitol solution, sucralose

Questions or comments?

call **1-877-895-3665** (toll-free) or **215-273-8755** (collect)

PRINCIPAL DISPLAY PANEL

NDC 50580-369-01

TYLENOL®

FOR ADULTS

COLD + FLU + COUGH

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,
Pain Reliever–Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant

- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE

NIGHT

WILD BERRY

BURST

Alcohol 0.7%

8 fl oz (240 mL)

30040371

NDC 50580-369-01

TYLENOL[®]

FOR ADULTS

COLD + FLU + COUGH

Acetaminophen, Dextromethorphan HBr, Doxylamine Succinate, Phenylephrine HCl,
Pain Reliever–Fever Reducer, Cough Suppressant, Antihistamine, Nasal Decongestant



- HEAD + BODY ACHES
- FEVER + SORE THROAT
- COUGH
- NASAL CONGESTION
- RUNNY NOSE



NIGHT



WILD BERRY
BURST



Alcohol 0.7%

8 fl oz (240 mL)

OPEN TO READ DRUG FACTS
(Warnings, Directions...)



PEEL



30040372 Do not use if neck band imprinted with
"TYLENOL COLD+FLU+ COUGH" or foil
inner seal imprinted with "SAFETY
SEAL[®]" is broken or missing.

**DO NOT USE WITH OTHER MEDICINES
CONTAINING ACETAMINOPHEN**

