

CBSE Class–10 Mathematics

NCERT solution

Chapter - 6

Triangles - Exercise 6.1

1. Fill in the blanks using the correct word given in brackets:

(i) All circles are _____. (congruent, similar)

(ii) All squares are _____. (similar, congruent)

(iii) All _____ triangles are similar. (isosceles, equilateral)

(iv) Two polygons of the same number of sides are similar, if (a) their corresponding angles are _____ and (b) their corresponding sides are _____. (equal, proportional)

Ans. (i) similar

(ii) similar

(iii) equilateral

(iv) equal, proportional

2. Give two different examples of pair of:

(i) similar figures

(ii) non-similar figures

Ans. (i) Two different examples of a pair of similar figures are:

(a) Any two rectangles



(b) Any two squares

(ii) Two different examples of a pair of non-similar figures are:

(a) A scalene and an equilateral triangle

(b) An equilateral triangle and a right angled triangle

3. State whether the following quadrilaterals are similar or not:



Ans. On looking at the given figures of the quadrilaterals, we can say that they are not similar because their angles are not equal.


CBSE Class-10 Mathematics

NCERT solution

Chapter - 6

Triangles - Exercise 6.2

1. In figure (i) and (ii), $DE \parallel BC$. Find EC in (i) and AD in (ii).



Ans. (i) Since $DE \parallel BC$,

$$\therefore \frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow \frac{1.5}{3} = \frac{1}{EC}$$

$$\Rightarrow EC = \frac{3}{1.5}$$

$$\Rightarrow EC = 2 \text{ cm}$$

(ii) Since $DE \parallel BC$,

$$\therefore \frac{AD}{DB} = \frac{AE}{EC}$$

$$\Rightarrow \frac{AD}{7.2} = \frac{1.8}{5.4}$$

$$\Rightarrow AD = \frac{1.8 \times 7.2}{5.4}$$

$$\Rightarrow EC = 2.4 \text{ cm}$$

2. E and F are points on the sides PQ and PR respectively of a $\triangle PQR$. For each of the following cases, state whether $EF \parallel QR$:


(i) $PE = 3.9 \text{ cm}$, $EQ = 4 \text{ cm}$, $PF = 3.6 \text{ cm}$ and $FR = 2.4 \text{ cm}$

(ii) $PE = 4 \text{ cm}$, $QE = 4.5 \text{ cm}$, $PF = 8 \text{ cm}$ and $RF = 9 \text{ cm}$

(iii) $PQ = 1.28 \text{ cm}$, $PR = 2.56 \text{ cm}$, $PE = 0.18 \text{ cm}$ and $PF = 0.36 \text{ cm}$

Ans. (i) Given: $PE = 3.9 \text{ cm}$, $EQ = 4 \text{ cm}$, $PF = 3.6 \text{ cm}$ and $FR = 2.4 \text{ cm}$

Now, $\frac{PE}{EQ} = \frac{3.9}{4} = 0.97 \text{ cm}$



And $\frac{PF}{FR} = \frac{3.6}{2.4} = 1.2 \text{ cm}$

$$\therefore \frac{PE}{EQ} \neq \frac{PF}{FR}$$

Therefore, EF does not divide the sides PQ and PR of $\triangle PQR$ in the same ratio.

\therefore EF is not parallel to QR.

(ii) Given: $PE = 4 \text{ cm}$, $QE = 4.5 \text{ cm}$, $PF = 8 \text{ cm}$ and $RF = 9 \text{ cm}$

Now, $\frac{PE}{EQ} = \frac{4}{4.5} = \frac{8}{9}$ cm

And $\frac{PF}{FR} = \frac{8}{9}$ cm

$\therefore \frac{PE}{EQ} = \frac{PF}{FR}$

Therefore, EF divides the sides PQ and PR of $\triangle PQR$ in the same ratio.

\therefore EF is parallel to QR.

(iii) Given: PQ = 1.28 cm, PR = 2.56 cm, PE = 0.18 cm and PF = 0.36 cm

$\Rightarrow EQ = PQ - PE = 1.28 - 0.18 = 1.10$ cm

And ER = PR - PF = 2.56 - 0.36 = 2.20 cm

Now, $\frac{PE}{EQ} = \frac{0.18}{1.10} = \frac{18}{110} = \frac{9}{55}$ cm


And $\frac{PF}{FR} = \frac{0.36}{2.20} = \frac{36}{220} = \frac{9}{55}$ cm

$\therefore \frac{PE}{EQ} = \frac{PF}{FR}$

Therefore, EF divides the sides PQ and PR of $\triangle PQR$ in the same ratio.

\therefore EF is parallel to QR.

3. In figure, if LM \parallel CB and LN \parallel CD, prove that $\frac{AM}{AB} = \frac{AN}{AD}$.



Ans. In $\triangle ABC$, $LM \parallel CB$

$$\therefore \frac{AM}{AB} = \frac{AL}{AC} \text{ [Basic Proportionality theorem](i)}$$


And in $\triangle ACD$, $LN \parallel CD$

$$\therefore \frac{AL}{AC} = \frac{AN}{AD} \text{ [Basic Proportionality theorem](ii)}$$

From eq. (i) and (ii), we have

$$\frac{AM}{AB} = \frac{AN}{AD}$$

4. In the given figure, $DE \parallel AC$ and $DF \parallel AE$. Prove that $\frac{BF}{FE} = \frac{BE}{EC}$.



Ans. In $\triangle BCA$, $DE \parallel AC$

$$\therefore \frac{BE}{EC} = \frac{BD}{DA} \text{ [Basic Proportionality theorem](i)}$$


And in $\triangle BEA$, $DF \parallel AE$

$$\therefore \frac{BE}{FE} = \frac{BD}{DA} \text{ [Basic Proportionality theorem](ii)}$$

From eq. (i) and (ii), we have

$$\frac{BF}{FE} = \frac{BE}{EC}$$

5. In the given figure, $DE \parallel OQ$ and $DF \parallel OR$. Show that $EF \parallel QR$.



Ans. In $\triangle PQO$, $DE \parallel OQ$

$$\therefore \frac{PE}{EQ} = \frac{PD}{DO} \text{ [Basic Proportionality theorem](i)}$$

And in $\triangle POR$, $DF \parallel OR$


$$\therefore \frac{PD}{DO} = \frac{PF}{FR} \text{ [Basic Proportionality theorem](ii)}$$

From eq. (i) and (ii), we have

$$\frac{PE}{EQ} = \frac{PF}{FR}$$

$\therefore EF \parallel QR$ [By the converse of BPT]

6. In the given figure, A, B, and C are points on OP, OQ and OR respectively such that $AB \parallel PQ$ and $AC \parallel PR$. Show that $BC \parallel QR$.



Ans. Given: O is any point in $\triangle PQR$, in which $AB \parallel PQ$ and $AC \parallel PR$.

To prove: $BC \parallel QR$

Construction: Join BC.

Proof: In $\triangle OPQ$, $AB \parallel PQ$

$$\therefore \frac{OA}{AP} = \frac{OB}{BQ} \text{ [Basic Proportionality theorem](i)}$$

And in $\triangle OPR$, $AC \parallel PR$

$$\therefore \frac{OA}{AP} = \frac{OC}{CR} \text{ [Basic Proportionality theorem](ii)}$$

From eq. (i) and (ii), we have

$$\frac{OB}{BQ} = \frac{OC}{CR}$$

\therefore In $\triangle OQR$, B and C are points dividing the sides OQ and OR in the same ratio.


\therefore By the converse of Basic Proportionality theorem,

$$\Rightarrow BC \parallel QR$$

7. Using Theorem 6.1, prove that a line drawn through the midpoint of one side of a triangle parallel to another side bisects the third side. (Recall that you have proved it in Class IX).

Ans. Given: A triangle ABC, in which D is the midpoint of side AB

and the line DE is drawn parallel to BC, meeting AC at E.



To prove: $AE = EC$

Proof: Since $DE \parallel BC$

$$\therefore \frac{AD}{DB} = \frac{AE}{EC} \text{ [Basic Proportionality theorem](i)}$$

But $AD = DB$ [Given]

$$\Rightarrow \frac{AD}{DB} = 1$$

$$\Rightarrow \frac{AE}{EC} = 1 \text{ [From eq. (i)]}$$


$$\Rightarrow AE = EC$$

Hence, E is the midpoint of the third side AC.

8. Using Theorem 6.2, prove that the line joining the midpoints of any two sides of a triangle is parallel to the third side. (Recall that you have done it in Class IX).

Ans. Given: A triangle ABC, in which D and E are the midpoints of

sides AB and AC respectively.



To Prove: $DE \parallel BC$

Proof: Since D and E are the midpoints of AB and AC respectively.

$$\therefore AD = DB \text{ and } AE = EC$$

Now, $AD = DB$

$$\Rightarrow \frac{AD}{DB} = 1 \text{ and } AE = EC$$

$$\Rightarrow \frac{AE}{EC} = 1$$

$$\therefore \frac{AD}{DB} = \frac{AE}{EC} = 1$$

$$\Rightarrow \frac{AD}{DB} = \frac{AE}{EC}$$

Thus, in triangle ABC, D and E are points dividing the sides AB and AC in the same ratio.


Therefore, by the converse of Basic Proportionality theorem, we have

$DE \parallel BC$

9. ABCD is a trapezium in which $AB \parallel DC$ and its diagonals intersect each other at the point O. Show that $\frac{AO}{BO} = \frac{CO}{DO}$.

Ans. Given: A trapezium ABCD, in which $AB \parallel DC$ and its diagonals

AC and BD intersect each other at O.



To Prove: $\frac{AO}{BO} = \frac{CO}{DO}$

Construction: Through O, draw $OE \parallel AB$, i.e. $OE \parallel DC$.

Proof: In $\triangle ADC$, we have $OE \parallel DC$

$$\therefore \frac{AE}{ED} = \frac{AO}{CO} \text{ [By Basic Proportionality theorem].....(i)}$$

Again, in $\triangle ABD$, we have $OE \parallel AB$ [Construction]

$$\therefore \frac{ED}{AE} = \frac{DO}{BO} \text{ [By Basic Proportionality theorem]}$$

$$\Rightarrow \frac{AE}{ED} = \frac{BO}{DO} \text{(ii)}$$

From eq. (i) and (ii), we get

$$\frac{AO}{CO} = \frac{BO}{DO}$$


$$\Rightarrow \frac{AO}{BO} = \frac{CO}{DO}$$

10. The diagonals of a quadrilateral ABCD intersect each other at the point O such that

$$\frac{AO}{BO} = \frac{CO}{DO} \text{ . Show that ABCD is a trapezium.}$$

Ans. Given: A quadrilateral ABCD, in which its diagonals AC and

BD intersect each other at O such that $\frac{AO}{BO} = \frac{CO}{DO}$, i.e.



$$\frac{AO}{CO} = \frac{BO}{DO}.$$

To Prove: Quadrilateral ABCD is a trapezium.

Construction: Through O, draw OE \parallel AB meeting AD at E.

Proof: In $\triangle ADB$, we have OE \parallel AB [By construction]

$$\therefore \frac{DE}{EA} = \frac{OD}{BO} \text{ [By Basic Proportionality theorem]}$$

$$\Rightarrow \frac{EA}{DE} = \frac{BO}{DO}$$

$$\Rightarrow \frac{EA}{DE} = \frac{BO}{DO} = \frac{AO}{CO}$$

$$\left[\because \frac{AO}{CO} = \frac{BO}{DO} \right]$$

$$\Rightarrow \frac{EA}{DE} = \frac{AO}{CO}$$

Thus in $\triangle ADC$, E and O are points dividing the sides AD and AC in the same ratio. Therefore by the converse of Basic Proportionality theorem, we have


$$EO \parallel DC$$

But EO \parallel AB [By construction]

$$\therefore AB \parallel DC$$

\therefore Quadrilateral ABCD is a trapezium

1. State which pairs of triangles in the given figure, are similar. Write the similarity criterion used by you for answering the question and also write the pairs of similar triangles in the symbolic form:



Ans. (i) In Δ s ABC and PQR, we observe that,

$$\angle A = \angle P = 60^\circ, \angle B = \angle Q = 80^\circ \text{ and } \angle C = \angle R = 40^\circ$$

\therefore By AAA criterion of similarity, $\Delta ABC \sim \Delta PQR$

(ii) In Δ s ABC and PQR, we observe that,

$$\frac{AB}{QR} = \frac{BC}{RP} = \frac{CA}{PQ} = \frac{1}{2}$$

\therefore By SSS criterion of similarity, $\Delta ABC \sim \Delta PQR$

(iii) In Δ s LMP and DEF, we observe that the ratio of the sides of these triangles is not equal.

Therefore, these two triangles are not similar.

(iv) In Δ s MNL and QPR, we observe that, $\angle M = \angle Q = 70^\circ$

But, $\frac{MN}{PQ} \neq \frac{ML}{QR}$

\therefore These two triangles are not similar as they do not satisfy SAS criterion of similarity.

(v) In Δ s ABC and FDE, we have, $\angle A = \angle F = 80^\circ$

But, $\frac{AB}{DE} \neq \frac{AC}{DF}$ [\because AC is not given]

\therefore These two triangles are not similar as they do not satisfy SAS criterion of similarity.


(vi) In Δ s DEF and PQR, we have, $\angle D = \angle P = 70^\circ$

$$[\because \angle P = 180^\circ - 80^\circ - 30^\circ = 70^\circ]$$

And $\angle E = \angle Q = 80^\circ$

\therefore By AAA criterion of similarity, $\Delta DEF \sim \Delta PQR$

2. In figure, $\triangle ODC \sim \triangle OBA$, $\angle BOC = 125^\circ$ and $\angle CDO = 70^\circ$. Find $\angle DOC$, $\angle DCO$ and $\angle OAB$.



Ans. Since BD is a line and OC is a ray on it.

$$\therefore \angle DOC + \angle BOC = 180^\circ$$

$$\Rightarrow \angle DOC + 125^\circ = 180^\circ$$

$$\Rightarrow \angle DOC = 55^\circ$$

In $\triangle CDO$, we have $\angle CDO + \angle DOC + \angle DCO = 180^\circ$

$$\Rightarrow 70^\circ + 55^\circ + \angle DCO = 180^\circ$$

$$\Rightarrow \angle DCO = 55^\circ$$

It is given that $\triangle ODC \sim \triangle OBA$


$$\therefore \angle OBA = \angle ODC, \angle OAB = \angle OCD$$

$$\Rightarrow \angle OBA = 70^\circ, \angle OAB = 55^\circ$$

Hence $\angle DOC = 55^\circ$, $\angle DCO = 55^\circ$ and $\angle OAB = 55^\circ$

3. Diagonals AC and BD of a trapezium ABCD with $AB \parallel CD$ intersect each other at the point O. Using a similarity criterion for two triangles, show that $\frac{OA}{OC} = \frac{OB}{OD}$.

Ans. Given: ABCD is a trapezium in which $AB \parallel DC$.



To Prove: $\frac{OA}{OC} = \frac{OB}{OD}$

Proof: In Δ s OAB and OCD, we have,

$\angle 5 = \angle 6$ [Vertically opposite angles]


$\angle 1 = \angle 2$ [Alternate angles]

And $\angle 3 = \angle 4$ [Alternate angles]

\therefore By AAA criterion of similarity, $\Delta OAB \sim \Delta ODC$

Hence, $\frac{OA}{OC} = \frac{OB}{OD}$

4. In figure, $\frac{QR}{QS} = \frac{QT}{PR}$ and $\angle 1 = \angle 2$. Show that $\Delta PQS \sim \Delta TQR$.



Ans. We have, $\frac{QR}{QS} = \frac{QT}{PR}$

$$\Rightarrow \frac{QT}{QR} = \frac{PR}{QS} \dots\dots\dots(1)$$

Also, $\angle 1 = \angle 2$ [Given]

∴ $PR = PQ$ (2) [∵ Sides opposite to equal \angle s are equal]

From eq.(1) and (2), we get

$$\frac{QT}{QR} = \frac{PR}{QS} \Rightarrow \frac{PQ}{QT} = \frac{QS}{QR}$$


In Δ s PQS and TQR, we have,

$$\frac{PQ}{QT} = \frac{QS}{QR} \text{ and } \angle PQS = \angle TQR = \angle Q$$

∴ By SAS criterion of similarity, $\Delta PQS \sim \Delta TQR$

5. S and T are points on sides PR and QR of a ΔPQR such that $\angle P = \angle RTS$. Show that $\Delta RPQ \sim \Delta RTS$.

Ans. In Δ s RPQ and RTS, we have




$$\angle RPQ = \angle RTS \text{ [Given]}$$

$$\angle PRQ = \angle TRS \text{ [Common]}$$

∴ By AA-criterion of similarity,

$$\Delta RPQ \sim \Delta RTS$$

6. In the given figure, if $\Delta ABE \cong \Delta ACD$, show that $\Delta ADE \sim \Delta ABC$.



Ans. It is given that $\triangle ABE \cong \triangle ACD$

$\therefore AB = AC$ and $AE = AD$

$$\Rightarrow \frac{AB}{AD} = \frac{AC}{AE}$$

$$\Rightarrow \frac{AB}{AC} = \frac{AD}{AE} \dots\dots\dots(1)$$


\therefore In \triangle s ADE and ABC, we have,

$$\frac{AB}{AC} = \frac{AD}{AE} \text{ [from eq.(1)]}$$

And $\angle BAC = \angle DAE$ [Common]

Thus, by SAS criterion of similarity, $\triangle ADE \sim \triangle ABC$

7. In figure, altitude AD and CE of a $\triangle ABC$ intersect each other at the point P. Show that:



(i) $\triangle AEP \sim \triangle CDP$

(ii) $\triangle ABD \sim \triangle CBE$

(iii) $\triangle AEP \sim \triangle ADB$

(iv) $\triangle PDC \sim \triangle BEC$

Ans. (i) In \triangle s AEP and CDP, we have,

$$\angle AEP = \angle CDP = 90^\circ \text{ [}\because \text{CE} \perp \text{AB, AD} \perp \text{BC]}$$

And $\angle APE = \angle CPD$ [Vertically opposite]

\therefore By AA-criterion of similarity, $\triangle AEP \sim \triangle CDP$

(ii) In \triangle s ABD and CBE, we have,

$$\angle ADB = \angle CEB = 90^\circ$$

And $\angle ABD = \angle CBE$ [Common]

\therefore By AA-criterion of similarity, $\triangle ABD \sim \triangle CBE$

(iii) In \triangle s AEP and ADB, we have,

$$\angle AEP = \angle ADB = 90^\circ \text{ [}\because \text{AD} \perp \text{BC, CE} \perp \text{AB]}$$

And $\angle PAE = \angle DAB$ [Common]

\therefore By AA-criterion of similarity, $\triangle AEP \sim \triangle ADB$

(iv) In \triangle s PDC and BEC, we have,


$$\angle PDC = \angle BEC = 90^\circ \text{ [}\because \text{CE} \perp \text{AB, AD} \perp \text{BC]}$$

And $\angle PCD = \angle BEC$ [Common]

\therefore By AA-criterion of similarity, $\triangle PDC \sim \triangle BEC$

8. E is a point on the side AD produced of a parallelogram ABCD and BE intersects CD at F. Show that $\triangle ABE \sim \triangle CFB$.

Ans. In \triangle s ABE and CFB, we have,




$$\angle AEB = \angle CBF [\text{Alt. } \angle \text{s}]$$

$$\angle A = \angle C [\text{opp. } \angle \text{s of a } \parallel \text{ gm}]$$

\therefore By AA-criterion of similarity, we have

$$\triangle ABE \sim \triangle CFB$$

9. In the given figure, ABC and AMP are two right triangles, right angles at B and M respectively. Prove that:



(i) $\triangle ABC \sim \triangle AMP$

(ii) $\frac{CA}{PA} = \frac{BC}{MP}$

Ans. (i) In \triangle s ABC and AMP, we have,

$$\angle ABC = \angle AMP = 90^\circ [\text{Given}]$$

$$\angle BAC = \angle MAP [\text{Common angles}]$$

\therefore By AA-criterion of similarity, we have

$$\triangle ABC \sim \triangle AMP$$

(ii) We have $\triangle ABC \sim \triangle AMP$ [As prove above]

$$\Rightarrow \frac{CA}{PA} = \frac{BC}{MP}$$


10. CD and GH are respectively the bisectors of $\angle ACB$ and $\angle EGF$ such that D and H lie on sides AB and FE at $\triangle ABC$ and $\triangle EFG$ respectively. If $\triangle ABC \sim \triangle FEG$, show that:

(i) $\frac{CD}{GH} = \frac{AC}{FG}$

(ii) $\triangle DCB \sim \triangle HE$

(iii) $\triangle DCA \sim \triangle HGF$

Ans. We have, $\triangle ABC \sim \triangle FEG$



$$\Rightarrow \angle A = \angle F \dots \dots (1)$$

And $\angle C = \angle G$

$$\Rightarrow \frac{1}{2} \angle C = \frac{1}{2} \angle G$$

$$\Rightarrow \angle 1 = \angle 3 \text{ and } \angle 2 = \angle 4 \dots \dots (2)$$

[\because CD and GH are bisectors of $\angle C$ and $\angle G$

respectively]

\therefore In \triangle s DCA and HGF, we have

$$\angle A = \angle F \text{ [From eq.(1)]}$$

$$\angle 2 = \angle 4 [\text{From eq.(2)}]$$

\therefore By AA-criterion of similarity, we have

$$\triangle DCA \sim \triangle HGF$$

Which proves the (iii) part

We have, $\triangle DCA \sim \triangle HGF$

$$\Rightarrow \frac{AG}{FG} = \frac{CD}{GH}$$

$$\Rightarrow \frac{CD}{GH} = \frac{AC}{FG}$$

Which proves the (i) part


In \triangle s DCA and HGF, we have

$$\angle 1 = \angle 3 [\text{From eq.(2)}]$$

$$\angle B = \angle E [\because \triangle DCB \sim \triangle HE]$$

Which proves the (ii) part

11. In the given figure, E is a point on side CB produced of an isosceles triangle ABC with $AB = AC$. If $AD \perp BC$ and $EF \perp AC$, prove that $\triangle ABD \sim \triangle ECF$.



Ans. Here $\triangle ABC$ is isosceles with $AB = AC$

$$\therefore \angle B = \angle C$$

In \triangle s ABD and ECF, we have


$$\angle ABD = \angle ECF [\because \angle B = \angle C]$$

$$\angle ABD = \angle ECF = 90^\circ [\because AD \perp BC \text{ and } EF \perp AC]$$

\therefore By AA-criterion of similarity, we have

$$\triangle ABD \sim \triangle ECF$$

12. Sides AB and BC and median AD of a triangle ABC are respectively proportional to sides PQ and QR and median PM of a \triangle PQR (see figure). Show that $\triangle ABC \sim \triangle PQR$.



Ans. Given: AD is the median of $\triangle ABC$ and PM is the median of $\triangle PQR$ such that

$$\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AD}{PM}$$

To prove: $\triangle ABC \sim \triangle PQR$

Proof: $BD = \frac{1}{2} BC$ [Given]

And $QM = \frac{1}{2} QR$ [Given]

Also $\frac{AB}{PQ} = \frac{BC}{QR} = \frac{AD}{PM}$ [Given]

$$\Rightarrow \frac{AB}{PQ} = \frac{2BD}{2QM} = \frac{AD}{PM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$$

$\therefore \triangle ABD \sim \triangle PQM$ [By SSS-criterion of similarity]

$\Rightarrow \angle B = \angle Q$ [Similar triangles have corresponding angles equal]


And $\frac{AB}{PQ} = \frac{BC}{QR}$ [Given]

\therefore By SAS-criterion of similarity, we have

$$\triangle ABC \sim \triangle PQR$$

13. D is a point on the side BC of a triangle ABC such that $\angle ADC = \angle BAC$. Show that $CA^2 = CB \cdot CD$.

ANS. In triangles ABC and DAC,



$$\angle ADC = \angle BAC \text{ [Given]}$$

$$\text{and } \angle C = \angle C \text{ [Common]}$$

\therefore By AA-similarity criterion,

$$\triangle ABC \sim \triangle DAC$$


$$\Rightarrow \frac{AB}{DA} = \frac{BC}{AC} = \frac{AC}{DC}$$

$$\Rightarrow \frac{CB}{CA} = \frac{CA}{CD}$$

$$\Rightarrow CA^2 = CB.CD$$

14. Sides AB and AC and median AD of a triangle ABC are respectively proportional to sides PQ and PR and median PM of another triangle PQR. Show that $\triangle ABC \sim \triangle PQR$.


ANS. Given: AD is the median of $\triangle ABC$ and PM is the median of $\triangle PQR$ such that



To prove: $\triangle ABC \sim \triangle PQR$

Proof:

Let us extend AD to point D such that AD = DE and PM up to point L such that PM = ML



Join B to E. C to E, and Q to L, and R to L

We know that medians is the bisector of opposite side

Hence

$$BD = DC$$

Also, AD = DE (by construction)

Hence in quadrilateral ABEC, diagonals AE and BC bisect each other at point D.

Therefore, quadrilateral ABEC is a parallelogram.

$$AC = BE$$

$$AB = EC \text{ (opposite sides of } \parallel \text{ gm are equal) (2)}$$

Similarly, we can prove that PQLR is a parallelogram

$$PR = QL$$

$$PQ = LR \text{ opposite sides of } \parallel \text{ gm are equal) (3)}$$

Given that


$$\frac{AB}{PQ} = \frac{AC}{PR} = \frac{AD}{PM}$$

$$\frac{AB}{PQ} = \frac{BE}{QL} = \frac{AD}{PM} \text{ [from (2) (3)]}$$

$$\frac{AB}{PQ} = \frac{BE}{QL} = \frac{2AD}{2PM}$$

$$\frac{AB}{PQ} = \frac{BE}{QL} = \frac{AE}{PL} \text{ [as } AD = DE, AE = AD + DE = 2AD \\ PM = ML. PL = PM + ML = 2PM]$$


$\triangle ABE \sim \triangle PQL$ (By SSS Similiarity Criteria)



We know that corresponding angles of similar triangles are equal.

$$\angle BAE = \angle QPL \text{ (4)}$$

Similarly, we can prove that $\triangle AEC \sim \triangle PLR$.



We know that corresponding angles of similar triangles are equal.


$$\angle CAE = \angle RPL \quad (5)$$

Adding (4) and (5),

$$\angle BAE + \angle CAE = \angle QPL + \angle RPL$$

$$\angle CAB = \angle RPQ$$

In $\triangle ABC$ and $\triangle PQR$,



$$\frac{AB}{PQ} = \frac{AC}{PR}$$


$$\angle CAB = \angle RPQ$$

$$\triangle ABC \sim \triangle PQR$$

Hence proved

15. A vertical pole of length 6 m casts a shadow 4 m long on the ground and at the same time a tower casts a shadow 28 m long. Find the height of the tower.

Ans. Let AB be the vertical pole and AC be its shadow. Also, let DE be the vertical tower and DF be its shadow. Joined BC and EF.




Let $DE = x$ meters

Here, $AB = 6$ m, $AC = 4$ m and $DF = 28$ m

In the triangles ABC and DEF,

$$\angle A = \angle D = 90^\circ$$

And $\angle C = \angle F$ [Each is the angular elevation of the sun]



\therefore By AA-similarity criterion,

$$\triangle ABC \sim \triangle DEF$$

$$\Rightarrow \frac{AB}{DE} = \frac{AC}{DF}$$

$$\Rightarrow \frac{6}{x} = \frac{4}{28}$$

$$\Rightarrow \frac{6}{x} = \frac{1}{7}$$


$$\Rightarrow x = 42 \text{ m}$$

Hence, the height of the tower is 42 meters.

16. If AD and PM are medians of triangles ABC and PQR respectively, where $\triangle ABC \sim \triangle PQR$, prove that $\frac{AB}{PQ} = \frac{AD}{PM}$.

Ans. Given: AD and PM are the medians of triangles

ABC and PQR respectively, where



$$\triangle ABC \sim \triangle PQR$$

To prove: $\frac{AB}{PQ} = \frac{AD}{PM}$

Proof: In triangles ABD and PQM,

$$\angle B = \angle Q \text{ [Given]}$$

$$\text{And } \frac{AB}{PQ} = \frac{\frac{1}{2}BC}{\frac{1}{2}QR} \text{ [}\because \text{ AD and PM are the medians of BC and QR respectively]}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM}$$

\therefore By SAS-criterion of similarity,

$$\triangle ABD \sim \triangle PQM$$

$$\Rightarrow \frac{AB}{PQ} = \frac{BD}{QM} = \frac{AD}{PM}$$

$$\Rightarrow \frac{AB}{PQ} = \frac{AD}{PM}$$

CBSE Class-10 Mathematics

NCERT solution

Chapter - 6

Triangles - Exercise 6.4

1. Let $\triangle ABC \sim \triangle DEF$ and their areas be, respectively, 64 cm^2 and 121 cm^2 . If $EF = 15.4 \text{ cm}$, find BC .

Ans. We have, $\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = \frac{BC^2}{EF^2}$


$$\Rightarrow \frac{64}{121} = \frac{BC^2}{(15.4)^2}$$

$$\Rightarrow \frac{8}{11} = \frac{BC}{15.4}$$

$$\Rightarrow BC = \left(\frac{8}{11} \times 15.4 \right) \text{ cm} = 11.2 \text{ cm}$$

2. Diagonals of a trapezium $ABCD$ with $AB \parallel DC$ intersect each other at the point O . If $AB = 2CD$, find the ratio of the areas of triangles AOB and COD .

Ans. In \triangle s AOB and COD , we have,



$$\angle AOB = \angle COD [\text{Vertically opposite angles}]$$

$$\angle OAB = \angle OCD [\text{Alternate angles}]$$

By AA-criterion of similarity,


$$\therefore \triangle AOB \sim \triangle COD$$

$$\therefore \frac{\text{Area}(\triangle AOB)}{\text{Area}(\triangle COD)} = \frac{AB^2}{DC^2}$$


$$\Rightarrow \frac{\text{Area}(\triangle AOB)}{\text{Area}(\triangle COD)} = \frac{(2DC)^2}{DC^2} = \frac{4}{1}$$

Hence, Area ($\triangle AOB$) : Area ($\triangle COD$) = 4 : 1

3. In the given figure, ABC and DBC are two triangles on the same base BC. If AD intersects BC at O, show that $\frac{\text{ar}(\triangle ABC)}{\text{ar}(\triangle DBC)} = \frac{AO}{DO}$.



Ans. Given: Two \triangle s ABC and DBC which stand on the same base but on the opposite sides of BC.



To Prove: $\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DBC)} = \frac{AO}{DO}$

Construction: Draw $AE \perp BC$ and $DF \perp BC$.

Proof: In \triangle s AOE and DOF, we have, $\angle AEO = \angle DFO = 90^\circ$

and $\angle AOE = \angle DOF$ [Vertically opposite]

$\therefore \triangle AOE \sim \triangle DOF$ [By AA-criterion]

$$\therefore \frac{AE}{DF} = \frac{AO}{OD} \dots\dots\dots(i)$$

$$\text{Now, } \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DBC)} = \frac{\frac{1}{2} \times BC \times AE}{\frac{1}{2} \times BC \times DF}$$


$$\Rightarrow \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DBC)} = \frac{AE}{DF}$$

$$\Rightarrow \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DBC)} = \frac{AO}{OD} \text{ [using eq. (i)]}$$

4. If the areas of two similar triangles are equal, prove that they are congruent.

Ans. Given: Two \triangle s ABC and DEF such that $\triangle ABC \sim \triangle DEF$

And $\text{Area}(\triangle ABC) = \text{Area}(\triangle DEF)$



To Prove: $\triangle ABC \cong \triangle DEF$

Proof: $\triangle ABC \sim \triangle DEF$

$$\therefore \angle A = \angle D, \angle B = \angle E, \angle C = \angle F$$

And $\frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF}$

To establish $\triangle ABC \cong \triangle DEF$, it is sufficient to prove that, $AB = DE$, $BC = EF$ and $AC = DF$

Now, $\text{Area}(\triangle ABC) = \text{Area}(\triangle DEF)$

$$\therefore \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle DEF)} = 1$$

$$\Rightarrow \frac{AB^2}{DE^2} = \frac{BC^2}{EF^2} = \frac{AC^2}{DF^2} = 1$$

$$\Rightarrow \frac{AB}{DE} = \frac{BC}{EF} = \frac{AC}{DF} = 1$$

$$\Rightarrow AB = DE, BC = EF, AC = DF$$

Hence, $\triangle ABC \cong \triangle DEF$


5. D, E and F are respectively the midpoints of sides AB, BC and CA of $\triangle ABC$. Find the ratio of the areas of $\triangle DEF$ and $\triangle ABC$.

Ans. Since D and E are the midpoints of the sides BC and CA of $\triangle ABC$ respectively.

$$\therefore DE \parallel BA \Rightarrow DE \parallel FA \dots\dots\dots(i)$$

Since D and F are the midpoints of the sides BC and AB of $\triangle ABC$ respectively.

$$\therefore DF \parallel CA \Rightarrow DF \parallel AE \dots\dots\dots(ii)$$



From (i) and (ii), we can say that AFDE is a parallelogram.

Similarly, BDEF is a parallelogram.

Now, in Δ s DEF and ABC, we have

$$\angle FDE = \angle A [\text{opposite angles of } \parallel \text{ gm AFDE}]$$

$$\text{And } \angle DEF = \angle B [\text{opposite angles of } \parallel \text{ gm BDEF}]$$

\therefore By AA-criterion of similarity, we have $\Delta DEF \sim \Delta ABC$


$$\Rightarrow \frac{\text{Area}(\Delta DEF)}{\text{Area}(\Delta ABC)} = \frac{DE^2}{AB^2} = \left(\frac{1}{2}AB\right)^2 \frac{1}{AB^2} = \frac{1}{4}$$

$$[\because DE = \frac{1}{2}AB]$$

Hence, Area (ΔDEF): Area (ΔABC) = 1 : 4

6. Prove that the ratio of the areas of two similar triangles is equal to the square of the ratio of their corresponding medians.

Ans. Given: $\Delta ABC \sim \Delta PQR$, AD and PM are the medians of Δ s ABC and PQR respectively.



$$\text{To Prove: } \frac{\text{Area}(\Delta ABC)}{\text{Area}(\Delta PQR)} = \frac{AD^2}{PM^2}$$

Proof: Since $\Delta ABC \sim \Delta PQR$

$$\therefore \frac{\text{Area}(\Delta ABC)}{\text{Area}(\Delta PQR)} = \frac{AB^2}{PQ^2} \dots\dots\dots(1)$$

But, $\frac{AB}{PQ} = \frac{AD}{PM}$ (2)


∴ From eq. (1) and (2), we have,

$$\frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle PQR)} = \frac{AD^2}{PM^2}$$

7. Prove that the area of an equilateral triangle described on one side of a square is equal to half the area of the equilateral triangle described on one of the diagonals.

Ans. Given: A square ABCD,

Equilateral \triangle s BCE and ACF have been drawn on side BC and the diagonal AC respectively.



To Prove: $\text{Area}(\triangle BCE) = \frac{1}{2} \text{Area}(\triangle ACF)$

Proof: $\triangle BCE \sim \triangle ACF$

[Being equilateral so similar by AAA criterion of similarity]

$$\Rightarrow \frac{\text{Area}(\triangle BCE)}{\text{Area}(\triangle ACF)} = \frac{BC^2}{AC^2}$$

$$\Rightarrow \frac{\text{Area}(\triangle BCE)}{\text{Area}(\triangle ACF)} = \frac{BC^2}{(\sqrt{2}BC)^2}$$

$$[\because \text{Diagonal} = \sqrt{2} \text{ side} \Rightarrow AC = \sqrt{2} BC]$$

$$\Rightarrow \frac{\text{Area}(\triangle BCE)}{\text{Area}(\triangle ACF)} = \frac{1}{2}$$

Tick the correct answer and justify:

8. ABC and BDE are two equilateral triangles such that D is the midpoint of BC. The ratio of the areas of triangles ABC and BDE is:

(A) 2: 1


(B) 1: 2

(C) 4: 1

(D) 1: 4

Ans. (C) Since $\triangle ABC$ and $\triangle BDE$ are equilateral, they are equiangular and hence,

$$\triangle ABC \sim \triangle BDE$$



$$\Rightarrow \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle BDE)} = \frac{BC^2}{BD^2}$$

$$\Rightarrow \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle BDE)} = \frac{(2BD)^2}{BD^2}$$

$[\because D \text{ is the midpoint of } BC]$

$$\Rightarrow \frac{\text{Area}(\triangle ABC)}{\text{Area}(\triangle BDE)} = \frac{4}{1}$$

∴ (C) is the correct answer.

9. Sides of two similar triangles are in the ratio 4: 9. Areas of these triangles are in the ratio:

(A) 2: 3

(B) 4: 9

(C) 81: 16

(D) 16: 81

Ans. (D) Since the ratio of the areas of two similar triangles is equal to the ratio of the squares of any two corresponding sides. Therefore,

$$\text{Ratio of areas} = \frac{(4)^2}{(9)^2} = \frac{16}{81}$$

∴ (D) is the correct answer.

CBSE Class-10 Mathematics

NCERT solution

Chapter - 6

Triangles - Exercise 6.5

1. Sides of triangles are given below. Determine which of them are right triangles. In case of a right triangle, write the length of its hypotenuse.

(i) 7 cm, 24 cm, 25 cm

(ii) 3 cm, 8 cm, 6 cm

(iii) 50 cm, 80 cm, 100 cm

(iv) 13 cm, 12 cm, 5 cm

Ans. (i) Let $a = 7$ cm, $b = 24$ cm and $c = 25$ cm

Here the larger side is $c = 25$ cm.

$$\text{We have, } a^2 + b^2 = 7^2 + 24^2 = 49 + 576 = 625 = c^2$$

So, the triangle with the given sides is a right triangle. Its hypotenuse = 25 cm

(ii) Let $a = 3$ cm, $b = 8$ cm and $c = 6$ cm

Here the larger side is $b = 8$ cm.

$$\text{We have, } a^2 + c^2 = 3^2 + 6^2 = 9 + 36 = 45 \neq b^2$$

So, the triangle with the given sides is not a right triangle.

(iii) Let $a = 50$ cm, $b = 80$ cm and $c = 100$ cm

Here the larger side is $c = 100$ cm.

$$\text{We have, } a^2 + b^2 = 50^2 + 80^2 = 2500 + 6400 = 8900 \neq c^2$$

So, the triangle with the given sides is not a right triangle.

(iv) Let $a = 13$ cm, $b = 12$ cm and $c = 5$ cm


Here the larger side is $a = 13$ cm.

We have, $b^2 + c^2 = 12^2 + 5^2 = 144 + 25 = 169 = a^2$

So, the triangle with the given sides is a right triangle. Its hypotenuse = 13 cm

2. PQR is a triangle right angled at P and M is a point on QR such that $PM \perp QR$. Show that $PM^2 = QM \times MR$.

Ans. Given: PQR is a triangle right angled at P and $PM \perp QR$



To Prove: $PM^2 = QM \cdot MR$

Proof: Since $PM \perp QR$

$\therefore \triangle QMP \sim \triangle PMR$

$$\Rightarrow \frac{QM}{PM} = \frac{PM}{RM}$$


$$\Rightarrow PM^2 = QM \cdot MR$$

3. In the given figure, ABD is a triangle right angled at A and $AC \perp BD$. Show that:

(i) $AB^2 = BC \cdot BD$

(ii) $AC^2 = BC \cdot DC$

(iii) $AD^2 = BD \cdot CD$




Ans. Given: ABD is a triangle right angled at A and $AC \perp BD$.

To Prove: (i) $AB^2 = BC \cdot BD$, (ii) $AC^2 = BC \cdot DC$, (iii) $AD^2 = BD \cdot CD$

Proof:(i) Since $AC \perp BD$

$\therefore \triangle CBA \sim \triangle CAD$ and each triangle is similar to $\triangle ABD$


$\therefore \triangle ABD \sim \triangle CBA$



$$\Rightarrow \frac{AB}{BD} = \frac{BC}{AB}$$

$$\Rightarrow AB^2 = BC \cdot BD$$

(ii) Since $\triangle ABC \sim \triangle DAC$



$$\Rightarrow \frac{AC}{BC} = \frac{DC}{AC}$$

$$\Rightarrow AC^2 = BC \cdot DC$$

(iii) Since $\triangle CAD \sim \triangle ABD$

$$\Rightarrow \frac{AD}{CD} = \frac{BD}{AD}$$

$$\Rightarrow AD^2 = BD \cdot CD$$

4. ABC is an isosceles triangle right angled at C. Prove that $AB^2 = 2AC^2$.

Ans. Since ABC is an isosceles right triangle, right angled at C.

$$\Rightarrow AB^2 = AC^2 + BC^2$$

$$\Rightarrow AB^2 = AC^2 + AC^2 \text{ [BC = AC, given]}$$

$$\Rightarrow AB^2 = 2AC^2$$

5. ABC is an isosceles triangle with AC = BC. If $AB^2 = 2AC^2$, prove that ABC is a right triangle.

Ans. Since ABC is an isosceles right triangle with AC = BC and $AB^2 = 2AC^2$


$$\Rightarrow AB^2 = AC^2 + AC^2$$

$$\Rightarrow AB^2 = AC^2 + BC^2 \text{ [BC = AC, given]}$$

$\therefore \triangle ABC$ is right angled at C.

6. ABC is an equilateral triangle of side $2a$. Find each of its altitudes.

Ans. Let ABC be an equilateral triangle of side $2a$ units.



Draw $AD \perp BC$. Then, D is the midpoint of BC.

$$\Rightarrow BD = \frac{1}{2} BC = \frac{1}{2} \times 2a = a$$

Since, ABD is a right triangle, right triangle at D.

$$\therefore AB^2 = AD^2 + BD^2$$

$$\Rightarrow (2a)^2 = AD^2 + (a)^2$$


$$\Rightarrow AD^2 = 4a^2 - a^2 = 3a^2$$

$$\therefore \text{Each of its altitude} = \sqrt{3}a$$

7. Prove that the sum of the squares of the sides of a rhombus is equal to the sum of squares of its diagonals.

Ans. Let the diagonals AC and BD of rhombus ABCD intersect each other at O. Since the diagonals of a rhombus bisect each other at right angles.

$$\therefore \angle AOB = \angle BOC = \angle COD = \angle DOA = 90^\circ \text{ and } AO = CO, BO = OD$$



Since AOB is a right triangle, right angled at O.

$$\therefore AB^2 = OA^2 + OB^2$$

$$\Rightarrow AB^2 = \left(\frac{1}{2}AC\right)^2 + \left(\frac{1}{2}BD\right)^2$$

[\because OA = OC and OB = OD]

$$\Rightarrow 4AB^2 = AC^2 + BD^2 \dots\dots\dots(1)$$

Similarly, we have $4BC^2 = AC^2 + BD^2 \dots\dots\dots(2)$

$$4CD^2 = AC^2 + BD^2 \dots\dots\dots(3)$$

$$4AD^2 = AC^2 + BD^2 \dots\dots\dots(4)$$


Adding all these results, we get

$$4(AB^2 + BC^2 + CD^2 + DA^2) = 4(AC^2 + BD^2)$$

$$\Rightarrow AB^2 + BC^2 + CD^2 + DA^2 = AC^2 + BD^2$$

8. In the given figure, O is a point in the interior of a triangle ABC, OD \perp BC,

OE \perp AC and OF \perp AB. Show that:




(i) $OA^2 + OB^2 + OC^2 - OD^2 - OE^2 - OF^2 = AF^2 + BD^2 + CE^2$

(ii) $AF^2 + BD^2 + CE^2 = AE^2 + CD^2 + BF^2$

Ans. Join AO, BO and CO.

(i) In right Δ s OFA, ODB and OEC, we have

$$OA^2 = AF^2 + OF^2, OB^2 = BD^2 + OD^2 \text{ and } OC^2 = CE^2 + OE^2$$



Adding all these, we get

$$OA^2 + OB^2 + OC^2 = AF^2 + BD^2 + CE^2 + OF^2 + OD^2 + OE^2$$

$$\Rightarrow OA^2 + OB^2 + OC^2 - OD^2 - OE^2 - OF^2 = AF^2 + BD^2 + CE^2$$

(ii) In right Δ s ODB and ODC, we have

$$OB^2 = BD^2 + OD^2 \text{ and } OC^2 = OD^2 + CD^2$$

$$\Rightarrow OB^2 - OC^2 = BD^2 - CD^2 \dots\dots\dots(1)$$

Similarly, we have $OB^2 - OC^2 = BD^2 - CD^2 \dots\dots\dots(2)$

and $OB^2 - OC^2 = BD^2 - CD^2 \dots\dots\dots(3)$

Adding equations (1), (2) and (3), we get

$$= (OB^2 - OC^2) + (OC^2 - OA^2) + (OA^2 - OB^2)$$

$$= (BD^2 - CD^2) + (CE^2 - AE^2) + (AF^2 - BF^2)$$


$$\Rightarrow (BD^2 + CE^2 + AF^2) - (AE^2 + CD^2 + BF^2) = 0$$

$$\Rightarrow AF^2 + BD^2 + CE^2 = AE^2 + BF^2 + CD^2$$

9. A ladder 10 m long reaches a window 8 m above the ground. Find the distance of the foot of the ladder from the base of the wall.

Ans. Let AB be the ladder, B be the window and CB be the wall. Then, ABC is a

right triangle, right angled at C.



$$\Rightarrow AB^2 = AC^2 + BC^2$$

$$\Rightarrow 10^2 = AC^2 + 8^2$$

$$\Rightarrow AC^2 = 100 - 64$$


$$\Rightarrow AC^2 = 36$$

$$\Rightarrow AC = 6$$

Hence, the foot of the ladder is at a distance 6 m from the base of the wall.

10. A guy wire attached to a vertical pole of height 18 m is 24 m long and has a stake attached to the other hand. How far from the base of the pole should the stake be driven so that the wire will be taut?

Ans. Let AB (= 24m) be a guy wire attached to a vertical pole. BC of height 18 m. To keep the wire taut, let it be fixed to a stake at A. Then, ABC is a right triangle, right angled at C.



$$\therefore AB^2 = AC^2 + BC^2$$

$$\Rightarrow 24^2 = AC^2 + 18^2$$

$$\Rightarrow AC^2 = 576 - 324$$

$$\Rightarrow AC^2 = 252$$


$$\Rightarrow AC = 6\sqrt{7}$$

Hence, the stake may be placed at a distance of $6\sqrt{7}$ m from the base of the pole.

11. An aeroplane leaves an airport and flies due north at a speed of 1000 km per hour. At the same time, another aeroplane leaves the same airport and flies due west at a speed of 1200 km per hour. How far apart will be the two planes after $1\frac{1}{2}$ hours?

Ans. Let the first aeroplane starts from O and goes upto A towards north where

$$OA = \left(1000 \times \frac{3}{2}\right) \text{ km} = 1500 \text{ km}$$



Let the second aeroplane starts from O at the same time and goes upto

B towards west where

$$OB = \left(1200 \times \frac{3}{2}\right) \text{ km} = 1800 \text{ km}$$

According to the question the required distance = BA

In right angled triangle ABC, by Pythagoras theorem, we have,

$$AB^2 = OA^2 + OB^2$$

$$= (1500)^2 + (1800)^2$$


$$= 2250000 + 3240000$$

$$= 5490000 = 9 \times 61 \times 100 \times 100$$

$$\Rightarrow AB = 300\sqrt{61} \text{ km}$$

12. Two poles of heights 6 m and 11 m stand on a plane ground. If the distance between the feet of the poles is 12 m, find the distance between their tops.

Ans. Let AB = 11 m and CD = 6 m be the two poles such that BD = 12 m



Draw $CE \perp AB$ and join AC.

$$\therefore CE = DB = 12 \text{ m}$$

$$AE = AB - BE = AB - CD = (11 - 6)\text{m} = 5 \text{ m}$$

In right angled triangle ACE, by Pythagoras theorem, we have

$$AC^2 = CE^2 + AE^2 = 12^2 + 5^2$$


$$= 144 + 25 = 169$$

$$\Rightarrow AC = 13 \text{ m}$$

Hence, the distance between the tops of the two poles is 13 m.

13. D and E are points on the sides CA and CB respectively of a triangle ABC right angled at C. Prove that $AE^2 + BD^2 = AB^2 + DE^2$.

Ans. In right angled Δ s ACE and DCB, we have



$$AE^2 = AC^2 + CE^2 \text{ and } BD^2 = DC^2 + BC^2$$


$$\Rightarrow AE^2 + BD^2 = (AC^2 + CE^2) + (DC^2 + BC^2)$$

$$\Rightarrow AE^2 + BD^2 = (AC^2 + BC^2) + (DC^2 + CE^2)$$

$$\Rightarrow AE^2 + BD^2 = AB^2 + DE^2$$

[By Pythagoras theorem, $AC^2 + BC^2 = AB^2$ and $DC^2 + CE^2 = DE^2$]

14. The perpendicular from A on side BC of a ΔABC intersects BC at D such that $DB = 3CD$ (see figure). Prove that $2AB^2 = 2AC^2 + BC^2$.



Ans. We have, $DB = 3CD$

Now, $BC = DB + CD$

$$\Rightarrow BC = 3CD + CD$$

$$\Rightarrow BC = 4CD$$

$$\therefore CD = \frac{1}{4} BC \text{ and } DB = 3CD = \frac{3}{4} BC \dots\dots\dots(1)$$

Since, ΔABD is a right triangle, right angled at D. Therefore by Pythagoras theorem, we have,

$$AB^2 = AD^2 + DB^2 \dots\dots\dots(2)$$

Similarly, from ΔACD , we have, $AC^2 = AD^2 + CD^2 \dots\dots\dots(3)$

From eq. (2) and (3) $AB^2 - AC^2 = DB^2 - CD^2$

$$\Rightarrow AB^2 - AC^2 = \left(\frac{3}{4}BC\right)^2 - \left(\frac{1}{4}BC\right)^2 \text{ [Using eq.(1)]}$$

$$\Rightarrow AB^2 - AC^2 = \left(\frac{9}{16} - \frac{1}{16}\right)BC^2$$


$$\Rightarrow AB^2 - AC^2 = \frac{1}{2}BC^2$$

$$\Rightarrow 2AB^2 - 2AC^2 = BC^2$$

$$\Rightarrow 2AB^2 = 2AC^2 + BC^2$$

15. In an equilateral triangle ABC, D is a point on side BC such that $BD = \frac{1}{3} BC$. Prove that $9AD^2 = 7AB^2$.

Ans. Let ABC be an equilateral triangle and let D be a point on BC such that $BD = \frac{1}{3} BC$



Draw $AE \perp BC$, Join AD .

In Δ s AEB and AEC , we have,

$$AB = AC [\because \Delta ABC \text{ is equilateral}]$$

$$\angle AEB = \angle AEC [\because \text{each } 90^\circ]$$

$$\text{And } AE = AE$$

\therefore By SAS-criterion of similarity, we have

$$\Delta AEB \sim \Delta AEC$$

$$\Rightarrow BE = EC$$

$$\text{Thus, we have, } BD = \frac{1}{3} BC, DC = \frac{2}{3} BC \text{ and } BE = EC = \frac{1}{3} BC \dots\dots\dots(1)$$

$$\text{Since, } \angle C = 60^\circ$$

$\therefore \Delta ADC$ is an acute angle triangle.

$$\therefore AD^2 = AC^2 + DC^2 - 2DC \times EC$$

$$= AC^2 + \left(\frac{2}{3}BC\right)^2 - 2 \times \frac{2}{3}BC \times \frac{1}{3}BC [\text{using eq.(1)}]$$

$$\Rightarrow AD^2 = AC^2 + \frac{4}{9}BC^2 - \frac{2}{3}BC^2$$


$$= AB^2 + \frac{4}{9}AB^2 - \frac{2}{3}AB^2 \quad [\because AB = BC = AC]$$

$$\Rightarrow AD^2 = \frac{(9+4-6)AB^2}{9} = \frac{7}{9}AB^2$$

$$\Rightarrow 9AD^2 = 7AB^2$$

16. In an equilateral triangle, prove that three times the square of one side is equal to four times the square of one of its altitudes.

Ans. Let ABC be an equilateral triangle and let $AD \perp BC$. In \triangle s ADB and ADC, we have,



$AB = AC$ [Given]

$\angle B = \angle C = 60^\circ$ [Given]

And $\angle ADB = \angle ADC$ [Each = 90°]

$\therefore \triangle ADB \cong \triangle ADC$ [By RHS criterion of congruence]

$\Rightarrow BD = DC$

$$\Rightarrow BD = DC = \frac{1}{2}BC$$

Since $\triangle ADB$ is a right triangle, right angled at D, by Pythagoras theorem, we have,

$$AB^2 = AD^2 + BD^2$$

$$\Rightarrow AB^2 = AD^2 + \left(\frac{1}{2}BC\right)^2$$

$$\Rightarrow AB^2 = AD^2 + \frac{1}{4}BC^2$$

$$\Rightarrow AB^2 = AD^2 + \frac{AB^2}{4} \quad [\because BC = AB]$$

$$\Rightarrow \frac{3}{4}AB^2 = AD^2$$

$$\Rightarrow 3AB^2 = 4AD^2$$

17. Tick the correct answer and justify: In $\triangle ABC$, $AB = 6\sqrt{3}$ cm, $AC = 12$ cm and $BC = 6$ cm. the angles A and B are respectively:

(A) 90° and 30°

(B) 90° and 60°

(C) 30° and 90°


(D) 60° and 90°

Ans. (C) In $\triangle ABC$, we have, $AB = 6\sqrt{3}$ cm, $AC = 12$ cm and $BC = 6$ cm.

$$\text{Now, } AB^2 + BC^2 = (6\sqrt{3})^2 + (6)^2 = 36 \times 3 + 36 = 108 + 36 = 144 = (AC)^2$$

Thus, $\triangle ABC$ is a right triangle, right angled at B.

$$\therefore \angle B = 90^\circ$$



Let D be the midpoint of AC. We know that the midpoint of the hypotenuse of a right triangle is equidistant from the vertices.

$$AD = BD = CD$$

$$\Rightarrow CD = BD = 6 \text{ cm } [\because CD = \frac{1}{2} AC]$$

$$\text{Also, } BC = 6 \text{ cm}$$

$$\therefore \text{ In } \triangle BDC, \text{ we have, } BD = CD = BC$$

$$\Rightarrow \triangle BDC \text{ is equilateral}$$

$$\Rightarrow \angle ACB = 60^\circ$$

$$\therefore \angle A = 180^\circ - (\angle B + \angle C) = 180^\circ - (90^\circ + 60^\circ) = 30^\circ$$

$$\text{Thus, } \angle A = 30^\circ \text{ and } \angle B = 90^\circ$$

CBSE Class-10 Mathematics


NCERT solution

Chapter - 6


Triangles - Exercise 6.6 (Optional)*

1. In the given figure, PS is the bisector of $\angle QPR$ of $\triangle PQR$. Prove that

$$\frac{QS}{SR} = \frac{PQ}{PR}.$$



Ans. Given: PQR is a triangle and PS is the internal bisector of $\angle QPR$ meeting QR at S.



$$\therefore \angle QPS = \angle SPR$$

To prove: $\frac{QS}{SR} = \frac{PQ}{PR}$

Construction: Draw $RT \parallel SP$ to cut QP produced at T .

Proof: Since $PS \parallel TR$ and PR cuts them, hence,

$$\angle SPR = \angle PRT \dots\dots\dots(i) \text{ [Alternate } \angle \text{ s]}$$

$$\text{And } \angle QPS = \angle PTR \dots\dots\dots(ii) \text{ [Corresponding } \angle \text{ s]}$$

$$\text{But } \angle QPS = \angle SPR \text{ [Given]}$$

$$\therefore \angle PRT = \angle PTR \text{ [From eq. (i) \& (ii)]}$$


$$\Rightarrow PT = PR \dots\dots\dots(iii)$$

[Sides opposite to equal angles are equal]

Now, in $\triangle QRT$,

$RT \parallel SP$ [By construction]


$$\therefore \frac{QS}{SR} = \frac{PQ}{PT} \text{ [Thales theorem]}$$



$$\Rightarrow \frac{QS}{SR} = \frac{PQ}{PR} \text{ [From eq. (iii)]}$$

2. In the given figure, D is a point on hypotenuse AC of $\triangle ABC$, $BD \perp AC$, $DM \perp BC$ and

DN \perp AB. Prove that:



(i) $DM^2 = DN \cdot MC$

(ii) $DN^2 = DM \cdot AN$

Ans. Since $AB \perp BC$ and $DM \perp BC$

$$\Rightarrow AB \parallel DM$$

Similarly, $BC \perp AB$ and $DN \perp AB$

$$\Rightarrow CB \parallel DN$$

\therefore quadrilateral BMDN is a rectangle.

$$\therefore BM = ND$$

(i) In $\triangle BMD$, $\angle 1 + \angle BMD + \angle 2 = 180^\circ$

$$\Rightarrow \angle 1 + 90^\circ + \angle 2 = 180^\circ$$

$$\Rightarrow \angle 1 + \angle 2 = 90^\circ$$

Similarly, in $\triangle DMC$, $\angle 3 + \angle 4 = 90^\circ$

Since $BD \perp AC$,

$$\therefore \angle 2 + \angle 3 = 90^\circ$$

Now, $\angle 1 + \angle 2 = 90^\circ$ and $\angle 2 + \angle 3 = 90^\circ$

$$\Rightarrow \angle 1 + \angle 2 = \angle 2 + \angle 3$$

$$\Rightarrow \angle 1 = \angle 3$$

Also, $\angle 3 + \angle 4 = 90^\circ$ and $\angle 2 + \angle 3 = 90^\circ$

$$\Rightarrow \angle 3 + \angle 4 = \angle 2 + \angle 3$$

$$\Rightarrow \angle 4 = \angle 2$$

Thus, in $\triangle BMD$ and $\triangle DMC$,

$$\angle 1 = \angle 3 \text{ and } \angle 4 = \angle 2$$

$$\therefore \triangle BMD \sim \triangle DMC$$

$$\Rightarrow \frac{BM}{DM} = \frac{MD}{MC}$$

$$\Rightarrow \frac{DN}{DM} = \frac{DM}{MC} \text{ [BM = ND]}$$

$$\Rightarrow DM^2 = DN.MC$$

(ii) Processing as in (i), we can prove that

$$\triangle BND \sim \triangle DNA$$


$$\Rightarrow \frac{BN}{DN} = \frac{ND}{NA}$$

$$\Rightarrow \frac{DM}{DN} = \frac{DN}{AN} \text{ [BN = DM]}$$

$$\Rightarrow DN^2 = DM.AN$$

3. In the given figure, ABC is a triangle in which $\angle ABC > 90^\circ$ and $AD \perp CB$ produced.

Prove that:



$$AC^2 = AB^2 + BC^2 + 2BC \cdot BD$$


Ans. Given: ABC is a triangle in which $\angle ABC > 90^\circ$ and $AD \perp CB$ produced.

To prove: $AC^2 = AB^2 + BC^2 + 2BC \cdot BD$

Proof: Since $\triangle ADB$ is a right triangle, right angled at D, therefore, by Pythagoras theorem,

$$AB^2 = AD^2 + DB^2 \dots\dots\dots(i)$$

Again, $\triangle ADC$ is a right triangle, right angled at D, therefore, by Pythagoras theorem,



$$AC^2 = AD^2 + DC^2$$

$$\Rightarrow AC^2 = AD^2 + (DB + BC)^2$$

$$\Rightarrow AC^2 = AD^2 + DB^2 + BC^2 + 2DB \cdot BC$$


$$\Rightarrow AC^2 = (AD^2 + DB^2) + BC^2 + 2DB \cdot BC$$

$$\Rightarrow AC^2 = AB^2 + BC^2 + 2DB \cdot BC$$

[Using eq. (i)]

4. In the given figure, ABC is a triangle in which $\angle ABC < 90^\circ$ and $AD \perp BC$ produced. Prove that:

$$AC^2 = AB^2 + BC^2 - 2BC \cdot BD$$



Ans. Given: ABC is a triangle in which $\angle ABC < 90^\circ$ and $AD \perp BC$ produced.

To prove: $AC^2 = AB^2 + BC^2 - 2BC \cdot BD$

Proof: Since $\triangle ADB$ is a right triangle, right angled at D, therefore, by Pythagoras theorem,

$$AB^2 = AD^2 + BD^2 \dots\dots(i)$$

Again, $\triangle ADC$ is a right triangle, right angled at D, therefore, by Pythagoras theorem,


$$AC^2 = AD^2 + DC^2$$

$$\Rightarrow AC^2 = AD^2 + (BC - BD)^2$$

$$\Rightarrow AC^2 = AD^2 + BC^2 + BD^2 - 2BC \cdot BD$$


$$\Rightarrow AC^2 = (AD^2 + DB^2) + BC^2 - 2DB \cdot BC$$

$$\Rightarrow AC^2 = AB^2 + BC^2 - 2DB \cdot BC$$



[Using eq. (i)]

5. In the given figure, AD is a median of a triangle ABC and $AM \perp BC$. Prove that:



$$(i) AC^2 = AD^2 + BC \cdot DM + \left(\frac{BC}{2}\right)^2$$

$$(ii) AB^2 = AD^2 - BC \cdot DM + \left(\frac{BC}{2}\right)^2$$

$$(iii) AC^2 + AB^2 = 2AD^2 + \frac{1}{2} BC$$

Ans. Since $\angle AMD = 90^\circ$, therefore $\angle ADM < 90^\circ$ and $\angle ADC > 90^\circ$


Thus, $\angle ADC$ is the acute angle and $\angle ADC$ is an obtuse angle.

(i) In $\triangle ADC$, $\angle ADC$ is an obtuse angle.

$$\therefore AC^2 = AD^2 + DC^2 + 2DC \cdot DM$$

$$\Rightarrow AC^2 = AD^2 + \left(\frac{BC}{2}\right)^2 + 2 \cdot \frac{BC}{2} \cdot DM$$

$$\Rightarrow AC^2 = AD^2 + \left(\frac{BC}{2}\right)^2 + BC \cdot DM$$



$$\Rightarrow AC^2 = AD^2 + BC \cdot DM + \left(\frac{BC}{2}\right)^2 \dots\dots\dots(i)$$

(ii) In $\triangle ABD$, $\angle ADM$ is an acute angle.

$$AB^2 = AD^2 + BD^2 - 2BD \cdot DM$$

$$\Rightarrow AB^2 = AD^2 + \left(\frac{BC}{2}\right)^2 - 2 \cdot \frac{BC}{2} \cdot DM$$

$$\Rightarrow AB^2 = AD^2 - BC \cdot DM + \left(\frac{BC}{2}\right)^2 \dots\dots\dots(ii)$$

(iii) From eq. (i) and eq. (ii),

$$AB^2 + AC^2 = 2AD^2 + \frac{1}{2}BC^2$$

6. Prove that the sum of the squares of the diagonals of a parallelogram is equal to the sum of the squares of its sides.

Ans. If AD is a median of $\triangle ABC$, then

$$AB^2 + AC^2 = 2AD^2 + \frac{1}{2}BC^2 \text{ [See Q.5 (iii)]}$$

Since the diagonals of a parallelogram bisect each other, therefore, BO and DO are medians of triangles ABC and ADC respectively.

$$\therefore AB^2 + BC^2 = 2BO^2 + \frac{1}{2} AC^2 \dots\dots\dots(i)$$

$$\text{And } AD^2 + CD^2 = 2DO^2 + \frac{1}{2} AC^2 \dots\dots\dots(ii)$$


Adding eq. (i) and (ii),

$$AB^2 + BC^2 + AD^2 + CD^2 = 2 (BO^2 + DO^2) + AC^2$$

$$\Rightarrow AB^2 + BC^2 + AD^2 + CD^2 = 2 \left(\frac{1}{4} BD^2 + \frac{1}{4} BD^2 \right) + AC^2 \left[DO = \frac{1}{2} BD \right]$$

$$\Rightarrow AB^2 + BC^2 + AD^2 + CD^2 = AC^2 + BD^2$$

7. In the given figure, two chords AB and CD intersect each other at the point P. Prove that:



(i) $\triangle APC \sim \triangle DPB$

(ii) $AP \cdot PB = CP \cdot DP$

Ans. (i) In the triangles APC and DPB,

$$\angle APC = \angle DPB \text{ [Vertically opposite angles]}$$

$$\angle CAP = \angle BDP \text{ [Angles in same segment of a circle are equal]}$$

∴ By AA-criterion of similarity,

$$\triangle APC \sim \triangle DPB$$


(ii) Since $\triangle APC \sim \triangle DPB$

$$\therefore \frac{AP}{DP} = \frac{CP}{PB} \Rightarrow AP \times PB = CP \times DP$$

8. In the give figure, two chords AB and CD of a circle intersect each other at the point P (when produced) outside the circle. Prove that:

(i) $\triangle PAC \sim \triangle PDB$

(ii) $PA.PB = PC.PD$



Ans. (i) In the triangles PAC and PDB,

$$\angle APC = \angle DPB \text{ [Common]}$$

$$\angle CAP = \angle BDP \text{ [}\because \angle BAC = 180^\circ - \angle PAC \text{ and } \angle PDB = \angle CDB]$$

$$= 180^\circ - \angle BAC = 180^\circ - (180^\circ - \angle PAC) = \angle PAC]$$

∴ By AA-criterion of similarity,


$$\triangle APC \sim \triangle DPB$$

(ii) Since $\triangle APC \sim \triangle DPB$


$$\therefore \frac{AP}{DP} = \frac{CP}{PB}$$

$$\Rightarrow PA.PB = PC.PD$$

9. In the given figure, D is a point on side BC of $\triangle ABC$ such that $\frac{BD}{CD} = \frac{AB}{AC}$. Prove that AD is the bisector of $\angle BAC$.



Ans. Given: ABC is a triangle and D is a point on BC such that $\frac{BD}{CD} = \frac{AB}{AC}$



To prove: AD is the internal bisector of $\angle BAC$.

Construction: Produce BA to E such that AE = AC. Join CE.

Proof: In $\triangle AEC$, since AE = AC

$$\therefore \angle AEC = \angle ACE \dots\dots\dots(i)$$

[Angles opposite to equal side of a triangle are equal]

$$\text{Now, } \frac{BD}{CD} = \frac{AB}{AC} \text{ [Given]}$$

$$\Rightarrow \frac{BD}{CD} = \frac{AB}{AE} [\because AE = AC, \text{ by construction}]$$

\therefore By converse of Basic Proportionality Theorem,

$$DA \parallel CE$$

Now, since CA is a transversal,

$$\therefore \angle BAD = \angle AEC \dots\dots\dots(ii) \text{ [Corresponding } \angle \text{ s]}$$


$$\text{And } \angle DAC = \angle ACE \dots\dots\dots(iii) \text{ [Alternate } \angle \text{ s]}$$

$$\text{Also } \angle AEC = \angle ACE \text{ [From eq. (i)]}$$


$$\text{Hence, } \angle BAD = \angle DAC \text{ [From eq. (ii) and (iii)]}$$

Thus, AD bisects \angle BAC internally.

10. Nazima is fly fishing in a stream. The tip of her fishing rod is 1.8 m above the surface of the water and the fly at the end of the string rests on the water 3.6 m away and 2.4 m from a point directly under the tip of the rod. Assuming that her string (from the tip of her rod to the fly) is taut, how much string does she have out (see Fig.)? If she pulls in the string at the rate of 5 cm per second, what will be the horizontal distance of the fly from her after 12 seconds?



Ans. I. To find The length of AC.



By Pythagoras theorem,


$$AC^2 = (2.4)^2 + (1.8)^2$$

$$\Rightarrow AC^2 = 5.76 + 3.24 = 9.00$$

$$\Rightarrow AC = 3 \text{ m}$$

\therefore Length of string she has out= 3 m

Length of the string pulled at the rate of 5 cm/sec in 12 seconds



$$= (5 \times 12) \text{ cm} = 60 \text{ cm} = 0.60 \text{ m}$$

$$\therefore \text{Remaining string left out} = 3 - 0.6 = 2.4 \text{ m}$$

II. To find: The length of PB

$$PB^2 = PC^2 - BC^2$$

$$= (2.4)^2 - (1.8)^2$$

$$= 5.76 - 3.24 = 2.52$$

$$\Rightarrow PB = \sqrt{2.52} = 1.59 \text{ (approx.)}$$

Hence, the horizontal distance of the fly from Nazima after 12 seconds

$$= 1.59 + 1.2 = 2.79 \text{ m (approx.)}$$