Physics 121: 1D Motion, Problem Solving

Cody Petrie

Mesa Community College

Questions

Any questions from the HW?

• If I know that a particle is at x(t = 0) = 5m, what is the position x at t = 5s?

- If I know that a particle is at x(t = 0) = 5m, what is the position x at t = 5s?
- If I know that a particle is at x(t = 0) = 5m and v(t = 0) = -2m/s, what is the position x at t = 5s?

- If I know that a particle is at x(t = 0) = 5m, what is the position x at t = 5s?
- If I know that a particle is at x(t = 0) = 5m and v(t = 0) = -2m/s, what is the position x at t = 5s?
- If the particle is changing velocity then this isn't enough... generally we need to know all three, \vec{r} , \vec{v} , and \vec{a} to fully describe the motion of an object.

 Those quantities are vectors and in three dimensions (3D) we would write them as

$$\vec{r} = x\hat{x} + y\hat{y} + z\hat{z} = (x, y, z)$$

 $\vec{v} = v_x\hat{x} + v_y\hat{y} + v_z\hat{z} = (v_x, v_y, v_z)$
 $\vec{a} = a_x\hat{x} + a_y\hat{y} + a_z\hat{z} = (a_x, a_y, a_z)$

 Those quantities are vectors and in three dimensions (3D) we would write them as


$$\vec{r} = x\hat{x} + y\hat{y} + z\hat{z} = (x, y, z)$$

 $\vec{v} = v_x\hat{x} + v_y\hat{y} + v_z\hat{z} = (v_x, v_y, v_z)$
 $\vec{a} = a_x\hat{x} + a_y\hat{y} + a_z\hat{z} = (a_x, a_y, a_z)$

• In 1D this can be written more simply as x, v_x , a_x , where each of these can be positive or negative. I might get lazy and just call these x, v, a when we know that the motion is 1D.

1D Motion - Signs of x, v_x, a_x

- Positive x means that the particle is on the positive side of the origin and negative x means that the particle is on the negative side of the origin... this is intuitive.
- Positive v means that the particle is moving in the positive direction as defined by the origin and visa-versa for negative v.
- What do you think positive and negative a mean? This is a bit more tricky.



 The sign of acceleration tells us which way the acceleration vector points, NOT whether the object is speeding up or slowing down.

Velocity	Acceleration	dwn/up
+	+	
-	+	
+	-	
-	-	

Velocity	Acceleration	dwn/up
+	+	up
-	+	
+	-	
-	-	

Velocity	Acceleration	dwn/up
+	+	up
-	+	dwn
+	-	
-	-	

Velocity	Acceleration	dwn/up
+	+	up
-	+	dwn
+	-	dwn
-	-	


Velocity	Acceleration	dwn/up
+	+	up
-	+	dwn
+	-	dwn
-	-	ир

Slowing down (dwn) or speeding up (up)? Do you see a pattern?


Velocity	Acceleration	dwn/up
+	+	up
-	+	dwn
+	-	dwn
-	-	up

Pattern: Same sign means speeding up, oppositite means slowing down.

A cyclist riding at 20 mph sees a stop sign and actually comes to a complete stop in 4 s. He then, in 6 s, returns to a speed of 15 mph. Which is his motion diagram? (Just think about what we just talked about)



A cyclist riding at 20 mph sees a stop sign and actually comes to a complete stop in 4 s. He then, in 6 s, returns to a speed of 15 mph. Which is his motion diagram? (Just think about what we just talked about)



An X-wing (not the USS Enterprise) is traveling to the left and is slowing down in preparation for his attack run on the Death Star.

The coordinate system is

$$\rightarrow +x$$

The velocity and acceleration are

- A. Velocity is positive, acceleration is positive.
- B. Velocity is negative, acceleration is positive.
- C. Velocity is positive, acceleration is negative.
- D. Velocity is negative, acceleration is negative.

An X-wing (not the USS Enterprise) is traveling to the left and is slowing down in preparation for his attack run on the Death Star.


The coordinate system is

$$\rightarrow +x$$

The velocity and acceleration are

- A Velocity is positive, acceleration is positive.
- ✓B. Velocity is negative, acceleration is positive.
 - C. Velocity is positive, acceleration is negative.
 - D. Velocity is negative, acceleration is negative.


• Another useful tool (in addition to a motion diagram) is an \times vs. t graph, like the ones you are generating in lab. Let's see if you are able to convert this motion diagram into an \times vs. t graph.



 $t = 0 \min$


• Another useful tool (in addition to a motion diagram) is an \times vs. t graph, like the ones you are generating in lab. Let's see if you are able to convert this motion diagram into an \times vs. t graph.

1 frame per minute




This is a graph of an object moving along a straight line. The most likely interpretation is:

- A. A person walking down a steep mountain.
- B. A car that drives and stops and drives and stops.
- C. An elevator descending.
- A rock that falls, bounces, and falls some more.
- A ball that is hit, caught, and thrown to someone else.



This is a graph of an object moving along a straight line. The most likely interpretation is:

- A. A person walking down a steep mountain.
- B. A car that drives and stopsand drives and stops.
- C. An elevator descending.
 - D. A rock that falls, bounces, and falls some more.
 - A ball that is hit, caught, and thrown to someone else.




- Uniform Motion is motion in which each each successive equal-time interval has the same displacement. This is a subset of 1D motion. You can only have motion in a straight line, but also you don't change velocity (speed or direction).
- For example, if you drive your car at a perfectly steady 60 mph, this means you change your position by 60 miles for every time interval of 1 hour.

$$v_{ave} = \frac{\Delta x}{\Delta t}$$

- What would uniform motion look like on a graph?
- How would you find the velocity of an object with uniform motion from a graph?

- What would uniform motion look like on a graph?
- How would you find the velocity of an object with uniform motion from a graph?




Cody Petrie

Chapters 1.6-7,2.1

 It turns out that the slope of the position vs. time graph is the velocity (We'll talk about this in terms of derivatives next time but since this is a constant slope it doesn't matter right now).


 It turns out that the slope of the position vs. time graph is the velocity (We'll talk about this in terms of derivatives next time but since this is a constant slope it doesn't matter right now).



If the position of a particle is given by s and the velocity is the slope then we can write

$$v_s = rac{\mathrm{rise}}{\mathrm{run}} = rac{\Delta s}{\Delta t} = rac{s_f - s_f}{t_f - t_f}$$

 It turns out that the slope of the position vs. time graph is the velocity (We'll talk about this in terms of derivatives next time but since this is a constant slope it doesn't matter right now).



If the position of a particle is given by s and the velocity is the slope then we can write

$$v_s = \frac{\mathrm{rise}}{\mathrm{run}} = \frac{\Delta s}{\Delta t} = \frac{s_f - s_i}{t_f - t_i}$$

$$s_f = s_i + v_s \Delta t$$
 (UniformMotionEquation) (1)

Problem Solving Strategy

- **1 Draw a motion diagram:** This developes intuition/visualize of the motion.
- Establish a coordinate system: Choose an axis and origin that seem "good" for the motion.
- Sketch the situation: Show the object at the beginning and end of the motion, and at any point where the motion changes in the middle.
- Oefine symbols: Place the symbols at relevant places on the sketch.
- **List known information:** Make a table of quantities that you know, or that are easily found out.
- **10 Identify the desired unknowns:** Make a table of quantities that you need to find to solve the problem.

Uniform Motion Example using the problem solving strategy

An hocker player is on a break away headed toward the other teams goal. He travels the whole time at 10 m/s. He is currently in the middle of the two goals which are about 52 m apart. How long does the goalie have to prepare for the shot? (Assume he shoots right as he gets to the goal)



- Draw a motion diagram
- Establish a coordinate system
- Sketch the situation
- Oefine symbols
- List known information
- Identify the desired unknowns

Uniform Motion Example using the problem solving strategy

An hocker player is on a break away headed toward the other teams goal. He travels the whole time at 10 m/s. He is currently in the middle of the two goals which are about 52 m apart. How long does the goalie have to prepare for the shot? (Assume he shoots right as he gets to the goal)



- Draw a motion diagram
- Establish a coordinate system
- Sketch the situation
- Oefine symbols
- List known information
- Identify the desired unknowns

t = 2.6 s

Uniform Motion Example using the problem solving strategy

Two billiard balls, labeled 1 and 2, are initially at rest on a smooth pool table and are separated by a distance of 50.0 m. Simultaneously, each ball is given a quick hit, and they begin to roll directly toward each other. Ball 1 moves with a speed of 5.23 m/s , and ball 2 moves with a speed of 2.67 m/s. What is the distance covered by ball 1 by the time the two balls collide?



Picture References

Hockey Player, accessed 28 Aug 2017: http://www.kevinneeld.com/breakaway-hockey-speed-qa/Billiard Balls, accessed 28 Aug 2017: http://www.healingtalks.com/wp-content/uploads/2011/12/billiard-balls.jpg