


Physics 121: Instantaneous Velocity, Kinematic Equations

Cody Petrie

Mesa Community College

Quiz

- 1 True/False: If an object has a negative acceleration it must be slowing down.
- 2 What is the velocity of the object whose $x(t)$ graph is shown below? Don't just write it, explain **briefly** how you got it.



How is HW going?

Instantaneous Velocity

- Instantaneous and average velocities are slightly different. Which is displayed on the speedometer of your car?

Instantaneous Velocity


- Instantaneous and average velocities are slightly different. Which is displayed on the speedometer of your car?
 - Instantaneous velocity

Instantaneous Velocity

- Instantaneous and average velocities are slightly different. Which is displayed on the speedometer of your car?
 - Instantaneous velocity
- Let's try to illustrate the difference and when it's appropriate to use average velocities.


This is an accelerating rocket (not uniform motion anymore)

(a) 30 frames per second




© 2013 Pearson Education, Inc.

(b) 3000 frames per second




The highly magnified section of the graph near point 2 is very nearly a straight line. The slope of this line is a good approximation to the instantaneous velocity at time t_2 . The slope is the instantaneous velocity in the limit $\Delta t \rightarrow 0$.

(c) The limiting case



Quick Check

Here is a motion diagram of a car moving along a straight road:




Which position-versus-time graph matches this motion diagram?




A.




B.



C.




D.



E.

Quick Check

Here is a motion diagram of a car moving along a straight road:




Which position-versus-time graph matches this motion diagram?




A.



B.



C.



D.




E.




Quick Check


Here is a motion diagram of a car moving along a straight road:




Which velocity-versus-time graph matches this motion diagram?




A.



B.



C.




D.


E. None of the above.

Quick Check


Here is a motion diagram of a car moving along a straight road:




Which velocity-versus-time graph matches this motion diagram?




A.



B.



C.



D.

E. None of the above.




Quick Check


Here is a motion diagram of a car moving along a straight road:




Which velocity-versus-time graph matches this motion diagram?




A.




B.



C.



D.




E.

Quick Check


Here is a motion diagram of a car moving along a straight road:




Which velocity-versus-time graph matches this motion diagram?




A.




B.



C.



D.



E.

Instantaneous Velocity

- As you zoom in more and more on the graph it's like taking smaller and smaller Δt steps. The truly instantaneous velocity is when you take $\Delta t \rightarrow 0$.

$$v_s \equiv \lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = \frac{ds}{dt} \quad (\text{Instantaneous Velocity})$$

v_s = slope of the position-versus-time graph at time t

Instantaneous Velocity

- As you zoom in more and more on the graph it's like taking smaller and smaller Δt steps. The truly instantaneous velocity is when you take $\Delta t \rightarrow 0$.

$$v_s \equiv \lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = \frac{ds}{dt} \quad (\text{Instantaneous Velocity})$$

v_s = slope of the position-versus-time graph at time t

- If v is the slope of $x(t)$ then how do we get x from $v(t)$?

Instantaneous Velocity

- As you zoom in more and more on the graph it's like taking smaller and smaller Δt steps. The truly instantaneous velocity is when you take $\Delta t \rightarrow 0$.

$$v_s \equiv \lim_{\Delta t \rightarrow 0} \frac{\Delta s}{\Delta t} = \frac{ds}{dt} \quad (\text{Instantaneous Velocity})$$

v_s = slope of the position-versus-time graph at time t

- If v is the slope of $x(t)$ then how to we get x from $v(t)$?
 - Some of you might recognize that as an integral.

$$x(t) = \int v(t) dt + x_0 \quad (1)$$

Derivatives Review

- $\frac{ds}{dt}$ is called *derivative of s with respect to t* .
- The derivative is the slope of the line that is tangent to the to the position vs. time graph.
- The book (and probably I) will only deal with derivatives of powers and polynomials so let's review those.

If $u(t) = ct^n$ then $\frac{du}{dt} =$

$$\text{If } u(t) = ct^n \text{ then } \frac{du}{dt} = nct^{n-1}$$

$$\text{If } u(t) = ct^n \text{ then } \frac{du}{dt} = nct^{n-1}$$

Let's try this with a specific example then. Let the trajectory of the particle be $s(t) = 2t^2$. Now solve for the velocity of the particle at any time along the path.

$$\frac{ds}{dt} =$$

$$\text{If } u(t) = ct^n \text{ then } \frac{du}{dt} = nct^{n-1}$$

Let's try this with a specific example then. Let the trajectory of the particle be $s(t) = 2t^2$. Now solve for the velocity of the particle at any time along the path.

$$\frac{ds}{dt} = 4t$$

Here are a couple more that might be useful to remember.

$$\frac{d}{dt}C =$$

$$\frac{d}{dt}(u(t) + w(t)) =$$

Here are a couple more that might be useful to remember.


$$\frac{d}{dt}C = 0$$

$$\frac{d}{dt}(u(t) + w(t)) = \frac{du}{dt} + \frac{dw}{dt}$$

Quick Check

Here is a position graph of an object:

At $t = 1.5$ s, the object's velocity is




- A. 40 m/s.
- B. 20 m/s.
- C. 10 m/s.
- D. -10 m/s.
- E. None of the above.

Quick Check

Here is a position graph of an object:

At $t = 1.5$ s, the object's velocity is




- A. 40 m/s.
- ✓ B. 20 m/s.
- C. 10 m/s.
- D. -10 m/s.
- E. None of the above.

Quick Check

Here is a position graph of an object:

At $t = 3.0$ s, the object's velocity is




- A. 40 m/s.
- B. 20 m/s.
- C. 10 m/s.
- D. -10 m/s.
- E. None of the above.

Quick Check

Here is a position graph of an object:

At $t = 3.0$ s, the object's velocity is




- A. 40 m/s.
- B. 20 m/s.
- C. 10 m/s.
- ✓ D. -10 m/s.
- E. None of the above.

Quick Check

When do objects 1 and 2 have the same velocity?


- A. At some instant before time t_0 .
- B. At time t_0 .
- C. At some instant after time t_0 .
- D. Both A and B.
- E. Never.



Quick Check

When do objects 1 and 2 have the same velocity?

- ✓ A. At some instant before time t_0 .
- B. At time t_0 .
- C. At some instant after time t_0 .
- D. Both A and B.
- E. Never.




Now Integrals

Position from Velocity


- Can we use our knowledge of the velocity to predict the position at a future time like we did with the uniform motion equation, $s_f = s_i + v_s \Delta t$?

Position from Velocity

- Can we use our knowledge of the velocity to predict the position at a future time like we did with the uniform motion equation, $s_f = s_i + v_s \Delta t$?
- Let's start by breaking up a $v(t)$ curve into N steps.




- Notice that the motion at each step is uniform motion so we can use ol' faithful, $s_f = s_i + v_s \Delta t \rightarrow \Delta s = v_s \Delta t$.




© 2015 Pearson Education, Inc.

Position from Velocity

- Can we use our knowledge of the velocity to predict the position at a future time like we did with the uniform motion equation, $s_f = s_i + v_s \Delta t$?
- Let's start by breaking up a $v(t)$ curve into N steps.





- Notice that the motion at each step is uniform motion so we can use ol' faithful, $s_f = s_i + v_s \Delta t \rightarrow \Delta s = v_s \Delta t$.
- So $v_s(t_k) \Delta t$ is the amount that is moved during between the steps k and $k + 1$.



© 2015 Pearson Education, Inc.

Position from Velocity

- Can we use our knowledge of the velocity to predict the position at a future time like we did with the uniform motion equation, $s_f = s_i + v_s \Delta t$?
- Let's start by breaking up a $v(t)$ curve into N steps.




- Notice that the motion at each step is uniform motion so we can use ol' faithful,
 $s_f = s_i + v_s \Delta t \rightarrow \Delta s = v_s \Delta t$.
- So $v_s(t_k) \Delta t$ is the amount that is moved during between the steps k and $k + 1$.
- So if I want $\Delta s = s_f - s_i$ then I want to sum over all of steps,

$$\Delta s \approx \Delta s_1 + \Delta s_2 + \dots + \Delta s_N = \sum_{k=1}^N (v_s)_k \Delta t.$$


© 2015 Pearson Education, Inc.

Position from Velocity

- We can then take this and from it determine the final position of a particle if we know it's initial position, s_i and it's velocity at each time, $v(t)$.



$$s_f \approx s_i + \sum_{k=1}^N (v_s)_k \Delta t$$





© 2013 Pearson Education, Inc.

Position from Velocity

- We can then take this and from it determine the final position of a particle if we know it's initial position, s_i and it's velocity at each time, $v(t)$.

$$s_f \approx s_i + \sum_{k=1}^N (v_s)_k \Delta t$$


- How do we get rid of the pesky approximate sign (\approx) though because we can't the true answer?



© 2013 Pearson Education, Inc.


Position from Velocity

- We can then take this and from it determine the final position of a particle if we know it's initial position, s_i and it's velocity at each time, $v(t)$.



$$s_f \approx s_i + \sum_{k=1}^N (v_s)_k \Delta t$$

- How do we get rid of the pesky approximate sign (\approx) though because we can't the true answer?




$$s_f = s_i + \lim_{\Delta t \rightarrow 0} \sum_{k=1}^N (v_s)_k \Delta t = s_i + \int_{t_i}^{t_f} v_s dt \quad (2)$$

Position from Velocity

- Just like we had a geometrical definition for a derivative let's find one for integrals.

Position from Velocity

- Just like we had a geometrical definition for a derivative let's find one for integrals.




During the interval t_i to t_f , the total displacement Δs is the "area under the curve."

© 2013 Pearson Education, Inc.

Position from Velocity

- Just like we had a geometrical definition for a derivative let's find one for integrals.




$$s_f = s_i + \text{area under the velocity curve } v_s \text{ between } t_i \text{ and } t_f \quad (3)$$

Quick Check

Here is the velocity graph of an object that is at the origin ($x = 0$ m) at $t = 0$ s.

At $t = 4.0$ s, the object's position is




- A. 20 m.
- B. 16 m.
- C. 12 m.
- D. 7 m.
- E. 4 m.

Quick Check

Here is the velocity graph of an object that is at the origin ($x = 0$ m) at $t = 0$ s.

At $t = 4.0$ s, the object's position is




- A. 20 m.
- B. 16 m.
- ✓ C. 12 m.
- D. 7 m.
- E. 4 m.

Quick Check

Here is the velocity graph of an object that is at the origin ($x = 0$ m) at $t = 0$ s.

What is Δs between $t=1$ s and $t=3$ s




- A. 20 m.
- B. 16 m.
- C. 12 m.
- D. 7 m.
- E. 4 m.

Quick Check

Here is the velocity graph of an object that is at the origin ($x = 0$ m) at $t = 0$ s.

What is Δs between $t=1$ s and $t=3$ s



- A. 20 m.
- B. 16 m.
- C. 12 m.
- ✓ D. 7 m.
- E. 4 m.

Integrals Review

- Again let's look at integrating polynomials. Think of undoing a derivative (multiply by power and subtract 1 from power).

$$\int_{t_i}^{t_f} ct^n dt =$$

- Again let's look at integrating polynomials. Think of undoing a derivative (multiply by power and subtract 1 from power).

$$\int_{t_i}^{t_f} ct^n dt = \left. \frac{ct^{n+1}}{n+1} \right|_{t_i}^{t_f} = \frac{ct_f^{n+1}}{n+1} - \frac{ct_i^{n+1}}{n+1} \quad (n \neq -1)$$

Integrals Review

- Again let's look at integrating polynomials. Think of undoing a derivative (multiply by power and subtract 1 from power).


$$\int_{t_i}^{t_f} ct^n dt = \left. \frac{ct^{n+1}}{n+1} \right|_{t_i}^{t_f} = \frac{ct_f^{n+1}}{n+1} - \frac{ct_i^{n+1}}{n+1} \quad (n \neq -1)$$

- Here is another identity for integrals

$$\int_{t_i}^{t_f} (u(t) + w(t)) dt = \int_{t_i}^{t_f} u(t) dt + \int_{t_i}^{t_f} w(t) dt$$

Quick Check


From the $v(t)$ graph find the position of this particle at time t .



© 2013 Pearson Education, Inc.

Quick Check

From the $v(t)$ graph find the position of this particle at time t .



- The first step might be to figure out $v(t)$, $v(t)=10-5t$ (where t is in seconds and v is in m/s).

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30m$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30\text{m}$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

$$x = x_i + \int_0^t v(t) dt = 30 \text{ m} + \int_0^t (10 - 5t) dt$$

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30\text{m}$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

$$\begin{aligned}x &= x_i + \int_0^t v(t) dt = 30 \text{ m} + \int_0^t (10 - 5t) dt \\&= 30 \text{ m} + \int_0^t 10 \, dt - \int_0^t 5t \, dt\end{aligned}$$

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30\text{m}$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

$$\begin{aligned}x &= x_i + \int_0^t v(t) dt = 30 \text{ m} + \int_0^t (10 - 5t) dt \\&= 30 \text{ m} + \int_0^t 10 dt - \int_0^t 5t dt \\&= 30 \text{ m} + 10t \Big|_0^t - \frac{5}{2}t^2 \Big|_0^t\end{aligned}$$

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30\text{m}$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

$$\begin{aligned}x &= x_i + \int_0^t v(t) dt = 30 \text{ m} + \int_0^t (10 - 5t) dt \\&= 30 \text{ m} + \int_0^t 10 dt - \int_0^t 5t dt \\&= 30 \text{ m} + 10t \Big|_0^t - \frac{5}{2}t^2 \Big|_0^t \\&= 30 \text{ m} + (10t - 10 * 0) - \left(\frac{5}{2}t^2 - \frac{5}{2}0^2 \right)\end{aligned}$$

Position from Velocity

$$v(t) = 10 - 5t \text{ and } x(t = 0) = 30\text{m}$$

- Then you might plug $v(t)$ and your initial conditions into the equation $s_f = s_i + \int_{t_i}^{t_f} v_s dt$.

$$\begin{aligned}x &= x_i + \int_0^t v(t) dt = 30 \text{ m} + \int_0^t (10 - 5t) dt \\&= 30 \text{ m} + \int_0^t 10 dt - \int_0^t 5t dt \\&= 30 \text{ m} + 10t \Big|_0^t - \frac{5}{2}t^2 \Big|_0^t \\&= 30 \text{ m} + (10t - 10 * 0) - \left(\frac{5}{2}t^2 - \frac{5}{2}0^2 \right) \\x &= 30 \text{ m} + 10t - \frac{5}{2}t^2\end{aligned}$$

Motion with Constant Acceleration

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?
 - $v = 7\text{m/s}$

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?
 - $v = 7\text{m/s}$
- What about after 2 seconds?

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?
 - $v = 7\text{m/s}$
- What about after 2 seconds?
 - $v = 9\text{m/s}$

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?
 - $v = 7\text{m/s}$
- What about after 2 seconds?
 - $v = 9\text{m/s}$
- What $a = -2\text{m/s}^2$ for 2 seconds?

Motion with Constant Acceleration

- **Acceleration:** Acceleration is the rate at which velocity changes.
- The SI units for acceleration are $(\text{m/s})/\text{s}$ or m/s^2 .
- If my velocity is 5m/s and I accelerate at $a = 2\text{m/s}^2$ for 1 second what is my new velocity?
 - $v = 7\text{m/s}$
- What about after 2 seconds?
 - $v = 9\text{m/s}$
- What $a = -2\text{m/s}^2$ for 2 seconds?
 - $v = 1\text{m/s}$

Motion with Constant Acceleration

- Just like the average velocity the **average acceleration** is the slope of $v(t)$.

$$a_{ave} = \frac{\Delta v}{\Delta t}$$

Motion with Constant Acceleration

- Just like the average velocity the **average acceleration** is the slope of $v(t)$.






$$a_{ave} = \frac{\Delta v}{\Delta t}$$

- If the acceleration is constant then a_{ave} and the instantaneous acceleration, a_s are the same.

$$a_s = \frac{dv}{dt}$$






Quick Check

A cart slows down while moving away from the origin. What do the position and velocity graphs look like?








Quick Check

A cart slows down while moving away from the origin. What do the position and velocity graphs look like?








Quick Check

A cart speeds up toward the origin.
What do the position and velocity graphs look like?








Quick Check

A cart speeds up toward the origin.
What do the position and velocity
graphs look like?








Quick Check

A cart *speeds up* while moving away from the origin. What do the velocity and acceleration graphs look like?








Quick Check

A cart *speeds up* while moving away from the origin. What do the velocity and acceleration graphs look like?








Quick Check

A cart *slows down* while moving away from the origin. What do the velocity and acceleration graphs look like?








Quick Check

A cart *slows down* while moving away from the origin. What do the velocity and acceleration graphs look like?








Quick Check

A cart *speeds up* while moving toward the origin. What do the velocity and acceleration graphs look like?



Quick Check

A cart *speeds up* while moving toward the origin. What do the velocity and acceleration graphs look like?



Kinematic Equations

- Remember with the initial position and the velocity we could calculate the position with $s_f = s_i + \int_{t_i}^{t_f} v_s dt$
- Since we know that the acceleration is the slope of $v(t)$, $a_{ave} = \frac{\Delta v}{\Delta t} \rightarrow v(t) = v_i + a\Delta t$ we can plug this into the integral.

Kinematic Equations

- Remember with the initial position and the velocity we could calculate the position with $s_f = s_i + \int_{t_i}^{t_f} v_s dt$
- Since we know that the acceleration is the slope of $v(t)$, $a_{ave} = \frac{\Delta v}{\Delta t} \rightarrow v(t) = v_i + a\Delta t$ we can plug this into the integral.
- Practice your integration by plugging $v(t)$ into the integral above to find an expression for the position $s(t)$ in terms of the initial velocity and the acceleration.

$$s_f = s_i + v_i \Delta t + \frac{1}{2} a (\Delta t)^2$$

Kinematic Equations

- If a particle is initially at s_i with a velocity of v_i you can figure out what the final velocity is by substituting in $\Delta t = \Delta v/a$ into the previous equation. This gives us the third kinematic equation.

$$v_f = v_i + a\Delta t \quad (4)$$

$$s_f = s_i + v_i\Delta t + \frac{1}{2}a(\Delta t)^2 \quad (5)$$

$$v_f^2 = v_i^2 + 2a\Delta s \quad (6)$$

Kinematic Equations - Practice


A rocket sled accelerates at 50 m/s^2 for 5.0 s , coasts for 3.0 s , then deploys a braking parachute and accelerated at -3.0 m/s^2 (you would call this deceleration) until it comes to a stop.

- a. What is the maximum velocity of the rocket sled?
- b. What is the total distance traveled?

Kinematic Equations - Practice

A rocket sled accelerates at 50 m/s^2 for 5.0 s , coasts for 3.0 s , then deploys a braking parachute and accelerated at -3.0 m/s^2 (you would call this deceleration) until it comes to a stop.

- What is the maximum velocity of the rocket sled?
- What is the total distance traveled?



© 2013 Pearson Education, Inc.

Known

$$x_0 = 0 \text{ m} \quad v_{0x} = 0 \text{ m/s} \quad t_0 = 0 \text{ s}$$

$$a_{0x} = 50 \text{ m/s}^2 \quad t_1 = 5.0 \text{ s}$$

$$a_{1x} = 0 \text{ m/s}^2 \quad t_2 = 8.0 \text{ s}$$

$$a_{2x} = -3.0 \text{ m/s}^2 \quad v_{3x} = 0 \text{ m/s}$$


Find

$$x_3 \text{ and } v_{1x}$$

Kinematic Equations - Practice

A rocket sled accelerates at 50 m/s^2 for 5.0 s , coasts for 3.0 s , then deploys a braking parachute and accelerated at -3.0 m/s^2 (you would call this deceleration) until it comes to a stop.

- What is the maximum velocity of the rocket sled?
- What is the total distance traveled?



Known

$$x_0 = 0 \text{ m} \quad v_{0x} = 0 \text{ m/s} \quad t_0 = 0 \text{ s}$$

$$a_{0x} = 50 \text{ m/s}^2 \quad t_1 = 5.0 \text{ s}$$

$$a_{1x} = 0 \text{ m/s}^2 \quad t_2 = 8.0 \text{ s}$$

$$a_{2x} = -3.0 \text{ m/s}^2 \quad v_{3x} = 0 \text{ m/s}$$

Find

$$x_3 \text{ and } v_{1x}$$

$$v_{1x} = 250 \text{ m/s}$$

$$x_3 = 12000 \text{ m}$$

Picture References

Nothing this time