

Physics 121: Newton's 1/2 Laws, Free Body Diagrams

Cody Petrie

Mesa Community College

- Next HW is due next Thursday (28th of September). Covers chapter 5 on Forces.

What Do Forced Do?

- We have established that a force causes an object to change its motion (acceleration). But by how much and how is that manifest?


What Do Forced Do?

- We have established that a force causes an object to change it's motion (acceleration). But by how much and how is that manifest?




What Do Forces Do?

Attach a rubber band to a block and stretch the rubber band (to a fixed amount) to pull the block across a frictionless surface. We know that the block will change its motion (accelerate).



What Do Forces Do?


Attach a rubber band to a block and stretch the rubber band (to a fixed amount) to pull the block across a frictionless surface. We know that the block will change its motion (accelerate).



- What happens to the acceleration if I add a second rubber band? Does it go up/down? By how much? The forces add so you get $F' = 2F$.

What Do Forces Do?

Attach a rubber band to a block and stretch the rubber band (to a fixed amount) to pull the block across a frictionless surface. We know that the block will change its motion (accelerate).




- What happens to the acceleration if I add a second rubber band? Does it go up/down? By how much? The forces add so you get $F' = 2F$.
- a also doubles. It turns out that the acceleration is proportional to the force applied.

$$a \propto F \quad \text{or} \quad a = cF$$

What does **proportional** mean?

What Do Forces Do?

Attach a rubber band to a block and stretch the rubber band (to a fixed amount) to pull the block across a frictionless surface. We know that the block will change its motion (accelerate).



- What happens to the acceleration if I add a second rubber band? Does it go up/down? By how much? The forces add so you get $F' = 2F$.
- a also doubles. It turns out that the acceleration is proportional to the force applied.

$$a \propto F \quad \text{or} \quad a = cF$$

What does **proportional** mean?
but what is c ?


What Do Forces Do?

- What if I were to double the mass of the block (say from 1kg to 2kg), what would happen to the acceleration? Think of pushing a broken down car vs a broken down semi ...

What Do Forces Do?

- What if I were to double the mass of the block (say from 1kg to 2kg), what would happen to the acceleration? Think of pushing a broken down car vs a broken down semi ...
- It's harder to change the motion of an object with large mass. In fact it turns out that the acceleration is **inversely proportional** to the mass of the object.



$$a = \frac{F}{m}$$




©2017 Pearson Education, Inc.

Quick Check


A cart is pulled to the right with a constant, steady force. How will its acceleration graph look?



A.





B.




C.

Quick Check


A cart is pulled to the right with a constant, steady force. How will its acceleration graph look?



A.



B.



C. ✓

- Alright, so if $a = F/m \rightarrow F = ma$ then what are the units of force?

Newton - units

- Alright, so if $a = F/m \rightarrow F = ma$ then what are the units of force?

$$\text{Force Unit} = \text{kg} \times \frac{\text{m}}{\text{s}^2} = \frac{\text{kg m}}{\text{s}^2} = \text{N}$$

- This is called a **newton**, so $1 \text{ N} = 1 \text{ kg m/s}^2$.

Inertial Mass

- 1 kg of mass is defined by a particular metal block in a vault in Paris (90% Platinum 10% Iridium). So it seems we could say that the mass of an object is related to how much stuff/matter it is made of.

Inertial Mass

- 1 kg of mass is defined by a particular metal block in a vault in Paris (90% Platinum 10% Iridium). So it seems we could say that the mass of an object is related to how much stuff/matter it is made of.
- However, now we have another way (more precise way) to define the mass of an object. If the same force causes one object to accelerate twice as much as another object then it must have half as much mass.

Inertial Mass

- 1 kg of mass is defined by a particular metal block in a vault in Paris (90% Platinum 10% Iridium). So it seems we could say that the mass of an object is related to how much stuff/matter it is made of.
- However, now we have another way (more precise way) to define the mass of an object. If the same force causes one object to accelerate twice as much as another object then it must have half as much mass.
- Mass *resists* acceleration due to a force, this is called **inertia** and so we call this mass **inertial mass**.


Newton's 2nd Law

- When there are multiple forces acting on an object the amount and direction of the acceleration is dictated by the net force.

$$\vec{a} = \frac{\vec{F}_{net}}{m}$$

more commonly seen as

$$\vec{F}_{net} = m\vec{a}$$



The acceleration is **NOT**
 $(T_1 + T_2)/m$ but

$$\vec{a} = \frac{\vec{T}_1 + \vec{T}_2}{m}$$

Newton's Laws

- Forces don't *overcome* each other, they simply add together to create a net force. Forces don't compete.
- One more interesting aspect of forces is that when an agent exerts a force on an object, the object exerts a force on agent as well. This is called **Newton's third law** and we'll talk more about it later. Think of hitting a solid brick wall . . . your hand exerted the force, but clearly the wall exerted one back.

A constant force causes an object to accelerate at 4 m/s^2 . What is the acceleration of an object with twice the mass that experiences the same force?

- A. 1 m/s^2
- B. 2 m/s^2
- C. 4 m/s^2
- D. 8 m/s^2
- E. 16 m/s^2

Quick Check

A constant force causes an object to accelerate at 4 m/s^2 . What is the acceleration of an object with twice the mass that experiences the same force?

- A. 1 m/s^2
- ✓ B. 2 m/s^2
- C. 4 m/s^2
- D. 8 m/s^2
- E. 16 m/s^2

Newton's 1st Law

- An object in motion tends to

Newton's 1st Law

- An object in motion tends to . . . stay in motion.

Newton's 1st Law

- An object in motion tends to . . . stay in motion.
- This is also called the law of inertia and is fully stated as

Newton's first law An object that is at rest will remain at rest, or an object that is moving will continue to move in a straight line with constant velocity, if and only if the net force acting on the object is zero.

© 2017 Pearson Education, Inc.


Newton's 1st Law

- An object on which the net force is zero and thus is either at rest or moving in a straight line with constant velocity is said to be in mechanical equilibrium and thus has zero acceleration.

An object at rest is in equilibrium: $\vec{F}_{\text{net}} = \vec{0}$.



$$\vec{v} = \vec{0} \quad \bullet \quad \vec{a} = \vec{0}$$




An object moving in a straight line at constant velocity is also in equilibrium: $\vec{F}_{\text{net}} = \vec{0}$.

© 2017 Pearson Education, Inc.

Quick Check


An object on a rope is lowered at constant speed.
Which is true?



- A. The rope tension is greater than the object's weight.
- B. The rope tension equals the object's weight.
- C. The rope tension is less than the object's weight.
- D. The rope tension can't be compared to the object's weight.

Quick Check


An object on a rope is lowered at constant speed.
Which is true?



- A. The rope tension is greater than the object's weight.
- ✓ B. The rope tension equals the object's weight.
- C. The rope tension is less than the object's weight.
- D. The rope tension can't be compared to the object's weight.

Quick Check


An object on a rope is lowered at a steadily decreasing speed. Which is true?



- A. The rope tension is greater than the object's weight.
- B. The rope tension equals the object's weight.
- C. The rope tension is less than the object's weight.
- D. The rope tension can't be compared to the object's weight.

Quick Check

An object on a rope is lowered at a steadily decreasing speed. Which is true?



- ✓A. The rope tension is greater than the object's weight.
- B. The rope tension equals the object's weight.
- C. The rope tension is less than the object's weight.
- D. The rope tension can't be compared to the object's weight.

Inertial Reference Frames



- Is there a force that throws this dummy into the windshield?

Inertial Reference Frames



- Is there a force that throws this dummy into the windshield?
- No, it just seems like it. What is the acceleration of the dummy relative to the car?

Inertial Reference Frames



- Is there a force that throws this dummy into the windshield?
- No, it just seems like it. What is the acceleration of the dummy relative to the car?
 - Large and positive

Inertial Reference Frames



- Is there a force that throws this dummy into the windshield?
- No, it just seems like it. What is the acceleration of the dummy relative to the car?
 - Large and positive
- What is the acceleration of the dummy relative to the earth?

Inertial Reference Frames




- Is there a force that throws this dummy into the windshield?
- No, it just seems like it. What is the acceleration of the dummy relative to the car?
 - Large and positive
- What is the acceleration of the dummy relative to the earth?
 - Zero

Inertial Reference Frames


- An **inertial reference frame** is a reference frame in which Newton's laws are valid . . . unlike the “car” reference frame in the dummy in the car situation.
- Inertial reference frames are non-accelerating reference frames.

(a) Cruising at constant speed.



The ball stays in place; the airplane is an inertial reference frame.

(b) Accelerating during takeoff.



The ball accelerates toward the back even though there are no horizontal forces; the airplane is *not* an inertial reference frame.

© 2017 Pearson Education, Inc.


Forces

- Every force has an agent which causes the force.
- Forces exist at the point of contact between the agent and the object (except for the few special cases of long-range forces).
- Forces exist due to interactions happening now, not due to what happened in the past.
- Consider a flying arrow.
 - A pushing force was required to accelerate the arrow as it was shot.
 - However, no force is needed to keep the arrow moving forward as it flies.
 - It continues to move because of inertia.




Quick Check

A hollow tube lies flat on a table.
A ball is shot through the tube.
As the ball emerges from the
other end, which path
does it follow?



Quick Check

A hollow tube lies flat on a table.
A ball is shot through the tube.
As the ball emerges from the
other end, which path
does it follow?



TACTICS BOX 5.3



Drawing a free-body diagram


- 1 **Identify all forces acting on the object.** This step was described in Tactics Box 5.2.
- 2 **Draw a coordinate system.** Use the axes defined in your pictorial representation.
- 3 **Represent the object as a dot at the origin of the coordinate axes.** This is the particle model.
- 4 **Draw vectors representing each of the identified forces.** This was described in Tactics Box 5.1. Be sure to label each force vector.
- 5 **Draw and label the *net force* vector \vec{F}_{net} .** Draw this vector beside the diagram, not on the particle. Or, if appropriate, write $\vec{F}_{\text{net}} = \vec{0}$. Then check that \vec{F}_{net} points in the same direction as the acceleration vector \vec{a} on your motion diagram.

Exercises 24–29




Quick Check

An elevator, lifted by a cable, is moving upward and slowing. Which is the correct free-body diagram?




Quick Check

An elevator, lifted by a cable, is moving upward and slowing. Which is the correct free-body diagram?




Quick Check

A ball has been tossed straight up. Which is the correct free-body diagram just after the ball has left the hand? Ignore air resistance.









Quick Check

A ball has been tossed straight up. Which is the correct free-body diagram just after the ball has left the hand? Ignore air resistance.





Quick Check

A ball, hanging from the ceiling by a string, is pulled back and released. Which is the correct free-body diagram just after its release?




Quick Check


A ball, hanging from the ceiling by a string, is pulled back and released. Which is the correct free-body diagram just after its release?




B.




C.



D.



E.



Picture References

Consider the Following (accessed 25 Sep 2017) <https://i.ytimg.com/vi/71VT6oTto1g/maxresdefault.jpg>