

Modern Physics and Quantum Mechanics

Fall 2001 Proficiency and Diagnostic

Useful constants:

- $e = 1.60 \times 10^{-19} \text{ C}$
- $hc = 1240 \text{ eV} \cdot \text{nm}$
- $c = 3.00 \times 10^8 \frac{\text{m}}{\text{s}}$
- $m_e = 0.511 \frac{\text{MeV}}{c^2}$

1. Cosmic ray photons from space are bombarding your laboratory and smashing massive objects to pieces. Your detectors indicate that two fragments, each of mass m_0 , depart such a collision moving at a speed of $0.6c$ at angles of 60° relative to the photon's original direction of motion.
 - a. In terms of m_0 and c , what is the energy of the cosmic ray photon?
 - b. In terms of m_0 , what is the mass M of the particle being struck (assumed to be initially stationary)?
2. The Pauli spin operators for a particle of spin $1/2$ are given by the matrices


$$\sigma_x = \begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}, \quad \sigma_y = \begin{pmatrix} 0 & -i \\ i & 0 \end{pmatrix}, \quad \sigma_z = \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}.$$

- a. Write the normalized eigenvectors of σ_z , $|+\rangle$ and $|-\rangle$ which are defined such that $\sigma_z|+\rangle = |+\rangle$ and $\sigma_z|-\rangle = -|-\rangle$, as column vectors in the same basis as the Pauli matrices given above. (You can assume, without loss of generality, that these vectors are real.)
 - b. Consider a wave vector $|\psi\rangle = a|+\rangle + b|-\rangle$. Assuming that a is a real number, $0 \leq a \leq 1$, show that $b = e^{i\phi}\sqrt{1-a^2}$, where $\phi \in [0, 2\pi)$ is an arbitrary angle.
 - c. Find the expectation values of σ_x , σ_y , and σ_z in the state $|\psi\rangle$ in terms of a and ϕ .
3. Neutron scattering is often done by cooling fast neutrons, which are among the fission products from a nuclear reactor, by thermalizing them in a moderator such as solid deuterium oxide (D_2O ice) held at some constant cryogenic temperature T . The average kinetic energy of neutrons from such a source is,

$$E = \frac{1}{2}m\langle v^2 \rangle = \frac{3}{2}kT,$$

where m is the neutron mass, $\langle v^2 \rangle$ is the mean-square velocity, and k is Boltzmann's constant.

- (a) What is the de Broglie wave length λ_n , for neutrons, in terms of their energy, mass, and Planck's constant?



- (b) What is the de Broglie wave length for neutrons with the average kinetic energy in terms of the temperature of the source?
- (c) Hydrogen freezes at 20.25 K and can be used to keep the D₂O moderator at a fixed temperature. What is the de Broglie wave length λ_n (in Å) for neutrons with the average energy corresponding to that temperature? What is the average neutron energy E (in eV)?
- (d) The neutrons are monochromatized at the 20.25 K thermal intensity maximum λ_n and diffracted off a crystalline target with lattice spacing $d = 4.21$ Å. What is the scattering angle θ that would give Bragg diffraction of the neutrons from this crystal? Use the convention for diffraction shown in the figure.

The neutron mass is $m_n = 1.68 \times 10^{-27}$ kg.

4. The time-independent wave function for a particle of mass m which moves in a one-dimensional potential $V(x)$ has the form $\psi(x) = A \exp[-a^2 x^2]$, where $a = \sqrt{m\omega/2\hbar}$ and A is a normalization constant.
 - a. Using the time-independent Schrödinger equation, find $V(x)$ and the energy eigenvalue for $\psi(x)$.
 - b. Identify the system. Which one of its quantum states is described by $\psi(x)$?
5. The Hamiltonian for the hydrogen atom is given by

$$H = -\frac{\hbar^2}{2m} \left[\frac{1}{r^2} \frac{\partial}{\partial r} \left(r^2 \frac{\partial}{\partial r} \right) + \frac{1}{r^2 \sin(\theta)} \frac{\partial}{\partial \theta} \left(\sin(\theta) \frac{\partial}{\partial \theta} \right) + \frac{1}{r^2 \sin^2(\theta)} \frac{\partial^2}{\partial \phi^2} \right] - \frac{e^2}{r},$$

where r , θ and ϕ are spherical coordinates of the position of the electron relative to the nucleus, m is the reduced mass, and e is the charge of the proton.

- (a) What are the quantum numbers characterizing the wave functions of the hydrogen atom?
- (b) Consider an s -wave state. Obtain the value of α for which the wave function

$$\psi(r) = C \exp(-\alpha r)$$

is a solution of Schrödinger's equation. Find the corresponding energy eigenvalue. Calculate the normalization constant C .

- (c) Obtain the expectation values of \vec{p}^2 and $1/r$ for the wave function in (b). Hint: the Hamiltonian above contains the expression for \vec{p}^2 in spherical coordinates.
- (d) Verify that the expectation value of the kinetic energy is $-1/2$ times that of the potential energy (virial theorem).
6. Consider a quantum mechanical system with two states, $|\alpha\rangle$ and $|\beta\rangle$. In this orthonormal basis of states the Hamiltonian is given by the matrix

$$H = \begin{pmatrix} W & V \\ V & -W \end{pmatrix}.$$

- (a) Obtain the exact energy eigenvalues.
- (b) Consider the Hamiltonian as $H = H_W + H_V$,

$$H_W = \begin{pmatrix} W & 0 \\ 0 & -W \end{pmatrix}, \quad H_V = \begin{pmatrix} 0 & V \\ V & 0 \end{pmatrix},$$

and assuming that $|V| \ll |W|$ obtain the energy eigenvalues to second order perturbation theory in V .

- (c) Compare your results in (a) and (b) and verify that they agree to second order in V .