


PHY6938 Proficiency Exam Spring 2002
April 5 , 2002
Mechanics

1. As shown in the diagram, two blocks with masses M_1 and M_2 are attached by an unstretchable rope around a frictionless pulley with radius r and moment of inertia I . There is no slipping between the rope and the pulley. The coefficient of kinetic friction between the blocks is the same as between block 1 and the surface, μ . A horizontal force F is applied to M_1 . Find the acceleration of M_1 .



2. The asteroid Toro was discovered in 1964. Its radius is about 5 km. Acceleration due to gravity on earth is $g = 9.81 \text{ m/s}^2$, the earth's radius is 6378 km, and the gravitational constant $G = 6.67 \times 10^{-11} \frac{\text{m}^3}{\text{kg s}^2}$.
- (a) Assuming the density of Toro is the same as that of Earth, find its total mass and the acceleration due to gravity at its surface.
 - (b) Suppose a body is placed in a circular orbit around Toro, with radius just slightly larger than the asteroid's radius. What is the speed of the body ?
 - (c) If a rock is thrown into a circular orbit around Toro at a height of 200 m above the surface, what is its period of revolution ?
3. Consider the off-center elastic collision of two objects of equal mass when one is initially at rest.
- (a) Show that the final velocity vectors of the two objects are perpendicular to each other.
 - (b) Show that the incoming object cannot have a backward scattered component.