

Flash Cards for Quantum/Nuclear Monte Carlo

Cody L. Petrie

October 21, 2015

Parameters in the Code

g2bval(d2b,sxz,fij) Given sxz this computes the d2b terms.

hspot(???) ???

op2val(d2b,sp,spx) ???

opmult(sp) This multiplies sp(s,i) by the 15 operators in this order 1-3 sx,sy,sz, 4-6 tx,ty,tz, 7-9 sx*(tx,ty,tz), 10-12 sy*(tx,ty,tz), 13-15 sz*(tx,ty,tz). This outputs opmult(s,kop,i)

szxupdate(sxznew(out),detrat(out),sxzold,i,opi,sp) Here the outputted detratt is simply di(i).

vnpsi2(w,dopot) This subroutine ...???

d15(kop) =di(i)= $\sum_k S_{ik}^{-1} \langle k | \mathcal{O}_i | \mathbf{r}, s_i \rangle = (S'/S)_{ii}$ for a specific kop (1 of 15)

d2b(s,s',ij) = $\frac{\langle \Phi | R, s_1, \dots, s_{i-1}, s, s_{i+1}, \dots, s_{j-1}, s', s_{j+1}, \dots, s_A \rangle}{\langle \Phi | RS \rangle}$

d2bip(s,s',s'',s''',ij,kl) = $\frac{\langle \Phi | R, s_1, \dots, s, \dots, s', \dots, s'', \dots, s''', \dots, s_A \rangle}{\langle \Phi | R, S \rangle}$ OR

di(m) = $\sum_k S_{mk}^{-1} \langle k | \mathcal{O}_i | \mathbf{r}_i, s_i \rangle = \sum_s \text{opi}(s, m) \langle s | s_i \rangle$

f2b(s,s',ij) = $\sum_{kop=1}^{15} f_{ij}^{kop} \langle s s' | \mathcal{O}_{ij}^{kop} | s_i s_j \rangle$

fst(3,3,ij) = f in front of specific operator

opi(s,m) = $\sum_k S_{mk}^{-1} \langle k | \mathcal{O}_i | \mathbf{r}_i, s \rangle = \sum_{s'} \text{sxz}(s', i, m) \langle s' | \mathcal{O}_i | s \rangle$

ph(i,4,j,idet) = $\sum_k S_{ik} \langle k | \mathcal{O}_j | s_j \rangle$

sigma(3,3,npair) I think the 3 is x,y,z.

sigtau(3,3,3,3,npair) I think the 3 is x,y,z.

$$\text{sp}(\mathbf{s}, \mathbf{i}) = \langle s | s_i \rangle$$

$$\text{spx}(\mathbf{s}, \mathbf{15}, \mathbf{i}) = \langle s | \mathcal{O}_i^p | s_i \rangle, \text{ where } p \text{ goes over the 15 cartesian coordinates.}$$

$$\text{stz}(\mathbf{s}, \mathbf{s}') = \text{spx}(\mathbf{s}, \sigma_\alpha \tau_\gamma, \mathbf{i})^* \text{spx}(\mathbf{s}', \sigma_\beta \tau_\gamma, \mathbf{j}) = \langle s, s' | \sigma_{\alpha i} \tau_{\gamma i} \sigma_{\beta j} \tau_{\gamma j} | s_i s_j \rangle$$

$$\text{sx15}(\mathbf{s}, \mathbf{15}, \mathbf{i}, \mathbf{j}) = \sum_k S_{jk}^{-1} \langle k | \sum_{kop=1}^{15} \mathcal{O}_{kop} | \mathbf{r}, s \rangle = \text{opmult}(\text{sxz0}), \text{ where } \text{sx15}(:, \text{kop}, :, k) = \text{opi}(\mathbf{s}, k)$$

$$\text{sxz}(\mathbf{s}, \mathbf{i}, \mathbf{j}) = \sum_k S_{jk}^{-1} \langle k | \mathbf{r}_i, s \rangle$$

$$\text{sxzi}(\mathbf{s}, \mathbf{i}, \mathbf{j}, \mathbf{iop}) = \sum_k S_{jk}^{\prime-1} S'_{ki}(s) = \sum_k S_{jk}^{\prime-1} \langle k | \mathcal{O}_i^{iop} | \mathbf{r}_i, s \rangle$$

$$\text{sxzj}(\mathbf{s}, \mathbf{i}, \mathbf{j}) = \sum_k S_{jk}^{\prime\prime-1} S''_{ki}(s) = \sum_k S_{jk}^{\prime-1} \langle k | \mathcal{O}_{ij}^{iopjop} | \mathbf{r}_i, s \rangle, \text{ each } iop \text{ and } jop \text{ is looped over and added to d2b using call g2bval(d2b, sxzj, fij).}$$

$$\text{sz}(\mathbf{s}, \mathbf{s}') = \text{spx}(\mathbf{s}, \sigma_\alpha, \mathbf{i})^* \text{spx}(\mathbf{s}', \sigma_\beta, \mathbf{j}) = \langle s, s' | \sigma_{\alpha i} \sigma_{\beta j} | s_i s_j \rangle$$

$$\text{tau}(\mathbf{3}, \mathbf{3}, \mathbf{npair}) \text{ I think the 3 is x,y,z.}$$

$$\text{tz}(\mathbf{s}, \mathbf{s}') = \text{spx}(\mathbf{s}, \tau_\alpha, \mathbf{i})^* \text{spx}(\mathbf{s}', \tau_\alpha, \mathbf{j}) = \langle s, s' | \tau_{\alpha i} \tau_{\alpha j} | s_i s_j \rangle$$

Variational Monte Carlo

Steps for Metropolis Algorithm:

1. Start with some random walker configuration \mathbf{R}
2. Propose a move to a new walker \mathbf{R}' from the distribution $T(\mathbf{R}' \leftarrow \mathbf{R})$
3. The probability of accepting the move is given by

$$A(\mathbf{R}' \leftarrow \mathbf{R}) = \min \left(1, \frac{T(\mathbf{R}' \leftarrow \mathbf{R}) P(\mathbf{R}')}{T(\mathbf{R} \leftarrow \mathbf{R}') P(\mathbf{R})} \right).$$

The move is accepted if $U[0, 1] < A(\mathbf{R}' \leftarrow \mathbf{R})$.

4. Repeat from step 2.

Variational Energy (In terms of $E_L(\mathbf{R})$ and $P(\mathbf{R})$), $\mathbf{x2} + E_L$ and P :

$$E_V = \frac{\int \Psi^*(\mathbf{R}) \hat{H} \Psi(\mathbf{R}) d\mathbf{R}}{\int \Psi^*(\mathbf{R}) \Psi(\mathbf{R}) d\mathbf{R}} = \int P(\mathbf{R}) E_L(\mathbf{R}) d\mathbf{R}$$

$$P(\mathbf{R}) = |\Psi_T(\mathbf{R})|^2 / \int |\Psi_T(\mathbf{R})|^2 d\mathbf{R}$$

$$E_L(\mathbf{R}) = \Psi_T^{-1}(\mathbf{R}) \hat{H} \Psi_T(\mathbf{R})$$

Sampled Variational Energy:

$$E_V \approx \frac{1}{N} \sum_{n=1}^N E_L(\mathbf{R}_n)$$

where \mathbf{R}_n are drawn from $P(\mathbf{R})$.

General Physics

Atomic Shell Model

- **Shells** are areas in which an electron can "orbit" a nucleus. They correspond to the principle quantum numbers ($n=1,2,3,4,\dots$) where $n=1$ is the closest shell.
- The number of allowed electrons is controlled by the spin statistics the other allowed quantum numbers being $l=0,1,\dots,n-1$ and $m_l=-l,\dots,l$. The number allowed for each shell is all the possible times two since $m_s=-1/2,1/2$.
- **Subshells** are given by the l quantum numbers where $l=1,2,3,4$ are s,p,d,f,g.
- List of how many each shell can hold

Shell name	Subshell name	Subshell max electrons	Shell max electrons
K	1s	2	2
L	2s	2	$2 + 6 = \mathbf{8}$
	2p	6	
M	3s	2	$2 + 6 + 10 = \mathbf{18}$
	3p	6	
	3d	10	
N	4s	2	$2 + 6 + 10 + 14 = \mathbf{32}$
	4p	6	
	4d	10	
	4f	14	

Nuclear Shell Model

The nuclear shell model is similar to the atomic shell model except that $n=1,2,3,4,\dots$ and $l=0,1,2,3,\dots$ independently from n . So you can have states like 1g