

---

Shorter Notes: The Roots of a Polynomial Vary Continuously as a Function of the Coefficients

Author(s): Gary Harris and Clyde Martin

Source: *Proceedings of the American Mathematical Society*, Jun., 1987, Vol. 100, No. 2 (Jun., 1987), pp. 390-392

Published by: American Mathematical Society

Stable URL: <https://www.jstor.org/stable/2045978>

---

JSTOR is a not-for-profit service that helps scholars, researchers, and students discover, use, and build upon a wide range of content in a trusted digital archive. We use information technology and tools to increase productivity and facilitate new forms of scholarship. For more information about JSTOR, please contact [support@jstor.org](mailto:support@jstor.org).

Your use of the JSTOR archive indicates your acceptance of the Terms & Conditions of Use, available at <https://about.jstor.org/terms>



American Mathematical Society is collaborating with JSTOR to digitize, preserve and extend access to *Proceedings of the American Mathematical Society*

JSTOR

## SHORTER NOTES

The purpose of this department is to publish very short papers of unusually polished character, for which there is no other outlet.

### THE ROOTS OF A POLYNOMIAL VARY CONTINUOUSLY AS A FUNCTION OF THE COEFFICIENTS

GARY HARRIS AND CLYDE MARTIN

**ABSTRACT.** We present an elementary topological proof that the roots of a polynomial vary continuously as a function of the coefficients.

It is, or should be, common knowledge that the roots of a polynomial over  $\mathbb{C}$  vary continuously as a function of the coefficients. Here we present an elementary topological proof of this important fact. Of course, the coefficients vary continuously as a function of the roots; indeed, the coefficients can be given as a symmetric polynomial mapping of the roots. It is natural to ask if this mapping is continuously invertible, and this is the question we answer in Theorem A.

Before becoming more precise, we should mention that proofs, probably in large numbers, exist in the literature in differing contexts. For example, for polynomials of degree 2, 3, or 4, Galois theory tells us the roots can be found by explicit formulas of the coefficients involving radicals. ([4] is a good source.) However, a general study of continuity can still be tricky. For polynomials without multiple roots, the Complex Implicit Function Theorem can be used to show the roots vary analytically with the coefficients [1]. For general polynomials our Theorem B follows from an application of Rouché's theorem [3].

As stated above, our purpose is to give a topological proof that the roots of a polynomial are given as a continuous function of the coefficients. We may assume all polynomials  $P$  of degree  $n$  are normalized so

$$P(z) = z^n + a_1 z^{n-1} + \cdots + a_n.$$

We will identify  $P$  with the vector  $a = (a_1, \dots, a_n)$  in  $\mathbb{C}^n$ . By the Fundamental Theorem of Algebra, we know  $P$  can be factored as

$$P(z) = \prod_{j=1}^n (z - \xi_j)$$

---

Received by the editors July 21, 1986 and, in revised form, November 5, 1986.

1980 *Mathematics Subject Classification* (1985 *Revision*). Primary 26C10, 30C15.

The second author was supported in part by NASA grants #NAG-82, #NAG-203 and NSA grant #MDA904-85-H-0009.

©1987 American Mathematical Society  
 0002-9939/87 \$1.00 + \$.25 per page

for a unique sequence  $\{\xi_1, \dots, \xi_n\}$  of elements of  $\mathbb{C}$ . We also know for each  $1 \leq j \leq n$ ,

$$a_j = \sigma_j(\xi_1, \dots, \xi_n)$$

for symmetric polynomials  $\sigma_1, \dots, \sigma_n$  in  $n$  variables. Let  $\sigma: \mathbb{C}^n \rightarrow \mathbb{C}^n$  be the continuous mapping defined by

$$\sigma(\xi) \doteq (\sigma_1(\xi), \dots, \sigma_n(\xi)).$$

We are assuming  $\mathbb{C}^n$  has the euclidean norm and are using the notation  $\xi \doteq (\xi_1, \dots, \xi_n)$ . It follows from the Fundamental Theorem of Algebra that  $\sigma$  is surjective.

The map  $\sigma$  is not injective since it is independent of the arrangement of the components of the vector  $\xi$ . For  $\mu \in S_n$ , the group of permutations on  $n$  elements, and  $\xi = (\xi_1, \dots, \xi_n)$  in  $\mathbb{C}^n$  let  $\xi_\mu \doteq (\xi_{\mu(1)}, \dots, \xi_{\mu(n)})$ . We define an equivalence relation on  $\mathbb{C}^n$  by

$$\xi \sim \beta \Leftrightarrow \exists \mu \in S_n \ni \xi_\mu = \beta.$$

Let  $\mathbb{C}^n / \sim$  be the quotient space endowed with the quotient topology induced by the canonical projection  $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^n / \sim$ , and let  $\hat{\sigma}$  be the unique mapping of  $\mathbb{C}^n / \sim$  onto  $\mathbb{C}^n$  for which  $\hat{\sigma} \circ \pi = \sigma$ . It follows that  $\hat{\sigma}$  is a continuous bijection. (All the facts we use about the quotient topology can be found in any good text on topology, for example [2, pp. 94–99].)

We can now prove our

**THEOREM A.** *The mapping  $\hat{\sigma}: \mathbb{C}^n / \sim \rightarrow \mathbb{C}^n$  is a homeomorphism.*

**PROOF.** Let  $d$  be the metric defined on  $\mathbb{C}^n / \sim$  by

$$d(\pi(\xi), \pi(\beta)) \doteq \min\{|\xi' - \beta'| : \xi' \in \pi(\xi) \text{ and } \beta' \in \pi(\beta)\}.$$

To see that  $d$  is a metric let  $\xi, \beta, \eta \in \mathbb{C}^n$ . Clearly

$$d(\pi(\xi), \pi(\beta)) = \min\{|\xi - \beta'| : \beta' \in \pi(\beta)\}.$$

We can choose  $\eta' \in \pi(\eta)$  so that  $d(\pi(\xi), \pi(\eta)) = |\xi - \eta'|$ . For each  $\beta' \in \pi(\beta)$ ,  $|\xi - \beta'| \leq |\xi - \eta'| + |\eta' - \beta'|$ . So

$$\begin{aligned} d(\pi(\xi), \pi(\beta)) &\leq \min\{|\xi - \eta'| + |\eta' - \beta'| : \beta' \in \pi(\beta)\} \\ &= d(\pi(\xi), \pi(\eta)) + d(\pi(\eta), \pi(\beta)). \end{aligned}$$

It follows immediately that  $\pi: \mathbb{C}^n \rightarrow \mathbb{C}^n / \sim$  is continuous if  $\mathbb{C}^n / \sim$  is endowed with the topology induced by  $d$ . Moreover for any set  $A$  in  $\mathbb{C}^n$  we have  $\pi^{-1}(\pi(A))$  is open (closed) if  $A$  is open (closed). Thus  $\pi$  is an open and closed map. Hence the topology induced by  $d$  is the quotient topology. Let  $M > 0$  be given and let  $B(0, M)$  denote the open ball about 0 in  $\mathbb{C}^n / \sim$  of radius  $M$ . We claim that  $\hat{\sigma}|_{\overline{B(0, M)}}$  is a homeomorphism. It suffices to show that  $\hat{\sigma}$  is closed. Let  $C \subset \overline{B(0, M)}$  be closed, then  $\pi^{-1}(C)$  is a closed and bounded subset of  $\mathbb{C}^n$ , hence compact. Thus  $\hat{\sigma}(C) = \sigma(\pi^{-1}(C))$  is compact, and hence closed in  $\mathbb{C}^n$ . We can now complete the proof by showing that  $\hat{\sigma}: \mathbb{C}^n / \sim \rightarrow \mathbb{C}^n$  is open. Let  $U$  be an open subset of  $\mathbb{C}^n / \sim$  and  $x \in U$ . Choose  $\varepsilon > 0$  and  $M > 0$  so that  $B(x, \varepsilon) \subset U$  and  $\overline{B(x, \varepsilon)} \subset B(0, M)$ . Since  $\hat{\sigma}$  is open on  $B(0, M)$  it follows that  $\hat{\sigma}(x)$  is in the interior of  $\hat{\sigma}(B(x, \varepsilon))$ . Since  $x$  was arbitrary this completes the proof.

Let  $\tau: \mathbb{C}^n \rightarrow \mathbb{C}^n/\sim$  denote the continuous inverse of  $\hat{\sigma}$ . Given  $a \in \mathbb{C}^n$  ( $a$  identified with a polynomial  $P$  as above) and  $\varepsilon > 0$ , Theorem A tells us there exists  $\delta > 0$  so that  $\tau(B(a, \delta)) \subset B(\tau(a), \varepsilon)$ . Writing out what this means in terms of the euclidean metric on  $\mathbb{C}^n$  and the metric  $d$  on  $\mathbb{C}^n/\sim$ , we obtain

**THEOREM B.** *Suppose*

$$P(z) = z^n + a_1 z^{n-1} + \cdots + a_n = \prod_{j=1}^s (z - \xi_j)^{m_j}$$

*for distinct  $\xi_1, \dots, \xi_s$ . Let  $\varepsilon > 0$  be given so that  $i \neq j$  implies that  $B(\xi_i, \varepsilon) \cap B(\xi_j, \varepsilon) = \emptyset$ . Then there exists  $\delta > 0$  so that  $b \in B(a, \delta)$  implies the polynomial*

$$Q(z) \doteq z^n + b_1 z^{n-1} + \cdots + b_n$$

*has exactly  $m_j$  roots (counting multiplicity) in  $B(\xi_j, \varepsilon)$ .*

#### REFERENCES

1. Lars Hörmander, *An introduction to complex analysis in several variables*, Van Nostrand, New York, 1966
2. John Kelley, *General topology*, Van Nostrand, New York, 1955
3. Morris Marden, *Geometry of polynomials*, Math. Surveys, no. 3, Amer. Math. Soc., Providence, R.I., 1966
4. B. L. van der Waerden, *Algebra*, I, Ungar, New York, 1970

DEPARTMENT OF MATHEMATICS, TEXAS TECH UNIVERSITY, LUBBOCK, TEXAS 79409