

# Functional Differential Geometry

merely in ClojureScript

Draft mode

# Functional Differential Geometry

merely in ClojureScript

Gerald Jay Sussman and Jack Wisdom

with Will Farr

The MIT Press  
Cambridge, Massachusetts  
London, England

This Typst edition was based on the source material published by [mentat-collective/fdg-book](https://mentat-collective/fdg-book). This version is licensed under CC BY NC SA 4.0. The text below is a reproduction of the published version of this book and does not represent the version you are looking at.

© 2013 Massachusetts Institute of Technology



This work is licensed under the Creative Commons Attribution-NonCommercial-ShareAlike 3.0 Unported License. To view a copy of this license, visit [creativecommons.org](https://creativecommons.org).

Other than as provided by this license, no part of this book may be reproduced, transmitted, or displayed by any electronic or mechanical means without permission from the MIT Press or as permitted by law.

MIT Press books may be purchased at special quantity discounts for business or sales promotional use. For information, please email [special\\_sales@mitpress.mit.edu](mailto:special_sales@mitpress.mit.edu) or write to Special Sales Department, The MIT Press, 55 Hayward Street, Cambridge, MA 02142.

This book was set in Computer Modern by the authors with the  $\text{\LaTeX}$  typesetting system and was printed and bound in the United States of America.

Library of Congress Cataloging-in-Publication Data

Sussman, Gerald Jay.

Functional Differential Geometry / Gerald Jay Sussman and Jack Wisdom; with Will Farr.  
p. cm.

Includes bibliographical references and index.

ISBN 978-0-262-01934-7 (hardcover : alk. paper)

1. Geometry, Differential. 2. Functional Differential Equations.  
3. Mathematical Physics.

I. Wisdom, Jack. II. Farr, Will. III. Title.

QC20.7.D52S87 2013

516.3'6—dc23

2012042107

*The author has spared himself no pains in his endeavour to present the main ideas in the simplest and most intelligible form, and on the whole, in the sequence and connection in which they actually originated. In the interest of clearness, it appeared to me inevitable that I should repeat myself frequently, without paying the slightest attention to the elegance of the presentation. I adhered scrupulously to the precept of that brilliant theoretical physicist L. Boltzmann, according to whom matters of elegance ought be left to the tailor and to the cobbler.*

*Albert Einstein, in Relativity, the Special and General Theory, (1961), p. v*

# Contents

|                                                                  |      |
|------------------------------------------------------------------|------|
| Preface to the Merely ClojureScript Edition .....                | ix   |
| Preface .....                                                    | x    |
| Acknowledgements .....                                           | x    |
| Prologue .....                                                   | xii  |
| Programming and Understanding .....                              | xii  |
| Functional Abstraction .....                                     | xiii |
| 1 Introduction .....                                             | 1    |
| 1.1 Lagrange Equations .....                                     | 2    |
| 1.2 The Metric .....                                             | 3    |
| 1.3 Euler-Lagrange Residuals .....                               | 4    |
| 1.4 Geodesic Equations .....                                     | 5    |
| 1.4.1 Exercise 1.1: Motion on a Sphere .....                     | 6    |
| 2 Manifolds .....                                                | 7    |
| 2.1 Coordinate Functions .....                                   | 7    |
| 2.2 Manifold functions .....                                     | 9    |
| 2.3 <i>Manifold Functions Are Coordinate Independent</i> .....   | 10   |
| 2.4 Naming Coordinate Functions .....                            | 10   |
| 2.5 Exercise 2.1: Curves .....                                   | 11   |
| 2.6 Exercise 2.2: Stereographic Projection .....                 | 11   |
| 3 Vector Fields and One-Form Fields .....                        | 13   |
| 3.1 Vector Fields .....                                          | 13   |
| 3.1.1 Coordinate Representation .....                            | 15   |
| 3.1.2 Vector Field Properties .....                              | 15   |
| 3.2 Coordinate-Basis Vector Fields .....                         | 16   |
| 3.2.1 Coordinate Transformations .....                           | 17   |
| 3.3 Integral Curves .....                                        | 17   |
| 3.4 One-Form Fields .....                                        | 19   |
| 3.4.1 Differential of a Function .....                           | 20   |
| 3.4.2 One-Form Fields .....                                      | 20   |
| 3.4.3 Coordinate-Basis One-Form Fields .....                     | 20   |
| 3.4.4 Not All One-Form Fields Are Differentials .....            | 22   |
| 3.4.5 Coordinate Transformations .....                           | 23   |
| 4 Basis Fields .....                                             | 24   |
| 4.1 Change of Basis .....                                        | 26   |
| 4.2 Rotation Basis .....                                         | 27   |
| 4.3 Commutators .....                                            | 28   |
| 4.4 Exercise 4.1: Alternate Angles .....                         | 32   |
| 4.5 Exercise 4.2: General Commutators .....                      | 32   |
| 4.6 Exercise 4.3: $SO(3)$ Basis and Angular Momentum Basis ..... | 32   |
| 5 Integration .....                                              | 33   |
| 5.1 Higher Dimensions .....                                      | 34   |
| 5.2 Wedge Product .....                                          | 34   |
| 5.3 3-Dimensional Euclidean Space .....                          | 35   |
| 5.4 Exercise 5.1: Wedge Product .....                            | 37   |
| 5.5 Exterior Derivative .....                                    | 37   |
| 5.6 Computing Exterior Derivatives .....                         | 38   |
| 5.7 Properties of Exterior Derivatives .....                     | 38   |
| 5.8 Stokes's Theorem .....                                       | 39   |
| 5.9 Vector Integral Theorems .....                               | 40   |
| 5.10 Exercise 5.2: Graded Formula .....                          | 41   |
| 5.11 Exercise 5.3: Iterated Exterior Derivative .....            | 41   |
| 6 Over a Map .....                                               | 42   |
| 6.1 Vector Fields Over a Map .....                               | 42   |
| 6.2 Restricted Vector Fields .....                               | 42   |
| 6.3 Differential of a Map .....                                  | 42   |

|        |                                                             |    |
|--------|-------------------------------------------------------------|----|
| 6.4    | Velocity at a Time .....                                    | 43 |
| 6.5    | One-Form Fields Over a Map .....                            | 43 |
| 6.6    | Basis Fields Over a Map .....                               | 43 |
| 6.7    | Walking on a Sphere .....                                   | 44 |
| 6.8    | Components of the Velocity .....                            | 44 |
| 6.9    | Pullbacks and Pushforwards .....                            | 45 |
| 6.10   | Pullback and Pushforward of a Function .....                | 45 |
| 6.11   | Pushforward of a Vector Field .....                         | 45 |
| 6.12   | Pushforward Along Integral Curves .....                     | 46 |
| 6.13   | Pullback of a Vector Field .....                            | 46 |
| 6.14   | Pullback of a Form Field .....                              | 46 |
| 6.15   | Properties of Pullback .....                                | 47 |
| 6.16   | Pushforward of a Form Field .....                           | 48 |
| 6.16.1 | Exercise 6.1: Velocities on a Globe .....                   | 48 |
| 7      | Directional Derivatives .....                               | 49 |
| 7.1    | Lie Derivative .....                                        | 50 |
| 7.2    | Functions .....                                             | 50 |
| 7.3    | Vector Fields .....                                         | 50 |
| 7.4    | An Alternate View .....                                     | 51 |
| 7.5    | Form Fields .....                                           | 52 |
| 7.6    | Uniform Interpretation .....                                | 52 |
| 7.7    | Properties of the Lie Derivative .....                      | 52 |
| 7.8    | Exponentiating Lie Derivatives .....                        | 53 |
| 7.9    | Interior Product .....                                      | 54 |
| 7.10   | Covariant Derivative .....                                  | 55 |
| 7.11   | Covariant Derivative of Vector Fields .....                 | 55 |
| 7.12   | An Alternate View .....                                     | 57 |
| 7.13   | Covariant Derivative of One-Form Fields .....               | 58 |
| 7.14   | Change of Basis .....                                       | 59 |
| 7.15   | Parallel Transport .....                                    | 62 |
| 7.16   | On a Sphere .....                                           | 63 |
| 7.17   | On a Great Circle .....                                     | 64 |
| 7.18   | Geodesic Motion .....                                       | 65 |
| 7.18.1 | Exercise 7.1: Hamiltonian Evolution .....                   | 67 |
| 7.18.2 | Exercise 7.2: Lie Derivative and Covariant Derivative ..... | 67 |
| 8      | Curvature .....                                             | 69 |
| 8.1    | Explicit Transport .....                                    | 70 |
| 8.1.1  | Verification in Two Dimensions .....                        | 71 |
| 8.1.2  | Geometrically .....                                         | 72 |
| 8.1.3  | Terms of the Riemann Curvature .....                        | 73 |
| 8.1.4  | Ricci Curvature .....                                       | 74 |
| 8.1.5  | Exercise 8.1: Ricci of a Sphere .....                       | 74 |
| 8.1.6  | Exercise 8.2: Pseudosphere .....                            | 74 |
| 8.2    | Torsion .....                                               | 75 |
| 8.2.1  | Torsion Doesn't Affect Geodesics .....                      | 75 |
| 8.3    | Geodesic Deviation .....                                    | 76 |
| 8.3.1  | Longitude Lines on a Sphere .....                           | 77 |
| 8.4    | Bianchi Identities .....                                    | 78 |
| 9      | Metrics .....                                               | 81 |
|        | Metric Music .....                                          | 81 |
| 9.1    | Metric Compatibility .....                                  | 82 |
| 9.1.1  | Exercise 9.1: Metric Compatibility .....                    | 83 |
| 9.2    | Metrics and Lagrange Equations .....                        | 83 |
| 9.3    | Kinetic Energy or Arc Length .....                          | 84 |
| 9.3.1  | For Two Dimensions .....                                    | 85 |
| 9.3.2  | Reparametrization .....                                     | 86 |

|        |                                                          |     |
|--------|----------------------------------------------------------|-----|
| 9.3.3  | Exercise 9.2: SO(3) Geodesics                            | 87  |
| 9.3.4  | Exercise 9.3: Curvature of a Spherical Surface           | 87  |
| 9.3.5  | Exercise 9.4: Curvature of a Pseudosphere                | 88  |
| 9.4    | General Relativity                                       | 88  |
| 9.4.1  | Exercise 9.5: Newton's Equations                         | 88  |
| 9.4.2  | Exercise 9.6: Curvature of Schwarzschild Spacetime       | 90  |
| 9.4.3  | Exercise 9.7: Circular Orbits in Schwarzschild Spacetime | 90  |
| 9.4.4  | Exercise 9.8: Stability of Circular Orbits               | 91  |
| 9.4.5  | Exercise 9.9: Friedmann-Lemaître-Robertson-Walker        | 91  |
| 9.4.6  | Exercise 9.10: Cosmology                                 | 92  |
| 10     | Hodge Star and Electrodynamics                           | 94  |
| 10.1   | Relationship to Vector Calculus                          | 94  |
| 10.2   | Spherical Coordinates                                    | 96  |
| 10.3   | The Wave Equation                                        | 97  |
| 10.4   | Electrodynamics                                          | 98  |
| 10.5   | Maxwell's Equations                                      | 100 |
| 10.6   | Lorentz Force                                            | 102 |
| 10.6.1 | Exercise 10.1: Relativistic Lorentz Force                | 103 |
| 11     | Special Relativity                                       | 104 |
| 11.1   | Invariance of the Wave Equation                          | 106 |
| 11.2   | Lorentz Transformations                                  | 106 |
| 11.3   | Simple Lorentz Transformations                           | 107 |
| 11.4   | More General Lorentz Transformations                     | 108 |
| 11.5   | Implementation                                           | 109 |
| 11.6   | Rotations                                                | 110 |
| 11.7   | General Lorentz Transformations                          | 110 |
| 11.7.1 | Exercise 11.1: Lorentz Decomposition                     | 111 |
| 11.8   | Special Relativity Frames                                | 111 |
| 11.8.1 | Velocity Addition Formula                                | 111 |
| 11.9   | Twin Paradox                                             | 112 |
| A      | Scheme                                                   | 115 |
| A.1    | Procedure Calls                                          | 115 |
| A.2    | Lambda Expressions                                       | 115 |
| A.3    | Definitions                                              | 116 |
| A.4    | Conditionals                                             | 116 |
| A.5    | Recursive Procedures                                     | 117 |
| A.6    | Local Names                                              | 117 |
| A.7    | Compound Data — Lists and Vectors                        | 118 |
| A.8    | Symbols                                                  | 119 |
| B      | Our Notation                                             | 121 |
| B.1    | Functions                                                | 121 |
| B.2    | Symbolic Values                                          | 122 |
| B.3    | Tuples                                                   | 123 |
| B.4    | Derivatives                                              | 125 |
| B.5    | Derivatives of Functions of Multiple Arguments           | 126 |
| B.6    | Structured Results                                       | 128 |
| B.6.1  | Exercise B.1: Chain Rule                                 | 129 |
| B.6.2  | Exercise B.2: Computing Derivatives                      | 129 |
| C      | Tensors                                                  | 130 |
| D      | ClojureScript                                            | 133 |
| D.1    | Function Calls                                           | 133 |
| D.2    | Function Expressions                                     | 133 |
| D.3    | Definitions                                              | 134 |
| D.3.1  | Names, functions, and local bindings                     | 134 |
| D.4    | Conditionals                                             | 135 |
| D.5    | Recursive Functions                                      | 135 |

|       |                                                       |     |
|-------|-------------------------------------------------------|-----|
| D.6   | Local Names .....                                     | 136 |
| D.7   | Compound Data — Lists and Vectors .....               | 136 |
| D.8   | Symbols .....                                         | 137 |
| D.9   | Scheme–ClojureScript Cheat Sheet .....                | 139 |
| E     | Our Notation in Emmy .....                            | 140 |
| E.1   | Functions .....                                       | 140 |
| E.2   | Symbolic Values .....                                 | 141 |
| E.3   | Tuples .....                                          | 142 |
| E.4   | Derivatives .....                                     | 144 |
| E.5   | Derivatives of Functions of Multiple Arguments .....  | 145 |
| E.6   | Structured Results .....                              | 147 |
| E.6.1 | Exercise E.1: Chain Rule .....                        | 148 |
| E.6.2 | Exercise E.2: Computing Derivatives .....             | 148 |
| F     | Tensors in Emmy .....                                 | 149 |
| G     | Running the Emmy Examples .....                       | 154 |
| G.1   | What the Book Runner Does .....                       | 154 |
| G.2   | Preparing and Starting the Runner .....               | 154 |
| G.3   | Automated Runs and Captured Results .....             | 155 |
| G.3.1 | Exact checks for expensive symbolic blocks .....      | 155 |
| G.4   | Running a Repository Block Manually .....             | 156 |
| G.5   | A Clean ClojureScript and Emmy Project .....          | 156 |
| H     | Edition-specific Prose Record .....                   | 159 |
| I     | Source and Edition Change Log .....                   | 160 |
| I.1   | Conversion contract .....                             | 160 |
| I.2   | Source/PDF fidelity corrections .....                 | 160 |
| I.3   | Source text, code, and math normalization .....       | 161 |
| I.4   | Imported setup and executable-claim repairs .....     | 163 |
| I.5   | Equation layout and counter changes .....             | 163 |
| I.6   | ClojureScript and Emmy edition transformations .....  | 164 |
| I.7   | Project-authored material and execution records ..... | 166 |
| I.8   | Annotated errata corrections .....                    | 166 |
| I.9   | Edition-specific prose records .....                  | 167 |
| I.10  | Presentation and policy changes .....                 | 168 |
| I.11  | Maintenance rule .....                                | 168 |
|       | Errata for FDG .....                                  | 169 |
|       | Chapter 1 .....                                       | 169 |
|       | Chapter 3 .....                                       | 169 |
|       | Chapter 4 .....                                       | 169 |
|       | Chapter 5 .....                                       | 169 |
|       | Chapter 6 .....                                       | 169 |
|       | Chapter 7 .....                                       | 169 |
|       | Chapter 8 .....                                       | 170 |
|       | Chapter 9 .....                                       | 171 |
|       | Chapter 10 .....                                      | 171 |
|       | Chapter 11 .....                                      | 172 |
|       | References .....                                      | 174 |



## Preface to the Merely ClojureScript Edition

This edition of the book is a vibe-coded monstrosity created by a mathematician who can program predominantly in Python but not (yet?) in Scheme or Clojure(Script). Starting from [mentat-collective/fdg-book](https://mentat-collective/fdg-book), ChatGPT 5.5 (and later ChatGPT 5.6) was used to create the PDF you are currently reading.

This book can be found in the Github repository [clvnkhr/fdgtyp](https://clvnkhr/fdgtyp). In this repository, one will also find the variant versions (Scheme only, ClojureScript, and both Scheme and ClojureScript), a copy of the org version, the code that produces variant books version, and a companion web-runner that can run the ClojureScript codeblocks in the book. This companion runner exists as a Github page at <https://clvnkhr.github.io/fdgtyp/> and can be run locally - see the emmy-runner folder.

There was no attempt made at matching the pagination or style of the original PDF book.

On the codeblocks: the .scm codeblocks are updated after an has ‘audited’ them. But I did not run any .scm code. The level of trust should be tempered against the following: my LLM(s) have read the code, and have converted the code to ClojureScript (i.e. there is a somewhat ad-hoc script that does this), and have ‘audited’ the ClojureScript code to compile and do what the surrounding text says it should do. This has led to discovering some errors in the original .scm codeblocks, such as outputs that were not commented out, typos in function names, or incorrectly saved outputs. The ClojureScript blocks compile and their outputs match the (audited) Scheme output comments. There were some changes that we had to put in to get to this point - apart from changing  $1/2$  to  $(/ 1 2)$  to avoid Javascript computing as floats, we needed to manually add calls to `simplify` to get outputs closer to `scmutils`’ outputs. Also, we (the AI, really) had to write a small compatability layer (`emmy-runner/src/fdg/compat.cljs`) to get closer to `scmutils`’ behavior.

I also (got the AI to) redo the diagrams and the cover page. The cover page is a real plot of trajectories, but is not a perfect recreation as I ran out of time to improve it further.

A comment on why this is a ‘mere’ translation. The name is a direct reference to [merely-true/merely-true](https://merely-true/), which is a “Lean project for AI generated mathematics, with very permissive rules about what can be contributed.” Although there seem to be other projects now in this vein, such as Mathlib’s [Formal Frontier](https://formal-frontier/) and the [TauCeti Project](https://tau-ceti-project/), they have avoided the beautifully evocative adjective ‘merely’. For the Lean community, AI contributions are at time of writing just not good enough for inclusion into the main mathlib project, even though they do certify correctness to the same extent. The merely-true project was meant as a suitable dumping ground for these less-than-stellar AI contributions. So it is for us, that I would not dare to call this a properly vetted translation, for I do not personally have the skills, and surely the result is inferior as a result. Yet AI has allowed me to convert the org and pdf book into a Typst book with ClojureScript codeblocks. The codeblocks all compile, and the book compiles. The diagrams are redone in Typst/ [CetZ](https://cet.z/). And now I am excited to read the book, and begin to learn some ClojureScript (and perhaps some differential geometry).

Last updated: 5th August 2026.

The page you just read is my sole human contribution in the book.

Calvin

## Preface

Learning physics is hard. Part of the problem is that physics is naturally expressed in mathematical language. When we teach we use the language of mathematics in the same way that we use our natural language. We depend upon a vast amount of shared knowledge and culture, and we only sketch an idea using mathematical idioms. We are insufficiently precise to convey an idea to a person who does not share our culture. Our problem is that since we share the culture we find it difficult to notice that what we say is too imprecise to be clearly understood by a student new to the subject. A student must simultaneously learn the mathematical language and the content that is expressed in that language. This is like trying to read *Les Misérables* while struggling with French grammar.

This book is an effort to ameliorate this problem for learning the differential geometry needed as a foundation for a deep understanding of general relativity or quantum field theory. Our approach differs from the traditional one in several ways. Our coverage is unusual. We do not prove the general Stokes's Theorem—this is well covered in many other books—instead, we show how it works in two dimensions. Because our target is relativity, we put lots of emphasis on the development of the covariant derivative, and we erect a common context for understanding both the Lie derivative and the covariant derivative. Most treatments of differential geometry aimed at relativity assume that there is a metric (or pseudometric). By contrast, we develop as much material as possible independent of the assumption of a metric. This allows us to see what results depend on the metric when we introduce it. We also try to avoid the use of traditional index notation for tensors. Although one can become very adept at “index gymnastics,” that leads to much mindless (though useful) manipulation without much thought to meaning. Instead, we use a semantically richer language of vector fields and differential forms.

But the single biggest difference between our treatment and others is that we integrate computer programming into our explanations. By programming a computer to interpret our formulas we soon learn whether or not a formula is correct. If a formula is not clear, it will not be interpretable. If it is wrong, we will get a wrong answer. In either case we are led to improve our program and as a result improve our understanding. We have been teaching advanced classical mechanics at MIT for many years using this strategy. We use precise functional notation and we have students program in a functional language. The students enjoy this approach and we have learned a lot ourselves. It is the experience of writing software for expressing the mathematical content and the insights that we gain from doing it that we feel is revolutionary. We want others to have a similar experience.

## Acknowledgements

We thank the people who helped us develop this material, and especially the students who have over the years worked through the material with us. In particular, Mark Tobenkin, William Throwe, Leo Stein, Peter Iannucci, and Micah Brodsky have suffered through bad explanations and have contributed better ones. Edmund Bertschinger, Norman Margolus, Tom Knight, Rebecca Frankel, Alexey Radul, Edwin Taylor, Joel Moses, Kenneth Yip, and Hal Abelson helped us with many thoughtful discussions and advice about physics and its relation to mathematics. We also thank Chris Hanson, Taylor Campbell, and the community of Scheme programmers for providing support and advice for the elegant language that we use. In particular, Gerald Jay Sussman wants to thank Guy Lewis Steele and Alexey Radul for many fun days of programming together—we learned much from each other's style. Matthew Halfant started us on the development of the Scmutils system. He encouraged us to get into scientific computation, using Scheme and functional style as an active way to explain the ideas, without the distractions of imperative languages such as C. In the 1980s he wrote some of the early Scheme procedures for numerical computation that we still use. Dan Zuras helped us with the invention of the unique organization of the Scmutils system. It is because of his insight that the system is organized around a generic extension of the chain rule for taking derivatives. He also helped in the heavy lifting that was required to make a really good polynomial GCD algorithm, based on ideas we learned from Richard Zippel. A special contribution that cannot be sufficiently acknowledged is from Seymour Papert and Marvin Minsky, who taught us that the practice of programming is a powerful way to develop a deeper understanding of any subject. Indeed, by the act of debugging we learn about our misconceptions, and by reflecting on our bugs and their resolutions we learn ways to learn more effectively. Indeed, *Turtle Geometry* [2], a beautiful book about discrete differential geometry at a more elementary level, was inspired by Papert's work on education. [13]

We acknowledge the generous support of the Computer Science and Artificial Intelligence Laboratory of the Massachusetts Institute of Technology. The laboratory provides a stimulating environment for efforts

to formalize knowledge with computational methods. We also acknowledge the Panasonic Corporation (formerly the Matsushita Electric Industrial Corporation) for support of Gerald Jay Sussman through an endowed chair.

Jack Wisdom thanks his wife, Cecile, for her love and support. Julie Sussman, PPA, provided careful reading and serious criticism that inspired us to reorganize and rewrite major parts of the text. She has also developed and maintained Gerald Jay Sussman over these many years.

Gerald Jay Sussman & Jack Wisdom

Cambridge, Massachusetts, USA

August 2012

## Prologue

### Programming and Understanding

One way to become aware of the precision required to unambiguously communicate a mathematical idea is to program it for a computer. Rather than using canned programs purely as an aid to visualization or numerical computation, we use computer programming in a functional style to encourage clear thinking. Programming forces us to be precise and unambiguous, without forcing us to be excessively rigorous. The computer does not tolerate vague descriptions or incomplete constructions. Thus the act of programming makes us keenly aware of our errors of reasoning or unsupported conclusions.<sup>1</sup>

Although this book is about differential geometry, we can show how thinking about programming can help in understanding in a more elementary context. The traditional use of Leibniz's notation and Newton's notation is convenient in simple situations, but in more complicated situations it can be a serious handicap to clear reasoning.

A mechanical system is described by a Lagrangian function of the system state (time, coordinates, and velocities). A motion of the system is described by a path that gives the coordinates for each moment of time. A path is allowed if and only if it satisfies the Lagrange equations. Traditionally, the Lagrange equations are written

$$\frac{d}{dt} \frac{\partial L}{\partial \dot{q}} - \frac{\partial L}{\partial q} = 0.$$

What could this expression possibly mean?

Let's try to write a program that implements Lagrange equations. What are Lagrange equations for? Our program must take a proposed path and give a result that allows us to decide if the path is allowed. This is already a problem; the equation shown above does not have a slot for a path to be tested.

So we have to figure out how to insert the path to be tested. The partial derivatives do not depend on the path; they are derivatives of the Lagrangian function and thus they are functions with the same arguments as the Lagrangian. But the time derivative  $d/dt$  makes sense only for a function of time. Thus we must be intending to substitute the path (a function of time) and its derivative (also a function of time) into the coordinate and velocity arguments of the partial derivative functions.

So probably we meant something like the following (assume that  $\omega$  is a path through the coordinate configuration space, and so  $w(t)$  specifies the configuration coordinates at time  $t$ ):

$$\frac{d}{dt} \left( \frac{\partial L(t, q, \dot{q})}{\partial \dot{q}} \Big|_{\substack{q=w(t) \\ \dot{q}=\frac{dw(t)}{dt}}} \right) - \frac{\partial L(t, q, \dot{q})}{\partial q} \Big|_{\substack{q=w(t) \\ \dot{q}=\frac{dw(t)}{dt}}} = 0.$$

In this equation we see that the partial derivatives of the Lagrangian function are taken, then the path and its derivative are substituted for the position and velocity arguments of the Lagrangian, resulting in an expression in terms of the time.

This equation is complete. It has meaning independent of the context and there is nothing left to the imagination. The earlier equations require the reader to fill in lots of detail that is implicit in the context. They do not have a clear meaning independent of the context.

By thinking computationally we have reformulated the Lagrange equations into a form that is explicit enough to specify a computation. We could convert it into a program for any symbolic manipulation program because it tells us *how* to manipulate expressions to compute the residuals of Lagrange's equations for a purported solution path.<sup>2</sup>

<sup>1</sup>The idea of using computer programming to develop skills of clear thinking was originally advocated by Seymour Papert. An extensive discussion of this idea, applied to the education of young children, can be found in Papert [13].

<sup>2</sup>The *residuals* of equations are the expressions whose value must be zero if the equations are satisfied. For example, if we know that for an unknown  $x$ ,  $x^3 - x = 0$  then the residual is  $x^3 - x$ . We can try  $x = -1$  and find a residual of 0, indicating that our purported solution satisfies the equation. A residual may provide information. For example, if we have the differential equation  $df(x)/dx - af(x) = 0$  and we plug in a test solution  $f(x) = Ae^{bx}$  we obtain the residual  $(b - a)Ae^{bx}$ , which can be zero only if  $b = a$ .

## Functional Abstraction

But this corrected use of Leibniz notation is ugly. We had to introduce extraneous symbols ( $q$  and  $\dot{q}$ ) in order to indicate the argument position specifying the partial derivative. Nothing would change here if we replaced  $q$  and  $\dot{q}$  by  $a$  and  $b$ .<sup>3</sup> We can simplify the notation by admitting that the partial derivatives of the Lagrangian are themselves new functions, and by specifying the particular partial derivative by the position of the argument that is varied

$$\frac{d}{dt} \left( (\partial_2 L) \left( t, w(t), \frac{d}{dt} w(t) \right) \right) - (\partial_1 L) \left( t, w(t), \frac{d}{dt} w(t) \right) = 0,$$

where  $\partial_i L$  is the function which is the partial derivative of the function  $L$  with respect to the  $i$ th argument.<sup>4</sup>

Two different notions of derivative appear in this expression. The functions  $\partial_2 L$   $\partial_1 L$ , constructed from the Lagrangian  $L$ , have the same arguments as  $L$ .

The derivative  $d/dt$  is an expression derivative. It applies to an expression that involves the variable  $t$  and it gives the rate of change of the value of the expression as the value of the variable  $t$  is varied.

These are both useful interpretations of the idea of a derivative. But functions give us more power. There are many equivalent ways to write expressions that compute the same value. For example  $1/(1/r_1 + 1/r_2) = (r_1 r_2)/(r_1 + r_2)$ . These expressions compute the same function of the two variables  $r_1$  and  $r_2$ . The first expression fails if  $r_1 = 0$  but the second one gives the right value of the function. If we abstract the function, say as  $\Pi(r_1, r_2)$ , we can ignore the details of how it is computed. The ideas become clearer because they do not depend on the detailed shape of the expressions.

So let's get rid of the expression derivative  $d/dt$  and replace it with an appropriate functional derivative. If  $f$  is a function then we will write  $Df$  as the new function that is the derivative of  $f$ :<sup>5</sup>

$$(Df)(t) = \frac{d}{dx} f(x)|_{x=t}.$$

To do this for the Lagrange equation we need to construct a function to take the derivative of.

Given a configuration-space path  $w$ , there is a standard way to make the state-space path. We can abstract this method as a mathematical function  $\Gamma$ :

$$\Gamma[w](t) = \left( t, w(t), \frac{d}{dt} w(t) \right).$$

Using  $\Gamma$  we can write:

$$\frac{d}{dt} ((\partial_2 L)(\Gamma[w](t))) - (\partial_1 L)(\Gamma[w](t)) = 0.$$

If we now define composition of functions  $(f \circ g)(x) = f(g(x))$ , we can express the Lagrange equations entirely in terms of functions:

$$D((\partial_2 L) \circ (\Gamma[w])) - (\partial_1 L) \circ (\Gamma[w]) = 0.$$

The functions  $\partial_1 L$  and  $\partial_2 L$  are partial derivatives of the function  $L$ . Composition with  $\Gamma[w]$  evaluates these partials with coordinates and velocities appropriate for the path  $w$ , making functions of time. Applying  $D$  takes the time derivative. The Lagrange equation states that the difference of the resulting functions of time must be zero. This statement of the Lagrange equation is complete, unambiguous, and functional. It is not encumbered with the particular choices made in expressing the Lagrangian. For example, it doesn't matter if the time is named  $t$  or  $\tau$ , and it has an explicit place for the path to be tested.

This expression is equivalent to a computer program:<sup>6</sup>

<sup>3</sup>That the symbols  $q$  and  $\dot{q}$  can be replaced by other arbitrarily chosen nonconflicting symbols without changing the meaning of the expression tells us that the partial derivative symbol is a logical quantifier, like forall and exists ( $\forall$  and  $\exists$ ).

<sup>4</sup>The argument positions of the Lagrangian are indicated by indices starting with zero for the time argument.

<sup>5</sup>An explanation of functional derivatives is in Appendix E, page 144.

<sup>6</sup>The programs in this edition are written in ClojureScript and use Emmy for generic arithmetic and symbolic mathematics. The details of the language are not germane to the points being made; what matters is that the expressions are mechanically

```
(defn Lagrange-equations                                     closure
  [Lagrangian]
  (fn [w]
    (- (D (compose ((partial 2) Lagrangian) (Gamma w)))
       (compose ((partial 1) Lagrangian) (Gamma w))))))
```

In `Lagrange-equations`, the parameter `Lagrangian` is a function implementing the Lagrangian. Its derivatives, for example `((partial 2) Lagrangian)`, are also functions. The state-space path `(Gamma w)` is constructed from the configuration-space path `w` by `Gamma`:

```
(defn Gamma [w] (fn [t] (up t (w t) ((D w) t))))          closure
```

where `up` is a constructor for a data structure that represents a state of the dynamical system (time, coordinates, velocities).

Applying `Lagrange-equations` to a Lagrangian function produces a function that accepts a configuration-space path `w` and returns a function giving the residual of the Lagrange equations at a time.

For example, consider the harmonic oscillator, with Lagrangian

$$L(t, q, v) = \frac{1}{2}mv^2 - \frac{1}{2}kq^2,$$

for mass  $m$  and spring constant  $k$ . This Lagrangian is implemented by

```
(defn L-harmonic                                           closure
  [m k]
  (fn [local]
    (let [q (coordinate local)
          v (velocity local)]
      (- (* (/ 1 2) m (square v)) (* (/ 1 2) k (square q))))))
```

We know that the motion of a harmonic oscillator is a sinusoid with a given amplitude  $a$ , frequency  $\omega$ , and phase  $\varphi$ :

$$x(t) = a \cos(\omega t + \varphi).$$

Suppose we have forgotten how the constants in the solution relate to the physical parameters of the oscillator. Let's plug in the proposed solution and look at the residual:

```
(defn proposed-solution [t] (* 'a (cos (+ (* 'omega t) 'phi))))          closure

(→tex-equation (((Lagrange-equations (L-harmonic 'm 'k)) proposed-solution) 't))
;; ⇒ "\begin{equation}n- a\\,m\\,,{\omega}^2\\,,\\cos\\left({\omega}\\,t + \\phi\\right) +
\\right) +
;; a\\,k\\,,\\cos\\left({\omega}\\,t + \\phi\\right)n\\end{equation}"
```

$$-am\omega^2 \cos(\omega t + \phi) + ak \cos(\omega t + \phi)$$

The residual here shows that for nonzero amplitude, the only solutions allowed are ones where  $(k - m\omega^2) = 0$  or  $\omega = \sqrt{k/m}$ .

But, suppose we had no idea what the solution looks like. We could propose a literal function for the path:

```
(→tex-equation (((Lagrange-equations (L-harmonic 'm 'k)) (literal-function 'x)) 't)) closure
;; ⇒ "\begin{equation}nk\\,,x\\left(t\\right) + m\\,,{D}^2x\\left(t\\right)n\\end{equation}"
```

---

interpretable and therefore unambiguous. Appendix D introduces the ClojureScript forms used here, Appendix E explains the Emmy notation, and Appendix G explains how the examples are run and checked.

$$kx(t) + mD^2x(t)$$

If this residual is zero we have the Lagrange equation for the harmonic oscillator.

Note that we can flexibly manipulate representations of mathematical functions. (See Appendices [D](#) and [E](#).)

We started out thinking that the original statement of Lagrange's equations accurately captured the idea. But we really don't know until we try to teach it to a naive student. If the student is sufficiently ignorant, but is willing to ask questions, we are led to clarify the equations in the way that we did. There is no dumber but more insistent student than a computer. A computer will absolutely refuse to accept a partial statement, with missing parameters or a type error. In fact, the original statement of Lagrange's equations contained an obvious type error: the Lagrangian is a function of multiple variables, but the  $d/dt$  is applicable only to functions of one variable.

# 1 Introduction

*Philosophy is written in that great book which ever lies before our eyes—I mean the Universe—but we cannot understand it if we do not learn the language and grasp the symbols in which it is written. This book is written in the mathematical language, and the symbols are triangles, circles, and other geometrical figures without whose help it is impossible to comprehend a single word of it, without which one wanders in vain through a dark labyrinth.*

*Galileo Galilei [8]*

Differential geometry is a mathematical language that can be used to express physical concepts. In this introduction we show a typical use of this language. Do not panic! At this point we do not expect you to understand the details of what we are showing. All will be explained as needed in the text. The purpose is to get the flavor of this material.

At the North Pole inscribe a line in the ice perpendicular to the Greenwich Meridian. Hold a stick parallel to that line and walk down the Greenwich Meridian keeping the stick parallel to itself as you walk. (The phrase “parallel to itself” is a way of saying that as you walk you keep its orientation unchanged. The stick will be aligned East-West, perpendicular to your direction of travel.) When you get to the Equator the stick will be parallel to the Equator. Turn East, and walk along the Equator, keeping the stick parallel to the Equator. Continue walking until you get to the  $90^\circ$  E meridian. When you reach the  $90^\circ$ E meridian turn North and walk back to the North Pole keeping the stick parallel to itself. Note that the stick is perpendicular to your direction of travel. When you get to the Pole note that the stick is perpendicular to the line you inscribed in the ice. But you started with that stick parallel to that line and you kept the stick pointing in the same direction on the Earth throughout your walk — how did it change orientation?

The answer is that you walked a closed loop on a curved surface. As seen in three dimensions the stick was actually turning as you walked along the Equator, because you always kept the stick parallel to the curving surface of the Earth. But as a denizen of a 2-dimensional surface, it seemed to you that you kept the stick parallel to itself as you walked, even when making a turn. Even if you had no idea that the surface of the Earth was embedded in a 3-dimensional space you could use this experiment to conclude that the Earth was not flat. This is a small example of intrinsic geometry. It shows that the idea of parallel transport is not simple. For a general surface it is necessary to explicitly define what we mean by parallel.

If you walked a smaller loop, the angle between the starting orientation and the ending orientation of the stick would be smaller. For small loops it would be proportional to the area of the loop you walked. This constant of proportionality is a measure of the curvature. The result does not depend on how fast you walked, so this is not a dynamical phenomenon.

Denizens of the surface may play ball games. The balls are constrained to the surface; otherwise they are free particles. The paths of the balls are governed by dynamical laws. This motion is a solution of the Euler-Lagrange equations<sup>1</sup> for the free-particle Lagrangian with coordinates that incorporate the constraint of living in the surface. There are coefficients of terms in the Euler-Lagrange equations that arise naturally in the description of the behavior of the stick when walking loops on the surface, connecting the static shape of the surface with the dynamical behavior of the balls. It turns out that the dynamical evolution of the balls may be viewed as parallel transport of the ball's velocity vector in the direction of the velocity vector. This motion by parallel transport of the velocity is called *geodesic motion*.

So there are deep connections between the dynamics of particles and the geometry of the space that the particles move in. If we understand this connection we can learn about dynamics by studying geometry and we can learn about geometry by studying dynamics. We enter dynamics with a Lagrangian and the associated Lagrange equations. Although this formulation exposes many important features of the system, such as how symmetries relate to conserved quantities, the geometry is not apparent. But when we express the Lagrangian and the Lagrange equations in differential geometry language, geometric properties become apparent. In the case of systems with no potential energy the Euler-Lagrange equations are equivalent to the geodesic equations on the configuration manifold. In fact, the coefficients of terms in the Lagrange

<sup>1</sup>It is customary to shorten “Euler-Lagrange equations” to “Lagrange equations.” We hope Leonhard Euler is not disturbed.



equations are Christoffel coefficients, which define parallel transport on the manifold. Let's look into this a bit.

## 1.1 Lagrange Equations

We write the Lagrange equations in functional notation<sup>2</sup> as follows:

$$D(\partial_2 L \circ \Gamma[q]) - \partial_1 L \circ \Gamma[q] = 0.$$

In SICM [19], Section 1.6.3, we showed that a Lagrangian describing the free motion of a particle subject to a coordinate-dependent constraint can be obtained by composing a free-particle Lagrangian with a function that describes how dynamical states transform given the coordinate transformation that describes the constraints.

A Lagrangian for a free particle of mass  $m$  and velocity  $v$  is just its kinetic energy,  $mv^2/2$ . The function `Lfree` implements the free Lagrangian:<sup>3</sup>

```
(defn Lfree [mass] (fn [state] (* (/ 1 2) mass (square (velocity state)))) closure
```

For us the dynamical state of a system of particles is a tuple of time, coordinates, and velocities. The free-particle Lagrangian depends only on the velocity part of the state.

For motion of a point constrained to move on the surface of a sphere the configuration space has two dimensions. We can describe the position of the point with the generalized coordinates colatitude and longitude. If the sphere is embedded in 3-dimensional space the position of the point in that space can be given by a coordinate transformation from colatitude and longitude to three rectangular coordinates.

For a sphere of radius  $R$  the function `sphere→R3` implements the transformation of coordinates from colatitude  $\theta$  and longitude  $\phi$  on the surface of the sphere to rectangular coordinates in the embedding space. (The  $\hat{z}$  axis goes through the North Pole, and the Equator is in the plane  $z = 0$ .)

```
(defn sphere→R3 closure
  [R]
  (fn [state]
    (let [q (coordinate state)]
      (let [theta (ref q 0)
            phi (ref q 1)]
        (up (* R (sin theta) (cos phi)) (* R (sin theta) (sin phi)) (* R (cos
theta)))))))
```

The coordinate transformation maps the generalized coordinates on the sphere to the 3-dimensional rectangular coordinates. Given this coordinate transformation we construct a corresponding transformation of velocities; these make up the state transformation. The function `F→C` implements the derivation of a transformation of states from a coordinate transformation:

```
(defn F→C closure
  [F]
  (fn [state]
    (up (time state)
        (F state)
        (+ (((partial 0) F) state) (* (((partial 1) F) state) (velocity state))))))
```

A Lagrangian governing free motion on a sphere of radius  $R$  is then the composition of the free Lagrangian with the transformation of states.

```
(defn Lsphere [m R] (compose (Lfree m) (F→C (sphere→R3 R)))) closure
```

So the value of the Lagrangian at an arbitrary dynamical state is:

<sup>2</sup>A short introduction to our functional notation, and why we have chosen it, is given in the prologue: Programming and Understanding. More details can be found in Appendix E

<sup>3</sup>The ClojureScript forms used in this edition are introduced in Appendix D.

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify ((Lsphere 'm 'R) (up 't (up 'theta 'phi) (up 'thetadot 'phidot))))
;; => (+ (* (/ 1 2) (expt R 2) m (expt phidot 2) (expt (sin theta) 2))
;;      (* (/ 1 2) (expt R 2) m (expt thetadot 2)))
```

or, in infix notation:

$$\frac{1}{2}R^2m\dot{\phi}^2(\sin(\theta))^2 + \frac{1}{2}R^2m\dot{\theta}^2 \quad (1.1)$$

## 1.2 The Metric

Let's now take a step into the geometry. A surface has a metric which tells us how to measure sizes and angles at every point on the surface. (Metrics are introduced in Chapter 9.)

The metric is a symmetric function of two vector fields that gives a number for every point on the manifold. (Vector fields are introduced in Chapter 3). Metrics may be used to compute the length of a vector field at each point, or alternatively to compute the inner product of two vector fields at each point. For example, the metric for the sphere of radius  $R$  is

$$g(u, v) = R^2 d\theta(u) d\theta(v) + R^2 (\sin \theta)^2 d\phi(u) d\phi(v), \quad (1.2)$$

where  $u$  and  $v$  are vector fields, and  $d\theta$  and  $d\phi$  are one-form fields that extract the named components of the vector-field argument. (One-form fields are introduced in Chapter 3.) We can think of  $d\theta(u)$  as a function of a point that gives the size of the vector field  $u$  in the  $\theta$  direction at the point. Notice that  $g(u, u)$  is a weighted sum of the squares of the components of  $u$ . In fact, if we identify

$$d\theta(v) = \dot{\theta} d\phi(v) = \dot{\phi},$$

then the coefficients in the metric are the same as the coefficients in the value of the Lagrangian, equation (1.1), apart from a factor of  $m/2$ .

We can generalize this result and write a Lagrangian for free motion of a particle of mass  $m$  on a manifold with metric  $g$ :

$$L_2(x, v) = \sum_{ij} \frac{1}{2} m g_{ij}(x) v^i v^j \quad (1.3)$$

This is written using indexed variables to indicate components of the geometric objects expressed with respect to an unspecified coordinate system. The metric coefficients  $g_{ij}$  are, in general, a function of the position coordinates  $x$ , because the properties of the space may vary from place to place.

We can capture this geometric statement as a program:

```
(defn L2 [mass metric] (fn [place velocity] (* (/ 1 2) mass ((metric velocity
velocity) place))))      clojure
```

This program gives the Lagrangian in a coordinate-independent, geometric way. It is entirely in terms of geometric objects, such as a place on the configuration manifold, the velocity at that place, and the metric that describes the local shape of the manifold. But to compute we need a coordinate system. We express the dynamical state in terms of coordinates and velocity components in the coordinate system. For each coordinate system there is a natural vector basis and the geometric velocity vectors can be constructed by contracting the basis with the components of the velocity. Thus, we can form a coordinate representation of the Lagrangian.

```
(defn Lc
  [mass metric coordsys]
  (fn [state]
    (let [x (coordinates state)
          v (velocities state)]
```

clojure

```
e (coordinate-system→vector-basis coordsys)]
((L2 mass metric) ((point coordsys) x) (* e v))))
```

The manifold point  $m$  represented by the coordinates  $x$  is given by `(define m ((point coordsys) x))`. The coordinates of  $m$  in a different coordinate system are given by `((chart coordsys2) m)`. The manifold point  $m$  is a geometric object that is the same point independent of how it is specified. Similarly, the velocity vector  $ev$  is a geometric object, even though it is specified using components  $v$  with respect to the basis  $e$ . Both  $v$  and  $e$  have as many components as the dimension of the space so their product is interpreted as a contraction.

Let's make a general metric on a 2-dimensional real manifold:<sup>4</sup>

```
(def the-metric (literal-metric 'g R2-rect))
```

closure

The metric is expressed in rectangular coordinates, so the coordinate system is `R2-rect`.<sup>5</sup> The component functions will be labeled as subscripted `gs`.

We can now make the Lagrangian for the system:

```
(def L (Lc 'm the-metric R2-rect))
```

closure

And we can apply our Lagrangian to an arbitrary state:

```
(L (up 't (up 'x 'y) (up 'vx 'vy)))
;; => (* (/ 1 2)
;;      m
;;      (+ (* (+ (* (g_00 (up x y)) vx) (* (g_01 (up x y)) vy)) vx)
;;          (* (+ (* (g_01 (up x y)) vx) (* (g_11 (up x y)) vy)) vy)))
```

closure

Compare this result with equation (1.3).

### 1.3 Euler-Lagrange Residuals

The Euler-Lagrange equations are satisfied on realizable paths. Let  $\gamma$  be a path on the manifold of configurations. (A path is a map from the 1-dimensional real line to the configuration manifold. We introduce maps between manifolds in Chapter 6.) Consider an arbitrary path:<sup>6</sup>

```
(def gamma (literal-manifold-map 'q R1-rect R2-rect))
```

closure

The values of  $\gamma$  are points on the manifold, not a coordinate representation of the points. We may evaluate `gamma` only on points of the real-line manifold; `gamma` produces points on the  $\mathbb{R}^2$  manifold. So to go from the literal real-number coordinate `'t` to a point on the real line we use `((point R1-rect) 't)` and to go from a point  $m$  in  $\mathbb{R}^2$  to its coordinate representation we use `((chart R2-rect) m)`. (The procedures `point` and `chart` are introduced in Chapter 2.) Thus

```
((chart R2-rect) (gamma ((point R1-rect) 't)))
;; => (up (q↑0 t) (q↑1 t))
```

closure

So, to work with coordinates we write:

<sup>4</sup>The function `literal-metric` provides a metric. It is a general symmetric function of two vector fields, with literal functions of the coordinates of the manifold points for its coefficients in the given coordinate system. The quoted symbol `'g` names the literal coefficient functions. Literal functions are discussed in Appendix E.

<sup>5</sup>`R2-rect` is the usual rectangular coordinate system on the 2-dimensional real manifold. (See Section 2.1, page 7.) We supply common coordinate systems for  $n$ -dimensional real manifolds. For example, `R2-polar` is a polar coordinate system on the same manifold.

<sup>6</sup>The procedure `literal-manifold-map` makes a map from the manifold implied by its second argument to the manifold implied by the third argument. These arguments must be coordinate systems. The quoted symbol that is the first argument is used to name the literal coordinate functions that define the map.

```
(def coordinate-path (compose (chart R2-rect) gamma (point R1-rect))) ;closure

(coordinate-path 't)
;; => (up (q↑0 t) (q↑1 t))
```

Now we can compute the residuals of the Euler-Lagrange equations, but we get a large messy expression that we will not show.<sup>7</sup> However, we will save it to compare with the residuals of the geodesic equations.

```
(def Lagrange-residuals (((Lagrange-equations L) coordinate-path) 't)) ;closure
```

## 1.4 Geodesic Equations

Now we get deeper into the geometry. The traditional way to write the geodesic equations is

$$\nabla_v v = 0 \quad (1.4)$$

where  $\nabla$  is a covariant derivative operator. Roughly,  $\nabla_v w$  is a directional derivative. It gives a measure of the variation of the vector field  $w$  as you walk along the manifold in the direction of  $v$ . (We will explain this in depth in Section 7.10, page 55.)  $\nabla_v v = 0$  is intended to convey that the velocity vector is parallel-transported by itself. When you walked East on the Equator you had to hold the stick so that it was parallel to the Equator. But the stick is constrained to the surface of the Earth, so moving it along the Equator required turning it in three dimensions. The  $\nabla$  thus must incorporate the 3-dimensional shape of the Earth to provide a notion of “parallel” appropriate for the denizens of the surface of the Earth. This information will appear as the “Christoffel coefficients” in the coordinate representation of the geodesic equations.

The trouble with the traditional way to write the geodesic equations (1.4) is that the arguments to the covariant derivative are vector fields and the velocity along the path is not a vector field. A more precise way of stating this relation is:

$$\nabla_{\partial/\partial t}^\gamma d\gamma(\partial/\partial t) = 0. \quad (1.5)$$

(We know that this may be unfamiliar notation, but we will explain it in Section 7.10, page 55.)

In coordinates, the geodesic equations are expressed

$$D^2 q^i(t) + \sum_{jk} \Gamma_{jk}^i(\gamma(t)) Dq^j(t) Dq^k(t) = 0, \quad (1.6)$$

where  $q(t)$  is the coordinate path corresponding to the manifold path  $\gamma$ , and  $\Gamma_{jk}^i(m)$  are Christoffel coefficients. The  $\Gamma_{jk}^i(m)$  describe the “shape” of the manifold close to the manifold point  $m$ . They can be derived from the metric  $g$ .

We can get and save the geodesic equation residuals by:

```
(def geodesic-equation-residuals ;closure
  (((((covariant-derivative Cartan gamma) d:dt) ((differential gamma) d:dt)) (chart
R2-rect))
  ((point R1-rect) 't)))
```

where  $d/dt$  is a vector field on the real line<sup>8</sup> and `Cartan` is a way of encapsulating the geometry, as specified by the Christoffel coefficients. The Christoffel coefficients are computed from the metric:

```
(def Cartan ;closure
  (Christoffel→Cartan (metric→Christoffel-2 the-metric (coordinate-system→basis R2-
rect))))
```

<sup>7</sup>For an explanation of equation residuals see page xii.

<sup>8</sup>The Emmy setup establishes the real-line coordinate objects with `(define-coordinates t R1-rect)`. The translated code uses `d:dt` for the coordinate vector field and `dt` for the one-form field.

The two messy residual results that we did not show are related by the metric. If we change the representation of the geodesic equations by “lowering” them using the mass and the metric, we see that the residuals are equal:

```
(def metric-components (metric→components the-metric (coordinate-system→basis R2-rect)))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (- Lagrange-residuals
  (* (* 'm (metric-components (gamma ((point R1-rect) 't))))
    geodesic-equation-residuals)))

;; ⇒ (down 0 0)
```

This establishes that for a 2-dimensional space the Euler-Lagrange equations are equivalent to the geodesic equations. The Christoffel coefficients that appear in the geodesic equation correspond to coefficients of terms in the Euler-Lagrange equations. This analysis will work for any number of dimensions (but will take your computer longer in higher dimensions, because the complexity increases).

#### 1.4.1 Exercise 1.1: Motion on a Sphere

The metric for a unit sphere, expressed in colatitude  $\theta$  and longitude  $\phi$ , is

$$g(u, v) = d\theta(u)d\theta(v) + (\sin \theta)^2 d\phi(u)d\phi(v).$$

Compute the Lagrange equations for motion of a free particle on the sphere and convince yourself that they describe great circles. For example, consider motion on the equator ( $\theta = \pi/2$ ) and motion on a line of longitude ( $\phi$  is constant).

## 2 Manifolds

A *manifold* is a generalization of our idea of a smooth surface embedded in Euclidean space. For an  $n$ -dimensional manifold, around every point there is a simply-connected open set, the *coordinate patch*, and a one-to-one continuous function, the *coordinate function* or *chart*, mapping every point in that open set to a tuple of  $n$  real numbers, the *coordinates*. In general, several charts are needed to label all points on a manifold. It is required that if a region is in more than one coordinate patch then the coordinates are consistent in that the function mapping one set of coordinates to another is continuous (and perhaps differentiable to some degree). A consistent system of coordinate patches and coordinate functions that covers the entire manifold is called an *atlas*.

An example of a 2-dimensional manifold is the surface of a sphere or of a coffee cup. The space of all configurations of a planar double pendulum is a more abstract example of a 2-dimensional manifold. A manifold that looks locally Euclidean may not look like Euclidean space globally: for example, it may not be simply connected. The surface of the coffee cup is not simply connected, because there is a hole in the handle for your fingers.

An example of a coordinate function is the function that maps points in a simply-connected open neighborhood of the surface of a sphere to the tuple of latitude and longitude<sup>1</sup>. If we want to talk about motion on the Earth, we can identify the space of configurations to a 2-sphere (the surface of a 3-dimensional ball). The map from the 2-sphere to the 3-dimensional coordinates of a point on the surface of the Earth captures the shape of the Earth.

Two angles specify the configuration of the planar double pendulum. The manifold of configurations is a torus, where each point on the torus corresponds to a configuration of the double pendulum. The constraints, such as the lengths of the pendulum rods, are built into the map between the generalized coordinates of points on the torus and the arrangements of masses in 3-dimensional space.

There are computational objects that we can use to model manifolds. For example, we can make an object that represents the plane<sup>2</sup>

```
(def R2 (make-manifold Rn 2))
```

closure

and give it the name `R2`. One useful patch of the plane is the one that contains the origin and covers the entire plane<sup>3</sup>.

```
(def U (patch 'origin R2))
```

closure

### 2.1 Coordinate Functions

A coordinate function  $\chi$  maps points in a coordinate patch of a manifold to a coordinate tuple<sup>4</sup>:

$$x = \chi(m), \quad (2.1)$$

where  $x$  may have a convenient tuple structure. Usually, the coordinates are arranged as an “up structure”; the coordinates are selected with superscripts:

$$x^i = \chi^i(m). \quad (2.2)$$

The number of independent components of  $x$  is the dimension of the manifold.


Assume we have two coordinate functions  $\chi$  and  $\chi'$ . The coordinate transformation from  $\chi'$  coordinates to  $\chi$  coordinates is just the composition  $\chi \circ \chi'^{-1}$ , where  $\chi'^{-1}$  is the functional inverse of  $\chi'$  (see figure 2.1).

<sup>1</sup>The open set for a latitude-longitude coordinate system cannot include either pole (because longitude is not defined at the poles) or the 180° meridian (where the longitude is discontinuous). Other coordinate systems are needed to cover these places.

<sup>2</sup>The expression `R^n` gives only one kind of manifold. We also have spheres `S^n` and `S03`.

<sup>3</sup>The word `origin` is an arbitrary symbol here. It labels a predefined patch in `R^n` manifolds.

<sup>4</sup>In the text that follows we will use sans-serif names, such as  $f$ ,  $v$ ,  $m$ , to refer to objects defined on the manifold. Objects that are defined on coordinates (tuples of real numbers) will be named with symbols like  $f$ ,  $v$ ,  $x$ .



**Figure 2.1:** Here there are two overlapping coordinate patches that are the domains of the two coordinate functions  $\chi$  and  $\chi'$ . It is possible to represent manifold points in the overlap using either coordinate system.

The coordinate transformation from  $\chi'$  coordinates to  $\chi$  coordinates is just the composition  $\chi \circ \chi'^{-1}$ .

We assume that the coordinate transformation is continuous and differentiable to any degree we require.

Given a coordinate system `coordsys` for a patch on a manifold the procedure that implements the function  $\chi$  that gives coordinates for a point is `(chart coordsys)`. The procedure that implements the inverse map that gives a point for coordinates is `(point coordsys)`.

We can have both rectangular and polar coordinates on a patch of the plane identified by the origin:<sup>56</sup>

```
;; Some charts on the patch U
(def R2-rect (coordinate-system 'rectangular U))

(def R2-polar (coordinate-system 'polar/cylindrical U))
```

For each of the coordinate systems above we obtain the coordinate functions and their inverses:

```
(def R2-rect-chi (chart R2-rect))

(def R2-rect-chi-inverse (point R2-rect))

(def R2-polar-chi (chart R2-polar))

(def R2-polar-chi-inverse (point R2-polar))
```

The coordinate transformations are then just compositions. The polar coordinates of a rectangular point are:

```
((compose R2-polar-chi R2-rect-chi-inverse) (up 'x0 'y0))
;; => (up (sqrt (+ (expt x0 2) (expt y0 2))) (atan y0 x0))
```

And the rectangular coordinates of a polar point are:

<sup>56</sup>The rectangular coordinates are good for the entire plane, but the polar coordinates are singular at the origin because the angle is not defined. Also, the patch for polar coordinates must exclude one ray from the origin, because of the angle variable.

<sup>6</sup>We can avoid explicitly naming the patch:

```
(def R2-rect (coordinate-system-at R2 :rectangular :origin))
```

```
((compose R2-rect-chi R2-polar-chi-inverse) (up 'r0 'theta0))
;; => (up (* r0 (cos theta0)) (* r0 (sin theta0)))
```


And we can obtain the Jacobian of the polar-to-rectangular transformation by taking its derivative<sup>7</sup>:

```
((D (compose R2-rect-chi R2-polar-chi-inverse)) (up 'r0 'theta0))
;; => (down (up (cos theta0) (sin theta0))
;;          (up (* r0 (- (sin theta0)) (* r0 (cos theta0))))
```

## 2.2 Manifold functions

Let  $f$  be a real-valued function on a manifold  $M$ : this function maps points  $m$  on the manifold to real numbers.

This function has a coordinate representation  $f_\chi$  with respect to the coordinate function  $\chi$  (see figure 2.2):



**Figure 2.2:** The coordinate function  $\chi$  maps points on the manifold in the coordinate patch to a tuple of coordinates. A function  $f$  on the manifold  $M$  can be represented in coordinates by a function  $f_\chi = f \circ \chi^{-1}$ .

$$f_\chi = f \circ \chi^{-1}. \quad (2.3)$$

Both the coordinate representation  $f_\chi$  and the tuple  $x$  depend on the coordinate system, but the value  $f_\chi(x)$  is independent of coordinates:

$$f_\chi(x) = (f \circ \chi^{-1})(\chi(m)) = f(m). \quad (2.4)$$

The subscript  $\chi$  may be dropped when it is unambiguous.

For example, in a 2-dimensional real manifold the coordinates of a manifold point  $m$  are a pair of real numbers,

$$(x, y) = \chi(m), \quad (2.5)$$

and the manifold function  $f$  is represented in coordinates by a function  $f$  that takes a pair of real numbers and produces a real number

$$f : R^2 \rightarrow R \quad f : (x, y) \rightarrow f(x, y). \quad (2.6)$$

We define our manifold function

<sup>7</sup>See Appendix E for an introduction to Emmy structures and a discussion of derivatives of functions with structured input or output.



$$f : M \rightarrow R \quad f : m \rightarrow (f \circ \chi)(m). \quad (2.7)$$

### 2.3 Manifold Functions Are Coordinate Independent

We can illustrate the coordinate independence with a program. We will show that an arbitrary manifold function  $f$ , when defined by its coordinate representation in rectangular coordinates, has the same behavior when applied to a manifold point independent of whether the point is specified in rectangular or polar coordinates.

We define a manifold function by specifying its behavior in rectangular coordinates<sup>8</sup>:

```
(def f (compose (literal-function 'f-rect R2→R) R2-rect-chi))
```

closure

where  $R2 \rightarrow R$  is a signature for functions that map an up structure of two reals to a real:

```
(def R2→R '(<math>\rightarrow</math> (UP Real Real) Real))
```

closure

We can specify a typical manifold point using its rectangular coordinates:

```
(def R2-rect-point (R2-rect-chi-inverse (up 'x0 'y0)))
```

closure

We can describe the *same point* using its polar coordinates:

```
(def corresponding-polar-point
  (R2-polar-chi-inverse (up (sqrt (+ (square 'x0) (square 'y0))) (atan 'y0 'x0))))
```

closure

`(f R2-rect-point)` and `(f corresponding-polar-point)` agree, even though the point has been specified in two different coordinate systems:

```
(f R2-rect-point)
;; => (f-rect (up x0 y0))

(f corresponding-polar-point)
;; => (f-rect (up x0 y0))
```

closure

### 2.4 Naming Coordinate Functions

To make things a bit easier, we can give names to the individual coordinate functions associated with a coordinate system. Here we name the coordinate functions for the `R2-rect` coordinate system `x` and `y` and for the `R2-polar` coordinate system `r` and `theta`.

```
(define-coordinates (up x y) R2-rect)

(define-coordinates (up r theta) R2-polar)
```

closure

This allows us to extract the coordinates from a point, independent of the coordinate system used to specify the point.

```
(x (R2-rect-chi-inverse (up 'x0 'y0)))
;; => x0

(x (R2-polar-chi-inverse (up 'r0 'theta0)))
;; => (* r0 (cos theta0))

(r (R2-polar-chi-inverse (up 'r0 'theta0)))
```

closure

<sup>8</sup>Alternatively, we can define the same function in a shorthand

```
(def f (literal-manifold-function 'f-rect R2-rect))
```

closure

```
;; => rθ

(r (R2-rect-chi-inverse (up 'x0 'y0)))
;; => (sqrt (+ (expt x0 2) (expt y0 2)))

(theta (R2-rect-chi-inverse (up 'x0 'y0)))
;; => (atan y0 x0)
```

We can work with the coordinate functions in a natural manner, defining new manifold functions in terms of them<sup>9</sup>:

```
(def h (+ (* x (square r)) (cube y))) closure

(h R2-rect-point)
;; => (+ (* x0 (+ (expt x0 2) (expt y0 2))) (expt y0 3))
```

We can also apply `h` to a point defined in terms of its polar coordinates:

```
(h (R2-polar-chi-inverse (up 'r0 'theta0))) closure
;; => (+ (* r0 (cos theta0) (expt r0 2)) (expt (* r0 (sin theta0)) 3))
```

## 2.5 Exercise 2.1: Curves

A curve may be specified in different coordinate systems. For example, a cardioid constructed by rolling a circle of radius  $a$  around another circle of the same radius is described in polar coordinates by the equation

$$r = 2a(1 + \cos(\theta)).$$

We can convert this to rectangular coordinates by evaluating the residual in rectangular coordinates.

```
(define-coordinates (up r theta) R2-polar) closure

((- r (* 2 'a (+ 1 (cos theta)))) ((point R2-rect) (up 'x 'y)))
;; => (- (sqrt (+ (expt x 2) (expt y 2))) (* 2 a (+ 1 (cos (atan y x)))))
```

The numerator of this expression is the equivalent residual in rectangular coordinates. If we rearrange terms and square it we get the traditional formula for the cardioid

$$(x^2 + y^2 - 2ax)^2 = 4a^2(x^2 + y^2).$$

**a.** The rectangular coordinate equation for the Lemniscate of Bernoulli is

$$(x^2 + y^2)^2 = 2a^2(x^2 - y^2).$$


Find the expression in polar coordinates.

**b.** Describe a helix space curve in both rectangular and cylindrical coordinates. Use the computer to show the correspondence. Note that we provide a cylindrical coordinate system on the manifold **R3** for you to use. It is called `R3-cyl`; with coordinates `(r, theta, z)`.

## 2.6 Exercise 2.2: Stereographic Projection

A stereographic projection is a correspondence between points on the unit sphere and points on the plane cutting the sphere at its equator. (See figure 2.3.)

<sup>9</sup>This is actually a nasty, but traditional, abuse of notation. An expression like  $\cos(r)$  can either mean the cosine of the angle  $r$  (if  $r$  is a number), or the composition  $\cos \circ r$  (if  $r$  is a function). In our system `(cos r)` behaves in this way—either computing the cosine of `r` or being treated as `(compose cos r)` depending on what `r` is.



**Figure 2.3:** For each point on the sphere (except for its north pole) a line is drawn from the north pole through the point and extending to the equatorial plane. The corresponding point on the plane is where the line intersects the plane. The rectangular coordinates of this point on the plane are the Riemann coordinates of the point on the sphere. The points on the plane can also be specified with polar coordinates  $(\rho, \theta)$  and the points on the sphere are specified both by Riemann coordinates and the traditional colatitude and longitude  $(\varphi, \lambda)$ .

The coordinate system for points on the sphere in terms of rectangular coordinates of corresponding points on the plane is `S2-Riemann`<sup>10</sup>. The procedure `(chart S2-Riemann)` gives the rectangular coordinates on the plane for every point on the sphere, except for the North Pole. The procedure `(point S2-Riemann)` gives the point on the sphere given rectangular coordinates on the plane. The usual spherical coordinate system on the sphere is `S2-spherical`.

We can compute the colatitude and longitude of a point on the sphere corresponding to a point on the plane with the following incantation:

```
((compose (chart S2-spherical) (point S2-Riemann) (chart R2-rect) (point R2-polar))) closure
  (up 'rho 'theta))
;; => (up (acos (/ (+ (expt rho 2) -1) (+ (expt rho 2) 1))) theta)
```

Perform an analogous computation to get the polar coordinates of the point on the plane corresponding to a point on the sphere given by its colatitude and longitude.

<sup>10</sup>The plane with the addition of a point at infinity is conformally equivalent to the sphere by this correspondence. This correspondence is called the Riemann sphere, in honor of the great mathematician Bernard Riemann (1826–1866), who made major contributions to geometry.

### 3 Vector Fields and One-Form Fields

We want a way to think about how a function varies on a manifold. Suppose we have some complex linkage, such as a multiple pendulum. The potential energy is an important function on the multi-dimensional configuration manifold of the linkage. To understand the dynamics of the linkage we need to know how the potential energy changes as the configuration changes. The change in potential energy for a step of a certain size in a particular direction in the configuration space is a real physical quantity; it does not depend on how we measure the direction or the step size. What exactly this means is to be determined: What is a step size? What is a direction? We cannot subtract two configurations to determine the distance between them. It is our job here to make sense of this idea.

So we would like something like a derivative, but there are problems. Since we cannot subtract two manifold points, we cannot take the derivative of a manifold function in the way described in elementary calculus. But we can take the derivative of a coordinate representation of a manifold function, because it takes real-number coordinates as its arguments. This is a start, but it is not independent of coordinate system. Let's see what we can build out of this.

#### 3.1 Vector Fields

In multiple dimensions the derivative of a function is the multiplier for the best linear approximation of the function at each argument point:<sup>1</sup>

$$f(x + \Delta x) \approx f(x) + (Df(x))\Delta x \quad (3.1)$$

The derivative  $Df(x)$  is independent of  $\Delta x$ . Although the derivative depends on the coordinates, the product  $(Df(x))\Delta x$  is invariant under change of coordinates in the following sense. Let  $\phi = \chi \circ \chi'^{-1}$  be a coordinate transformation, and  $x = \phi(y)$ . Then  $\Delta x = D\phi(y)\Delta y$  is the linear approximation to the change in  $x$  when  $y$  changes by  $\Delta y$ . If  $f$  and  $g$  are the representations of a manifold function in the two coordinate systems,  $g(y) = f(\phi(y)) = f(x)$ , then the linear approximations to the increments in  $f$  and  $g$  are equal:

$$Dg(y)\Delta y = Df(\phi(y))(D\phi(y)\Delta y) = Df(x)\Delta x. \quad (3.2)$$

The invariant product  $(Df(x))\Delta x$  is the *directional derivative* of  $f$  at  $x$  with respect to the vector specified by the tuple of components  $\Delta x$  in the coordinate system. We can generalize this idea to allow the vector at each point to depend on the point, making a *vector field*. Let  $b$  be a function of coordinates. We then have a directional derivative of  $f$  at each point  $x$ , determined by  $b$

$$D_b(f)(x) = (Df(x))b(x). \quad (3.3)$$

Now we bring this back to the manifold and develop a useful generalization of the idea of directional derivative for functions on a manifold, rather than functions on  $\mathbb{R}^n$ . A *vector field* on a manifold is an assignment of a vector to each point on the manifold. In elementary geometry, a vector is an arrow anchored at a point on the manifold with a magnitude and a direction. In differential geometry, a vector is an operator that takes directional derivatives of manifold functions at its anchor point. The direction and magnitude of the vector are the direction and scale factor of the directional derivative.

Let  $m$  be a point on a manifold,  $v$  be a vector field on the manifold, and  $f$  be a real-valued function on the manifold. Then  $v(f)$  is the directional derivative of the function  $f$  and  $v(f)(m)$  is the directional derivative of the function  $f$  at the point  $m$ . The vector field is an operator that takes a real-valued manifold function and a manifold point and produces a number. The order of arguments is chosen to make  $v(f)$  be a new manifold function that can be manipulated further. Directional derivative operators, unlike ordinary derivative operators, produce a result of the same type as their argument. Note that there is no mention here of any coordinate system. The vector field specifies a direction and magnitude at each manifold point that is independent of how it is described using any coordinate system.

A useful way to characterize a vector field in a particular coordinate system is by applying it to the coordinate functions. The resulting functions  $b_{\chi, v}^i$  are called the *coordinate component functions* or *coefficient functions* of the vector field; they measure how quickly the coordinate functions change in the direction of the vector field, scaled by the magnitude of the vector field:

<sup>1</sup>In multiple dimensions the derivative  $Df(x)$  is a down tuple structure of the partial derivatives and the increment  $\Delta x$  is an up tuple structure, so the indicated product is to be interpreted as a contraction. (See equation (B.8).)

$$b_{\chi,v}^i = v(\chi^i) \circ \chi^{-1}. \quad (3.4)$$

Note that we have chosen the coordinate components to be functions of the coordinate tuple, not of a manifold point.

A vector with coordinate components  $b_{\chi,v}$  applies to a manifold function  $f$  via

$$\begin{aligned} v(f)(m) &= ((D(f \circ \chi^{-1})b_{\chi,v}) \circ \chi)(m) \\ &= D(f \circ \chi^{-1})(\chi(m))b_{\chi,v}(\chi(m)) \\ &= \sum_i \partial_i(f \circ \chi^{-1})(\chi(m))b_{\chi,v}^i(\chi(m)). \end{aligned} \quad (3.5)$$

In equation (3.4), the quantity  $f \circ$  is the coordinate representation of the manifold function  $f$ . We take its derivative, and weight the components of the derivative with the coordinate components  $b_{\chi,v}$  of the vector field that specify its direction and magnitude. Since this product is a function of coordinates we use  $\chi$  to extract the coordinates from the manifold point  $m$ . In equation (3.5), the composition of the product with the coordinate chart  $\chi$  is replaced by function evaluation. In equation (3.6) the tuple multiplication is expressed explicitly as a sum of products of corresponding components. So the application of the vector is a linear combination of the partial derivatives of  $f$  in the coordinate directions weighted by the vector components. This computes the rate of change of  $f$  in the direction specified by the vector.

Equations (3.3) and (3.5) are consistent:

$$\begin{aligned} v(x)(\chi^{-1}(x)) &= D(\chi \circ \chi^{-1})(x)b_{\chi,v}(x) \\ &= D(I)(x)b_{\chi,v}(x) \\ &= b_{\chi,v}(x). \end{aligned} \quad (3.6)$$

The coefficient tuple  $b_{\chi,v}(x)$  is an up structure compatible for addition to the coordinates. Note that for any vector field  $v$  the coefficients  $b_{\chi,v}(x)$  are different for different coordinate functions  $\chi$ . In the text that follows we will usually drop the subscripts on  $b$ , understanding that it is dependent on the coordinate system and the vector field.

We implement the definition of a vector field (3.4) as:

```
(defn components→vector-field
  [components coordsys]
  (letfn [(v [f] (compose (* (D (compose f (point coordsys))) components) (chart
    coordsys)))]
    (procedure→vector-field v)))
```

The vector field is an operator, like derivative.<sup>2</sup>

Given a coordinate system and coefficient functions that map coordinates to real values, we can make a vector field. For example, a general vector field can be defined by giving components relative to the coordinate system `R2-rect` by

```
(def v
  (components→vector-field (up (literal-function 'b^0 R2→R) (literal-function 'b^1
    R2→R))
    R2-rect))
```

To make it convenient to define literal vector fields we provide a shorthand: `(define v (literal-vector-field 'b R2-rect))`. This makes a vector field with component functions named `b^0` and `b^1` and names the result `v`. When this vector field is applied to an arbitrary manifold function it gives the directional derivative of that manifold function in the direction specified by the components `b^0` and `b^1`:

<sup>2</sup>An operator is just like a procedure except that multiplication is interpreted as composition. For example, the derivative procedure is made into an operator `D` so that we can say `(expt D 2)` and expect it to compute the second derivative. The procedure `procedure→vector-field` makes a vector-field operator.

```
((v (literal-manifold-function 'f-rect R2-rect)) R2-rect-point)
;; => (+ (* ((partial 0) f-rect) (up x0 y0)) (b↑0 (up x0 y0)))
;;      (* ((partial 1) f-rect) (up x0 y0)) (b↑1 (up x0 y0)))
```

This result is what we expect from equation (3.6).

We can recover the coordinate components of the vector field by applying the vector field to the coordinate chart:

```
((v (chart R2-rect)) R2-rect-point)
;; => (up (b↑0 (up x0 y0)) (b↑1 (up x0 y0)))
```

### 3.1.1 Coordinate Representation

The vector field  $\mathbf{v}$  has a coordinate representation  $v$ :

$$\begin{aligned} \mathbf{v}(f)(m) &= D(f \circ \chi^{-1})(\chi(m))b(\chi(m)) \\ &= Df(x)b(x) \\ &= v(f)(x), \end{aligned} \tag{3.7}$$

with the definitions  $f = f \circ \chi^{-1}$  and  $x = \chi(m)$ . The function  $b$  is the coefficient function for the vector field  $\mathbf{v}$ . It provides a scale factor for the component in each coordinate direction. However,  $v$  is the coordinate representation of the vector field  $\mathbf{v}$  in that it takes directional derivatives of coordinate representations of manifold functions.

Given a vector field  $\mathbf{v}$  and a coordinate system `coordsys` we can construct the coordinate representation of the vector field.<sup>3</sup>

```
(defn coordinatize
  [v coordsys]
  (letfn [(coordinatized-v [f]
            (fn [x] (let [b (compose (v (chart coordsys)) (point coordsys))] (* ((D f)
x) (b x)))))]
    (make-operator coordinatized-v)))
```

We can apply a coordinatized vector field to a function of coordinates to get the same answer as before.

```
((coordinatize v R2-rect) (literal-function 'f-rect R2→R)) (up 'x0 'y0))
;; => (+ (* ((partial 0) f-rect) (up x0 y0)) (b↑0 (up x0 y0)))
;;      (* ((partial 1) f-rect) (up x0 y0)) (b↑1 (up x0 y0)))
```

### 3.1.2 Vector Field Properties

The vector fields on a manifold form a vector space over the field of real numbers and a module over the ring of real-valued manifold functions. A module is like a vector space except that there is no multiplicative inverse operation on the scalars of a module. Manifold functions that are not the zero function do not necessarily have multiplicative inverses, because they can have isolated zeros. So the manifold functions form a ring, not a field, and vector fields must be a module over the ring of manifold functions rather than a vector space.

Vector fields have the following properties. Let  $\mathbf{u}$  and  $\mathbf{v}$  be vector fields and let  $\alpha$  be a real-valued manifold function. Then

$$(\mathbf{u} + \mathbf{v})(f) = \mathbf{u}(f) + \mathbf{v}(f) \quad (\alpha f)(f) = \alpha(\mathbf{u}(f)). \tag{3.8}$$

Vector fields are linear operators. Assume  $f$  and  $g$  are functions on the manifold,  $a$  and  $b$  are real constants.<sup>4</sup> The constants  $a$  and  $b$  are not manifold functions, because vector fields take derivatives. See equation (3.13).

$$\mathbf{v}(af + bg)(m) = a\mathbf{v}(f)(m) + b\mathbf{v}(g)(m) \quad \mathbf{v}(af)(m) = a\mathbf{v}(f)(m) \tag{3.9}$$

<sup>3</sup>The `make-operator` procedure takes a procedure and returns an operator.

<sup>4</sup>If  $f$  has structured output then  $\mathbf{v}(f)$  is the structure resulting from  $\mathbf{v}$  being applied to each component of  $f$ .

Vector fields satisfy the product rule (Leibniz rule).

$$v(fg)(m) = v(f)(m)g(m) + f(m)v(g)(m) \quad (3.10)$$

Vector fields satisfy the chain rule. Let  $F$  be a function on the range of  $f$ .

$$v(F \circ f)(m) = DF(f(m))v(f)(m) \quad (3.11)$$

### 3.2 Coordinate-Basis Vector Fields

For an  $n$ -dimensional manifold any set of  $n$  linearly independent vector fields<sup>5</sup> form a *basis* in that any vector field can be expressed as a linear combination of the basis fields with manifold-function coefficients. Given a coordinate system we can construct a basis as follows: we choose the component tuple  $b_i(x)$  (see equation (3.5)) to be the  $i$ th unit tuple  $u_i(x)$ —an up tuple with one in the  $i$ th position and zeros in all other positions—selecting the partial derivative in that direction. Here  $u_i$  is a constant function. Like  $b$ , it formally takes coordinates of a point as an argument, but it ignores them. We then define the basis vector field  $X_i$  by

$$X_i(f)(m) = D(f \circ \chi^{-1})(\chi(m))u_i(\chi(m)) = \partial_i(f \circ \chi^{-1})(\chi(m)). \quad (3.12)$$

In terms of  $X_i$  the vector field of equation (3.6) is

$$v(f)(m) = \sum_i X_i(f)(m)b^i(\chi(m)). \quad (3.13)$$

We can also write

$$v(f)(m) = X(f)(m)b(\chi(m)), \quad (3.14)$$

letting the tuple algebra do its job.

The basis vector field is often written

$$\frac{\partial}{\partial x^i} = X_i, \quad (3.15)$$

to call to mind that it is an operator that computes the directional derivative in the  $i$ th coordinate direction.

In addition to making the coordinate functions, the procedure `define-coordinates` also makes the traditional named basis vectors. Using these we can examine the application of a rectangular basis vector to a polar coordinate function:

```
(define-coordinates (up x y) R2-rect) closure

(define-coordinates (up r theta) R2-polar)

((d:dx (square r)) R2-rect-point)
;; => (* 2
;;      (sqrt (+ (expt x0 2) (expt y0 2)))
;;      (/ 1 (* (sqrt (+ (expt x0 2) (expt y0 2))) 2))
;;      2
;;      x0)
```

More general functions and vectors can be made as combinations of these simple pieces:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify (((+ d:dx (* 2 d:dy)) (+ (square r) (* 3 x))) R2-rect-point))
;; => (+ (* 2 x0) (* 4 y0) 3)
```

<sup>5</sup>A set of vector fields,  $\{v_i\}$ , is linearly independent with respect to manifold functions if we cannot find nonzero manifold functions,  $\{a_i\}$ , such that

$$\sum_i a_i v_i(f) = 0(f),$$

where  $0$  is the vector field such that  $0(f)(m) = 0$  for all  $f$  and  $m$ .

### 3.2.1 Coordinate Transformations

Consider a coordinate change from the chart  $\chi$  to the chart  $\chi'$ .

$$\begin{aligned} X(f)(m) &= D(f \circ \chi^{-1})(\chi(m)) \\ &= D(f \circ (\chi')^{-1} \circ \chi' \circ \chi^{-1})(\chi(m)) \\ &= D(f \circ (\chi')^{-1})(\chi'(m))(D(\chi' \circ \chi^{-1})(\chi(m))) \\ &= X'(f)(m)(D(\chi' \circ \chi^{-1})(\chi(m))). \end{aligned} \quad (3.16)$$

This is the rule for the transformation of basis vector fields. The second factor can be recognized as “ $\partial x' / \partial x$ ,” the Jacobian.<sup>6</sup>

The vector field does not depend on coordinates. So, from equation (3.17), we have

$$v(f)(m) = X(f)(m)b(\chi(m)) = X'(f)(m)b'(\chi'(m)). \quad (3.17)$$

Using equation (3.19) with  $x = \chi(m)$  and  $x' = \chi'(m)$ , we deduce

$$D(\chi' \circ \chi^{-1})(x)b(x) = b'(x'). \quad (3.18)$$

Because  $\chi' \circ \chi^{-1}$  is the inverse function of  $\chi \circ (\chi')^{-1}$ , their derivatives are multiplicative inverses,

$$D(\chi' \circ \chi^{-1})(x) = \left( D(\chi \circ (\chi')^{-1})(x') \right)^{-1}, \quad (3.19)$$

and so

$$b(x) = D(\chi \circ (\chi')^{-1})(x')b'(x'), \quad (3.20)$$

as expected.<sup>7</sup>

It is traditional to express this rule by saying that the basis elements transform *covariantly* and the coefficients of a vector in terms of a basis transform contravariantly; their product is invariant under the transformation.

### 3.3 Integral Curves

A vector field gives a direction and rate for every point on a manifold. We can start at any point and go in the direction specified by the vector field, tracing out a parametric curve on the manifold. This curve is an *integral curve* of the vector field.

More formally, let  $v$  be a vector field on the manifold  $M$ . An integral curve  $\gamma_m^v : R \rightarrow M$  of  $v$  is a parametric path on  $M$  satisfying

$$D(f \circ \gamma_m^v)(t) = v(f)(\gamma_m^v(t)) = (v(f) \circ \gamma_m^v)(t)\gamma_m^v(0) = m, \quad (3.21)$$

for arbitrary functions  $f$  on the manifold, with real values or structured real values. The rate of change of a function along an integral curve is the vector field applied to the function evaluated at the appropriate place along the curve. Often we will simply write  $\gamma$ , rather than  $\gamma_m^v$ . Another useful variation is  $\phi_t^v(m) = \gamma_m^v(t)$ .

We can recover the differential equations satisfied by a coordinate representation of the integral curve by letting  $f = \chi$ , the coordinate function, and letting  $\sigma = \chi \circ \gamma$  be the coordinate path corresponding to the curve  $\gamma$ . Then the derivative of the coordinate path  $\sigma$  is

<sup>6</sup>This notation helps one remember the transformation rule:

$$\frac{\partial f}{\partial x^i} = \sum_j \frac{\partial f}{\partial x'^j} \frac{\partial x'^j}{\partial x^i},$$

which is the relation in the usual Leibniz notation. As Spivak pointed out in *Calculus on Manifolds*, p.45,  $f$  means something different on each side of the equation.

<sup>7</sup>For coordinate paths  $q$  and  $q'$  related by  $q(t) = (\chi \circ (\chi')^{-1})(q'(t))$  the velocities are related by  $Dq(t) = D(\chi \circ (\chi')^{-1})(q'(t))Dq'(t)$ . Abstracting off paths, we get  $v = D(\chi \circ (\chi')^{-1})(x')v'$ .



$$\begin{aligned}
D\sigma(t) &= D(\chi \circ \gamma)(t) \\
&= (\mathbf{v}(\chi) \circ \gamma)(t) \\
&= (\mathbf{v}(\chi) \circ \chi^{-1} \circ \chi \circ \gamma)(t) \\
&= (\mathbf{b} \circ \sigma)(t)
\end{aligned} \tag{3.22}$$

where  $\mathbf{b} = \mathbf{v}(\chi) \circ \chi^{-1}$  is the coefficient function for the vector field  $\mathbf{v}$  for coordinates  $\chi$  (see equation (3.7)). So the coordinate path  $\sigma$  satisfies the differential equations

$$D\sigma = \mathbf{b} \circ \sigma. \tag{3.23}$$

Differential equations for the integral curve can be expressed only in a coordinate representation, because we cannot go from one point on the manifold to another by addition of an increment. However, we can do this by adding the coordinates to an increment of coordinates and then finding the corresponding point on the manifold.

Iterating the process described by equation (3.24) we can compute higher-order derivatives of functions along the integral curve:

$$\begin{aligned}
D(f \circ \gamma) &= \mathbf{v}(f) \circ \gamma \\
D^2(f \circ \gamma) &= D(\mathbf{v}(f) \circ \gamma) = \mathbf{v}(\mathbf{v}(f)) \circ \gamma \\
&\dots \\
D^n(f \circ \gamma) &= \mathbf{v}^n(f) \circ \gamma
\end{aligned} \tag{3.24}$$

Thus, the evolution of  $f \circ \gamma$  can be written formally as a Taylor series in the parameter:

$$\begin{aligned}
(f \circ \gamma)(t) &= (f \circ \gamma)(0) + tD(f \circ \gamma)(0) + \frac{1}{2}t^2D^2(f \circ \gamma)(0) + \dots \\
&= (e^{tD}(f \circ \gamma))(0) \\
&= e^{t\mathbf{v}f}(\gamma(0)).
\end{aligned} \tag{3.25}$$

Using  $\phi$  rather than  $\gamma$

$$(f \circ \gamma_m^{\mathbf{v}})(t) = (f \circ \phi_t^{\mathbf{v}})(\mathbf{m}), \tag{3.26}$$

so, when the series converges,

$$(e^{t\mathbf{v}}f)(\mathbf{m}) = (f \circ \phi_t^{\mathbf{v}})(\mathbf{m}). \tag{3.27}$$

In particular, let  $f = \chi$ , then

$$\sigma(t) = (\chi \circ \gamma)(t) = (e^{tD}(\chi \circ \gamma))(0) = (e^{t\mathbf{v}}\chi)(\gamma(0)), \tag{3.28}$$

a Taylor series representation of the solution to the differential equation (3.27).

For example, a vector field circular that generates a rotation about the origin is:<sup>8</sup>

```
(def circular (- (* x d:dy) (* y d:dx))) closure
```

We can exponentiate the circular vector field, to generate an evolution in a circle around the origin starting at  $(1, 0)$ :

```
(series-for-each print-expression
  (((exp (* 't circular)) (chart R2-rect)) ((point R2-rect) (up 1 0)))
6) closure
```

These are the first six terms of the series expansion of the coordinates of the position for parameter  $t$ .

---

<sup>8</sup>In this expression  $d/dx$  and  $d/dy$  are vector fields that take directional derivatives of manifold functions and evaluate them at manifold points;  $x$  and  $y$  are manifold functions. `define-coordinates` was used to create these operators and functions, see page 10.

Note that `circular` is an operator—a property inherited from  $d/dx$  and  $d/dy$ .

We can define an evolution operator  $E_{\Delta t, \mathbf{v}}$  using equation (3.31)

$$(E_{\Delta t, \mathbf{v}} \mathbf{f})(\mathbf{m}) = (e^{\Delta t \mathbf{v}} \mathbf{f})(\mathbf{m}) = (\mathbf{f} \circ \phi_{\Delta t}^{\mathbf{v}})(\mathbf{m}). \quad (3.29)$$

We can approximate the evolution operator by summing the series up to a given order:

```
(defn evolution                                     clojure
  [order]
  (fn [delta-t v] (fn [f] (fn [m] (series:sum (((exp (* delta-t v)) f) m) order))))))
```

We can evolve circular from the initial point up to the parameter  $t$ , and accumulate the first six terms as follows:

```
((((evolution 6) 'delta-t circular) (chart R2-rect)) ((point R2-rect) (up 1 0)))      clojure
;; => (up (+ 1
;;       (* (/ 1 2) delta-t delta-t -1)
;;       (* (/ 1 24) delta-t delta-t -1 delta-t delta-t -1)
;;       (* (/ 1 720) delta-t delta-t -1 delta-t delta-t -1 delta-t delta-t -1))
;;       (+ delta-t
;;       (* (/ 1 6) delta-t delta-t -1 delta-t)
;;       (* (/ 1 120) delta-t delta-t -1 delta-t delta-t -1 delta-t)))
```

Note that these are just the series for  $\cos \Delta t$  and  $\sin \Delta t$ , so the coordinate tuple of the evolved point is  $(\cos \Delta t, \sin \Delta t)$ .

For functions whose series expansions have finite radius of convergence, evolution can progress beyond the point at which the Taylor series converges because evolution is well defined whenever the integral curve is defined.

### Exercise 3.1: State Derivatives

Newton's equations for the motion of a particle in a plane, subject to a force that depends only on the position in the plane, are a system of second-order differential equations for the rectangular coordinates  $(X, Y)$  of the particle:

$$D^2 X(t) = A_x(X(t), Y(t)) \text{ and } D^2 Y(t) = A_y(X(t), Y(t)), \quad (3.30)$$

where  $A$  is the acceleration of the particle.

These are equivalent to a system of first-order equations for the coordinate path  $\sigma = \chi \circ \gamma$ , where  $\chi = (t, x, y, v_x, v_y)$  is a coordinate system on the manifold  $\mathbb{R}^5$ . Then our equations are:

$$\begin{aligned} D(t \circ \gamma) &= 1 \\ D(x \circ \gamma) &= v_x \circ \gamma \\ D(y \circ \gamma) &= v_y \circ \gamma \\ D(v_x \circ \gamma) &= A_x(x \circ \gamma, y \circ \gamma) \\ D(v_y \circ \gamma) &= A_y(x \circ \gamma, y \circ \gamma) \end{aligned} \quad (3.31)$$

Construct a vector field on  $\mathbb{R}^5$  corresponding to this system of differential equations. Derive the first few terms in the series solution of this problem by exponentiation.

## 3.4 One-Form Fields

A vector field that gives a velocity for each point on a topographic map of the surface of the Earth can be applied to a function, such as one that gives the height for each point on the topographic map, or a map that gives the temperature for each point. The vector field then provides the rate of change of the height or temperature as one moves in the way described by the vector field. Alternatively, we can think of a topographic map, which gives the height at each point, as measuring a velocity field at each point. For example, we may be interested in the velocity of the wind or the trajectories of migrating birds. The topographic map gives the rate of change of height at each point for each velocity vector field. The rate of

change of height can be thought of as the number of equally-spaced (in height) contours that are pierced by each velocity vector in the vector field.

### 3.4.1 Differential of a Function

For example, consider the *differential*<sup>9</sup>  $df$  of a manifold function  $f$ , defined as follows. If  $df$  is applied to a vector field  $\mathbf{v}$  we obtain

$$df(\mathbf{v}) = \mathbf{v}(f), \quad (3.32)$$

which is a function of a manifold point.

The differential of the height function on the topographic map is a function that gives the rate of change of height at each point for a velocity vector field. This gives the same answer as the velocity vector field applied to the height function.

The differential of a function is linear in the vector fields. The differential is also a linear operator on functions: if  $f_1$  and  $f_2$  are manifold functions, and if  $c$  is a real constant, then

$$d(f_1 + f_2) = df_1 + df_2 \quad (3.33)$$

and

$$d(cf) = cdf. \quad (3.34)$$

Note that  $c$  is not a manifold function.

### 3.4.2 One-Form Fields

A one-form field is a generalization of this idea; it is something that measures a vector field at each point.

*One-form fields* are linear functions of vector fields that produce real-valued functions on the manifold. A one-form field is linear in vector fields: if  $\omega$  is a one-form field,  $\mathbf{v}$  and  $\mathbf{w}$  are vector fields, and  $c$  is a manifold function, then

$$\omega(\mathbf{v} + \mathbf{w}) = \omega(\mathbf{v}) + \omega(\mathbf{w}) \quad (3.35)$$

and

$$\omega(c\mathbf{v}) = c\omega(\mathbf{v}). \quad (3.36)$$

Sums and scalar products of one-form fields on a manifold have the following properties. If  $\omega$  and  $\theta$  are one-form fields, and if  $f$  is a real-valued manifold function, then:

$$(\omega + \theta)(\mathbf{v}) = \omega(\mathbf{v}) + \theta(\mathbf{v}), (f\omega)(\mathbf{v}) = f\omega(\mathbf{v}). \quad (3.37)$$

### 3.4.3 Coordinate-Basis One-Form Fields

Given a coordinate function  $\chi$ , we define the coordinate-basis one-form fields  $\tilde{X}^i$  by

$$\tilde{X}^i(\mathbf{v})(m) = \mathbf{v}(\chi^i)(m) \quad (3.38)$$

or collectively

$$\tilde{X}(\mathbf{v})(m) = \mathbf{v}(\chi)(m). \quad (3.39)$$

With this definition the coordinate-basis one-form fields are dual to the coordinate-basis vector fields in the following sense (see equation (3.15)):<sup>10</sup>

$$\tilde{X}^i(X_j)(m) = X_j(\chi^i)(m) = \partial_j(\chi^i \circ \chi^{-1})(\chi(m)) = \delta_j^i. \quad (3.40)$$

The tuple of basis one-form fields  $\tilde{X}(\mathbf{v})(m)$  is an up structure like that of  $\chi$ .

The general one-form field  $\omega$  is a linear combination of coordinate-basis one-form fields:

$$\omega(\mathbf{v}) = (a \circ \chi)\tilde{X}(\mathbf{v}) \quad (3.41)$$

<sup>9</sup>The differential of a manifold function will turn out to be a special case of the exterior derivative, which will be introduced later.

<sup>10</sup>The Kronecker delta  $\delta_j^i$  is one if  $i = j$  and zero otherwise.

with coefficient-function tuple  $a(x)$ , for  $x = \chi(m)$ . We can write this more simply as

$$\omega(v) = (a \circ \chi)\tilde{X}(v), \quad (3.42)$$

because everything is evaluated at  $m$ .

The coefficient tuple can be recovered from the one-form field:<sup>11</sup>

$$a_i(x) = \omega(\tilde{X}_i)(\chi^{-1}(x)). \quad (3.43)$$

This follows from the dual relationship (3.41). We can see this as a program:<sup>12</sup>

```
(def omega
  (components→oneform-field (down (literal-function 'a_0 R2→R) (literal-function
    'a_1 R2→R))
    R2-rect))
;; ⇒ (down (down d:dx d:dy) R2-rect-point)
;; ⇒ (down (a_0 (up x0 y0)) (a_1 (up x0 y0)))
```

We provide a shortcut for this construction:

```
(def omega (literal-oneform-field 'a R2-rect))
```

A differential can be expanded in a coordinate basis:

$$df(v) = \sum_i c_i \tilde{X}^i(v). \quad (3.44)$$

The coefficients  $c_i = df(X_i) = X_i(f) = \partial_i(f \circ \chi^{-1}) \circ \chi$  are the partial derivatives of the coordinate representation of  $f$  in the coordinate system of the basis:

```
((d (literal-manifold-function 'f-rect R2-rect)) (coordinate-system→vector-basis R2-closure
  rect))
  R2-rect-point)
;; ⇒ (down ((partial 0) f-rect) (up x0 y0)) ((partial 1) f-rect) (up x0 y0)))
```

However, if the coordinate system of the basis differs from the coordinates of the representation of the function, the result is complicated by the chain rule:

```
(simplify ((d (literal-manifold-function 'f-polar R2-polar))
  (coordinate-system→vector-basis R2-rect))
  ((point R2-polar) (up 'r 'theta)))
;; ⇒ (down (/ (+ (* r (cos theta) ((partial 0) f-polar) (up r theta)))
  (* -1 (sin theta) ((partial 1) f-polar) (up r theta))))
  r)
;;
;; (/ (+ (* r (sin theta) ((partial 0) f-polar) (up r theta)))
  (* (cos theta) ((partial 1) f-polar) (up r theta))))
  r))
```

) The coordinate-basis one-form fields can be used to find the coefficients of vector fields in the corresponding coordinate vector-field basis:

$$\tilde{X}^i(v) = v(\chi^i) = b^i \circ \chi \quad (3.45)$$

or collectively,

$$\tilde{X}(v) = v(\chi) = b \circ \chi. \quad (3.46)$$

<sup>11</sup>The analogous recovery of coefficient tuples from vector fields is equation (3.3):  $b^i_{\chi,v} = v(\chi^i) \circ \chi^{-1}$ .

<sup>12</sup>The procedure `components→1form-field` is analogous to the procedure `components→vector-field` introduced earlier.

A coordinate-basis one-form field is often written  $dx^i$ . This traditional notation for the coordinate-basis one-form fields is justified by the relation:

$$dx^i = \tilde{X}^i = d(\chi^i). \quad (3.47)$$

The `define-coordinates` procedure also makes the basis one-form fields with these traditional names inherited from the coordinates.

We can illustrate the duality of the coordinate-basis vector fields and the coordinate-basis one-form fields:

```
(define-coordinates (up x y) R2-rect) closure

((dx d:dy) R2-rect-point)
;; => 0

((dx d:dx) R2-rect-point)
;; => 1
```

We can use the coordinate-basis one-form fields to extract the coefficients of `circular` on the rectangular vector basis:

```
((dx circular) R2-rect-point) closure
;; => (- yθ)

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((dy circular) R2-rect-point))
;; => xθ
```

But we can also find the coefficients on the polar vector basis:

```
(simplify ((dr circular) R2-rect-point)) closure
;; => 0

(simplify ((dtheta circular) R2-rect-point))
;; => 1
```

So `circular` is the same as `d/dtheta`, as we can see by applying them both to the general function `f`:

```
(def f (literal-manifold-function 'f-rect R2-rect)) closure

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((- circular d:dtheta) f) R2-rect-point))
;; => 0
```

### 3.4.4 Not All One-Form Fields Are Differentials

Although all one-form fields can be constructed as linear combinations of basis one-form fields, not all one-form fields are differentials of functions.

The coefficients of a differential are (see equation (3.45)):

$$c_i = X_i(f) = df(X_i) \quad (3.48)$$

and partial derivatives of functions commute

$$X_i(X_j(f)) = X_j(X_i(f)). \quad (3.49)$$

As a consequence, the coefficients of a differential are constrained

$$X_i(c_j) = X_j(c_i), \quad (3.50)$$

but a one-form field can be constructed with arbitrary coefficient functions. For example:

$$x dx + x dy \quad (3.51)$$

is not a differential of any function. This is why we started with the basis one-form fields and built the general one-form fields in terms of them.

### 3.4.5 Coordinate Transformations

Consider a coordinate change from the chart  $\chi$  to the chart  $\chi'$ .

$$\begin{aligned}\tilde{X}(v) &= v(\chi) \\ &= v(\chi \circ (\chi')^{-1} \circ \chi') \\ &= (D(\chi \circ (\chi')^{-1}) \circ \chi') v(\chi') \\ &= (D(\chi \circ (\chi')^{-1}) \circ \chi') \circ \tilde{X}'(v),\end{aligned}\tag{3.52}$$

where the third line follows from the chain rule for vector fields.

One-form fields are independent of coordinates. So,

$$\omega(v) = (a \circ \chi) \tilde{X}(v) = (a' \circ \chi') \tilde{X}'(v).\tag{3.53}$$

Eqs. (3.54) and (3.53) require that the coefficients transform under coordinate transformations as follows:

$$a(\chi(m)) D(\chi \circ (\chi')^{-1})(\chi'(m)) = a'(\chi'(m)),\tag{3.54}$$

or

$$a(\chi(m)) = a'(\chi'(m)) (D(\chi \circ (\chi')^{-1})(\chi'(m)))^{-1}.\tag{3.55}$$

The coefficient tuple  $a(x)$  is a down structure compatible for contraction with  $b(x)$ . Let  $v$  be the vector with coefficient tuple  $b(x)$ , and  $\omega$  be the one-form with coefficient tuple  $a(x)$ . Then, by equation (3.43),

$$\omega(v) = (a \circ \chi)(b \circ \chi).\tag{3.56}$$

As a program:

```
(def omega (literal-oneform-field 'a R2-rect)) ;clojure

(def v (literal-vector-field 'b R2-rect))

((omega v) R2-rect-point)
;; => (+ (* (a_0 (up x0 y0)) (b^0 (up x0 y0))) (* (a_1 (up x0 y0)) (b^1 (up x0 y0))))
```

Comparing equation (3.56) with equation (3.23) we see that one-form components and vector components transform oppositely, so that

$$a(x)b(x) = a'(x')b'(x'),\tag{3.57}$$

as expected because  $\omega(v)(m)$  is independent of coordinates.

#### Exercise 3.2: Verification

Verify that the coefficients of a one-form field transform as described in equation (3.56). You should use equation (3.44) in your derivation.

#### Exercise 3.3: Hill Climbing

The topography of a region on the Earth can be specified by a manifold function  $h$  that gives the altitude at each point on the manifold. Let  $v$  be a vector field on the manifold, perhaps specifying a direction and rate of walking at every point on the manifold.

- Form an expression that gives the power that must be expended to follow the vector field at each point.
- Write this as a computational expression.

## 4 Basis Fields

A vector field may be written as a linear combination of basis vector fields. If  $n$  is the dimension, then any set of  $n$  linearly independent vector fields may be used as a basis. The coordinate basis  $X$  is an example of a basis.<sup>1</sup> We will see later that not every basis is a coordinate basis: in order to be a coordinate basis, there must be a coordinate system such that each basis element is the directional derivative operator in a corresponding coordinate direction.

Let  $e$  be a tuple of basis vector fields, such as the coordinate basis  $X$ . The general vector field  $v$  applied to an arbitrary manifold function  $f$  can be expressed as a linear combination


$$v(f)(m) = e(f)(m)b(m) = \sum_i e_i(f)(m)b^i(m), \quad (4.1)$$

where  $b$  is a tuple-valued coefficient function on the manifold. When expressed in a coordinate basis, the coefficients that specify the direction of the vector are naturally expressed as functions  $b^i$  of the coordinates of the manifold point. Here, the coefficient function  $b$  is more naturally expressed as a tuple-valued function on the manifold. If  $b$  is the coefficient function expressed as a function of coordinates, then  $b = b \circ \chi$  is the coefficient function as a function on the manifold.

The coordinate-basis forms have a simple definition in terms of the coordinate-basis vectors and the coordinates (equation (3.40)). With this choice, the dual property, equation (3.41), holds without further fuss. More generally, we can define a basis of one-forms  $\tilde{e}$  that is dual to  $e$  in that the property

$$\tilde{e}^i(e_j)(m) = \delta_j^i \quad (4.2)$$

is satisfied, analogous to property (3.41). Figure 4.1 illustrates the duality of basis fields.



**Figure 4.1:** Let arrows  $e_0$  and  $e_1$  depict the vectors of a basis vector field at a particular point. Then the foliations shown by the parallel lines depict the dual basis one-form fields at that point. The dotted lines represent the field  $\tilde{e}^0$  and the dashed lines represent the field  $\tilde{e}^1$ . The spacings of the lines are  $\frac{1}{3}$  unit. That the vectors pierce three of the lines representing their duals and do not pierce any of the lines representing the other basis elements is one way to see the relationship  $\tilde{e}^i(e_j)(m) = \delta_j^i$ .

To solve for the dual basis  $\tilde{e}$  given the basis  $e$ , we express the basis vectors  $e$  in terms of a coordinate basis<sup>2</sup>

$$e_j(f) = \sum_k X_k(f)c_j^k, \quad (4.3)$$

and the dual one-forms  $\tilde{e}$  in terms of the dual coordinate one-forms

<sup>1</sup>We cannot say if the basis vectors are orthogonal or normalized until we introduce a metric.

<sup>2</sup>We write the vector components on the right and the tuple of basis vectors on the left because if we think of the basis vectors as organized as a row and the components as organized as a column then the formula is just a matrix multiplication.

$$\tilde{e}^i(v) = \sum_l d_l^i \tilde{X}^l(v), \quad (4.4)$$

then

$$\begin{aligned} \tilde{e}^i(e_j) &= \sum_l d_l^i \tilde{X}^l(e_j) \\ &= \sum_l d_l^i e_j(\chi^l) \\ &= \sum_l d_l^i \sum_k \chi_k(\chi^l) c_j^k \\ &= \sum_{kl} d_l^i \delta_k^l c_j^k \\ &= \sum_k d_k^i c_j^k. \end{aligned} \quad (4.5)$$

Applying this at  $m$  we get

$$\tilde{e}^i(e_j)(m) = \delta_j^i = \sum_k d_k^i(m) c_j^k(m). \quad (4.6)$$

So the  $d$  coefficients can be determined from the  $c$  coefficients (essentially by matrix inversion).

A set of vector fields  $\{e_i\}$  may be linearly independent in the sense that a weighted sum of them may not be identically zero over a region, yet it may not be a basis in that region. The problem is that there may be some places in the region where the vectors are not independent. For example, two of the vectors may be parallel at a point but not parallel elsewhere in the region. At such a point  $m$  the determinant of the matrix  $c(m)$  is zero. So at these points we cannot define the dual basis forms.<sup>3</sup>

The dual form fields can be used to determine the coefficients  $b$  of a vector field  $v$  relative to a basis  $e$ , by applying the dual basis form fields  $\tilde{e}$  to the vector field. Let

$$v(f) = \sum_i e_i(f) b^i. \quad (4.7)$$

Then

$$\tilde{e}^j(v) = b^j. \quad (4.8)$$

Define two general vector fields:

```
(def e0
  (+ (* (literal-manifold-function 'e0x R2-rect) d:dx)
    (* (literal-manifold-function 'e0y R2-rect) d:dy)))

(def e1
  (+ (* (literal-manifold-function 'e1x R2-rect) d:dx)
    (* (literal-manifold-function 'e1y R2-rect) d:dy)))
```

We use these as a vector basis and compute the dual:

```
(def e-vector-basis (down e0 e1))

(def e-dual-basis (vector-basis→dual e-vector-basis R2-polar))
```

The procedure `vector-basis→dual` requires an auxiliary coordinate system (here `R2-polar`) to get the  $c_j^k$  coefficient functions from which we compute the  $d_i^k$  coefficient functions. However, the final result is independent of this coordinate system. Then we can verify that the bases  $e$  and  $\tilde{e}$  satisfy the dual relationship (equation (3.41)) by applying the dual basis to the vector basis:

<sup>3</sup>This is why the set of vector fields and the set of one-form fields are modules rather than vector spaces.



```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify ((e-dual-basis e-vector-basis) R2-rect-point))
;; => (up (down 1 0) (down 0 1))
```

Note that the dual basis was computed relative to the polar coordinate system: the resulting objects are independent of the coordinates in which they were expressed!

Or we can make a general vector field with this basis and then pick out the coefficients by applying the dual basis:

```
(def v                                          clojure
  (* (up (literal-manifold-function 'b↑0 R2-rect) (literal-manifold-function 'b↑1 R2-
rect))
    e-vector-basis))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((e-dual-basis v) R2-rect-point))
;; => (up (b↑0 (up x0 y0)) (b↑1 (up x0 y0)))
```

## 4.1 Change of Basis

Suppose that we have a vector field  $v$  expressed in terms of one basis  $e$  and we want to reexpress it in terms of another basis  $e'$ . We have

$$v(f) = \sum_i e_i(f) b^i = \sum_i e'_j(f) b'^j. \quad (4.9)$$

The coefficients  $b'$  can be obtained from  $v$  by applying the dual basis

$$b'^j = \tilde{e}'^j(v) = \sum_i \tilde{e}'^j(e_i) b^i. \quad (4.10)$$

Let

$$J_i^j = \tilde{e}'^j(e_i), \quad (4.11)$$

then

$$b'^j = \sum_i J_i^j b^i, \quad (4.12)$$

and

$$e_i(f) = \sum_j e'_j(f) J_i^j. \quad (4.13)$$

The Jacobian  $J$  is a structure of manifold functions. Using tuple arithmetic, we can write

$$b' = Jb \quad (4.14)$$

and

$$e(f) = e'(f)J. \quad (4.15)$$

We can write

```
(defn Jacobian                                clojure
  [to-basis from-basis]
  (mapr (basis→oneform-basis to-basis) (basis→vector-basis from-basis)))
```

The polar components are:

```
(def b-polar                                clojure
  (* (Jacobian (coordinate-system→basis R2-polar) (coordinate-system→basis R2-rect))
```

```

b-rect))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (b-polar ((point R2-rect) (up 'x0 'y0))))
;; => (up (/ (+ (* x0 (b↑0 (up x0 y0))) (* y0 (b↑1 (up x0 y0))))
;;      (sqrt (+ (expt x0 2) (expt y0 2))))
;;      (/ (+ (* x0 (b↑1 (up x0 y0))) (* -1 y0 (b↑0 (up x0 y0))))
;;      (+ (expt x0 2) (expt y0 2))))

```

We can also get the polar components directly:

```

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((coordinate-system→oneform-basis R2-polar) (literal-vector-field 'b R2-
rect)))
((point R2-rect) (up 'x0 'y0)))
;; => (up (/ (+ (* x0 (b↑0 (up x0 y0))) (* y0 (b↑1 (up x0 y0))))
;;      (sqrt (+ (expt x0 2) (expt y0 2))))
;;      (/ (+ (* x0 (b↑1 (up x0 y0))) (* -1 y0 (b↑0 (up x0 y0))))
;;      (+ (expt x0 2) (expt y0 2))))

```

We see that they are the same.

If  $K$  is the Jacobian that relates the basis vectors in the other direction

$$e'(f) = e(f)K \quad (4.16)$$

then

$$KJ = I = JK \quad (4.17)$$

where  $I$  is a manifold function that returns the multiplicative identity.

The dual basis transforms oppositely. Let

$$\omega = \sum_i a_i \tilde{e}'^i. \quad (4.18)$$

The coefficients are<sup>4</sup>

$$a_i = \omega(e_i) = \sum_j a'_j \tilde{e}'^j(e_i) = \sum_j a'_j J_i^j \quad (4.19)$$

or, in tuple arithmetic,

$$a = a' J. \quad (4.20)$$

Because of equation (4.18) we can deduce

$$\tilde{e} = K \tilde{e}'. \quad (4.21)$$

## 4.2 Rotation Basis

One interesting basis for rotations in 3-dimensional space is not a coordinate basis.

Rotations are the actions of the special orthogonal group  $SO(3)$ , which is a 3-dimensional manifold. The elements of this group may be represented by the set of  $3 \times 3$  orthogonal matrices with determinant +1.

We can use a coordinate patch on this manifold with Euler angle coordinates: each element has three coordinates,  $\theta$ ,  $\phi$ ,  $\psi$ . A manifold point may be represented by a rotation matrix. The rotation matrix for Euler angles is a product of three simple rotations:  $M(\theta, \phi, \psi) = R_z(\phi)R_x(\theta)R_z(\psi)$ , where  $R_x$  and  $R_z$  are functions that take an angle and produce the matrices representing rotations about the  $x$  and  $z$  axes, respectively. We can visualize  $\theta$  as the colatitude of the pole from the  $\hat{z}$ -axis,  $\phi$  as the longitude, and  $\psi_a$  as the rotation around the pole.

<sup>4</sup>We see from equations (4.15) and (4.16) that  $J$  and  $K$  are inverses. We can obtain their coefficients by:  $J_i^j = \tilde{e}'^j(e_i)$  and  $K_i^j = \tilde{e}^j(e'_i)$ .

Given a rotation specified by Euler angles, how do we change the Euler angle to correspond to an incremental rotation of size  $\epsilon$  about the  $\hat{x}$ -axis? The direction  $(a, b, c)$  is constrained by the equation

$$R_x(\epsilon)M(\theta, \phi, \psi) = M(\theta + a\epsilon, \phi + b\epsilon, \psi + c\epsilon). \quad (4.22)$$

Linear equations for  $(a, b, c)$  can be found by taking the derivative of this equation with respect to  $\epsilon$ . We find

$$0 = c \cos \theta + b, \quad (4.23)$$

$$0 = a \sin \phi - c \cos \phi \sin \theta, \quad (4.24)$$

$$1 = c \sin \phi \sin \theta + a \cos \phi, \quad (4.25)$$

with the solution

$$a = \cos \phi, \quad (4.26)$$

$$b = -\frac{\sin \phi \cos \theta}{\sin \theta}, \quad (4.27)$$

$$c = \frac{\sin \phi}{\sin \theta}. \quad (4.28)$$

Therefore, we can write the basis vector field that takes directional derivatives in the direction of incremental  $x$  rotations as

$$\mathbf{e}_x = a \frac{\partial}{\partial \theta} + b \frac{\partial}{\partial \phi} + c \frac{\partial}{\partial \psi} = \cos \phi \frac{\partial}{\partial \theta} - \frac{\sin \phi \cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} + \frac{\sin \phi}{\sin \theta} \frac{\partial}{\partial \psi}. \quad (4.29)$$

Similarly, vector fields for the incremental  $y$  and  $z$  rotations are

$$\mathbf{e}_y = \frac{\cos \phi \cos \theta}{\sin \theta} \frac{\partial}{\partial \phi} + \sin \phi \frac{\partial}{\partial \theta} - \frac{\cos \phi}{\sin \theta} \frac{\partial}{\partial \psi} \quad (4.30)$$

$$\mathbf{e}_z = \frac{\partial}{\partial \phi}. \quad (4.31)$$

### 4.3 Commutators

The commutator of two vector fields is defined as

$$[\mathbf{v}, \mathbf{w}](f) = \mathbf{v}(\mathbf{w}(f)) - \mathbf{w}(\mathbf{v}(f)). \quad (4.32)$$

In the special case that the two vector fields are coordinate basis fields, the commutator is zero:

$$\begin{aligned} [X_i, X_j](f) &= X_i(X_j(f)) - X_j(X_i(f)) \\ &= \partial_i \partial_j (f \circ \chi^{-1}) \circ \chi - \partial_j \partial_i (f \circ \chi^{-1}) \circ \chi \\ &= 0, \end{aligned} \quad (4.33)$$

because the individual partial derivatives commute. The vanishing commutator is telling us that we get to the same manifold point by integrating from a point along first one basis vector field and then another as from integrating in the other order. If the commutator is zero we can use the integral curves of the basis vector fields to form a coordinate mesh.

More generally, the commutator of two vector fields is a vector field. Let  $\mathbf{v}$  be a vector field with coefficient function  $\mathbf{c} = \mathbf{c} \circ \chi$ , and  $\mathbf{u}$  be a vector field with coefficient function  $\mathbf{b} = \mathbf{b} \circ \chi$ , both with respect to the coordinate basis  $\mathbf{X}$ . Then

$$\begin{aligned}
[u, v](f) &= u(v(f)) - v(u(f)) \\
&= u\left(\sum_i X_i(f) c^i\right) - v\left(\sum_j X_j(f) b^j\right) \\
&= \sum_j X_j\left(\sum_i X_i(f) c^i\right) b^j - \sum_i X_i\left(\sum_j X_j(f) b^j\right) c^i \\
&= \sum_{ij} [X_j, X_i](f) c^i b^j \\
&\quad + \sum_i X_i(f) \sum_j (X_j(c^i) b^j - X_j(b^i) c^j) \\
&= \sum_i X_i(f) a^i,
\end{aligned} \tag{4.34}$$

where the coefficient function  $\mathbf{a}$  of the commutator vector field is

$$\begin{aligned}
a^i &= \sum_j (X_j(c^i) b^j \\
&\quad - X_j(b^i) c^j) \\
&= u(c^i) - v(b^i).
\end{aligned} \tag{4.35}$$

We used the fact, shown above, that the commutator of two coordinate basis fields is zero.

We can check this formula for the commutator for the general vector fields `e0` and `e1` in polar coordinates:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (let [polar-basis (coordinate-system→basis R2-polar)
                polar-vector-basis (basis→vector-basis polar-basis)
                polar-dual-basis (basis→oneform-basis polar-basis)
                f (literal-manifold-function 'f-rect R2-rect)]
  ((- ((commutator e0 e1) f)
    (* (- (e0 (polar-dual-basis e1)) (e1 (polar-dual-basis e0)))
      (polar-vector-basis f)))
  R2-rect-point)))

;; ⇒ 0
```

Let  $\mathbf{e}$  be a tuple of basis vector fields. The commutator of two basis fields can be expressed in terms of the basis vector fields:

$$[e_i, e_j](f) = \sum_k d_{ij}^k e_k(f), \tag{4.36}$$

where  $d_{ij}^k$  are functions of  $\mathbf{m}$ , called the *structure constants* for the basis vector fields. The coefficients are

$$d_{ij}^k = \tilde{e}^k([e_i, e_j]). \tag{4.37}$$

The commutator  $[u, v]$  with respect to a non-coordinate basis  $\mathbf{e}_i$  is

$$[u, v](f) = \sum_k e_k(f) \left( u(c^k) - v(b^k) + \sum_{ij} c^i b^j d_{ji}^k \right) \tag{4.38}$$

Define the vector fields `Jx`, `Jy`, and `Jz` that generate rotations about the three rectangular axes in three dimensions:<sup>5</sup>

<sup>5</sup>Using

```
(def R3-rect (coordinate-system-at R3 :rectangular :origin))      clojure

(define-coordinates (up x y z) R3-rect)
```

```

(define-coordinates (up x y z) R3-rect)                                clojure

(def Jz (- (* x d:dy) (* y d:dx)))

(def Jx (- (* y d:dz) (* z d:dy)))

(def Jy (- (* z d:dx) (* x d:dz)))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((+ (commutator Jx Jy) Jz) g) R3-rect-point))
;; ⇒ 0

(simplify (((+ (commutator Jy Jz) Jx) g) R3-rect-point))
;; ⇒ 0

(simplify (((+ (commutator Jz Jx) Jy) g) R3-rect-point))
;; ⇒ 0

```

We see that

$$\begin{aligned}
 [J_x, J_y] &= -J_z \\
 [J_y, J_z] &= -J_x \\
 [J_z, J_x] &= -J_y
 \end{aligned} \tag{4.39}$$

We can also compute the commutators for the basis vector fields  $\mathbf{e}_x$ ,  $\mathbf{e}_y$ , and  $\mathbf{e}_z$  in the  $\text{SO}(3)$  manifold (see equations (4.29) – (4.31)) that correspond to rotations about the  $x$ ,  $y$ , and  $z$  axes, respectively:<sup>6</sup>

```

(def Euler-angles (coordinate-system-at S03 :Euler :Euler-patch))    clojure

(define-coordinates (up theta phi psi) Euler-angles)

(def S03-point ((point Euler-angles) (up 'theta 'phi 'psi)))

(def f (literal-manifold-function 'f-Euler Euler-angles))

(def e_x
  (+ (* (cos phi) d:dtheta)
    (* -1 (/ (* (sin phi) (cos theta)) (sin theta)) d:dphi)
    (* (/ (sin phi) (sin theta)) d:dpsi)))

(def e_y
  (+ (/ (* (cos phi) (cos theta) d:dphi) (sin theta))
    (* (sin phi) d:dtheta)
    (* -1 (/ (cos phi) (sin theta)) d:dpsi)))

```

```

(def R3-rect-point ((point R3-rect) (up 'x0 'y0 'z0)))

(def g (literal-manifold-function 'g-rect R3-rect))

```

<sup>6</sup>Using

```

(def Euler-angles (coordinate-system-at S03 :Euler :Euler-patch))    clojure

(def Euler-angles-chi-inverse (point Euler-angles))

(define-coordinates (up theta phi psi) Euler-angles)

(def S03-point ((point Euler-angles) (up 'theta 'phi 'psi)))

(def f (literal-manifold-function 'f-Euler Euler-angles))

```

```

(def e_z d:dphi)

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((+ (commutator e_x e_y) e_z) f) S03-point))
;; => 0

(simplify (((+ (commutator e_y e_z) e_x) f) S03-point))
;; => 0

(simplify (((+ (commutator e_z e_x) e_y) f) S03-point))
;; => 0

```

You can tell if a set of basis vector fields is a coordinate basis by calculating the commutators. If they are nonzero, then the basis is not a coordinate basis. If they are zero then the basis vector fields can be integrated to give the coordinate system.

Recall equation (3.31)

$$(e^{tv})(m) = (f \circ \phi_t^v)(m). \quad (4.40)$$

Iterating this equation, we find

$$(e^{sw}e^{tv})(m) = (f \circ \phi_t^v \circ \phi_s^w)(m). \quad (4.41)$$

Notice that the evolution under  $w$  occurs before the evolution under  $v$ .

To illustrate the meaning of the commutator, consider the evolution around a small loop with sides made from the integral curves of two vector fields  $v$  and  $w$ . We will first follow  $v$ , then  $w$ , then  $-v$ , and then  $-w$ :

$$(e^{\epsilon v} e^{\epsilon w} e^{-\epsilon v} e^{-\epsilon w} f)(m). \quad (4.42)$$

To second order in  $\epsilon$  the result is<sup>7</sup>

$$(e^{\epsilon^2 [v, w]} f)(m) \quad (4.43)$$


This result is illustrated in figure 4.2.

---

<sup>7</sup>For non-commuting operators  $A$  and  $B$ ,

$$e^A e^B e^{-A} e^{-B} = \left(1 + A + \frac{A^2}{2} + \dots\right) \left(1 + B + \frac{B^2}{2} + \dots\right) \times \left(1 - A + \frac{A^2}{2} + \dots\right) \left(1 - B + \frac{B^2}{2} + \dots\right) = 1 + [A, B] + \dots,$$

to second order in  $A$  and  $B$ . All higher-order terms can be written in terms of higher-order commutators of  $A$  and  $B$ . An example of a higher-order commutator is  $[A, [A, B]]$ .



**Figure 4.2:** The commutator of two vector fields computes the residual of a small loop following their integral curves.

Take a point  $0$  in  $M$  as the origin. Then, presuming  $[\mathbf{e}_i, \mathbf{e}_j] = 0$ , the coordinates  $x$  of the point  $m$  in the coordinate system corresponding to the  $\mathbf{e}$  basis satisfy<sup>8</sup>

$$m = \phi_1^{xe}(0) = \chi^{-1}(x), \quad (4.44)$$

where  $\chi$  is the coordinate function being defined. Because the elements of  $\mathbf{e}$  commute, we can translate separately along the integral curves in any order and reach the same point; the terms in the exponential can be factored into separate exponentials if needed.

#### 4.4 Exercise 4.1: Alternate Angles

Note that the Euler angles are singular at  $\theta = 0$  (where  $\phi$  and  $\psi$  become degenerate), so the representations of  $\mathbf{e}_x$ ,  $\mathbf{e}_y$ , and  $\mathbf{e}_z$  (defined in equations (4.29) – (4.31)) have problems there. An alternate coordinate system avoids this problem, while introducing a similar problem elsewhere in the manifold. Consider the “alternate angles”  $(\theta_a, \phi_a, \psi_a)$  which define a rotation matrix via  $M(\theta_a, \phi_a, \psi_a) = R_z(\phi_a)R_x(\theta_a)R_y(\psi_a)$ .

a. Where does the singularity appear in these alternate coordinates? Do you think you could define a coordinate system for rotations that has no singularities?

b. What do the  $\mathbf{e}_x$ ,  $\mathbf{e}_y$ , and  $\mathbf{e}_z$  basis vector fields look like in this coordinate system?

#### 4.5 Exercise 4.2: General Commutators

Verify equation (4.38).

#### 4.6 Exercise 4.3: SO(3) Basis and Angular Momentum Basis

How are  $J_x$ ,  $J_y$ , and  $J_z$  related to  $\mathbf{e}_x$ ,  $\mathbf{e}_y$ , and  $\mathbf{e}_z$  in equations (4.29) – (4.31)?

<sup>8</sup>Here  $x$  is an up-tuple structure of components, and  $\mathbf{e}$  is down-tuple structure of basis vectors. The product of the two contracts to make a scaled vector, along which we translate by one unit.

## 5 Integration

We know how to integrate real-valued functions of a real variable. We want to extend this idea to manifolds, in such a way that the integral is independent of the coordinate system used to compute it.

The integral of a real-valued function of a real variable is the limit of a sum of products of the values of the function on subintervals and the lengths of the increments of the independent variable in those subintervals:

$$\int_a^b f = \int_a^b f(x)dx = \lim_{\Delta x_i \rightarrow 0} \sum_i f(x_i) \Delta x_i \quad (5.1)$$

If we change variables ( $x = g(y)$ ), then the form of the integral changes:

$$\begin{aligned} \int_a^b f &= \int_a^b f(x)dx \\ &= \int_{g^{-1}(a)}^{g^{-1}(b)} f(g(y))Dg(y)dy \\ &= \int_{g^{-1}(a)}^{g^{-1}(b)} (f \circ g)Dg \end{aligned} \quad (5.2)$$

We can make a coordinate-independent notion of integration in the following way. An interval of the real line is a 1-dimensional manifold with boundary. We can assign a coordinate chart  $\chi$  to this manifold. Let  $x = \chi(m)$ . The coordinate basis is associated with a coordinate-basis vector field, here  $\partial/\partial x$ . Let  $\omega$  be a one-form on this manifold. The application of  $\omega$  to  $\partial/\partial x$  is a real-valued function on the manifold. If we compose this with the inverse chart, we get a real-valued function of a real variable. We can then write the usual integral of this function

$$I = \int_a^b \omega(\partial/\partial x) \circ \chi^{-1} \quad (5.3)$$

It turns out that the value of this integral is independent of the coordinate chart used in its definition. Consider a different coordinate chart  $x' = \chi'(m)$ , with associated basis vector field  $\partial/\partial x'$ . Let  $g = \chi' \circ \chi^{-1}$ . We have

$$\begin{aligned} \int_{a'}^{b'} \omega(\partial/\partial x') \circ \chi'^{-1} &= \int_{a'}^{b'} \omega(\partial/\partial x (D(\chi \circ \chi'^{-1}) \circ \chi')) \circ \chi'^{-1} \\ &= \int_{a'}^{b'} (\omega(\partial/\partial x) D(\chi \circ \chi'^{-1}) \circ \chi') \circ \chi'^{-1} \\ &= \int_{a'}^{b'} (\omega(\partial/\partial x) \circ \chi'^{-1}) D(\chi \circ \chi'^{-1}) \\ &= \int_a^b (((\omega(\partial/\partial x) \circ \chi^{-1}) D(\chi \circ \chi'^{-1})) \circ g) Dg \\ &= \int_a^b \omega(\partial/\partial x) \circ \chi^{-1}, \end{aligned} \quad (5.4)$$

where we have used the rule for coordinate transformations of basis vectors (equation (3.19)), linearity of forms in the first two lines, and the rule for change-of-variables under an integral in the last line.<sup>1</sup>

Because the integral is independent of the coordinate chart, we can write simply

$$I = \int_M \omega, \quad (5.5)$$

where  $M$  is the 1-dimensional manifold with boundary corresponding to the interval.

<sup>1</sup>Note  $(D(\chi \circ \chi'^{-1}) \circ (\chi' \circ \chi^{-1})) D(\chi' \circ \chi^{-1}) = 1$ . With  $g = \chi' \circ \chi^{-1}$  this is  $(D(g^{-1} \circ g)(Dg) = 1$ .



We are exploiting the fact that coordinate basis vectors in different coordinate systems are related by a Jacobian (see equation (3.19)), which cancels the Jacobian that appears in the change-of-variables formula for integration (see equation (5.2)).

### 5.1 Higher Dimensions

We have seen that we can integrate one-forms on 1-dimensional manifolds. We need higher-rank forms that we can integrate on higher-dimensional manifolds in a coordinate-independent manner.

Consider the integral of a real-valued function,  $f : R^n \rightarrow R$ , over a region  $U$  in  $R^n$ . Under a coordinate transformation  $g : R^n \rightarrow R^n$ , we have<sup>2</sup>

$$\int_U f = \int_{g^{-1}(U)} (f \circ g) \det(Dg). \quad (5.6)$$

A rank  $n$  form field takes  $n$  vector field arguments and produces a real-valued manifold function:  $\omega(v, w, \dots, u)(m)$ . By analogy with the 1-dimensional case, higher-rank forms are linear in each argument. Higher-rank forms must also be antisymmetric under interchange of any two arguments in order to make a coordinate-free definition of integration analogous to equation (5.3).

Consider an integral in the coordinate system  $\chi$ :

$$\int_{\chi(U)} \omega(X_0, X_1, \dots) \circ \chi^{-1}. \quad (5.7)$$

Under coordinate transformations  $g = \chi \circ \chi'^{-1}$ , the integral becomes

$$\int_{\chi'(U)} \omega(X_0, X_1, \dots) \circ \chi'^{-1} \det(Dg). \quad (5.8)$$

Using the change-of-basis formula, equation (3.19):

$$X(f) = X'(f)(D(\chi' \circ \chi^{-1})) \circ \chi = X'(f)(D(g^{-1}) \circ \chi). \quad (5.9)$$

If we let  $M = (D(g^{-1})) \circ \chi$  then

$$\begin{aligned} (\omega(X_0, X_1, \dots) \circ \chi'^{-1}) \det(Dg) &= (\omega(X'_0 M_0, X'_1 M_1, \dots) \circ \chi'^{-1}) \det(Dg) \\ &= (\omega(X'_0, X'_1, \dots) \circ \chi'^{-1}) \alpha(M_0, M_1, \dots) \det(Dg), \end{aligned} \quad (5.10)$$

using the multilinearity of  $\omega$ , where  $M_i$  is the  $i^{\text{th}}$  column of  $M$ . The function  $\alpha$  is multilinear in the columns of  $M$ . To make a coordinate-independent integration we want the expression (5.10) to be the same as the integrand in

$$I' = \int_{\chi'(U)} \omega(X'_0, X'_1, \dots) \circ \chi'^{-1}. \quad (5.11)$$

For this to be the case,  $\alpha(M_0, M_1, \dots)$  must be  $(\det(D(g))^{-1} = \det(M))$ . So  $\alpha$  is an antisymmetric function, and thus so is  $\omega$ .

Thus higher-rank form fields must be antisymmetric multilinear functions from vector fields to manifold functions. So we have a coordinate-independent definition of integration of form fields on a manifold and we can write

$$I = I' = \int_U \omega \quad (5.12)$$

### 5.2 Wedge Product

There are several ways we can construct antisymmetric higher-rank forms. Given two one-form fields  $\omega$  and  $\tau$  we can form a two-form field  $\omega \wedge \tau$  as follows:

---

<sup>2</sup>The determinant is the unique function of the rows of its argument that i) is linear in each row, ii) changes sign under any interchange of rows, and iii) is one when applied to the identity multiplier.

$$(\omega \wedge \tau)(v, w) = \omega(v)\tau(w) - \omega(w)\tau(v). \quad (5.13)$$

More generally we can form the wedge of higher-rank forms. Let  $\omega$  be a  $k$ -form field and  $\tau$  be an  $l$ -form field. We can form a  $(k+l)$ -form field  $\omega \wedge \tau$  as follows:

$$\omega \wedge \tau = \frac{(k+l)!}{k!l!} \text{Alt}(\omega \otimes \tau) \quad (5.14)$$

where, if  $\eta$  is a function on  $m$  vectors,

$$\text{Alt}(\eta)(v_0, \dots, v_{m-1}) = \frac{1}{m!} \sum_{\sigma \in \text{Perm}(m)} \text{Parity}(\sigma) \eta(v_{\sigma(0)}, \dots, v_{\sigma(m-1)}), \quad (5.15)$$

and where

$$\omega \otimes \tau(v_0, \dots, v_{k-1}, v_k, \dots, v_{k+l-1}) = \omega(v_0, \dots, v_{k-1}) \tau(v_k, \dots, v_{k+l-1}). \quad (5.16)$$

The wedge product is associative, and thus we need not specify the order of a multiple application. The factorial coefficients of these formulas are chosen so that


$$(dx \wedge dy \wedge \dots)(\partial/\partial x, \partial/\partial y, \dots) = 1. \quad (5.17)$$

This is true independent of the coordinate system.

Equation (5.17) gives us

$$\int_U dx \wedge dy \wedge \dots = \text{Volume}(U) \quad (5.18)$$

where  $\text{Volume}(U)$  is the ordinary volume of the region corresponding to  $U$  in the Euclidean space of  $\mathbb{R}^n$  with the orthonormal coordinate system  $(x, y, \dots)$ .<sup>3</sup>



**Figure 5.1:** The area of the parallelogram in the  $(x, y)$  coordinate plane is given by  $A(u, v)(m)$ .

An example two-form (see figure 5.1) is the oriented area of a parallelogram in the  $(x, y)$  coordinate plane at the point  $m$  spanned by two vectors  $u = u^0 \partial/\partial x + u^1 \partial/\partial y$  and  $v = v^0 \partial/\partial x + v^1 \partial/\partial y$ , which is given by

$$A(u, v)(m) = u^0(m)v^1 - v^0(m)u^1(m). \quad (5.19)$$

Note that this is the area of the parallelogram in the coordinate plane, which is the range of the coordinate function. It is not the area on the manifold. To define that, we need more structure — the metric. We will put a metric on the manifold in Chapter 9.

### 5.3 3-Dimensional Euclidean Space

Let's specialize to 3-dimensional Euclidean space. Following equation (5.18) we can write the coordinate-area two-form in another way:  $A = dx \wedge dy$ . As code:

<sup>3</sup>By using the word “orthonormal” here we are assuming that the range of the coordinate chart is an ordinary Euclidean space with the usual Euclidean metric. The coordinate basis in that chart is orthonormal. Under these conditions we can usefully use words like “length,” “area,” and “volume” in the coordinate space.

```

(define-coordinates (up x y z) R3-rect)
;; closure

(def u (+ (* 'u↑0 d:dx) (* 'u↑1 d:dy)))

(def v (+ (* 'v↑0 d:dx) (* 'v↑1 d:dy)))

(((wedge dx dy) u v) R3-rect-point)
;; ⇒ (+ (* u↑0 v↑1) (* -1 v↑0 u↑1))

```

If we use cylindrical coordinates and define cylindrical vector fields we get the analogous answer in cylindrical coordinates:

```

(define-coordinates (up r theta z) R3-cyl)
;; closure

(def a (+ (* 'a↑0 d:dr) (* 'a↑1 d:dtheta)))

(def b (+ (* 'b↑0 d:dr) (* 'b↑1 d:dtheta)))

(((wedge dr dtheta) a b) ((point R3-cyl) (up 'r0 'theta0 'z0)))
;; ⇒ (+ (* a↑0 b↑1) (* -1 b↑0 a↑1))

```

The moral of this story is that this is the area of the parallelogram in the coordinate plane. It is not the area of the manifold!

There is a similar story with volumes. The wedge product of the elements of the coordinate basis is a three-form that measures our usual idea of coordinate volumes in  $R^3$  with a Euclidean metric:

```

(def u (+ (* 'u↑0 d:dx) (* 'u↑1 d:dy) (* 'u↑2 d:dz)))
;; closure

(def v (+ (* 'v↑0 d:dx) (* 'v↑1 d:dy) (* 'v↑2 d:dz)))

(def w (+ (* 'w↑0 d:dx) (* 'w↑1 d:dy) (* 'w↑2 d:dz)))

(simplify (((wedge dx dy dz) u v w) R3-rect-point))
;; ⇒ (+ (* u↑0 v↑1 w↑2)
;;      (* -1 u↑0 v↑2 w↑1)
;;      (* -1 u↑1 v↑0 w↑2)
;;      (* u↑1 v↑2 w↑0)
;;      (* u↑2 v↑0 w↑1)
;;      (* -1 u↑2 v↑1 w↑0))

```

This last expression is the determinant of a  $3 \times 3$  matrix:

```

(simplify (- (((wedge dx dy dz) u v w) R3-rect-point)
  (determinant
    (matrix-by-rows (list 'u↑0 'u↑1 'u↑2) (list 'v↑0 'v↑1 'v↑2) (list 'w↑0
'w↑1 'w↑2)))))
;; ⇒ 0

```

If we did the same operations in cylindrical coordinates we would get the analogous formula, showing that what we are computing is volume in the coordinate space, not volume on the manifold.

Because of antisymmetry, if the rank of a form is greater than the dimension of the manifold then the form is identically zero. The  $k$ -forms on an  $n$ -dimensional manifold form a module of dimension  $\binom{n}{k}$ . We can write a coordinate-basis expression for a  $k$ -form as

$$\omega = \sum_{i_0, \dots, i_{k-1}}^n \omega_{i_0, \dots, i_{k-1}} dx^{i_0} \wedge \dots \wedge dx^{i_{k-1}}. \quad (5.20)$$

The antisymmetry of the wedge product implies that

$$\omega_{i_{\sigma(0)}, \dots, i_{\sigma(k-1)}} = \text{Parity}(\sigma) \omega_{i_0, \dots, i_{k-1}}, \quad (5.21)$$

from which we see that there are only  $\binom{n}{k}$  independent components of  $\omega$ .

### 5.4 Exercise 5.1: Wedge Product

Pick a coordinate system and use the computer to verify that

- the wedge product is associative for forms in your coordinate system
- formula (5.17) is true in your coordinate system.

### 5.5 Exterior Derivative

The intention of introducing the exterior derivative is to capture all of the classical theorems of “vector analysis” into one unified Stokes's Theorem, which asserts that the integral of a form on the boundary of a manifold is the integral of the exterior derivative of the form on the interior of the manifold:<sup>4</sup>

$$\int_{\partial M} \omega = \int_M d\omega. \quad (5.22)$$

As we have seen in equation (3.34), the differential of a function on a manifold is a one-form field. If a function on a manifold is considered to be a form field of rank zero,<sup>5</sup> then the differential operator increases the rank of the form by one. We can generalize this to  $k$ -form fields with the exterior derivative operation.

Consider a one-form  $\omega$ . We define<sup>6</sup>

$$d\omega(v_1, v_2) = v_1(\omega(v_2)) - v_2(\omega(v_1)) - \omega([v_1, v_2]). \quad (5.23)$$

More generally, the exterior derivative of a  $k$ -form field is a  $k+1$ -form field, given by:<sup>7</sup>

$$\begin{aligned} d\omega(v_0, \dots, v_k) = & \sum_{i=0}^k \left\{ \left( (-1)^i v_i(\omega(v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_k)) \right. \right. \\ & \left. \left. + \sum_{j=i+1}^k (-1)^{i+j} \omega([v_i, v_j], v_0, \dots, v_{i-1}, v_{i+1}, \dots, v_{j-1}, v_{j+1}, \dots, v_k) \right) \right\}. \end{aligned} \quad (5.24)$$

This formula is coordinate-system independent. This is the way we compute the exterior derivative in our software.

If the form field  $\omega$  is represented in a coordinate basis

$$\omega = \sum_{i_0=0, \dots, i_{k-1}=0}^{n-1} a_{i_0, \dots, i_{k-1}} dx^{i_0} \wedge \dots \wedge dx^{i_{k-1}} \quad (5.25)$$

then the exterior derivative can be expressed as

$$d\omega = \sum_{i_0=0, \dots, i_{k-1}=0}^{n-1} da_{i_0, \dots, i_{k-1}} dx^{i_0} \wedge \dots \wedge dx^{i_{k-1}}. \quad (5.26)$$

Though this formula is expressed in terms of a coordinate basis, the result is independent of the choice of coordinate system.

<sup>4</sup>This is a generalization of the Fundamental Theorem of Calculus.

<sup>5</sup>A manifold function  $f$  induces a form field  $\hat{f}$  of rank 0 as follows:

$$\hat{f}()(m) = f(m).$$

<sup>6</sup>The definition is chosen to make Stokes's Theorem pretty.

<sup>7</sup>See Spivak, *Differential Geometry*, Volume 1, p.289.

## 5.6 Computing Exterior Derivatives

We can test that the computation indicated by equation (5.24) is equivalent to the computation indicated by equation (5.26) in three dimensions with a general one-form field:

```
(def a (literal-manifold-function 'alpha R3-rect))
(def b (literal-manifold-function 'beta R3-rect))
(def c (literal-manifold-function 'gamma R3-rect))
(def theta (+ (* a dx) (* b dy) (* c dz)))
```

The test will require two arbitrary vector fields

```
(def X (literal-vector-field 'X-rect R3-rect))
(def Y (literal-vector-field 'Y-rect R3-rect))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((- (d theta) (+ (wedge (d a) dx) (wedge (d b) dy) (wedge (d c) dz))) X Y)
            R3-rect-point))
;; => 0
```

We can also try a general two-form field in 3-dimensional space:

Let

$$\omega = a dy \wedge dz + b dz \wedge dx + c dx \wedge dy, \quad (5.27)$$

where  $a = \alpha^\wedge \chi$ ,  $b = \beta^\wedge \chi$ ,  $c = \gamma^\wedge \chi$ , and  $\alpha$ ,  $\beta$ , and  $\gamma$  are real-valued functions of three real arguments. As a program,

```
(def omega (+ (* a (wedge dy dz)) (* b (wedge dz dx)) (* c (wedge dx dy))))
```

Here we need another vector field because our result will be a three-form field.

```
(def Z (literal-vector-field 'Z-rect R3-rect))

(simplify (((- (d omega) (+ (wedge (d a) dy dz) (wedge (d b) dz dx) (wedge (d c) dx
dy))) X Y Z)
            R3-rect-point))
;; => 0
```

## 5.7 Properties of Exterior Derivatives

The exterior derivative of the wedge of two form fields obeys the graded Leibniz rule. It can be written in terms of the exterior derivatives of the component form fields:

$$d(\omega \wedge \tau) = d\omega \wedge \tau + (-1)^k \omega \wedge d\tau, \quad (5.28)$$

where  $k$  is the rank of  $\omega$ .

A form field  $\omega$  that is the exterior derivative of another form field  $\omega = d\theta$  is called exact. A form field whose exterior derivative is zero is called closed.

Every exact form field is a closed form field: applying the exterior derivative operator twice always yields zero:

$$d^2\omega = 0 \quad (5.29)$$

This is equivalent to the statement that partial derivatives with respect to different variables commute.<sup>8</sup>

<sup>8</sup>See Spivak, *Calculus on Manifolds*, p.92

It is easy to show equation (5.29) for manifold functions:

$$\begin{aligned}
 d^2f(u, v) &= d(df)(u, v) \\
 &= u(df(v)) - v(df(u)) - df([u, v]) \\
 &= u(v(f)) - v(u(f)) - [u, v](f) \\
 &= 0
 \end{aligned} \tag{5.30}$$

Consider the general one-form field  $\theta$  defined on 3-dimensional rectangular space. Taking two exterior derivatives of  $\theta$  yields a three-form field. It is zero:

```
// scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (((d (d theta)) X Y Z) R3-rect-point))
// => 0
```

Not every closed form field is an exact form field. Whether a closed form field is exact depends on the topology of a manifold.

## 5.8 Stokes's Theorem

The proof of the general Stokes's Theorem for  $n$ -dimensional orientable manifolds is quite complicated, but it is easy to see how it works for a 2-dimensional region  $M$  that can be covered with a single coordinate patch.<sup>9</sup>

Given a coordinate chart  $\chi(m) = (x(m), y(m))$  we can obtain a pair of coordinate-basis vectors  $\partial/\partial x = X_0$  and  $\partial/\partial y = X_1$ .

The coordinate image of  $M$  can be divided into small rectangular areas in the  $(x, y)$  coordinate plane. The union of the rectangular areas gives the coordinate image of  $M$ . The clockwise integrals around the boundaries of the rectangles cancel on neighboring rectangles, because the boundary is traversed in opposite directions. But on the boundary of the coordinate image of  $M$  the boundary integrals do not cancel, yielding an integral on the boundary of  $M$ . Area integrals over the rectangular areas add to produce an integral over the entire coordinate image of  $M$ .

So, consider Stokes's Theorem on a small patch  $P$  of the manifold for which the coordinates form a rectangular region ( $x_{\min} < x < x_{\max}$  and  $y_{\min} < y < y_{\max}$ ). Stokes's Theorem on  $P$  states

$$\int_{\partial P} \omega = \int_P d\omega. \tag{5.31}$$

The area integral on the right can be written as an ordinary multidimensional integral using the coordinate basis vectors (recall that the integral is independent of the choice of coordinates):

$$\int_{\chi(P)} d\omega(\partial/\partial x, \partial/\partial y) \circ \chi^{-1} = \int_{x_{\min}}^{x_{\max}} \int_{y_{\min}}^{y_{\max}} (\partial/\partial x(\omega(\partial/\partial y)) - \partial/\partial y(\omega(\partial/\partial x))) \circ \chi^{-1}. \tag{5.32}$$

We have used equation (5.23) to expand the exterior derivative.

Consider just the first term of the right-hand side of equation (5.32). Then using the definition of basis vector field  $\partial/\partial x$  we obtain

$$\begin{aligned}
 \int_{x_{\min}}^{x_{\max}} \int_{y_{\min}}^{y_{\max}} (\partial/\partial x(\omega(\partial/\partial y)) \circ \chi^{-1}) &= \int_{x_{\min}}^{x_{\max}} \int_{y_{\min}}^{y_{\max}} (X_0(\omega(\partial/\partial y)) \circ \chi^{-1}) \\
 &= \int_{x_{\min}}^{x_{\max}} \int_{y_{\min}}^{y_{\max}} \partial_0((\omega(\partial/\partial y)) \circ \chi^{-1}).
 \end{aligned} \tag{5.33}$$

This integral can now be evaluated using the Fundamental Theorem of Calculus. Accumulating the results for both integrals

<sup>9</sup>We do not develop the machinery for integration on chains that is usually needed for a full proof of Stokes's Theorem. This is adequately done in other books. A beautiful treatment can be found in Spivak, *Calculus on Manifolds* [17].

$$\begin{aligned}
\int_{\chi(P)} d\omega(\partial/\partial x, \partial/\partial y) \circ \chi^{-1} &= \int_{x_{\min}}^{x_{\max}} ((\omega(\partial/\partial x)) \circ \chi^{-1})(x, y_{\min}) dx \\
&\quad + \int_{y_{\min}}^{y_{\max}} ((\omega(\partial/\partial y)) \circ \chi^{-1})(x_{\max}, y) dy \\
&\quad - \int_{x_{\min}}^{x_{\max}} ((\omega(\partial/\partial x)) \circ \chi^{-1})(x, y_{\max}) dx \\
&\quad - \int_{y_{\min}}^{y_{\max}} ((\omega(\partial/\partial y)) \circ \chi^{-1})(x_{\min}, y) dy \\
&= \int_{\partial P} \omega,
\end{aligned} \tag{5.34}$$

as was to be shown.

## 5.9 Vector Integral Theorems

Green's Theorem states that for an arbitrary compact set  $M \subset \mathbb{R}^2$ , a 2-dimensional Euclidean space:

$$\int_{\partial M} ((\alpha \circ \chi) dx + (\beta \circ \chi) dy) = \int_M ((\partial_0 \beta - \partial_1 \alpha) \circ \chi) dx \wedge dy. \tag{5.35}$$

We can test this. By Stokes's Theorem, the integrands are related by an exterior derivative. We need some vectors to test our forms:

```
(def v (literal-vector-field 'v-rect R2-rect)) closure
(def w (literal-vector-field 'w-rect R2-rect))
```

We can now test our integrands:<sup>10</sup>

```
(def alpha (literal-function 'alpha R2→R)) closure
(def beta (literal-function 'beta R2→R))

(simplify
  (let [dx (ref (basis→oneform-basis R2-rect-basis) 0)
        dy (ref (basis→oneform-basis R2-rect-basis) 1)]
    (((- (d (+ (* (compose alpha (chart R2-rect)) dx) (* (compose beta (chart R2-rect)) dy)))
      (* (compose (- ((partial 0) beta) ((partial 1) alpha)) (chart R2-rect))
      (wedge dx dy)))
      v
      w)
      R2-rect-point)))
;; ⇒ 0
```

We can also compute the integrands for the Divergence Theorem: For an arbitrary compact set  $M \subset \mathbb{R}^3$  and a vector field  $w$

$$\int_M \operatorname{div}(w) dV = \int_{\partial M} w \cdot n dA \tag{5.36}$$

where  $n$  is the outward-pointing normal to the surface  $\partial M$ . Again, the integrands should be related by an exterior derivative, if this is an instance of Stokes's Theorem.

<sup>10</sup>Using `(define R2-rect-basis (coordinate-system→basis R2-rect))`.

Here we extract  $dx$  and  $dy$  from `R2-rect-basis` to avoid globally installing coordinates.

Note that even the statement of this theorem cannot be made with the machinery we have developed at this point. The concepts “outward-pointing normal,” area  $A$ , and volume  $V$  on the manifold are not definable without using a metric

(see Chapter 9). However, for orthonormal rectangular coordinates in  $\mathbb{R}^3$  we can interpret the integrands in terms of forms.

Let the vector field describing the flow of stuff be

$$\mathbf{w} = a \frac{\partial}{\partial x} + b \frac{\partial}{\partial y} + c \frac{\partial}{\partial z}. \quad (5.37)$$

The rate of leakage of stuff through each element of the boundary is  $\mathbf{w} \cdot \mathbf{n} dA$ . We interpret this as the two-form

$$a \, dy \wedge dz + b \, dz \wedge dx + c \, dx \wedge dy, \quad (5.38)$$

because any part of the boundary will have  $y - z$ ,  $z - x$ , and  $x - y$  components, and each such component will pick up contributions from the normal component of the flux  $\mathbf{w}$ . Formalizing this as code we have

```
(def a (literal-manifold-function 'a-rect R3-rect)) closure
(def b (literal-manifold-function 'b-rect R3-rect))
(def c (literal-manifold-function 'c-rect R3-rect))

(def flux-through-boundary-element (+ (* a (wedge dy dz)) (* b (wedge dz dx)) (* c
(wedge dx dy))))
```

The rate of production of stuff in each element of volume is  $\text{div}(\mathbf{w})dV$ . We interpret this as the three-form

$$\left( \frac{\partial}{\partial x} a + \frac{\partial}{\partial y} b + \frac{\partial}{\partial z} c \right) dx \wedge dy \wedge dz. \quad (5.39)$$

or:

```
(def production-in-volume-element (* (+ (d:dx a) (d:dy b) (d:dz c)) (wedge dx dy dz))) closure
```

Assuming Stokes's Theorem, the exterior derivative of the leakage of stuff per unit area through the boundary must be the rate of production of stuff per unit volume in the interior. We check this by applying the difference to arbitrary vector fields at an arbitrary point:

```
(def X (literal-vector-field 'X-rect R3-rect)) closure
(def Y (literal-vector-field 'Y-rect R3-rect))
(def Z (literal-vector-field 'Z-rect R3-rect))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((- production-in-volume-element (d flux-through-boundary-element)) X Y Z)
  R3-rect-point))

;; => 0
```

as expected.

## 5.10 Exercise 5.2: Graded Formula

Derive equation (5.28).

## 5.11 Exercise 5.3: Iterated Exterior Derivative

We have shown that the equation (5.29) is true for manifold functions. Show that it is true for any form field.



## 6 Over a Map


To deal with motion on manifolds we need to think about paths on manifolds and vectors along these paths. Tangent vectors along paths are not vector fields on the manifold because they are defined only on the path. And the path may even cross itself, which would give more than one vector at a point. Here we introduce the concept of a *vector field over a map*.<sup>1</sup> A vector field over a map assigns a vector to each image point of the map. In general the map may be a function from one manifold to another. If the domain of the map is the manifold of the real line, the range of the map is a 1-dimensional path on the target manifold. One possible way to define a vector field over a map is to assign a tangent vector to each image point of a path, allowing us to work with tangent vectors to paths. A *one-form field over the map* allows us to extract the components of a vector field over the map.

### 6.1 Vector Fields Over a Map

Let  $\mu$  be a map from points  $n$  in the manifold  $N$  to points  $m$  in the manifold  $M$ . A vector over the map  $\mu$  takes directional derivatives of functions on  $M$  at points  $m = \mu(n)$ . The vector over the map applied to the function on  $M$  is a function on  $N$ .

### 6.2 Restricted Vector Fields

One way to make a vector field over a map is to restrict a vector field on  $M$  to the image of  $N$  over  $\mu$ , as illustrated in figure 6.1.



**Figure 6.1:** The vector field  $v$  on  $M$  is indicated by arrows. The solid arrows are  $v_\mu$ , the restricted vector field over the map  $\mu$ . The vector field over the map is restricted to the image of  $N$  in  $M$ .

Let  $v$  be a vector field on  $M$ , and  $f$  a function on  $M$ . Then

$$v_\mu(f) = v(f) \circ \mu, \quad (6.1)$$

is a vector over the map  $\mu$ . Note that  $v_\mu(f)$  is a function on  $N$ , not  $M$ :

$$v_\mu(f)(n) = v(f)(\mu(n)). \quad (6.2)$$

We can implement this definition as:

```
(defn vector-field→vector-field-over-map                                clojure
  [mu:N→M]
  (fn [v-on-M] (procedure→vector-field (fn [f-on-M] (compose (v-on-M f-on-M) mu:N-
    >M))))))
```

### 6.3 Differential of a Map

Another way to construct a vector field over a map  $\mu$  is to transport a vector field from the source manifold  $N$  to the target manifold  $M$  with the *differential* of the map

$$d\mu(v)(f)(n) = v(f \circ \mu)(n), \quad (6.3)$$

<sup>1</sup>See Bishop and Goldberg, *Tensor Analysis on Manifolds* [3].

which takes its argument in the source manifold  $N$ . The differential of a map  $\mu$  applied to a vector field  $v$  on  $N$  is a vector field over the map. A procedure to compute the differential is:

```
(defn differential [mu] (fn [v] (fn [f] (v (compose f mu)))))) closure
```

The nomenclature of this subject is confused. The “differential of a map between manifolds,”  $d\mu$ , takes one more argument than the “differential of a real-valued function on a manifold,”  $df$ , but when the target manifold of  $\mu$  is the reals and  $I$  is the identity function on the reals,

$$d\mu(v)(I)(n) = (v(I \circ \mu))(n) = (v(\mu))(n) = d\mu(v)(n). \quad (6.4)$$

We avoid this problem in our notation by distinguishing  $d$  and  $d$ . In our programs we encode  $d$  as differential and  $d$  as d.

## 6.4 Velocity at a Time

Let  $\mu$  be the map from the time line to the manifold  $M$ , and  $\partial/\partial t$  be a basis vector on the time line. Then  $d\mu(\partial/\partial t)$  is the vector over the map  $\mu$  that computes the rate of change of functions on  $M$  along the path that is the image of  $\mu$ . This is the velocity vector. We can use the differential to assign a velocity vector to each moment, solving the problem of multiple vectors at a point if the path crosses itself.

## 6.5 One-Form Fields Over a Map

Given a one-form  $\omega$  on the manifold  $M$ , the one-form over the map  $\mu : N \rightarrow M$  is constructed as follows:

$$\omega^\mu(v_\mu)(n) = \omega(u)(\mu(n)), \text{ where } u(f)(m) = v_\mu(f)(n). \quad (6.5)$$

The object  $u$  is not really a vector field on  $M$  even though we have given it that shape so that the dual vector can apply to it;  $u(f)$  is evaluated only at images  $m = \mu(n)$  of points  $n$  in  $N$ . If we were defining  $u$  as a vector field we would need the inverse of  $\mu$  to find the point  $n = \mu^{-1}(m)$ , but this is not required to define the object  $u$  in a context where there is already an  $m$  associated with the  $n$  of interest. To extend this idea to  $k$ -forms, we carry each vector argument over the map.

The procedure that constructs a  $k$ -form over the map from a  $k$ -form is:

```
(defn form-field→form-field-over-map closure
  [mu:N→M]
  (fn [w-on-M]
    (letfn [(make-fake-vector-field [V-over-mu n]
              (letfn [(u [f] (fn [m] ((V-over-mu f) n))))
                (procedure→vector-field u)))]
      (procedure→nform-field (fn [& vectors-over-map]
                                (fn [n]
                                  ((apply w-on-M
                                        (map (fn [V-over-mu] (make-fake-vector-field V-
over-mu n))
                                              vectors-over-map))
                                   (mu:N→M n))))
                              (get-rank w-on-M))))))
```

The internal procedure `make-fake-vector-field` counterfeits a vector field  $u$  on  $M$  from the vector field over the map  $\mu : N \rightarrow M$ . This works here because the only value that is ever passed as  $m$  is  $(\mu:N \rightarrow M \ n)$ .

## 6.6 Basis Fields Over a Map

Let  $e$  be a tuple of basis vector fields, and  $\tilde{e}$  be the tuple of basis one-forms that is dual to  $e$ :

$$\tilde{e}^i(e_j)(m) = \delta_j^i. \quad (6.6)$$

The *basis vectors* over the map,  $e^\mu$ , are particular cases of vectors over a map:

$$e^\mu(f) = e(f) \circ \mu. \quad (6.7)$$

And the elements of the *dual basis over the map*,  $\tilde{e}_\mu$ , are particular cases of one-forms over the map. The basis and dual basis over the map satisfy

$$\tilde{e}_\mu^i(e_j^\mu)(n) = \delta_j^i. \quad (6.8)$$

## 6.7 Walking on a Sphere

For example, let  $\mu$  map the time line to the unit sphere.<sup>2</sup> We use colatitude  $\theta$  and longitude  $\phi$  as coordinates on the sphere:

```
(def S2 (make-manifold S2-type 2 3)) closure

(def S2-spherical (coordinate-system-at S2 :spherical :north-pole))

(define-coordinates (up theta phi) S2-spherical)

(def S2-basis (coordinate-system→basis S2-spherical))
```

A general path on the sphere is:<sup>3</sup>

```
(def mu closure
  (compose (point S2-spherical)
            (up (literal-function 'theta) (literal-function 'phi))
            (chart R1-rect)))
```

The basis over the map is constructed from the basis on the sphere:

```
(def S2-basis-over-mu (basis→basis-over-map mu S2-basis)) closure

(def h (literal-manifold-function 'h-spherical S2-spherical))

(((basis→vector-basis S2-basis-over-mu) h) ((point R1-rect) 't0))
;; ⇒ (down ((partial 0) h-spherical) (up (theta t0) (phi t0)))
;;      ((partial 1) h-spherical) (up (theta t0) (phi t0)))
```

The basis vectors over the map compute derivatives of the function  $h$  evaluated on the path at the given time.

We can check that the dual basis over the map does the correct thing:

```
((basis→oneform-basis S2-basis-over-mu) (basis→vector-basis S2-basis-over-mu)) closure
((point R1-rect) 't0)
;; ⇒ (up (down 1 0) (down 0 1))
```

## 6.8 Components of the Velocity

Let  $\chi$  be a tuple of coordinates on  $M$ , with associated basis vectors  $X_i$ , and dual basis elements  $dx^i$ . The vector basis and dual basis over the map  $\mu$  are  $X_i^\mu$  and  $dx_\mu^i$ . The components of the velocity (rates of change of coordinates along the path  $\mu$ ) are obtained by applying the dual basis over the map to the velocity

$$v^i(t) = dx_\mu^i(d\mu(\partial/\partial t))(t), \quad (6.9)$$

where  $t$  is the coordinate for the point  $t$ .

For example, the coordinate velocities on a sphere are

<sup>2</sup>We execute `(define-coordinates t R1-rect)` to establish the real-line coordinate objects; the translated code names its coordinate vector field `d:dt`.

<sup>3</sup>We provide a shortcut to make literal manifold maps:

```
(def mu (literal-manifold-map 'mu R1-rect S2-spherical)) closure
```

```
((basis→oneform-basis S2-basis-over-mu) ((differential mu) d:dt)) ((point R1-rect) closure
't0))
;; ⇒ (up ((D theta) t0) ((D phi) t0))
```

as expected.

## 6.9 Pullbacks and Pushforwards

Maps from one manifold to another can also be used to relate the vector fields and one-form fields on one manifold to those on the other. We have introduced two such relations: restricted vector fields and the differential of a function. However, there are other ways to relate the vector fields and form fields on different manifolds that are connected by a map.

## 6.10 Pullback and Pushforward of a Function

The *pullback* of a function  $f$  on  $M$  over the map  $\mu$  is defined as

$$\mu^*f = f \circ \mu. \quad (6.10)$$

This allows us to take a function defined on  $M$  and use it to define a new function on  $N$ .

For example, the integral curve of  $v$  evolved for time  $t$  as a function of the initial manifold point  $m$  generates a map  $\phi_t^v$  of the manifold onto itself. This is a simple currying<sup>4</sup> of the integral curve of  $v$  from  $m$  as a function of time:  $\phi_t^v(m) = \gamma_m^v(t)$ . The evolution of the function  $f$  along an integral curve, equation (3.33), can be written in terms of the pullback over  $\phi_t^v$ :

$$(E_{t,v}f)(m) = f(\phi_t^v(m)) = ((\phi_t^v)^*f)(m). \quad (6.11)$$

This is implemented as:

```
(defn pullback-function [mu:N→M] (fn [f-on-M] (compose f-on-M mu:N→M)))
```

A vector field over the map that was constructed by restriction (equation (6.1)) can be seen as the pullback of the function constructed by application of the vector field to a function:

$$v_\mu(f) = v(f) \circ \mu = \mu^*(v(f)). \quad (6.12)$$

A vector field over the map that was constructed by a differential (equation (6.3)) can be seen as the vector field applied to the pullback of the function:

$$d\mu(v)(f)(n) = v(f \circ \mu)(n) = v(\mu^*f)(n). \quad (6.13)$$

If we have an inverse for the map  $\mu$  we can also define a *push-forward* of the function  $g$ , defined on the source manifold of the map:<sup>5</sup>

$$\mu_*g = g \circ \mu^{-1}. \quad (6.14)$$

## 6.11 Pushforward of a Vector Field

We can also define the *pushforward* of a vector field over the map  $\mu$ . The pushforward takes a vector field  $v$  defined on  $N$ . The result takes directional derivatives of functions on  $M$  at a place determined by a point in  $M$ :

$$\mu_*v(f)(m) = v(\mu^*f)(\mu^{-1}(m)) = v(f \circ \mu)(\mu^{-1}(m)), \quad (6.15)$$

or

$$\mu_*v(f) = \mu_*(v(\mu^*f)). \quad (6.16)$$

<sup>4</sup>A function of two arguments may be seen as a function of one argument whose value is a function of the other argument. This can be done in two different ways, depending on which argument is supplied first. The general process of specifying a subset of the arguments to produce a new function of the others is called *currying* the function, in honor of the logician Haskell Curry (1900-1982) who, with Moses Schönfinkel (1889-1942), developed combinatory logic.

<sup>5</sup>Notation note: superscript asterisk indicates pullback, subscript asterisk indicates pushforward. Pullbacks and pushforwards are tightly binding operators, so, for example  $\mu^*f(n) = (\mu^*f)(n)$ .

Here we expressed the pushforward of the vector field in terms of pullbacks and pushforwards of functions. Note that the pushforward requires the inverse of the map.

If the map is from time to some configuration manifold and represents the time evolution of a process, we can think of the pushforward of a vector field as a velocity measured at a point on the trajectory in the configuration manifold. By contrast, the differential of the map applied to the vector field gives us the velocity vector at each moment in time. Because a trajectory may cross itself, the pushforward is not defined at any point where the crossing occurs, but the differential is always defined.

## 6.12 Pushforward Along Integral Curves

We can push a vector field forward over the map generated by an integral curve of a vector field  $\mathbf{w}$ , because the inverse is always available.<sup>6</sup>

$$((\phi_t^{\mathbf{w}})_* \mathbf{v})(f)(m) = \mathbf{v}((\phi_t^{\mathbf{w}})^* f)(\phi_{-t}^{\mathbf{w}}(m)) = \mathbf{v}(f \circ \phi_t^{\mathbf{w}})(\phi_{-t}^{\mathbf{w}}(m)). \quad (6.17)$$

This is implemented as:

```
(defn pushforward-vector                                     clojure
  [mu:N→M mu-inverse:M→N]
  (fn [v-on-N]
    (procedure→vector-field (fn [f] (compose (v-on-N (compose f mu:N→M)) mu-
                                              inverse:M→N))))))
```

## 6.13 Pullback of a Vector Field

Given a vector field  $\mathbf{v}$  on a manifold  $M$  we can pull the vector field back through the map  $\mu : N \rightarrow M$  as follows:

$$\mu^* \mathbf{v}(f)(n) = (\mathbf{v}(f \circ \mu^{-1}))(\mu(n)) \quad (6.18)$$

or

$$\mu^* \mathbf{v}(f) = \mu^*(\mathbf{v}(\mu_* f)). \quad (6.19)$$

This may be useful when the map is invertible, as in the flow generated by a vector field.

This is implemented as:

```
(defn pullback-vector-field [mu:N→M mu-inverse:M→N] (pushforward-vector mu-
  inverse:M→N mu:N→M))                                     clojure
```

## 6.14 Pullback of a Form Field

We can also pull back a one-form field  $\omega$  defined on  $M$ , but an honest definition is rarely written. The pullback of a one-form field applied to a vector field is intended to be the same as the one-form field applied to the pushforward of the vector field.

The pullback of a one-form field is often described by the relation

$$\mu^* \omega(\mathbf{v}) = \omega(\mu_* \mathbf{v}), \quad (6.20)$$

but this is wrong, because the two sides are not functions of points in the same manifold. The one-form field  $\omega$  applies to a vector field on the manifold  $M$ , which takes a directional derivative of a function defined on  $M$  and is evaluated at a point on  $M$ , but the left-hand side is evaluated at a point on the manifold  $N$ .

A more precise description would be

$$\mu^* \omega(\mathbf{v})(n) = \omega(\mu_* \mathbf{v})(\mu(n)) \quad (6.21)$$

or

$$\mu^* \omega(\mathbf{v}) = \mu^*(\omega(\mu_* \mathbf{v})). \quad (6.22)$$

<sup>6</sup>The map  $\phi_t^{\mathbf{w}}$  is always invertible:  $(\phi_t^{\mathbf{w}})^{-1} = \phi_{-t}^{\mathbf{w}}$  because of the uniqueness of the solutions of the initial-value problem for ordinary differential equations.

Although this is accurate, it may not be effective, because computing the pushforward requires the inverse of the map  $\mu$ . But the inverse is available when the map is the flow generated by a vector field.

In fact it is possible to compute the pullback of a one-form field without having the inverse of the map. Instead we can use `form-field→form-field-over-map` to avoid needing the inverse:

$$\mu^*\omega(v)(n) = \omega^\mu(d\mu(v))(n). \quad (6.23)$$

The pullback of a  $k$ -form generalizes equation (6.21):

$$\mu^*\omega(u, v, \dots)(n) = \omega(\mu_*, u, \mu_*, v, \dots)(\mu(n)). \quad (6.24)$$

This is implemented as follows:<sup>7</sup>

```
(defn pullback-form
  [mu:N→M]
  (fn [omega-on-M]
    (let [k (get-rank omega-on-M)]
      (if (= k 0)
        ((pullback-function mu:N→M) omega-on-M)
        (procedure→nform-field (fn [& vectors-on-N]
                                   (apply ((form-field→form-field-over-map mu:N→M)
                                           (map (differential mu:N→M) vectors-on-N)))
                                   k))))))
  omega-on-M)
```

## 6.15 Properties of Pullback

The pullback through a map has many nice properties: it distributes through addition and through wedge product:

$$\mu^*(\theta + \phi) = \mu^*\theta + \mu^*\phi, \quad (6.25)$$

$$\mu^*(\theta \wedge \phi) = \mu^*\theta \wedge \mu^*\phi. \quad (6.26)$$

The pullback also commutes with the exterior derivative:

$$d(\mu^*\theta) = \mu^*(d\theta), \quad (6.27)$$

for  $\theta$  a function or  $k$ -form field.

We can verify this by computing an example. Let  $\mu$  map the rectangular plane to rectangular 3-space:

```
(def mu (literal-manifold-map 'MU R2-rect R3-rect))
```

First, let's compare the pullback of the exterior derivative of a function with the exterior derivative of the pullback of the function:

```
(def f (literal-manifold-function 'f-rect R3-rect))
(def X (literal-vector-field 'X-rect R2-rect))
(((pullback mu) (d f)) (d ((pullback mu) f))) X ((point R2-rect) (up 'x0 'y0)))
;; => 0
```

More generally, we can consider what happens to a form field. For a one-form field the result is as expected:

```
(def theta (literal-oneform-field 'THETA R3-rect))
(def Y (literal-vector-field 'Y-rect R2-rect))
```

<sup>7</sup>There is a generic pullback procedure that operates on any kind of manifold object. However, to pull a vector field back requires providing the inverse map.

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((- ((pullback mu) (d theta)) (d ((pullback mu) theta))) X Y)
  ((point R2-rect) (up 'x0 'y0))))
;; => 0
```

## 6.16 Pushforward of a Form Field

By symmetry, it is possible to define the pushforward of a one-form field as

$$\mu_*\omega(v) = \mu_*(\omega(\mu^*v)), \quad (6.28)$$

but this is rarely useful.

### 6.16.1 Exercise 6.1: Velocities on a Globe

We can use manifold functions, vector fields, and one-forms over a map to understand how paths behave.

a. Suppose that a vehicle is traveling east on the Earth at a given rate of change of longitude. What is the actual ground speed of the vehicle?

b. Stereographic projection is useful for navigation because it is conformal (it preserves angles). For the situation of part a, what is the speed measured on a stereographic map? Remember that the stereographic projection is implemented with [S2-Riemann](#).

But if we used this shortcut, the component functions would be named  $\mu^0$  and  $\mu^1$ . Here we wanted to use more mnemonic names for the component functions.

## 7 Directional Derivatives

The vector field was a generalization of the directional derivative to functions on a manifold. When we want to generalize the directional derivative idea to operate on other manifold objects, such as directional derivatives of vector fields or of form fields, there are several useful choices. In the same way that a vector field applies to a function to produce a function, we will build directional derivatives so that when applied to any object it will produce another object of the same kind. All directional derivatives require a vector field to give the direction and scale factor.

We will have a choice of directional derivative operators that give different results for the rate of change of vector and form fields along integral curves. But all directional derivative operators must agree when computing rates of change of functions along integral curves. When applied to functions, all directional derivative operators give:

$$\mathcal{D}_v(f) = v(f). \quad (7.1)$$

Next we specify the directional derivative of a vector field  $u$  with respect to a vector field  $v$ . Let an integral curve of the vector field  $v$  be  $\gamma$ , parameterized by  $t$ , and let  $m = \gamma(t)$ . Let  $u'$  be a vector field that results from transporting the vector field  $u$  along  $\gamma$  for a parameter increment  $\delta$ . How  $u$  is transported to make  $u'$  determines the type of derivative. We formulate the method of transport by:

$$u' = F_\delta^v u. \quad (7.2)$$

We can assume without loss of generality that  $F_\delta^v u$  is a linear transformation over the reals on  $u$ , because we care about its behavior only in an incremental region around  $\delta = 0$ .

Let  $g$  be the comparison of the original vector field at a point with the transported vector field at that point:

$$g(\delta) = u(f)(m) - (F_\delta^v u)(f)(m). \quad (7.3)$$

So we can compute the directional derivative operator using only ordinary derivatives:

$$\mathcal{D}_v u(f)(m) = Dg(0). \quad (7.4)$$

The result  $\mathcal{D}_v u$  is of type vector field.

The general pattern of constructing a directional derivative operator from a transport operator is given by the following schema:<sup>1</sup>

```
(defn F→directional-derivative
  [F]
  (fn [v]
    (fn [u]
      (fn [f]
        (fn [m]
          (letfn [(g [delta] (- ((u f) m) (((F v) delta) u) f) m)))]
            ((D g) 0)))))))
```

clojure

The linearity of transport implies that

$$\mathcal{D}_v(\alpha O + \beta P) = \alpha \mathcal{D}_v O + \beta \mathcal{D}_v P, \quad (7.5)$$

for any real  $\alpha$  and  $\beta$  and manifold objects  $O$  and  $P$ .

The directional derivative obeys superposition in its vector-field argument:

$$\mathcal{D}_{v+w} = \mathcal{D}_v + \mathcal{D}_w. \quad (7.6)$$

The directional derivative is homogeneous over the reals in its vector-field argument:

$$\mathcal{D}_{\alpha v} = \alpha \mathcal{D}_v, \quad (7.7)$$

<sup>1</sup>The directional derivative of a vector field must itself be a vector field. Thus the real program for this must make the function of  $f$  into a vector field. However, we leave out this detail here to make the structure clear.



for any real  $\alpha$ .<sup>2</sup> This follows from the fact that for evolution along integral curves: when  $\alpha$  is a real number,

$$\phi_t^{\alpha v}(m) = \phi_{\alpha t}^v(m). \quad (7.8)$$

When applied to products of functions, directional derivative operators satisfy Leibniz's rule:

$$\mathcal{D}_v(fg) = f(\mathcal{D}_v g) + (\mathcal{D}_v f)g. \quad (7.9)$$

The Leibniz rule is extended to applications of one-form fields to vector fields:

$$\mathcal{D}_v(\omega(y)) = \omega(\mathcal{D}_v y) + (\mathcal{D}_v \omega)(y). \quad (7.10)$$

The extension of the Leibniz rule, combined with the choice of transport of a vector field, determines the action of the directional derivative on form fields.<sup>3</sup>

## 7.1 Lie Derivative

The Lie derivative is one kind of directional derivative operator. We write the Lie derivative operator with respect to a vector field  $v$  as  $\mathcal{L}_v$ .

## 7.2 Functions

The Lie derivative of the function  $f$  with respect to the vector field  $v$  is given by:

$$\mathcal{L}_v f = v(f). \quad (7.11)$$

The tangent vector  $v$  measures the rate of change of  $f$  along integral curves.

## 7.3 Vector Fields

For the Lie derivative of a vector field  $y$  with respect to a vector field  $v$  we choose the transport operator  $F_\delta^v y$  to be the pushforward of  $y$  along the integral curves of  $v$ . Recall equation (6.15). So the Lie derivative of  $y$  with respect to  $v$  at the point  $m$  is

$$(\mathcal{L}_v y)(f)(m) = Dg(0), \quad (7.12)$$

where

$$g(\delta) = y(f)(m) - ((\phi_\delta^v)_* y)(f)(m). \quad (7.13)$$

We can construct a procedure that computes the Lie derivative of a vector field by supplying an appropriate transport operator (F-Lie phi) for F in our schema F→directional-derivative. In this first stab at the Lie derivative, we introduce a coordinate system and we expand the integral curve to a given order. Because in the schema we evaluate the derivative of  $g$  at 0, the dependence on the order and the coordinate system disappears. They will not be needed in the final version.

```
(defn Lie-directional
  [coordsys order]
  (let [Phi (phi coordsys order)] (F→directional-derivative (F-Lie Phi))))

(defn F-Lie [phi] (fn [v] (fn [delta] (pushforward-vector ((phi v) delta) ((phi v) (-
delta))))))

(defn phi
  [coordsys order]
  (fn [v]
    (fn [delta]
      (fn [m] ((point coordsys) (series:sum (((exp (* delta v)) (chart coordsys)) m)
order))))))
```

clojure

Expand the quantities in equation (7.13) to first order in  $\delta$ :

<sup>2</sup>For some derivative operators  $\alpha$  can be a real-valued manifold function.

<sup>3</sup>The action on functions, vector fields, and one-form fields suffices to define the action on all tensor fields. See Appendix F.

$$\begin{aligned}
g(\delta) &= y(f)(m) - (\phi_{\delta*}^\vee y)(f)(m) \\
&= y(f)(m) - y(f \circ \phi_\delta^\vee)(\phi_{-\delta}^\vee(m)) \\
&= (y(f) - y(f + \delta v(f) + \dots) + \delta v(y(f + \delta v(f) + \dots)))(m) + \dots \\
&= (-\delta y(v(f)) + \delta v(y(f)))(m) + \dots \\
&= \delta[v, y](f)(m) + \mathcal{O}(\delta^2).
\end{aligned} \tag{7.14}$$

So the Lie derivative of a vector field  $y$  with respect to a vector field  $v$  is a vector field that is defined by its behavior when applied to an arbitrary manifold function  $f$ :

$$(\mathcal{L}_v y)(f) = [v, y](f) \tag{7.15}$$

Verifying this computation

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (let [v (literal-vector-field 'v-rect R3-rect)
               w (literal-vector-field 'w-rect R3-rect)
               f (literal-manifold-function 'f-rect R3-rect)]
  ((- (((Lie-directional R3-rect 2) v) w) f) ((commutator v w) f))
  ((point R3-rect) (up 'x0 'y0 'z0))))
;; => 0
```

Although this is tested to second order, evaluating the derivative at zero ensures that first order is enough. So we can safely define:

```
(defn Lie-derivative-vector [V] (fn [Y] (commutator V Y)))      clojure
```

We can think of the Lie derivative as the rate of change of the manifold function  $y(f)$  as we move in the  $v$  direction, adjusted to take into account that some of the variation is due to the variation of  $f$ :

$$\begin{aligned}
(\mathcal{L}_v y)(f) &= [v, y](f) \\
&= v(y(f)) - y(v(f)) \\
&= v(y(f)) - y(\mathcal{L}_v f).
\end{aligned} \tag{7.16}$$

The first term in the commutator,  $v(y(f))$ , measures the rate of change of the combination  $y(f)$  along the integral curves of  $v$ . The change in  $y(f)$  is due to both the intrinsic change in  $y$  along the curve and the change in  $f$  along the curve; the second term in the commutator subtracts this latter quantity. The result is the intrinsic change in  $y$  along the integral curves of  $v$ .

Additionally, we can extend the product rule, for any manifold function  $g$  and any vector field  $u$ :

$$\begin{aligned}
\mathcal{L}_v(gu)(f) &= [v, gu](f) \\
&= v(g)u(f) + g[v, u](f) \\
&= (\mathcal{L}_v g)u(f) + g(\mathcal{L}_v u)(f).
\end{aligned} \tag{7.17}$$

## 7.4 An Alternate View

We can write the vector field

$$y(f) = \sum_i y^i e_i(f). \tag{7.18}$$

By the extended product rule (equation (7.17)) we get

$$\mathcal{L}_v y(f) = \sum_i (v(y^i) e_i(f) + y^i \mathcal{L}_v e_i(f)). \tag{7.19}$$

Because the Lie derivative of a vector field is a vector field, we can extract the components of  $\mathcal{L}_v e_i$  using the dual basis. We define  $\Delta_j^i(v)$  to be those components:

$$\Delta_j^i(v) = \tilde{e}^i(\mathcal{L}_v e_j) = \tilde{e}^i([v, e_j]). \tag{7.20}$$

So the Lie derivative can be written

$$(\mathcal{L}_\nu y)(f) = \sum_i \left( \nu(y^i) + \sum_j \Delta_j^i(\nu) y^j \right) e_i(f). \quad (7.21)$$

The components of the Lie derivatives of the basis vector fields are the structure constants for the basis vector fields. (See equation (4.37).) The structure constants are antisymmetric in the lower indices:

$$\tilde{e}^i(\mathcal{L}_{e_k} e_j) = \tilde{e}^i([e_k, e_j]) = d_{kj}^i. \quad (7.22)$$

Resolving  $\nu$  into components and applying the product rule, we get

$$(\mathcal{L}_\nu y)(f) = \sum_k (\nu^k [e_k, y](f) - y(\nu^k) e_k(f)). \quad (7.23)$$

So  $\Delta_j^i$  is related to the structure constants by

$$\begin{aligned} \Delta_j^i(\nu) &= \tilde{e}^i(\mathcal{L}_\nu e_j) \\ &= \sum_k (\nu^k \tilde{e}^i([e_k, e_j]) - e_j(\nu^k) \tilde{e}^i(e_k)) \\ &= \sum_k (\nu^k d_{kj}^i - e_j(\nu^k) \delta_k^i) \\ &= \sum_k \nu^k d_{kj}^i - e_j(\nu^i). \end{aligned} \quad (7.24)$$

Note: Despite their appearance, the  $\Delta_j^i$  are not form fields because  $\Delta_j^i(f\nu) \neq f\Delta_j^i(\nu)$ .

## 7.5 Form Fields

We can also define the Lie derivative of a form field  $\omega$  with respect to the vector field  $\nu$  by its action on an arbitrary vector field  $y$ , using the extended Leibniz rule (see equation (7.10)):

$$(\mathcal{L}_\nu(\omega))(y) \equiv \nu(\omega(y)) - \omega(\mathcal{L}_\nu y). \quad (7.25)$$

The first term computes the rate of change of the combination  $\omega(y)$  along the integral curve of  $\nu$ , while the second subtracts  $\omega$  applied to the change in  $y$ . The result is the change in  $\omega$  along the curve.

The Lie derivative of a  $k$ -form field  $\omega$  with respect to a vector field  $\nu$  is a  $k$ -form field that is defined by its behavior when applied to  $k$  arbitrary vector fields  $w_0, \dots, w_{k-1}$ . We generalize equation (7.25):

$$\mathcal{L}_\nu \omega(w_0, \dots, w_{k-1}) = \nu(\omega(w_0, \dots, w_{k-1})) - \sum_{i=0}^{k-1} \omega(w_0, \dots, \mathcal{L}_\nu w_i, \dots, w_{k-1}). \quad (7.26)$$

## 7.6 Uniform Interpretation

Consider abstracting equations (7.16), (7.25), and (7.27). The Lie derivative of an object,  $a$ , that can apply to other objects,  $b$ , to produce manifold functions,  $a(b) : M \rightarrow R^n$ , is

$$(\mathcal{L}_\nu a)(b) = \nu(a(b)) - a(\mathcal{L}_\nu b). \quad (7.27)$$

The first term in this expression computes the rate of change of the compound object  $a(b)$  along integral curves of  $\nu$ , while the second subtracts the change in  $a$  due to the change in  $b$  along the curves. The result is a measure of the “intrinsic” change in  $a$  along integral curves of  $\nu$ , with  $b$  held “fixed.”

## 7.7 Properties of the Lie Derivative

As required by properties 7.7-7.5, the Lie derivative is linear in its arguments:

$$\mathcal{L}_{\alpha\nu + \beta w} = \alpha\mathcal{L}_\nu + \beta\mathcal{L}_w, \quad (7.28)$$

and

$$\mathcal{L}_\nu(\alpha a + \beta b) = \alpha\mathcal{L}_\nu a + \beta\mathcal{L}_\nu b, \quad (7.29)$$

with  $\alpha, \beta \in R$  and vector fields or one-form fields  $a$  and  $b$ .

For any  $k$ -form field  $\omega$  and any vector field  $\mathbf{v}$  the exterior derivative commutes with the Lie derivative with respect to the vector field:

$$\mathcal{L}_{\mathbf{v}}(d\omega) = d(\mathcal{L}_{\mathbf{v}}\omega). \quad (7.30)$$

If  $\omega$  is an element of surface then  $d\omega$  is an element of volume. The Lie derivative computes the rate of change of its argument under a deformation described by the vector field. The answer is the same whether we deform the surface before computing the volume or compute the volume and then deform it.

We can verify this in 3-dimensional rectangular space for a general one-form field:<sup>4</sup>

```
(simplify (((- ((Lie-derivative V) (d theta))) (d ((Lie-derivative V) theta))) X Y) R3-closure
rect-point))
;; => 0
```

and for the general two-form field:

```
(simplify (((- ((Lie-derivative V) (d omega))) (d ((Lie-derivative V) omega))) X Y Z) closure
R3-rect-point))
;; => 0
```

The Lie derivative satisfies another nice elementary relationship. If  $\mathbf{v}$  and  $\mathbf{w}$  are two vector fields, then

$$[\mathcal{L}_{\mathbf{v}}, \mathcal{L}_{\mathbf{w}}] = \mathcal{L}_{[\mathbf{v}, \mathbf{w}]}. \quad (7.31)$$

Again, for our general one-form field  $\theta$ :

```
(simplify
  ((((- (commutator (Lie-derivative X) (Lie-derivative Y)) (Lie-derivative (commutator
X Y))) theta)
    Z)
  R3-rect-point))
;; => 0
```

and for the two-form field  $\omega$ :

```
(simplify
  ((((- (commutator (Lie-derivative X) (Lie-derivative Y)) (Lie-derivative (commutator
X Y))) omega)
    Z
    V)
  R3-rect-point))
;; => 0
```

## 7.8 Exponentiating Lie Derivatives

The Lie derivative computes the rate of change of objects as they are advanced along integral curves. The Lie derivative of an object produces another object of the same type, so we can iterate Lie derivatives. This gives us Taylor series for objects along the curve.

The operator  $e^{t\mathcal{L}_{\mathbf{v}}} = 1 + t\mathcal{L}_{\mathbf{v}} + \frac{t^2}{2!}\mathcal{L}_{\mathbf{v}}^2 + \dots$  evolves objects along the curve by parameter  $t$ . For example, the exponential of a Lie derivative applied to a vector field is

---

<sup>4</sup>In these experiments we need some setup.

```
(def a (literal-manifold-function 'alpha R3-rect))
(def b (literal-manifold-function 'beta R3-rect))
(def c (literal-manifold-function 'gamma R3-rect))
```

$$e^{t\mathcal{L}_v}y = y + t\mathcal{L}_v y + \frac{t^2}{2}\mathcal{L}_v^2 y + \dots = y + t[v, y] + \frac{t^2}{2}[v, [v, y]] + \dots \quad (7.32)$$

Consider a simple case. We advanced the coordinate-basis vector field  $\partial/\partial y$  by an angle  $a$  around the circle. Let  $J_z = x\partial/\partial y - y\partial/\partial x$ , the circular vector field. We recall

```
(define-coordinates (up x y z) R3-rect) closure

(def Jz (- (* x d:dy) (* y d:dx)))
```

We can apply the exponential of the Lie derivative with respect to  $J_z$  to  $\partial/\partial y$ . We examine how the result affects a general function on the manifold:

```
(series:for-each print-expression closure
  (((exp (* 'a (Lie-derivative Jz))) d:dy)
   (literal-manifold-function 'f-rect R3-rect))
  ((point R3-rect) (up 1 0 0)))
5)
```

Apparently the result is

$$\exp\left(\alpha\mathcal{L}_{(x\partial/\partial y - y\partial/\partial x)}\right)\frac{\partial}{\partial y} = -\sin(a)\frac{\partial}{\partial x} + \cos(a)\frac{\partial}{\partial y}. \quad (7.33)$$

## 7.9 Interior Product

There is a simple but useful operation available between vector fields and form fields called *interior product*. This is the substitution of a vector field  $v$  into the first argument of a  $p$ -form field  $\omega$  to produce a  $p-1$ -form field:

$$(i_v\omega)(v_1, \dots, v_{p-1}) = \omega(v, v_1, \dots, v_{p-1}). \quad (7.34)$$

There is a mundane identity corresponding to the product rule for the Lie derivative of an interior product:

$$\mathcal{L}_v(i_y\omega) = i_{\mathcal{L}_v y}\omega + i_y(\mathcal{L}_v\omega). \quad (7.35)$$

And there is a rather nice identity for the Lie derivative in terms of the interior product and the exterior derivative, called *Cartan's formula*:

$$\mathcal{L}_v\omega = i_v(d\omega) + d(i_v\omega). \quad (7.36)$$

We can verify Cartan's formula in a simple case with a program:

```
(def X (literal-vector-field 'X-rect R3-rect)) closure

(def Y (literal-vector-field 'Y-rect R3-rect))

(def Z (literal-vector-field 'Z-rect R3-rect))

(def a (literal-manifold-function 'alpha R3-rect))

(def b (literal-manifold-function 'beta R3-rect))

(def c (literal-manifold-function 'gamma R3-rect))

(def omega (+ (* a (wedge dx dy)) (* b (wedge dy dz)) (* c (wedge dz dx))))

(defn L1 [X] (fn [omega] (+ ((interior-product X) (d omega)) (d ((interior-product X)
omega)))))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((- ((Lie-derivative X) omega) Y Z) (((L1 X) omega) Y Z)))
```

```
;; => 0
((point R3-rect) (up 'x0 'y0 'z0)))
```

Note that  $i_v \circ i_u + i_u \circ i_v = 0$ . One consequence of this is that  $i_v \circ i_v = 0$ .

## 7.10 Covariant Derivative

The covariant derivative is another kind of directional derivative operator. We write the covariant derivative operator with respect to a vector field  $\mathbf{v}$  as  $\nabla_{\mathbf{v}}$ . This is pronounced “covariant derivative with respect to  $\mathbf{v}$ ” or “nabla  $\mathbf{v}$ .”

## 7.11 Covariant Derivative of Vector Fields

We may also choose our  $F_{\delta}^{\mathbf{v}} \mathbf{u}$  to define what we mean by “parallel” transport of the vector field  $\mathbf{u}$  along an integral curve of the vector field  $\mathbf{v}$ . This may correspond to our usual understanding of parallel in situations where we have intuitive insight.

The notion of parallel transport is path dependent. Remember our example from the Introduction, [page 1](#): Start at the North Pole carrying a stick along a line of longitude to the Equator, always pointing it south, parallel to the surface of the Earth. Then proceed eastward for some distance, still pointing the stick south. Finally, return to the North Pole along this new line of longitude, keeping the stick pointing south all the time. At the pole the stick will not point in the same direction as it did at the beginning of the trip, and the discrepancy will depend on the amount of eastward motion.<sup>5</sup>

So if we try to carry a stick parallel to itself and tangent to the sphere, around a closed path, the stick generally does not end up pointing in the same direction as it started. The result of carrying the stick from one point on the sphere to another depends on the path taken. However, the direction of the stick at the endpoint of a path does not depend on the rate of transport, just on the particular path on which it is carried. Parallel transport over a zero-length path is the identity.

A vector may be resolved as a linear combination of other vectors. If we parallel-transport each component, and form the same linear combination, we get the transported original vector. Thus parallel transport on a particular path for a particular distance is a linear operation.

So the transport function  $F_{\delta}^{\mathbf{v}}$  is a linear operator on the components of its argument, and thus:

$$F_{\delta}^{\mathbf{v}} \mathbf{u}(f)(m) = \sum_{i,j} \left( A_j^i(\delta) (u^j \circ \phi_{-\delta}^{\mathbf{v}}) \mathbf{e}_i(f) \right) (m) \quad (7.37)$$

for some functions  $A_j^i$  that depend on the particular path (hence its tangent vector  $\mathbf{v}$ ) and the initial point. We reach back along the integral curve to pick up the components of  $\mathbf{u}$  and then parallel-transport them forward by the matrix  $A_j^i(\delta)$  to form the components of the parallel-transported vector at the advanced point.

As before, we compute

$$\nabla_{\mathbf{v}} \mathbf{u}(f)(m) = Dg(0), \quad (7.38)$$

where

$$g(\delta) = \mathbf{u}(f)(m) - (F_{\delta}^{\mathbf{v}} \mathbf{u})(f)(m). \quad (7.39)$$

Expanding with respect to a basis  $\{\mathbf{e}_i\}$  we get

$$g(\delta) = \sum_i \left( u^i \mathbf{e}_i(f) - \sum_j A_j^i(\delta) (u^j \circ \phi_{-\delta}^{\mathbf{v}}) \mathbf{e}_i(f) \right) (m). \quad (7.40)$$

By the product rule for derivatives,

$$Dg(\delta) = \sum_{ij} \left( A_j^i(\delta) ((\mathbf{v}(u^j)) \circ \phi_{-\delta}^{\mathbf{v}}) \mathbf{e}_i(f) - DA_j^i(\delta) (u^j \circ \phi_{-\delta}^{\mathbf{v}}) \mathbf{e}_i(f) \right) (m). \quad (7.41)$$

<sup>5</sup>In the introduction the stick was always kept east-west rather than pointing south, but the phenomenon is the same!



So

$$Dg(0) = \sum_i \left( v(u^i)(m) + \sum_j \varpi_j^i(v)(m) u^j(m) \right) e_i(f)(m). \quad (7.48)$$

Thus the covariant derivative is

$$\nabla_v u(f) = \sum_i \left( v(u^i) + \sum_j \varpi_j^i(v) u^j \right) e_i(f). \quad (7.49)$$

The one-form fields  $\varpi_j^i$  are called the *Cartan one-forms*, or the *connection one-forms*. They are defined with respect to the basis  $e$ .

As a program, the covariant derivative is:<sup>6</sup>

```
(defn covariant-derivative-vector                                     clojure
  [Cartan]
  (fn [V]
    (fn [U]
      (fn [f]
        (let [basis (Cartan→basis Cartan)
              Cartan-forms (Cartan→forms Cartan)]
          (let [vector-basis (basis→vector-basis basis)
                oneform-basis (basis→oneform-basis basis)]
            (let [u-components (oneform-basis U)]
              (* (vector-basis f) (+ (V u-components) (* (Cartan-forms V) u-
components))))))))))
```

An important property of  $\nabla_v u$  is that it is linear over manifold functions  $g$  in the first argument

$$\nabla_{gv} u(f) = g \nabla_v u(f), \quad (7.50)$$

consistent with the fact that the Cartan forms  $\varpi_j^i$  share the same property.

Additionally, we can extend the product rule, for any manifold function  $g$  and any vector field  $u$ :

$$\begin{aligned} \nabla_v(gu)(f) &= \sum_i \left( v(gu^i) + \sum_j \varpi_j^i(v) g u^j \right) e_i(f) \\ &= \sum_i v(g) u^i e_i(f) + g \nabla_v u(f) \\ &= (\nabla_v g) u(f) + g \nabla_v u(f). \end{aligned} \quad (7.51)$$

## 7.12 An Alternate View

As we did with the Lie derivative (equations (7.18) - (7.21)), we can write the vector field

$$u(f)(m) = \sum_i u^i(m) e_i(f)(m). \quad (7.52)$$

By the extended product rule, equation (7.51), we get:

$$\nabla_v u(f) = \sum_i (v(u^i) e_i(f) + u^i \nabla_v e_i(f)). \quad (7.53)$$

Because the covariant derivative of a vector field is a vector field we can extract the components of  $\nabla_v e_i$  using the dual basis:

$$\varpi_j^i(v) = \tilde{e}^i(\nabla_v e_j). \quad (7.54)$$

This gives an alternate expression for the Cartan one forms. So

---

<sup>6</sup>This program is incomplete. It must construct a vector field; it must make a differential operator; and it does not apply to functions or forms.



$$\nabla_v u(f) = \sum_i \left( v(u^i) + \sum_j \varpi_j^i(v) u^j \right) e_i(f). \quad (7.55)$$

This analysis is parallel to the analysis of the Lie derivative, except that here we have the Cartan form fields  $\varpi_j^i$  and there we had  $\Delta_j^i$ , which are not form fields.

Notice that the Cartan forms appear here (equation (7.53)) in terms of the covariant derivatives of the basis vectors. By contrast, in the first derivation (see equation (7.42)) the Cartan forms appear as the derivatives of the linear forms that accomplish the parallel transport of the coefficients.

The Cartan forms can be constructed from the dual basis one-forms:

$$\varpi_j^i(v)(m) = \sum_k \Gamma_{jk}^i(m) \tilde{e}^k(v)(m). \quad (7.56)$$

The connection coefficient functions  $\Gamma_{jk}^i$  are called the *Christoffel coefficients* (traditionally called *Christoffel symbols*).<sup>7</sup> Making use of the structures,<sup>8</sup> the Cartan forms are

$$\varpi(v) = \Gamma \tilde{e}(v). \quad (7.57)$$

Conversely, the Christoffel coefficients may be obtained from the Cartan forms

$$\Gamma_{jk}^i = \varpi_j^i(e_k). \quad (7.58)$$

### 7.13 Covariant Derivative of One-Form Fields

The covariant derivative of a vector field induces a compatible covariant derivative for a one-form field. Because the application of a one-form field to a vector field yields a manifold function, we can evaluate the covariant derivative of such an application. Let  $\tau$  be a one-form field and  $w$  be a vector field. Then

$$\begin{aligned} \nabla_v(\tau(w)) &= v \left( \sum_j \tau_j w^j \right) \\ &= \sum_j (v(\tau_j) w^j + \tau_j v(w^j)) \\ &= \sum_j \left( v(\tau_j) w^j + \tau_j \left( \tilde{e}^j(\nabla_v w) - \sum_k (\varpi_k^j(v) w^k) \right) \right) \\ &= \sum_j \left( v(\tau_j) w^j - \tau_j \sum_k (\varpi_k^j(v) w^k) \right) + \tau(\nabla_v w) \\ &= \sum_j \left( v(\tau_j) \tilde{e}^j - \tau_j \sum_k (\varpi_k^j(v) \tilde{e}^k) \right) (w) + \tau(\nabla_v w). \end{aligned}$$

So if we define the covariant derivative of a one-form field to be

$$\nabla_v(\tau) = \sum_k \left( v(\tau_k) - \sum_j \tau_j \varpi_k^j(v) \right) \tilde{e}^k, \quad (7.59)$$

then the generalized product rule holds:

$$\nabla_v(\tau(u)) = (\nabla_v \tau)(u) + \tau(\nabla_v u). \quad (7.60)$$

Alternatively, assuming the generalized product rule forces the definition of covariant derivative of a one-form field.

As a program this is

<sup>7</sup>This terminology may be restricted to the case in which the basis is a coordinate basis.

<sup>8</sup>The structure of the Cartan forms  $\varpi$  together with this equation forces the shape of the Christoffel coefficient structure.

```

(defn covariant-derivative-oneform                                     closure
  [Cartan]
  (fn [V]
    (fn [tau]
      (fn [U]
        (let [nabla_V ((covariant-derivative-vector Cartan) V)]
          (- (V (tau U)) (tau (nabla_V U))))))))

```

This program extends naturally to higher-rank form fields:

```

(defn covariant-derivative-form                                     closure
  [Cartan]
  (fn [V]
    (fn [tau]
      (fn [vs]
        (let [k (get-rank tau)
              nabla_V ((covariant-derivative-vector Cartan) V)]
          (- (V (apply tau vs))
             (sigma (fn [i] (apply tau (list-with-substituted-coord vs i (nabla_V (nth
vs i))))))
              0
              (- k 1))))))))

```

## 7.14 Change of Basis

The basis-independence of the covariant derivative implies a relationship between the Cartan forms in one basis and the equivalent Cartan forms in another basis. Recall (equation (4.13)) that the basis vector fields of two bases are always related by a linear transformation. Let  $J$  be the matrix of coefficient functions and let  $\mathbf{e}$  and  $\mathbf{e}'$  be down tuples of basis vector fields. then

$$\mathbf{e}(f) = \mathbf{e}'(f)J. \quad (7.61)$$

We want the covariant derivative to be independent of basis. This will determine how the connection transforms with a change of basis:

$$\begin{aligned}
\nabla_{\mathbf{v}} u(f) &= \sum_i \mathbf{e}_i(f) \left( \mathbf{v}(u^i) + \sum_j \varpi_j^i(\mathbf{v}) u^j \right) \\
&= \sum_{ijk} \mathbf{e}'_i(f) J_j^i \left( \mathbf{v} \left( (J^{-1})_k^j (u')^k \right) + \sum_l \varpi_k^j(\mathbf{v}) (J^{-1})_l^k (u')^l \right) \\
&= \sum_i \mathbf{e}'_i(f) \left( \mathbf{v} \left( (u')^i \right) + \sum_{jk} J_j^i \mathbf{v} \left( (J^{-1})_k^j \right) (u')^k \right. \\
&\quad \left. + \sum_{jkl} J_j^i \varpi_k^j(\mathbf{v}) (J^{-1})_l^k (u')^l \right) \\
&= \sum_i \mathbf{e}'_i(f) \left( \mathbf{v} \left( (u')^i \right) + \sum_j (\varpi')_j^i(\mathbf{v}) (u')^j \right).
\end{aligned} \quad (7.62)$$

The last line of equation (7.62) gives the formula for the covariant derivative we would have written down naturally in the primed coordinates; comparing with the next-to-last line, we see that

$$\varpi'(\mathbf{v}) = J\mathbf{v}(J^{-1}) + J\varpi(\mathbf{v})J^{-1}. \quad (7.63)$$

This transformation rule is weird. It is not a linear transformation of  $\varpi$  because the first term is an offset that depends on  $\mathbf{v}$ . So it is not required that  $\varpi' = 0$  when  $\varpi = 0$ . Thus  $\varpi$  is not a tensor field. See Appendix F.

We can write equation (7.61) in terms of components

$$e_i(f) = \sum_j e'_j(f) J_i^j. \quad (7.64)$$

Let  $K = J^{-1}$ , so  $\sum_j K_j^i(m) J_k^j(m) = \delta_k^i$ . Then

$$\varpi_l^i(v) = \sum_j J_j^i v(K_l^j) + \sum_{jk} J_j^i \varpi_k^j(v) K_l^k. \quad (7.65)$$

The transformation rule for  $\varpi$  is implemented in the following program:

```
(defn Cartan-transform                                     closure
  [Cartan basis-prime]
  (let [basis (Cartan→basis Cartan)
        forms (Cartan→forms Cartan)
        prime-dual-basis (basis→oneform-basis basis-prime)
        prime-vector-basis (basis→vector-basis basis-prime)]
    (let [vector-basis (basis→vector-basis basis)
          oneform-basis (basis→oneform-basis basis)]
      (let [J-inv (mapr oneform-basis prime-vector-basis)
            J (mapr prime-dual-basis vector-basis)]
        (let [omega-prime-forms (procedure→oneform-field
                                (fn [v] (+ (* J (v J-inv)) (* J (* (forms v) J-
inv))))))]
          (make-Cartan omega-prime-forms basis-prime))))))
```

Emmy's `mapr` function recursively maps its first argument over the Emmy structure supplied as its second argument, preserving that structure's shape.

We can illustrate that the covariant derivative is independent of the coordinate system in a simple case, using rectangular and polar coordinates in the plane.<sup>9</sup> We can choose Christoffel coefficients for rectangular coordinates that are all zero:<sup>10</sup>

```
(def R2-rect-Christoffel                                   closure
  (make-Christoffel (let [zero (fn [m] 0)]
    (down (down (up zero zero) (up zero zero))
          (down (up zero zero) (up zero zero))))
  R2-rect-basis))
```

With these Christoffel coefficients, parallel transport preserves the components relative to the rectangular basis. This corresponds to our usual notion of parallel in the plane. We will see later in Chapter 9 that these Christoffel coefficients are a natural choice for the plane. From these we obtain the Cartan form:<sup>11</sup>

```
(def R2-rect-Cartan (Christoffel→Cartan R2-rect-Christoffel)) closure
```

<sup>9</sup>We will need a few definitions:

```
(def R2-rect-basis (coordinate-system→basis R2-rect)) closure
(def R2-polar-basis (coordinate-system→basis R2-polar))
(define-coordinates (up x y) R2-rect)
(define-coordinates (up r theta) R2-polar)
```

<sup>10</sup>Since the Christoffel coefficients are basis-dependent they are packaged with the basis.

<sup>11</sup>The code for making the Cartan forms is as follows:

```
(defn Christoffel→Cartan                                   closure
  [Christoffel]
  (let [basis (Christoffel→basis Christoffel)
        Christoffel-symbols (Christoffel→symbols Christoffel)]
    (make-Cartan (* Christoffel-symbols (basis→oneform-basis basis)) basis)))
```

And from equation (7.63) we can get the corresponding Cartan form for polar coordinates:

```
(def R2-polar-Cartan (Cartan-transform R2-rect-Cartan R2-polar-basis))
```

closure

The vector field  $\partial/\partial\theta$  generates a rotation in the plane (the same as circular). The covariant derivative with respect to  $\partial/\partial x$  of  $\partial/\partial\theta$  applied to an arbitrary manifold function is:

```
(def circular (- (* x d:dy) (* y d:dx)))
```

closure

```
(def f (literal-manifold-function 'f-rect R2-rect))
```

```
(def R2-rect-point ((point R2-rect) (up 'x0 'y0)))
```

```
(((((covariant-derivative R2-rect-Cartan) d:dx) circular) f) R2-rect-point)
```

```
;; => (((partial 1) f-rect) (up x0 y0))
```


Note that this is the same thing as  $\partial/\partial y$  applied to the function:

```
((d:dy f) R2-rect-point)
```

closure

```
;; => (((partial 1) f-rect) (up x0 y0))
```

In rectangular coordinates, where the Christoffel coefficients are zero, the covariant derivative  $\nabla_u v$  is the vector whose coefficients are obtained by applying  $u$  to the coefficients of  $v$ . Here, only one coefficient of  $\partial/\partial\theta$  depends on  $x$ , the coefficient of  $\partial/\partial y$ , and it depends linearly on  $x$ . So  $\nabla_{\partial/\partial x} \partial/\partial\theta = \partial/\partial y$ . (See figure 7.1.)



**Figure 7.1:** If  $v$  and  $v'$  are “arrow” representations of vectors in the circular field and we parallel-transport  $v$  in the  $\partial/\partial x$  direction, then the difference between  $v'$  and the parallel transport of  $v$  is in the  $\partial/\partial y$  direction.

Note that we get the same answer if we use polar coordinates to compute the covariant derivative:

```
(simplify (((((covariant-derivative R2-polar-Cartan) d:dx) circular) f) R2-rect-  
point))
```

closure

```
;; => (((partial 1) f-rect) (up x0 y0))
```

In rectangular coordinates the Christoffel coefficients are all zero; in polar coordinates there are nonzero coefficients, but the value of the covariant derivative is the same. In polar coordinates the basis elements vary with position, and the Christoffel coefficients compensate for this.

Of course, this is a pretty special situation. Let's try something more general:

```
(def V (literal-vector-field 'V-rect R2-rect))

(def W (literal-vector-field 'W-rect R2-rect))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((((- (covariant-derivative R2-rect-Cartan) (covariant-derivative R2-polar-
Cartan)) V) W)
           f)
          R2-rect-point))

;; => 0
```

## 7.15 Parallel Transport

We have defined parallel transport of a vector field along integral curves of another vector field. But not all paths are integral curves of a vector field. For example, paths that cross themselves are not integral curves of any vector field.

Here we extend the idea of a parallel transport of a stick to make sense for arbitrary paths on the manifold. Any path can be written as a map  $\gamma$  from the real-line manifold to the manifold  $M$ . We construct a vector field over the map  $u_\gamma$  by parallel-transporting the stick to all points on the path  $\gamma$ .

For any path  $\gamma$  there are locally directional derivatives of functions on  $M$  defined by tangent vectors to the curve. The vector over the map  $w_\gamma = d\gamma(\partial/\partial t)$  is a directional derivative of functions on the manifold  $M$  along the path  $\gamma$ .

Our goal is to determine the equations satisfied by the vector field over the map  $u_\gamma$ . Consider the parallel-transport  $F_\delta^{w_\gamma} u_\gamma$ .<sup>12</sup> So a vector field  $u_\gamma$  is parallel-transported to itself if and only if  $u_\gamma = F_\delta^{w_\gamma} u_\gamma$ . Restricted to a path, the equation analogous to equation (7.40) is

$$g(\delta) = \sum_i \left( u^i(t) - \sum_j A_j^i(\delta) u^j(t - \delta) \right) e_i^\gamma(f)(t), \quad (7.66)$$

where the coefficient function  $u^i$  is now a function on the real-line parameter manifold and where we have rewritten the basis as a basis over the map  $\gamma$ .<sup>13</sup> Here  $g(\delta) = 0$  if  $u_\gamma$  is parallel-transported into itself.

Taking the derivative and setting  $\delta = 0$  we find

$$0 = \sum_i \left( Du^i(t) + \sum_j^\gamma \varpi_j^i(w_\gamma)(t) u^j(t) \right) e_i^\gamma(f)(t). \quad (7.67)$$

But this implies that

$$0 = Du^i(t) + \sum_j^\gamma \varpi_j^i(w_\gamma)(t) u^j(t), \quad (7.68)$$

an ordinary differential equation in the coefficients of  $u_\gamma$ .

We can abstract these equations of parallel transport by inventing a covariant derivative over a map. We also generalize the time line to a source manifold  $N$ .

<sup>12</sup>The argument  $w_\gamma$  makes sense because our parallel-transport operator never depended on the vector field tangent to the integral curve existing off of the curve. Because the connection is a form field (see equation (7.47)), it does not depend on the value of its vector argument anywhere except at the point where it is being evaluated.

The argument  $u_\gamma$  is more difficult. We must modify equation (7.37):

$$F_\delta^{w_\gamma} u_\gamma(f)(t) = \sum_{i,j} A_j^i(\delta) u^j(t - \delta) e_i^\gamma(f)(t).$$

<sup>13</sup>You may have noticed that  $t$  and  $\mathbf{t}$  appear here. The real-line manifold point  $\mathbf{t}$  has coordinate  $t$ .

$$\nabla_v^\gamma u_\gamma(f)(n) = \sum_i \left( v(u^i)(n) + \sum_j^\gamma \varpi_j^i(d\gamma(v))(n) u^j(n) \right) e_i^\gamma(f)(n), \quad (7.69)$$

where the map  $\gamma : N \rightarrow M$ ,  $v$  is a vector on  $N$ ,  $u_\gamma$  is a vector over the map  $\gamma$ ,  $f$  is a function on  $M$ , and  $n$  is a point in  $N$ . Indeed, if  $w$  is a vector field on  $M$ ,  $f$  is a manifold function on  $M$ , and if  $d\gamma(v) = w_\gamma$  then

$$\nabla_v^\gamma u_\gamma(f)(n) = \nabla_w u(f)(\gamma(n)). \quad (7.70)$$

This is why we are justified in calling  $\nabla_v^\gamma$  a covariant derivative.

Respecializing the source manifold to the real line, we can write the equations governing the parallel transport of  $u_\gamma$  as

$$\nabla_{\partial/\partial t}^\gamma u_\gamma = 0. \quad (7.71)$$

We obtain the set of differential equations (7.68) for the coordinates of  $u_\gamma$ , the vector over the map  $\gamma$ , that is parallel-transported along the curve  $\gamma$ :

$$Du^i(t) + \sum_j^\gamma \varpi_j^i(d\gamma(\partial/\partial t))(t) u^j(t) = 0. \quad (7.72)$$

Expressing the Cartan forms in terms of the Christoffel coefficients we obtain

$$Du^i(t) + \sum_{j,k} \Gamma_{jk}^i(\gamma(t)) D\sigma^k(t) u^j(t) = 0 \quad (7.73)$$

where  $\sigma = \chi_M \circ \gamma \circ \chi_R^{-1}$  are the coordinates of the path ( $\chi_M$  and  $\chi_R$  are the coordinate functions for  $M$  and the real line).

## 7.16 On a Sphere

Let's figure out what the equations of parallel transport of  $u_\gamma$ , an arbitrary vector over the map  $\gamma$ , along an arbitrary path  $\gamma$  on a sphere are. We start by constructing the necessary manifold.

```
(def sphere (make-manifold S2-type 2 3)) closure

(def S2-spherical (coordinate-system-at sphere :spherical :north-pole))

(def S2-basis (coordinate-system→basis S2-spherical))

(define-coordinates (up theta phi) S2-spherical)
```

We need the path  $\gamma$ , which we represent as a map from the real line to  $M$ , and  $w$ , the parallel-transported vector over the map:

```
(def gamma closure
  (compose (point S2-spherical)
    (up (literal-function 'alpha) (literal-function 'beta))
    (chart R1-rect)))
```

where  $\alpha$  is the colatitude and  $\beta$  is the longitude.

We also need an arbitrary vector field  $u_{\text{gamma}}$  over the map  $\gamma$ . To make this we multiply the structure of literal component functions by the vector basis structure.

```
(def basis-over-gamma (basis→basis-over-map gamma S2-basis)) closure

(def u_gamma
  (* (up (compose (literal-function 'u↑0) (chart R1-rect))
    (compose (literal-function 'u↑1) (chart R1-rect)))
    (basis→vector-basis basis-over-gamma)))
```

We specify a connection by giving the Christoffel coefficients.<sup>14</sup>

```
(def S2-Christoffel
  (make-Christoffel (let [zero (fn [point] 0)]
    (down (down (up zero zero) (up zero (/ 1 (tan theta))))
      (down (up zero (/ 1 (tan theta)))
        (up (- (* (sin theta) (cos theta)) zero))))
    S2-basis)))

(def sphere-Cartan (Christoffel→Cartan S2-Christoffel))
```

Finally, we compute the residual of the equation (7.71) that governs parallel transport for this situation:<sup>15</sup>

```
(define-coordinates t R1-rect)

(mapr (fn [omega]
  ((omega ((covariant-derivative sphere-Cartan gamma) d:dt) u_gamma))
    ((point R1-rect) 'tau)))
(basis→oneform-basis basis-over-gamma))
;; => (up (+ ((D u↑0) tau)
;;          (* (- (* (sin (alpha tau)) (cos (alpha tau))) ((D beta) tau) (u↑1 tau)))
;;          (+ ((D u↑1) tau)
;;            (* (/ 1 (tan (alpha tau))) ((D beta) tau) (u↑0 tau))
;;            (* (/ 1 (tan (alpha tau))) ((D alpha) tau) (u↑1 tau))))))
```

Thus the equations governing the evolution of the components of the transported vector are:

$$Du^0(\tau) = \sin(\alpha(\tau)) \cos(\alpha(\tau)) D\beta(\tau) u^1(\tau),$$

$$Du^1(\tau) = -\frac{\cos(\alpha(\tau))}{\sin(\alpha(\tau))} (D\beta(\tau) u^0(\tau) + D\alpha(\tau) u^1(\tau)). \quad (7.74)$$

These equations describe the transport on a sphere, but more generally they look like

$$Du(\tau) = f(\sigma(\tau), D\sigma(\tau))u(\tau), \quad (7.75)$$

where  $\sigma$  is the tuple of the coordinates of the path on the manifold and  $u$  is the tuple of the components of the vector. The equation is linear in  $u$  and is driven by the path  $\sigma$ , as in a variational equation.

We now set this up for numerical integration. Let  $s(t) = (t, u(t))$  be a state tuple, combining the time and the coordinates of  $u_\gamma$  at that time. Then we define  $g$ :

$$g(s(t)) = Ds(t) = (1, Du(t)), \quad (7.76)$$

where  $Du(t)$  is the tuple of right-hand sides of equation (7.72).

## 7.17 On a Great Circle

We illustrate parallel transport in a case where we should know the answer: we carry a vector along a great circle of a sphere. Given a path and Cartan forms for the manifold we can produce a state derivative suitable for numerical integration. Such a state derivative takes a state and produces the derivative of the state.

```
(defn g
  [gamma Cartan]
  (let [omega ((Cartan→forms (Cartan→Cartan-over-map Cartan gamma)) ((differential
gamma) d:dt))]
    (letfn
      [(the-state-derivative []
        (fn [state]
```

<sup>14</sup>We will show later that these Christoffel coefficients are a natural choice for the sphere.

<sup>15</sup>If we give `covariant-derivative` an extra argument, in addition to the Cartan form, the covariant derivative treats the extra argument as a map and transforms the Cartan form to work over the map.

```

    (let [t ((point R1-rect) (ref state 0)) u (ref state 1)] (up 1 (* -1 (omega
t) u))))))
    the-state-derivative)))

```

The path on the sphere will be the target of a map from the real line. We choose one that starts at the origin of longitudes on the equator and follows the great circle that makes a given tilt angle with the equator.

```

(defn transform
  [tilt]
  (fn [coords]
    (let [colat (ref coords 0)
          long (ref coords 1)]
      (let [x (* (sin colat) (cos long))
            y (* (sin colat) (sin long))
            z (cos colat)]
        (let [vp ((rotate-x tilt) (up x y z))]
          (let [colatp (acos (ref vp 2)) longp (atan (ref vp 1) (ref vp 0))] (up
colatp longp)))))))

(defn tilted-path
  [tilt]
  (letfn [(coords [t] ((transform tilt) (up (/ pi 2) t)))]
    (compose (point S2-spherical) coords (chart R1-rect))))

```

A southward pointing vector, with components (up 1 0), is transformed to an initial vector for the tilted path by multiplying by the derivative of the tilt transform at the initial point. We then parallel transport this vector by numerically integrating the differential equations. In this example we tilt by 1 radian, and we advance for  $\pi/2$  radians. In this case we know the answer: by advancing by  $\pi/2$  we walk around the circle a quarter of the way and at that point the transported vector points south:

```

((state-advancer (g (tilted-path 1) sphere-Cartan))
 (up 0 (* ((D (transform 1)) (up (/ pi 2) 0)) (up 1 0)))
 (/ pi 2))
;; => (up 1.5707963267948957 (up 1.0000000005999354 -2.3386485677699687e-8))

```

However, if we transport by 1 radian rather than  $\pi/2$ , the numbers are not so pleasant, and the transported vector no longer points south:

```

((state-advancer (g (tilted-path 1) sphere-Cartan))
 (up 0 (* ((D (transform 1)) (up (/ pi 2) 0)) (up 1 0)))
 1)
;; => (up 1.0000000000000004 (up 0.76515026455619 0.9117920267996223))

```

But the transported vector can be obtained by tilting the original southward-pointing vector after parallel-transporting along the equator:<sup>16</sup>

```

(* ((D (transform 1)) (up (/ pi 2) 1)) (up 1 0))
;; => (up 0.7651502649370374 0.9117920272004736)

```

## 7.18 Geodesic Motion

In geodesic motion the velocity vector is parallel-transported by itself. Recall (equation (6.9)) that the velocity is the differential of the vector  $\partial/\partial t$  over the map  $\gamma$ . The equation of geodesic motion is<sup>17</sup>

<sup>16</sup>A southward-pointing vector remains southward-pointing when it is parallel-transported along the equator. To do this we do not have to integrate the differential equations, because we know the answer.

<sup>17</sup>The equation of a geodesic path is often said to be

$$\nabla_v v = 0, \quad (7.77)$$



$$\nabla_{\partial/\partial t}^\gamma d\gamma(\partial/\partial t) = 0. \quad (7.78)$$

In coordinates, this is

$$D^2\sigma^i(t) + \sum_{jk} \Gamma_{jk}^i(\gamma(t)) D\sigma^j(t) D\sigma^k(t) = 0, \quad (7.79)$$

where  $\sigma(t)$  is the coordinate path corresponding to the manifold path  $\gamma$ .

For example, let's consider geodesic motion on the surface of a unit sphere. We let  $\gamma$  be a map from the real line to the sphere, with colatitude  $\alpha$  and longitude  $\beta$ , as before. The geodesic equation is:

```
(show-expression (((covariant-derivative sphere-Cartan gamma) d:dt) ((differential
gamma) d:dt)))
      (chart S2-spherical))
      ((point R1-rect) 't0)))
;; => (up (+ ((expt D 2) alpha) t0)
;;        (* (- (* (sin (alpha t0)) (cos (alpha t0)))) ((D beta) t0) ((D beta)
t0)))
;;      (+ ((expt D 2) beta) t0)
;;      (* (/ 1 (tan (alpha t0))) ((D beta) t0) ((D alpha) t0))
;;      (* (/ 1 (tan (alpha t0))) ((D alpha) t0) ((D beta) t0)))
```

$$\left( \begin{aligned} & -\cos(\alpha(t_0)) \sin(\alpha(t_0)) (D\beta(t_0))^2 + D^2\alpha(t_0) \\ & \frac{2D\beta(t_0) \cos(\alpha(t_0)) D\alpha(t_0)}{\sin(\alpha(t_0))} + D^2\beta(t_0) \end{aligned} \right)$$

The geodesic equation is the same as the Lagrange equation for free motion constrained to the surface of the unit sphere. The Lagrangian for motion on the sphere is the composition of the free-particle Lagrangian and the state transformation induced by the coordinate constraint:<sup>18</sup>

```
(defn Lfree [s] (* (/ 1 2) (square (velocity s))))
;; closure

(defn sphere→R3
  [s]
  (let [q (coordinate s)]
    (let [theta (ref q 0)
          phi (ref q 1)]
      (up (* (sin theta) (cos phi)) (* (sin theta) (sin phi)) (cos theta)))))

(def Lsphere (compose Lfree (F→C sphere→R3)))
```

Then the Lagrange equations are:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (show-expression ((Lagrange-equations Lsphere)
  (up (literal-function 'alpha) (literal-function 'beta))
  't)))
;; => (down
;;      (+ (* #emmy/bigint -1 (cos (alpha t)) (sin (alpha t)) (expt ((D beta) t) 2))
;;        ((expt D 2) alpha) t))
;;      (+ (* #emmy/bigint 2 (cos (alpha t)) (sin (alpha t)) ((D beta) t) ((D alpha)
t))
;;          (* (expt (sin (alpha t)) 2) ((expt D 2) beta) t)))
```

but this is nonsense. The geodesic equation is a constraint on the path, but the path does not appear in this equation. Further, the velocity along a path is not a vector field, so it cannot appear in either argument to the covariant derivative.

What is true is that a vector field  $\mathbf{v}$  all of whose integral curves are geodesics satisfies equation (7.77).

<sup>18</sup>The method of formulating a system with constraints by composing a free system with the state-space coordinate transformation that represents the constraints can be found in [19], section 1.6.3. The function  $F \rightarrow C$  takes a coordinate transformation and produces a corresponding transformation of Lagrangian state.

$$[-(D\beta(t))^2 \sin(\alpha(t)) \cos(\alpha(t)) + D^2\alpha(t) 2D\alpha(t)D\beta(t) \sin(\alpha(t)) \cos(\alpha(t)) + D^2\beta(t)(\sin(\alpha(t)))^2]$$

The Lagrange equations are true of the same paths as the geodesic equations. The second Lagrange equation is the second geodesic equation multiplied by  $(\sin(\alpha(t)))^2$ , and the Lagrange equations are arranged in a down tuple, whereas the geodesic equations are arranged in an up tuple.<sup>19</sup> The two systems are equivalent unless  $\alpha(t) = 0$ , where the coordinate system is singular.

### 7.18.1 Exercise 7.1: Hamiltonian Evolution

We have just seen that the Lagrange equations for the motion of a free particle constrained to the surface of a sphere determine the geodesics on the sphere. We can investigate the phenomenon in the Hamiltonian formulation. The Hamiltonian is obtained from the Lagrangian by a Legendre transformation:

```
;; Prove that Emmy's generic Legendre transform agrees with the equivalent
;; closed form at a fully symbolic state. The following example differentiates
;; the closed form, avoiding a much slower differentiation through the generic
;; transform while retaining an exact symbolic check.
(def Hsphere
  (let [via-Legendre (Lagrangian→Hamiltonian Lsphere)
        closed-form (fn [_ [theta _] [p-theta p-phi]]
                      (* (/ 1 2) (+ (square p-theta) (/ (square p-phi) (square (sin
theta))))))]
    symbolic-state (up 't (up 'theta 'phi) (down 'p_theta 'p_phi))
    residual (simplify (- (via-Legendre symbolic-state) (closed-form symbolic-
state))))]
  (verified-zero closed-form (up residual))))
```

We can get the coordinate representation of the Hamiltonian vector field as follows:

```
(simplify ((phase-space-derivative Hsphere) (up 't (up 'theta 'phi) (down 'p_theta
'p_phi))))
;; => (up 1
;;      (up p_theta (/ p_phi (expt (sin theta) 2)))
;;      (down (/ (* (expt p_phi 2) (cos theta)) (expt (sin theta) 3)) -0.0))
```

The state space for Hamiltonian evolution has five dimensions: time, two dimensions of position on the sphere, and two dimensions of momentum:

```
(def state-space (make-manifold Rn 5))
(def states (coordinate-system-at state-space :rectangular :origin))
(define-coordinates (up t (up theta phi) (down p_theta p_phi)) states)
```

So now we have coordinate functions and the coordinate-basis vector fields and coordinate-basis one-form fields.

a. Define the Hamiltonian vector field as a linear combination of these fields.

b. Obtain the first few terms of the Taylor series for the evolution of the coordinates  $(\theta, \phi)$  by exponentiating the Lie derivative of the Hamiltonian vector field.

### 7.18.2 Exercise 7.2: Lie Derivative and Covariant Derivative

How are the Lie derivative and the covariant derivative related?

a. Prove that for every vector field there exists a connection such that the covariant derivative for that connection and the given vector field is equivalent to the Lie derivative with respect to that vector field.

b. Show that there is no connection that for every vector field makes the Lie derivative the same as the covariant derivative with the chosen connection.

<sup>19</sup>The geodesic equations and the Lagrange equations are related by a contraction with the metric.

```
(define-coordinates (up x y z) R3-rect)

(def theta (+ (* a dx) (* b dy) (* c dz)))

(def omega (+ (* a (wedge dy dz)) (* b (wedge dz dx)) (* c (wedge dx dy))))

(def X (literal-vector-field 'X-rect R3-rect))

(def Y (literal-vector-field 'Y-rect R3-rect))

(def Z (literal-vector-field 'Z-rect R3-rect))

(def V (literal-vector-field 'V-rect R3-rect))

(def R3-rect-point ((point R3-rect) (up 'x0 'y0 'z0)))
```

## 8 Curvature

If the intrinsic curvature of a manifold is not zero, a vector parallel-transported around a small loop will end up different from the vector that started. We saw the consequence of this before, on [page 1](#) and [page 63](#). The Riemann tensor encapsulates this idea.

The Riemann curvature operator is

$$\mathcal{R}(\mathbf{w}, \mathbf{v}) = [\nabla_{\mathbf{w}}, \nabla_{\mathbf{v}}] - \nabla_{[\mathbf{w}, \mathbf{v}]}.$$
 (8.1)

The traditional Riemann tensor is

$$\mathcal{R}(\omega, u, v, w) = \omega((\mathcal{R}(\mathbf{w}, \mathbf{v}))(u)),$$
 (8.2)

where  $\omega$  is a one-form field that measures the incremental change in the vector field  $u$  caused by parallel-transporting it around the loop defined by the vector fields  $\mathbf{w}$  and  $\mathbf{v}$ .  $\mathcal{R}$  allows us to compute the *intrinsic curvature* of a manifold at a point.

The Riemann curvature is computed by

```
(defn Riemann-curvature                                     clojure
  [nabla]
  (fn [w v] (- (commutator (nabla w) (nabla v)) (nabla (commutator w v)))))
```

The `Riemann-curvature` procedure is parameterized by the relevant `covariant-derivative` operator `nabla`, which implements  $\nabla$ . The `nabla` is itself dependent on the connection, which provides the details of the local geometry. The same `Riemann-curvature` procedure works for ordinary covariant derivatives and for covariant derivatives over a map. Given two vector fields, the result of `((Riemann-curvature nabla) w v)` is a procedure that takes a vector field and produces a vector field so we can implement the Riemann tensor as

```
(defn Riemann [nabla] (fn [omega u w v] (omega (((Riemann-curvature nabla) w v) u)))) clojure
```

So, for example,<sup>1</sup>

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((Riemann (covariant-derivative sphere-Cartan)) dphi d:dtheta d:dphi
  d:dtheta)
  ((point S2-spherical) (up 'theta0 'phi0))))
;; => 1                                     clojure
```

Here we have computed the  $\varphi$  component of the result of carrying a  $\partial/\partial\theta$  basis vector around the parallelogram defined by  $\partial/\partial\phi$  and  $\partial/\partial\theta$ . The result shows a net rotation in the  $\phi$  direction.

Most of the sixteen coefficients of the Riemann tensor for the sphere are zero. The following are the nonzero coefficients:

<sup>1</sup>The connection specified by `sphere-Cartan` is defined in Section 7.16, [page 63](#).

$$\begin{aligned}
R\left(d\theta, \frac{\partial}{\partial\phi}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}\right) \\
(\chi^{-1}(q^\theta, q^\phi)) &= (\sin(q^\theta))^2, \\
R\left(d\theta, \frac{\partial}{\partial\phi}, \frac{\partial}{\partial\phi}, \frac{\partial}{\partial\theta}\right) \\
(\chi^{-1}(q^\theta, q^\phi)) &= -(\sin(q^\theta))^2, \\
R\left(d\phi, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}\right) \\
(\chi^{-1}(q^\theta, q^\phi)) &= -1, \\
R\left(d\phi, \frac{\partial}{\partial\theta}, \frac{\partial}{\partial\phi}, \frac{\partial}{\partial\theta}\right) \\
(\chi^{-1}(q^\theta, q^\phi)) &= 1.
\end{aligned} \tag{8.3}$$

### 8.1 Explicit Transport

We will show that the result of the Riemann calculation of the change in a vector, as we traverse a loop, is what we get by explicitly calculating the transport. The coordinates of the vector to be transported are governed by the differential equations (see equation (7.72))

$$Du^i(t) = - \sum_j \varpi_j^i(\nu)(\chi^{-1}(\sigma(t)))u^j(t) \tag{8.4}$$

and the coordinates as a function of time,  $\sigma = \chi \circ \gamma \circ \chi_R^{-1}$ , of the path  $\gamma$ , are governed by the differential equations<sup>2</sup>

$$D\sigma(t) = \nu(\chi)(\chi^{-1}(\sigma(t))). \tag{8.5}$$

We have to integrate these equations (8.4), (8.5) together to transport the vector over the map  $u_\gamma$  a finite distance along the vector field  $\nu$ .

Let  $s(t) = (\sigma(t), u(t))$  be a state tuple, combining  $\sigma$  the coordinates of  $\gamma$ , and  $u$  the coordinates of  $u_\gamma$ . Then

$$Ds(t) = (D\sigma(t), Du(t)) = g(s(t)), \tag{8.6}$$

where  $g$  is the tuple of right-hand sides of equations (8.4), (8.5).

The differential equations describing the evolution of a function  $h$  of state  $s$  along the state path are

$$D(h \circ s) = (Dh \circ s)(g \circ s) = L_g h \circ s, \tag{8.7}$$

defining the operator  $L_g$ .

Exponentiation gives a finite evolution:<sup>3</sup>

$$h(s(t + \epsilon)) = (e^{\epsilon L_g} h)(s(t)). \tag{8.8}$$

The finite parallel transport of the vector with components  $u$  is

$$u(t + \epsilon) = (e^{\epsilon L_g} U)(s(t)), \tag{8.9}$$

where the selector  $U(\sigma, u) = u$ , and the initial state is  $s(t) = (\sigma(t), u(t))$ .

Consider parallel-transporting a vector  $u$  around a parallelogram defined by two coordinate-basis vector fields  $w$  and  $v$ . The vector  $u$  is really a vector over a map, where the map is the parametric curve describing our parallelogram. This map is implicitly defined in terms of the vector fields  $w$  and  $v$ . Let  $g_w$  and  $g_v$  be

<sup>2</sup>The map  $\gamma$  takes points on the real line to points on the target manifold. The chart  $\chi$  gives coordinates of points on the target manifold while  $\chi_R$  gives a time coordinate on the real line.

<sup>3</sup>The series may not converge for large increments in the independent variable. In this case it is appropriate to numerically integrate the differential equations directly.

the right-hand sides of the differential equations for parallel transport along  $\mathbf{w}$  and  $\mathbf{v}$  respectively. Then evolution along  $\mathbf{w}$  for interval  $\epsilon$ , then along  $\mathbf{v}$  for interval  $\epsilon$ , then reversing  $\mathbf{w}$ , and reversing  $\mathbf{v}$ , brings  $\sigma$  back to where it started to second order in  $\epsilon$ .

The state  $s = (\sigma, u)$  after transporting  $s_0$  around the loop is<sup>4</sup>

$$\begin{aligned} & (e^{-\epsilon L_{g_v}} I) \circ (e^{-\epsilon L_{g_w}} I) \circ (e^{\epsilon L_{g_v}} I) \\ & \quad \circ (e^{\epsilon L_{g_w}} I)(s_0) \\ & = (e^{\epsilon L_{g_w}} e^{\epsilon L_{g_v}} e^{-\epsilon L_{g_w}} e^{-\epsilon L_{g_v}} I)(s_0) \\ & = (e^{\epsilon^2 [L_{g_w}, L_{g_v}] + \dots} I)(s_0). \end{aligned} \tag{8.10}$$

So the lowest-order change in the transported vector is

$$\epsilon^2 U\left(\left([L_{g_w}, L_{g_v}] I\right)(s_0)\right), \tag{8.11}$$

where  $U(\sigma, u) = u$ .

However, if  $\mathbf{w}$  and  $\mathbf{v}$  do not commute, the indicated loop does not bring  $\sigma$  back to the starting point, to second order in  $\epsilon$ . We must account for the commutator. (See figure 4.2.) In the general case the lowest order change in the transported vector is

$$\epsilon^2 U\left(\left(\left([L_{g_w}, L_{g_v}] - L_{g_{[\mathbf{w}, \mathbf{v}]}}\right) I\right)(s_0)\right), \tag{8.12}$$

This is what the Riemann tensor computation gives, scaled by  $\epsilon^2$ .

### 8.1.1 Verification in Two Dimensions

We can verify this in two dimensions. We need to make the structure representing a state:

```
(defn make-state [sigma u] (vector sigma u)) ;closure
(defn Sigma [state] (ref state 0))
(defn U-select [state] (ref state 1))
```

And now we get to the meat of the matter: First we find the rate of change of the components of the vector  $\mathbf{u}$  as we carry it along the vector field  $\mathbf{v}$ .<sup>5</sup>

```
(defn Du ;closure
  [v]
  (fn [state]
    (let [CF (Cartan→forms general-Cartan-2)]
      (* -1 ((CF v) (Chi-inverse (Sigma state))) (U-select state))))))
```

We also need to determine the rate of change of the coordinates of the integral curve of  $\mathbf{v}$ .

<sup>4</sup>The parallel-transport operators are evolution operators, and therefore descend into composition:

$$e_A(F \circ G) = F \circ (e^A G),$$

for any state function  $G$  and any compatible  $F$ . As a consequence, we have the following identity:

$$e^A e^B I = e^A ((e^B I) \circ I) = (e^B I) \circ (e^A I),$$

where  $I$  is the identity function on states.

<sup>5</sup>The setup for this experiment is a bit complicated. We need to make a manifold with a general connection.

```
(def Chi-inverse (point R2-rect)) ;closure
(def Chi (chart R2-rect))
```

```
(defn Dsigma [v] (fn [state] ((v Chi) (Chi-inverse (Sigma state))))))
```

closure

Putting these together to make the derivative of the state vector

```
(defn g [v] (fn [state] (make-state ((Dsigma v) state) ((Du v) state))))
```

closure

gives us just what we need to construct the differential operator for evolution of the combined state:

```
(defn L
  [v]
  (letfn [(l [h] (fn [state] (* ((D h) state) ((g v) state))))]
    (make-operator l)))
```

closure

So now we can demonstrate that the lowest-order change resulting from explicit parallel transport of a vector around an infinitesimal loop is what is computed by the Riemann curvature.

```
(simplify (let [U (literal-vector-field 'U-rect R2-rect)
                W (literal-vector-field 'W-rect R2-rect)
                V (literal-vector-field 'V-rect R2-rect)
                sigma (up 'sigma0 'sigma1)]
  (let [nabla (covariant-derivative general-Cartan-2)
        m (Chi-inverse sigma)]
    (let [s (make-state sigma ((U Chi) m))]
      (- (((- (commutator (L V) (L W)) (L (commutator V W))) U-select) s)
          (((Riemann-curvature nabla) W V) U) Chi) m))))))
;; => (up 0 0)
```

closure

### 8.1.2 Geometrically

The explicit transport above was done with differential equations operating on a state consisting of coordinates and components of the vector being transported. We can simplify this so that it is entirely built on manifold objects, eliminating the state. After a long algebraic story we find that

$$\begin{aligned}
 & ((\mathcal{R}(w, v))(u))(f) \\
 &= e(f) \{ (w(\varpi(v)) - v(\varpi(w)) - \varpi([w, v])) \tilde{e}(u) \\
 & \quad + \varpi(w) \varpi(v) \tilde{e}(u) - \varpi(v) \varpi(w) \tilde{e}(u) \}
 \end{aligned} \tag{8.13}$$

or as a program:

```
(defn curvature-from-transport
  [Cartan]
  (fn [w v]
    (fn [u]
      (fn [f]
        (let [CF (Cartan→forms Cartan)
              basis (Cartan→basis Cartan)
              fi (basis→oneform-basis basis)
              ei (basis→vector-basis basis)]
          (* (ei f)
              (+ (* (- (- (w (CF v)) (v (CF w))) (CF (commutator w v))) (fi u))
                  (- (* (CF w) (* (CF v) (fi u))) (* (CF v) (* (CF w) (fi u))))))))))
```

closure

This computes the same operator as the traditional Riemann curvature operator:

```
(defn test
  [coordsys Cartan]
  (let [m (typical-point coordsys)
        u (literal-vector-field 'u-coord coordsys)
        w (literal-vector-field 'w-coord coordsys)
        v (literal-vector-field 'v-coord coordsys)]
```

closure

```

    f (literal-manifold-function 'f-coord coordsys)]
  (let [nabla (covariant-derivative Cartan)]
    (- (((curvature-from-transport Cartan) w v) u) f) m)
    (((Riemann-curvature nabla) w v) u) f) m))))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (test R2-rect general-Cartan-2))
;; ⇒ 0

;; Full rational simplification enters an expensive polynomial GCD here.
;; Polynomial simplification proves the same exact zero without that GCD.
(polynomially-verified-zero (test R2-polar general-Cartan-2))
;; ⇒ 0

```

### 8.1.3 Terms of the Riemann Curvature

Since the Riemann curvature is defined as in equation (8.1),

$$\mathcal{R}(w, v) = [\nabla_w, \nabla_v] - \nabla_{[w, v]}, \quad (8.14)$$

it is natural<sup>6</sup> to identify these terms with the corresponding terms in

$$\left( \left( [L_{g_w}, L_{g_v}] - L_{g_{[w, v]}} \right) U \right) (s_0). \quad (8.15)$$

Unfortunately, this does not work, as demonstrated below:

```

(simplify (let [U (literal-vector-field 'U-rect R2-rect)
                V (literal-vector-field 'V-rect R2-rect)
                W (literal-vector-field 'W-rect R2-rect)
                nabla (covariant-derivative general-Cartan-2)
                sigma (up 'sigma0 'sigma1)]
  (let [m (Chi-inverse sigma)]
    (let [s (make-state sigma ((U Chi) m))]
      (- (((commutator (L W) (L V)) U-select) s)
          (((commutator (nabla W) (nabla V)) U) Chi) m))))))
;; ⇒ (up (+ (* 2 (U-rect↑1 (up sigma0 sigma1)) (W-rect↑1 (up sigma0 sigma1))
;;      (Gamma_11↑0 (up sigma0 sigma1)) (Gamma_00↑0 (up sigma0 sigma1)) (V-rect↑0
;;      (up sigma0 sigma1))) (* -2 (U-rect↑1 (up s...
;; [full result retained in chapter008/013.cljs]

```

The obvious identification does not work, but neither does the other one!

```

(let [U (literal-vector-field 'U-rect R2-rect)
      V (literal-vector-field 'V-rect R2-rect)
      W (literal-vector-field 'W-rect R2-rect)
      nabla (covariant-derivative general-Cartan-2)
      sigma (up 'sigma0 'sigma1)]
  (let [m (Chi-inverse sigma)]
    (let [s (make-state sigma ((U Chi) m))]
      (- (((commutator (L W) (L V)) U-select) s) (((nabla (commutator W V)) U) Chi)
          m))))))
;; ⇒ (up (- (- (+ (* (+ (* (U-rect↑1 (up sigma0 sigma1)) -1 (V-rect↑1 (up sigma0
;;      sigma1)) (((partial 0) Gamma_11↑0) (up sigma0 sigma1))) (* (U-rect↑1 (up
;;      sigma0 sigma1)) -1 (V-rect↑0 (up sigma0...
;; [full result retained in chapter008/014.cljs]

```

Let's compute the two parts of the Riemann curvature operator and see how this works out. First, recall

<sup>6</sup>People often say “Geodesic evolution is exponentiation of the covariant derivative.” But this is wrong. The evolution is by exponentiation of  $L_g$ .



$$\begin{aligned}
\nabla_v u(f) &= \sum_i e_i(f) \left( v(\tilde{e}^i(u)) \right. \\
&\quad \left. + \sum_j \varpi_j^i(v) \tilde{e}^j(u) \right) \\
&= e(f)(v(\tilde{e}(u)) \\
&\quad + \varpi(v)\tilde{e}(u)),
\end{aligned} \tag{8.16}$$

where the second form uses tuple arithmetic. Now let's consider the first part of the Riemann curvature operator:

$$\begin{aligned}
[\nabla_w, \nabla_v]u &= \nabla_w \nabla_v u - \nabla_v \nabla_w u \\
&= e\{w(v(\tilde{e}(u)) + \varpi(v)\tilde{e}(u)) \\
&\quad + \varpi(w)(v(\tilde{e}(u)) + \varpi(v)\tilde{e}(u))\} \\
&\quad - e\{v(w(\tilde{e}(u)) + \varpi(w)\tilde{e}(u)) \\
&\quad + \varpi(v)(w(\tilde{e}(u)) + \varpi(w)\tilde{e}(u))\} \\
&= e\{[w, v]\tilde{e}(u) \\
&\quad + w(\varpi(v))\tilde{e}(u) - v(\varpi(w))\tilde{e}(u) \\
&\quad + \varpi(w)\varpi(v)\tilde{e}(u) - \varpi(v)\varpi(w)\tilde{e}(u)\}.
\end{aligned} \tag{8.17}$$

The second term of the Riemann curvature operator is

$$\nabla_{[w, v]}u = e\{[w, v]\tilde{e}(u) + \varpi([w, v])\tilde{e}(u)\}. \tag{8.18}$$

The difference of these is the Riemann curvature operator. Notice that the first term in each cancels, and the rest gives equation (8.13).

#### 8.1.4 Ricci Curvature

One measure of the curvature is the Ricci tensor, which is computed from the Riemann tensor by

$$R(u, v) = \sum_i R(\tilde{e}^i, u, e_i, v). \tag{8.19}$$

Expressed as a program:

```
(defn Ricci [nabla basis] (fn [u v] (contract (fn [ei wi] ((Riemann nabla) wi u ei v)) closure
basis))))
```

Einstein's field equation (9.27) for gravity, which we will encounter later, is expressed in terms of the Ricci tensor.

#### 8.1.5 Exercise 8.1: Ricci of a Sphere

Compute the components of the Ricci tensor of the surface of a sphere.

#### 8.1.6 Exercise 8.2: Pseudosphere

A pseudosphere is a surface in 3-dimensional space. It is a surface of revolution of a tractrix about its asymptote (along the  $\hat{z}$ -axis). We can make coordinates for the surface  $(t, \theta)$  where  $t$  is the coordinate along the asymptote and  $\theta$  is the angle of revolution. We embed the pseudosphere in rectangular 3-dimensional space with

```
(defn pseudosphere
  [q]
  (let [t (ref q 0)
        theta (ref q 1)]
    (up (* (sech t) (cos theta)) (* (sech t) (sin theta)) (- t (tanh t)))))
```

The structure of Christoffel coefficients for the pseudosphere is



where  $d_{ij}^k$  are the structure constants of the basis. See equations (4.37), (4.38). For a commuting basis the structure constants are zero, and the components of the torsion are the antisymmetric part of  $\Gamma$  with respect to the lower indices.

Recall the geodesic equation (7.79):

$$D^2\sigma^i(t) = \sum_{jk} \Gamma_{jk}^i(\gamma(t)) D\sigma^j(t) D\sigma^k(t) = 0. \quad (8.22)$$

Observe that the lower indices of  $\Gamma$  are contracted with two copies of the velocity. Because the use of  $\Gamma$  is symmetrical here, any asymmetry of  $\Gamma$  in its lower indices is irrelevant to the geodesics. Thus one can study the geodesics of any connection by first symmetrizing the connection, eliminating torsion. The resulting equations will be simpler.

### 8.3 Geodesic Deviation

Geodesics may converge and intersect (as in the lines of longitude on a sphere) or they may diverge (for example, on a saddle). To capture this notion requires some measure of the convergence or divergence, but this requires metrics (see Chapter 9). But even in the absence of a metric we can define a quantity, the *geodesic deviation*, that can be interpreted in terms of relative acceleration of neighboring geodesics from a reference geodesic.

Let there be a one-parameter family of geodesics, with parameter  $s$ , and let  $T$  be the vector field of tangent vectors to those geodesics:

$$\nabla_T T = 0. \quad (8.23)$$

We can parameterize travel along the geodesics with parameter  $t$ : a geodesic curve  $\gamma_s(t) = \phi_t^T(m_s)$  where

$$f \circ \phi_t^T(m_s) = (e^{tT} f)(m_s). \quad (8.24)$$

Let  $U = \partial/\partial s$  be the vector field corresponding to the displacement of neighboring geodesics. Locally,  $(t, s)$  is a coordinate system on the 2-dimensional submanifold formed by the family of geodesics. The vector fields  $T$  and  $U$  are a coordinate basis for this coordinate system, so  $[T, U] = 0$ .

The geodesic deviation vector field is defined as:

$$\nabla_T(\nabla_T U). \quad (8.25)$$

If the connection has zero torsion, the geodesic deviation can be related to the Riemann curvature:

$$\nabla_T(\nabla_T U) = -\mathcal{R}(U, T)(T), \quad (8.26)$$

as follows, using equation (8.21),

$$\nabla_T(\nabla_T U) = \nabla_T(\nabla_U T), \quad (8.27)$$

because both the torsion is zero and  $[T, U] = 0$ . Continuing

$$\begin{aligned} \nabla_T(\nabla_T U) &= \nabla_T(\nabla_U T) \\ &= \nabla_T(\nabla_U T) + \nabla_U(\nabla_T T) - \nabla_U(\nabla_T T) \\ &= \nabla_U(\nabla_T T) - \mathcal{R}(U, T)(T) \\ &= -\mathcal{R}(U, T)(T). \end{aligned} \quad (8.28)$$

In the last line the first term was dropped because  $T$  satisfies the geodesic equation (8.24).

The geodesic deviation is defined without using a metric, but it helps to have a metric (see Chapter 9) to interpret the geodesic deviation. Consider two neighboring geodesics, with parameters  $s$  and  $s + \Delta s$ . Given a metric we can assume that  $t$  is proportional to path length along each geodesic, and we can define a distance  $\delta(s, t, \Delta s)$  between the geodesics at the same value of the parameter  $t$ . So the velocity of separation of the two geodesics is

$$(\nabla_T U) = \partial_1 \delta(s, t, \Delta s) \hat{s} \quad (8.29)$$

where  $\hat{s}$  is a unit vector in the direction of increasing  $s$ . So  $\nabla_T U$  is the factor of increase of velocity with increase of separation. Similarly, the geodesic deviation can be interpreted as the factor of increase of acceleration with increase of separation:

$$\nabla T(\nabla_T U) = \partial_1 \partial_1 \delta(s, t, \Delta s) \hat{s}. \quad (8.30)$$

### 8.3.1 Longitude Lines on a Sphere

Consider longitude lines on the unit sphere.<sup>7</sup> Let `theta` be colatitude and `phi` be longitude. These are the parameters  $s$  and  $t$ , respectively. Then let `T` be the vector field `d/dtheta` that is tangent to the longitude lines.

We can verify that every longitude line is a geodesic:

```
((omega ((covariant-derivative Cartan) T) T)) m)
;; => 0
```

closure

where `omega` is an arbitrary one-form field.

Now let `U` be `d/dphi`, then `U` commutes with `T`:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((commutator U T) f) m))
;; => 0
```

closure

The torsion for the usual connection for the sphere is zero:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (let [X (literal-vector-field 'X-sphere S2-spherical)
                Y (literal-vector-field 'Y-sphere S2-spherical)]
            (((torsion-vector nabla) X Y) f) m)))
;; => 0
```

closure

So we can compute the geodesic deviation using `Riemann`

```
(simplify ((+ (omega ((nabla T) ((nabla T) U))) ((Riemann nabla) omega T U T)) m))
;; => 0
```

closure

confirming equation (8.29).

Lines of longitude are geodesics. How do the lines of longitude behave? As we proceed from the North Pole, the lines of constant longitude diverge. At the Equator they are parallel and they converge towards the South Pole.

Let's compute  $\nabla_T U$  and  $\nabla_T(\nabla_T U)$ . We know that the distance is purely in the  $\phi$  direction, so

---

<sup>7</sup>The setup for this example is:

```
(define-coordinates (up theta phi) S2-spherical)
(def T d:dtheta)
(def U d:dphi)
(def m ((point S2-spherical) (up 'theta0 'phi0)))
(def Cartan (Christoffel→Cartan S2-Christoffel))
(def nabla (covariant-derivative Cartan))
(def omega (literal-oneform-field 'omega-sphere S2-spherical))
(def f (literal-manifold-function 'f S2-spherical))
```

closure

```
((dphi ((nabla T) U)) m)
;; ⇒ (* (/ 1 (expt (sin theta0) 2)) (/ 1 2) 2 (sin theta0) (cos theta0))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((dphi ((nabla T) ((nabla T) U))) m))
;; ⇒ -1
```

Let's interpret these results. On a sphere of radius  $R$  the distance at colatitude  $\theta$  between two geodesics separated by  $\Delta\phi$  is  $d(\phi, \theta, \Delta\phi) = R \sin(\theta) \Delta\phi$ . Assuming that  $\theta$  is uniformly increasing with time, the magnitude of the velocity is just the  $\theta$ -derivative of this distance:

```
(defn delta [R] (fn [phi theta Delta-phi] (* R (sin theta) Delta-phi)))
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((partial 1) (delta 'R)) 'phi0 'theta0 'Delta-phi))
;; ⇒ (* Delta-phi R (cos theta0))
```

The direction of the velocity is the unit vector in the  $\phi$  direction:

```
(def phi-hat (* (/ 1 (sin theta)) d:dphi))
```

This comes from the fact that the separation of lines of longitude is proportional to the sine of the colatitude. So the velocity vector field is the product.

We can measure the  $\phi$  component with  $d\phi$ :

```
((dphi (* (((partial 1) (delta 'R)) 'phi0 'theta0 'Delta-phi) phi-hat)) m)
;; ⇒ (* Delta-phi R (cos theta0) (/ 1 (sin theta0)))
```

This agrees with  $\nabla_T U \Delta\phi$  for the unit sphere. Indeed, the lines of longitude diverge until they reach the Equator and then they converge.

Similarly, the magnitude of the acceleration is

```
((partial 1) ((partial 1) (delta 'R))) 'phi0 'theta0 'Delta-phi
;; ⇒ (* Delta-phi R (- (sin theta0)))
```

and the acceleration vector is the product of this result with  $\hat{\phi}$ . Measuring this with  $d\phi$  we get:

```
((dphi (* (((partial 1) ((partial 1) (delta 'R))) 'phi0 'theta0 'Delta-phi) phi-hat)) m)
;; ⇒ (* Delta-phi R (- (sin theta0)) (/ 1 (sin theta0)))
```

And this agrees with the calculation of  $\nabla_T \nabla_T U \Delta\phi$  for the unit sphere. We see that the separation of the lines of longitude are uniformly decelerated as they progress from pole to pole.

## 8.4 Bianchi Identities

There are some important mathematical properties of the Riemann curvature. These identities will be used to constrain the possible geometries that can occur.

A system with a symmetric connection,  $\Gamma_{jk}^i = \Gamma_{kj}^i$ , is torsion free.<sup>8</sup>

---

<sup>8</sup>Setup for this section:

```
(def omega (literal-oneform-field 'omega-rect R4-rect))
(def X (literal-vector-field 'X-rect R4-rect))
(def Y (literal-vector-field 'Y-rect R4-rect))
```

```
(def nabla
  (covariant-derivative (Christoffel→Cartan (symmetrize-Christoffel
                                            (literal-Christoffel-2 'C R4-rect)))))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((torsion nabla) omega X Y) (typical-point R4-rect)))
;; ⇒ 0
```

The Bianchi identities are defined in terms of a cyclic-summation operator, which is most easily described as a ClojureScript function:

```
(defn cyclic-sum [f] (fn [x y z] (+ (f x y z) (f y z x) (f z x y)))))
```

The first Bianchi identity is

$$R(\omega, x, y, z) + R(\omega, y, z, x) + R(\omega, z, x, y) = 0, \quad (8.31)$$

or, as a program:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((cyclic-sum (fn [x y z] ((Riemann nabla) omega x y z))) X Y Z) (typical-
point R4-rect)))
;; ⇒ 0
```

The second Bianchi identity is

$$\nabla_x R(\omega, v, y, z) + \nabla_y R(\omega, v, z, x) + \nabla_z R(\omega, v, x, y) = 0 \quad (8.32)$$

or, as a program:

```
((cyclic-sum (fn [x y z] (((nabla x) (Riemann nabla)) omega V y z))) X Y Z)
(typical-point R4-rect))
```

Things get more complicated when there is torsion. We can make a general connection, which has torsion:

```
(def nabla (covariant-derivative (Christoffel→Cartan (literal-Christoffel-2 'C R4-
rect))))

(def R (Riemann nabla))

(def T (torsion-vector nabla))

(defn TT [omega x y] (omega (T x y)))
```

The first Bianchi identity is now:<sup>9</sup>

```
((cyclic-sum (fn [x y z] (- (R omega x y z) (+ (omega (T (T x y) z)) (((nabla x) TT)
omega y z)))))
X
Y
Z)
(typical-point R4-rect))
```

and the second Bianchi identity for a general connection is

```
(def Z (literal-vector-field 'Z-rect R4-rect))

(def V (literal-vector-field 'V-rect R4-rect))
```

<sup>9</sup>The Bianchi identities are much nastier to write in traditional mathematical notation than as ClojureScript programs.

```
(((cyclic-sum (fn [x y z] (+ (((nabla x) R) omega V y z) (R omega V (T x y) z)))) X Y closure
Z)
(typical-point R4-rect))
```

We now make the Cartan forms from the most general 2-dimensional Christoffel coefficient structure:

```
(def general-Cartan-2 (Christoffel→Cartan (literal-Christoffel-2 'Gamma R2-rect))) closure
```

[11], [4], and [14] use our definition. [20] uses a different convention for the order of arguments and a different sign. See Appendix F for a definition of tensors in Emmy.

## 9 Metrics

We often want to impose further structure on a manifold to allow us to define lengths and angles. This is done by generalizing the idea of the Euclidean dot product, which allows us to compute lengths of vectors and angles between vectors in traditional vector algebra.

For vectors  $\vec{u} = u^x \hat{x} + u^y \hat{y} + u^z \hat{z}$  and  $\vec{v} = v^x \hat{x} + v^y \hat{y} + v^z \hat{z}$  the dot product is  $\vec{u} \cdot \vec{v} = u^x v^x + u^y v^y + u^z v^z$ . The generalization is to provide coefficients for these terms and to include cross terms, consistent with the requirement that the function of two vectors is symmetric. This symmetric, bilinear, real-valued function of two vector fields is called a *metric field*.

For example, the natural metric on a sphere of radius  $R$  is

$$g(u, v) = R^2(d\theta(u)d\theta(v) + (\sin \theta)^2 d\phi(u)d\phi(v)), \quad (9.1)$$

and the Minkowski metric on the 4-dimensional space of special relativity is

$$g(u, v) = dx(u)dx(v) + dy(u)dy(v) + dz(u)dz(v) - c^2 dt(u)dt(v). \quad (9.2)$$

Although these examples are expressed in terms of a coordinate basis, the value of the metric on vector fields does not depend on the coordinate system that is used to specify the metric.

Given a metric field  $g$  and a vector field  $v$  the scalar field  $g(v, v)$  is the squared length of the vector at each point of the manifold.

### Metric Music

The metric can be used to construct a one-form field  $\omega_u$  from a vector field  $u$ , such that for any vector field  $v$  we have

$$\omega_u(v) = g(v, u). \quad (9.3)$$

The operation of constructing a one-form field from a vector field using a metric is called “lowering” the vector field. It is sometimes notated as

$$\omega_u = g^\flat(u). \quad (9.4)$$

There is also an inverse metric that takes two one-form fields. It is defined by the relation

$$\delta_k^i = \sum_j g^{-1}(\tilde{e}^i, \tilde{e}^j) g(e_j, e_k). \quad (9.5)$$

where  $e$  and  $\tilde{e}$  are any basis and its dual basis.

The inverse metric can be used to construct a vector field  $v_\omega$  from a one-form field  $\omega$ , such that for any one-form field  $\tau$  we have

$$\tau(v_\omega) = g^{-1}(\omega, \tau). \quad (9.6)$$

This definition is implicit, but the vector field can be explicitly computed from the one-form field with respect to a basis as follows:

$$v_\omega = \sum_i g^{-1}(\omega, \tilde{e}^i) e_i. \quad (9.7)$$

The operation of constructing a vector field from a one-form field using a metric is called “raising” the one-form field. It is sometimes notated

$$v_\omega = g^\sharp(\omega). \quad (9.8)$$

The raising and lowering operations allow one to interchange the vector fields and the one-form fields. However they should not be confused with the dual operation that allows one to construct a dual one-form basis from a vector basis or construct a vector basis from a one-form basis. The dual operation that interchanges bases is defined without assigning a metric structure on the space.

Lowering a vector field with respect to a metric is a simple program:



```
(defn lower
  [metric]
  (fn [u]
    (letfn [(omega [v] (metric v u))]
      (procedure→oneform-field omega))))
```

clojure

But raising a one-form field to make a vector field is a bit more complicated:

```
(defn raise
  [metric basis]
  (let [gi (invert metric basis)]
    (fn [omega] (contract (fn [e i wfi] (* (gi omega wfi) e i)) basis))))
```

clojure

where `contract` is the trace over a basis of a two-argument function that takes a vector field and a one-form field as its arguments.<sup>1</sup>

```
(defn contract
  [proc basis]
  (let [vector-basis (basis→vector-basis basis)
        oneform-basis (basis→oneform-basis basis)]
    (sumr proc vector-basis oneform-basis)))
```

clojure

## 9.1 Metric Compatibility

A connection is said to be compatible with a metric  $g$  if the covariant derivative for that connection obeys the “product rule”:

$$\Delta_X(g(Y, Z)) = g(\Delta_X(Y), Z) + g(Y, \Delta_X(Z)). \quad (9.9)$$

For a metric there is a unique torsion-free connection that is compatible with it. The Christoffel coefficients of the first kind are computed from the metric by the following:

$$\bar{\Gamma}_{ijk} = \frac{1}{2} (e_k(g(e_i, e_j)) + e_j(g(e_i, e_k)) - e_i(g(e_j, e_k))) \quad (9.10)$$

for the coordinate basis  $\mathbf{e}$ . We can then construct the Christoffel coefficients of the second kind (the ones used previously to define a connection) by “raising the first index.” To do this we define a function of three vectors, with a weird currying:

$$\sum_{ijk} \bar{\Gamma}_{ijk} \tilde{\mathbf{e}}^i(u) \tilde{\mathbf{e}}^j(v) \tilde{\mathbf{e}}^k(w). \quad (9.11)$$

This function takes two vector fields and produces a one-form field. We can use it with equation (9.7) to construct a new function that takes two vector fields and produces a vector field:

$$\hat{\Gamma}(v, w) = \sum_i g^{-1}(\tilde{\Gamma}(v, w), \tilde{\mathbf{e}}^i) \mathbf{e}_i. \quad (9.12)$$

We can now construct the Christoffel coefficients of the second kind:

$$\Gamma_{jk}^i = \tilde{\mathbf{e}}^i(\hat{\Gamma}(\mathbf{e}_j, \mathbf{e}_k)) = \sum_m \bar{\Gamma}_{mjk} g^{-1}(\tilde{\mathbf{e}}^m, \tilde{\mathbf{e}}^i) \quad (9.13)$$

The Cartan forms are then just

$$\varpi_j^i = \sum_k \Gamma_{jk}^i \tilde{\mathbf{e}}^k = \sum_k \tilde{\mathbf{e}}^i(\hat{\Gamma}(\mathbf{e}_j, \mathbf{e}_k)) \tilde{\mathbf{e}}^k. \quad (9.14)$$

So, for example, we can compute the Christoffel coefficients for the sphere from the metric for the sphere. First, we need the metric:

---

<sup>1</sup>Notice that `raise` and `lower` are not symmetrical. This is because vector fields and form fields are not symmetrical: a vector field takes a manifold function as its argument, whereas a form field takes a vector field as its argument. This asymmetry is not apparent in traditional treatments based on index notation.

```
(defn g-sphere
  [R]
  (fn [u v]
    (* (square R)
      (+ (* (dtheta u) (dtheta v)) (* (compose (square sin) theta) (dphi u) (dphi
v))))))
```

closure

The Christoffel coefficients of the first kind are a complex structure with all three indices down:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((Christoffel→symbols (metric→Christoffel-1 (g-sphere 'R) S2-basis))
  ((point S2-spherical) (up 'theta0 'phi0))))
;; ⇒ (down (down (down 0 0) (down 0 (* (expt R 2) (sin theta0) (cos theta0))))
;;      (down (down 0 (* (expt R 2) (sin theta0) (cos theta0))))
;;      (down (* #emmy/bigint -1 (expt R 2) (sin theta0) (cos theta0)) 0)))
```

closure

And the Christoffel coefficients of the second kind have the innermost index up:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((Christoffel→symbols (metric→Christoffel-2 (g-sphere 'R) S2-basis))
  ((point S2-spherical) (up 'theta0 'phi0))))
;; ⇒ (down (down (up 0 0) (up 0 (/ (cos theta0) (sin theta0))))
;;      (down (up 0 (/ (cos theta0) (sin theta0))))
;;      (up (* #emmy/bigint -1 (sin theta0) (cos theta0)) 0)))
```

closure

### 9.1.1 Exercise 9.1: Metric Compatibility

The connections constructed from a metric by equation (9.13) are “metric compatible,” as described in equation (9.9). Demonstrate that this is true for a literal metric, as described in Section 1.2, page 3, in  $\mathbf{R}^4$ . Your program should produce a zero.

## 9.2 Metrics and Lagrange Equations

In the Introduction (Chapter 1) we showed that the Lagrange equations for a free particle constrained to a 2-dimensional surface are equivalent to the geodesic equations for motion on that surface. We illustrated that in detail in Section 7.4 for motion on a sphere.

Here we expand this understanding to show that the Christoffel symbols can be derived from the Lagrange equations. Specifically, if we solve the Lagrange equations for the acceleration (the highest-order derivatives) we find that the Christoffel symbols are the symmetrized coefficients of the quadratic velocity terms.

Consider the Lagrange equations for a free particle, with Lagrangian

$$L_2(t, x, v) = \frac{1}{2}g(x)(v, v). \quad (9.15)$$

If we solve the Lagrange equations for the accelerations, the accelerations can be expressed with the geodesic equations (7.79):

$$D^2q^i + \sum_{jk} (\Gamma_{jk}^i \circ \chi^{-1} \circ q) Dq^j Dq^k = 0. \quad (9.16)$$

We can verify this computationally. Given a metric, we can construct a Lagrangian where the kinetic energy is the metric applied to the velocity twice: The kinetic energy is proportional to the squared length of the velocity vector.

```
(defn metric→Lagrangian
  [metric coordsys]
  (letfn [(L [state]
    (let [q (ref state 1)
          qd (ref state 2)]
      (let [v (components→vector-field (fn [m] qd) coordsys)]
```

closure

```

L))
      ((* (/ 1 2) (metric v v)) ((point coordsys) q))))]]

```

The following code compares the Christoffel symbols with the coefficients of the terms of second order in velocity appearing in the accelerations, determined by solving the Lagrange equations for the highest-order derivative.<sup>2</sup> We extract these terms by taking two partials with respect to the structure of velocities. Because the elementary partials commute we get two copies of each coefficient, requiring a factor of 1/2.

```

(simplify
  (let [metric (literal-metric 'g R3-rect)
        q (typical-coords R3-rect)
        L2 (metric→Lagrangian metric R3-rect)]
    (+ (* (/ 1 2)
          ((expt (partial 2) 2) (Lagrange-explicit L2)) (up 't q (corresponding-
            velocities q))))
        ((Christoffel→symbols (metric→Christoffel-2 metric (coordinate-system→basis
          R3-rect)))
          ((point R3-rect) q))))))
;; ⇒ (down (down (up 0 0 0) (up 0 0 0) (up 0 0 0))
;;      (down (up 0 0 0) (up 0 0 0) (up 0 0 0))
;;      (down (up 0 0 0) (up 0 0 0) (up 0 0 0)))

```

We get a structure of zeros, demonstrating the correspondence between Christoffel symbols and coefficients of the Lagrange equations.

Thus, if we have a metric specifying an inner product, the geodesic equations are equivalent to the Lagrange equations for the Lagrangian that is equal to the inner product of the generalized velocities with themselves

### 9.3 Kinetic Energy or Arc Length

A geodesic is a path of stationary length with respect to variations in the path that keep the endpoints fixed. On the other hand, the solutions of the Lagrange equations are paths of stationary action that keep the endpoints fixed. How are these solutions related?

The integrand of the traditional action is the Lagrangian, which is in this case the Lagrangian  $L_2$ , the kinetic energy. The integrand of the arc length is

$$L_1(t, x, v) = \sqrt{g(x)(v, v)} = \sqrt{2L_2(t, x, v)} \quad (9.17)$$

and the path length is

$$\tau = \int_{t_1}^{t_2} L_1(t, q(t), Dq(t)) dt. \quad (9.18)$$

If we compute the Lagrange equations for  $L_2$  we get the Lagrange equations for  $L_1$  with a correction term. Since

$$L_2(t, x, v) = \frac{1}{2}(L_1(t, x, v))^2, \quad (9.19)$$

and the Lagrange operator for  $L_2$  is<sup>3</sup>

```

(defn Lagrange-explicit
  [L]
  (let [P ((partial 2) L)
        F ((partial 1) L)]
    (/ (- F (+ ((partial 0) P) (* ((partial 1) P) velocity))) ((partial 2) P))))

```

<sup>2</sup>The procedure `Lagrange-explicit` produces the accelerations of the coordinates. In this code the division operator `(/)` multiplies its first argument on the left by the inverse of its second argument.

<sup>3</sup> $\mathbf{E}$  is the Euler-Lagrange operator, which gives the residuals of the Lagrange equations for a Lagrangian.  $\mathbf{\Gamma}$  extends a configuration-space path  $q$  to make a state-space path, with as many terms as needed:  $\mathbf{\Gamma}[q](t) = (t, q(t), Dq(t), \dots)$ . The total

$$\mathbf{E}[L_2] = D_t \partial_2 L_2 - \partial_1 L_2,$$

we find

$$\mathbf{E}[L_2] = L_1 \mathbf{E}[L_1] + \partial_2 L_1 D_t L_1. \quad (9.20)$$

$L_2$  is the kinetic energy. It is conserved along solution paths, since there is no explicit time dependence. Because of the relation between  $L_1$  and  $L_2$ ,  $L_1$  is also a conserved quantity. Let  $L_1$  take the constant value  $a$  on the geodesic coordinate path  $q$  we are considering. Then  $\tau = a(t_2 - t_1)$ . Since  $L_1$  is conserved,  $(D_t L_1) \circ \Gamma[q] = 0$  on the geodesic path  $q$ , and both  $\mathbf{E}[L_1] \circ \Gamma[q] = 0$  and  $\mathbf{E}[L_2] \circ \Gamma[q] = 0$ , as required by equation (9.20).

Since  $L_2$  is homogeneous of degree 2 in the velocities,  $L_1$  is homogeneous of degree 1. So we cannot solve for the highest-order derivative in the Lagrange-Euler equations derived from  $L_1$ : The Lagrange equations of the Lagrangian  $L_1$  are dependent. But although they do not uniquely specify the evolution, they do specify the geodesic path.

On the other hand, we can solve for the highest-order derivative in  $\mathbf{E}[L_2]$ . This is because  $L_1 \mathbf{E}[L_1]$  is homogeneous of degree 2. So the equations derived from  $L_2$  uniquely

### 9.3.1 For Two Dimensions

We can show this is true for a 2-dimensional system with a general metric. We define the Lagrangians in terms of this metric:

```
(def L2 (metric→Lagrangian (literal-metric 'm R2-rect) R2-rect)) ;closure
(defn L1 [state] (sqrt (* 2 (L2 state))))
```

Although the mass matrix of  $L_2$  is nonsingular

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. ;closure
(simplify (determinant (((partial 2) ((partial 2) L2)) (up 't (up 'x 'y) (up 'vx
'vy')))))
;; ⇒ (+ (* (m_00 (up x y)) (m_11 (up x y)))
;;      (* #emmy/bigint -1 (expt (m_01 (up x y)) 2)))
```

the mass matrix of  $L_1$  has determinant zero

```
(simplify (determinant (((partial 2) ((partial 2) L1)) (up 't (up 'x 'y) (up 'vx
'vy'))))) ;closure
;; ⇒ 0
```

showing that these Lagrange equations are dependent.

We can show this dependence explicitly, for a simple system. Consider the simplest possible system, a geodesic (straight line) in a plane:

```
(defn L1 [state] (sqrt (square (velocity state)))) ;closure

;; Emmy expands speed^2^(3/2) into speed^2 sqrt(speed^2). Cancel that common
;; denominator before simplifying the numerators, then put the compact common
;; denominator back so the displayed result remains readable.
(let [x (literal-function 'x)
      y (literal-function 'y)
      equations (((Lagrange-equations L1) (up x y)) 't)
      speed2 (+ (square ((D x) 't)) (square ((D y) 't)))
      denominator (* speed2 (sqrt speed2))]
  (/ (simplify (* equations denominator)) denominator))
;; ⇒ (down (* (+ (* -1 ((D x) t) ((D y) t) ((expt D 2) y) t))
```

---

time derivative  $D_t$  is defined by  $D_t F \circ \Gamma[q] = D(F \circ \Gamma[q])$  for any state function  $F$  and path  $q$ . The Lagrange equations are  $\mathbf{E}[L] \circ \Gamma[q] = 0$ . See [19] for more details.

```

;;      (* (((expt D 2) x) t) (expt ((D y) t) 2)))
;;      (/ 1
;;      (* (+ (expt ((D x) t) 2) (expt ((D y) t) 2))
;;          (sqrt (+ (expt ((D x) t) 2) (expt ((D y) t) 2))))))
;;      (* (+ (* (expt ((D x) t) 2) (((expt D 2) y) t))
;;          (* -1 ((D x) t) (((expt D 2) x) t) ((D y) t)))
;;      (/ 1
;;      (* (+ (expt ((D x) t) 2) (expt ((D y) t) 2))
;;          (sqrt (+ (expt ((D x) t) 2) (expt ((D y) t) 2))))))

```

These residuals must be zero; so the numerators must be zero.<sup>4</sup> They are:

$$D^2x(Dy)^2 = Dx Dy D^2y D^2x Dx Dy = (Dx)^2 D^2y$$

Note that the only constraint is  $D^2x Dy = Dx D^2y$ , so the resulting Lagrange equations are dependent.

This is enough to determine that the result is a straight line, without specifying the rate along the line. Suppose  $y = f(x)$ , for path  $(x(t), y(t))$ . Then

$$Dy = Df(x) Dx \text{ and } D^2y = D^2f(x) Dx + Df(x) D^2x.$$

Substituting, we get

$$Df(x) Dx D^2x = Dx(D^2f(x) Dx + Df(x) D^2x)$$

or

$$Df(x) D^2x = D^2f(x) Dx + Df(x) D^2x,$$

so  $D^2f(x) = 0$ . Thus  $f$  is a straight line, as required.

### 9.3.2 Reparametrization

More generally, a differential equation system  $F[q](t) = 0$  is said to be *reparameterized* if the coordinate path  $q$  is replaced with a new coordinate path  $q \circ f$ . For example, we may change the scale of the independent variable. The system  $F[q \circ f] = 0$  is said to be independent of the parameterization if and only if  $F[q] \circ f = 0$ . So the differential equation system is satisfied by  $q \circ f$  if and only if it is satisfied by  $q$ .

The Lagrangian  $L_1$  is homogeneous of degree 1 in the velocities; so

$$E[L_1] \circ \Gamma[q \circ f] - (E[L_1] \circ \Gamma[q] \circ f) Df = 0. \quad (9.21)$$

We can check this in a simple case. For two dimensions  $q = (x, y)$ , the condition under which a reparameterization  $f$  of the geodesic paths with coordinates  $q$  satisfies the Lagrange equations for  $L_1$  is:

```

;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (let [x (literal-function 'x)
               y (literal-function 'y)
               f (literal-function 'f)
               E1 (Euler-Lagrange-operator L1)]
  ((- (compose E1 (Gamma (up (compose x f) (compose y f)) 4))
      (* (compose E1 (Gamma (up x y) 4) f) (D f)))
    't)))
;; => (down 0 0)

```

This residual is identically satisfied, showing that the Lagrange equations for  $L_1$  are independent of the parameterization of the independent variable.

The Lagrangian  $L_2$  is homogeneous of degree 2 in the velocities; so

$$E[L_2][q \circ f] - (E[L_2][q] \circ f)(Df)^2 = (\partial_2 L_2 \circ \Gamma[q] \circ f)(D^2f). \quad (9.22)$$

Although the Euler-Lagrange equations for  $L_1$  are invariant under an arbitrary reparameterization ( $Df \neq 0$ ), the Euler-Lagrange equations for  $L_2$  are invariant only for a restricted set of  $f$ . The conditions under

<sup>4</sup>We cheated: We hand-simplified the denominator to make the result more obvious.

which a reparameterization  $f$  of geodesic paths with coordinates  $q$  satisfies the Lagrange equations for  $L_2$  are:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (let [q (up (literal-function 'x) (literal-function 'y))
                f (literal-function 'f)]
  ((- (compose (Euler-Lagrange-operator L2) (Gamma (compose q f) 4))
      (* (compose (Euler-Lagrange-operator L2) (Gamma q 4) f) (expt (D f)
2))))
  't)))
;; => (down (+ (* ((D x) (f t)) (((expt D 2) f) t) (m_00 (up (x (f t)) (y (f t))))))
;;          (* ((D y) (f t)) (((expt D 2) f) t) (m_01 (up (x (f t)) (y (f t))))))
;;          (+ (* ((D x) (f t)) (((expt D 2) f) t) (m_01 (up (x (f t)) (y (f t))))))
;;          (* ((D y) (f t)) (((expt D 2) f) t) (m_11 (up (x (f t)) (y (f t)))))))
```

We see that if these expressions must be zero, then  $D^2f = 0$ . This tells us that  $f$  is at most affine in  $t$ :  $f(t) = at + b$ .

### 9.3.3 Exercise 9.2: SO(3) Geodesics

We have derived a basis for SO(3) in terms of incremental rotations around the rectangular axes. See equations (4.29), (4.30), (4.31). We can use the dual basis to define a metric on SO(3).

```
(defn S03-metric [v1 v2] (+ (* (e-x v1) (e-x v2)) (* (e-y v1) (e-y v2)) (* (e-z v1)
(e-z v2))))      clojure
```

This metric determines a connection. Show that uniform rotation about an arbitrary axis traces a geodesic on SO(3).

### 9.3.4 Exercise 9.3: Curvature of a Spherical Surface

The 2-dimensional surface of a 3-dimensional sphere can be embedded in three dimensions with a metric that depends on the radius:

```
(def M (make-manifold S2-type 2 3))      clojure

(def spherical (coordinate-system-at M :spherical :north-pole))

(define-coordinates (up theta phi) spherical)

(def spherical-basis (coordinate-system->basis spherical))

(defn spherical-metric
  [r]
  (fn [v1 v2]
    (* (square r) (+ (* (dtheta v1) (dtheta v2)) (* (square (sin theta)) (dphi v1)
(dphi v2))))))
```

If we raise one index of the Ricci tensor (see equation (8.20)) by contracting it with the inverse of the metric tensor we can further contract it to obtain a scalar manifold function:

$$R = \sum_{ij} g(\tilde{e}^i, \tilde{e}^j) r(e^i, e^j). \quad (9.23)$$

The `trace2down` procedure converts a tensor that takes two vector fields into a tensor that takes a vector field and a one-form field, and then it contracts the result over a basis to make a trace. It is useful for getting the Ricci scalar from the Ricci tensor, given a metric and a basis.

```
(defn trace2down      clojure
  [metric basis]
  (fn [tensor]
    (let [inverse-metric-tensor (invert metric basis)]
      (contract (fn [v1 w1]
        (contract (fn [v w] (* (inverse-metric-tensor w1 w) (tensor v v1))))
```

basis))

basis)))

Evaluate the Ricci scalar for a sphere of radius  $r$  to obtain a measure of its intrinsic curvature. You should obtain the answer  $2/r^2$ .

### 9.3.5 Exercise 9.4: Curvature of a Pseudosphere

Compute the scalar curvature of the pseudosphere (see exercise 8.2). You should obtain the value  $-2$ .

## 9.4 General Relativity

By analogy to Newtonian mechanics, relativistic mechanics has two parts. There are equations of motion that describe how particles move under the influence of “forces” and there are field equations that describe how the forces arise. In general relativity the only force considered is gravity. However, gravity is not treated as a force. Instead, gravity arises from curvature in the spacetime, and the equations of motion are motion along geodesics of that space.

The geodesic equations for a spacetime with the metric

$$g(v_1, v_2) = -c^2 1 + \frac{2V}{c^2} dt(v_1)dt(v_2) + dx(v_1)dx(v_2) + dy(v_1)dy(v_2) + dz(v_1)dz(v_2) \quad (9.24)$$

are Newton's equations to lowest order in  $V/c^2$ :

$$D^2 \vec{x}(t) = -\text{grad } V(\vec{x}(t)). \quad (9.25)$$

### 9.4.1 Exercise 9.5: Newton's Equations

Verify that Newton's equations (9.25) are indeed the lowest-order terms of the geodesic equations for the metric (9.24).

Einstein's field equations tell how the local energy-momentum distribution determines the local shape of the spacetime, as described by the metric tensor  $g$ . The equations are traditionally written

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} + \Lambda g_{\mu\nu} = \frac{8\pi G}{c^4}T_{\mu\nu} \quad (9.26)$$

where  $R_{\mu\nu}$  are the components of the Ricci tensor (equation (8.20)),  $R$  is the Ricci scalar (equation (9.23)),<sup>5</sup> and  $\Lambda$  is the cosmological constant.

$T_{\mu\nu}$  are the components of the stress-energy tensor describing the energy-momentum distribution. Equivalently, one can write

$$R_{\mu\nu} = \frac{8\pi G}{c^4} \left( T_{\mu\nu} - \frac{1}{2}Tg_{\mu\nu} \right) - \Lambda g_{\mu\nu} \quad (9.27)$$

where  $T = T_{\mu\nu}g^{\mu\nu}$ .<sup>6</sup>

Einstein's field equations arise from a heuristic derivation by analogy to the Poisson equation for a Newtonian gravitational field:

$$\text{Lap}(V) = 4\pi G\rho \quad (9.28)$$

where  $V$  is the gravitational potential field at a point,  $\rho$  is the mass density at that point, and  $\text{Lap}$  is the Laplacian operator.

The time-time component of the Ricci tensor derived from the metric (9.24) is the Laplacian of the potential, to lowest order.

<sup>5</sup>The tensor with components  $G_{\mu\nu} = R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu}$  is called the Einstein tensor. In his search for an appropriate field equation for gravity, Einstein demanded *general covariance* (independence of coordinate system) and local Lorentz invariance (at each point transformations must preserve the line element). These considerations led Einstein to look for a tensor equation (see Appendix F).

<sup>6</sup>Start with equation (9.26). Raise one index of both sides, and then contract. Notice that the trace  $g^\mu_\mu = 4$ , the dimension of spacetime. This gets  $R = -(\frac{8\pi G}{c^4})T$ , from which we can deduce equation (9.27).

```

(define-coordinates (up t x y z) spacetime-rect)

(def spacetime-rect-basis (coordinate-system→basis spacetime-rect))

(defn Newton-metric
  [M G c V]
  (let [a (+ 1 (* (/ 2 (square c)) (compose V (up x y z))))]
    (letfn [(g [v1 v2]
              (+ (* -1 (square c) a (dt v1) (dt v2))
                 (* (dx v1) (dx v2))
                 (* (dy v1) (dy v2))
                 (* (dz v1) (dz v2))))]
      g)))

(defn Newton-connection
  [M G c V]
  (Christoffel→Cartan (metric→Christoffel-2 (Newton-metric M G c V) spacetime-rect-basis)))

(def V (literal-function 'V '(→ (UP Real Real Real) Real)))

(def nabla (covariant-derivative (Newton-connection 'M 'G 'c V)))

(((Ricci nabla (coordinate-system→basis spacetime-rect)) d:dt d:dt)
 (point spacetime-rect) (up 't 'x 'y 'z)))
;; ⇒ (+
;;      (- (+ (* (/ (* -1 (expt c 2) (+ 1 (* (/ 2 (expt c 2)) (V (up x y z)))))
;;            (* -1 (expt c 2) (+ 1 (* (/ 2 (expt c 2)) (V (up x y z)))))
;;          (/ 1 2)
;;      ""
;; [full result retained in chapter009/019.cljs]

```

The leading terms of the mess are

```

(+ (((partial 0) ((partial 0) V)) (up 'x 'y 'z))
    (((partial 1) ((partial 1) V)) (up 'x 'y 'z))
    (((partial 2) ((partial 2) V)) (up 'x 'y 'z)))
;; ⇒ (+ (((partial 0) ((partial 0) V)) (up x y z))
;;      (((partial 1) ((partial 1) V)) (up x y z))
;;      (((partial 2) ((partial 2) V)) (up x y z)))

```

which is the Laplacian of  $V$ . The other terms are smaller by  $V/c^2$ .

Now consider the right-hand side of equation (9.27). In the Poisson equation the source of the gravitational potential is the density of matter. Let the time-time component of the stress-energy tensor  $T_{00}$  be the matter density  $\rho$ . Here is a program for the stress-energy tensor:

```

(defn Tdust
  [rho]
  (letfn [(T [w1 w2] (* rho (w1 d:dt) (w2 d:dt)))]
    T))

```

If we evaluate the right-hand side expression we obtain<sup>7</sup>

<sup>7</sup>The procedure `trace2down` is defined in Section 9.3.4, page 87. This expression also uses `drop2`, which converts a tensor field that takes two one-form fields into a tensor field that takes two vector fields. Its definition is

```

(defn drop2
  [metric-tensor basis]
  (fn [tensor]
    (fn [v1 v2]
      (contract (fn [e1 w1]
                   (contract (fn [e2 w2]

```



```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (let [g (Newton-metric 'M 'G 'c V)]
  (let [T_ij ((drop2 g spacetime-rect-basis) (Tdust 'rho))]
    (let [T ((trace2down g spacetime-rect-basis) T_ij)]
      ((- (T_ij d:dt d:dt) (* (/ 1 2) T (g d:dt d:dt)))
        ((point spacetime-rect) (up 't 'x 'y 'z))))))
;; => (+ (* (/ 1 2) (expt c 4) rho)
;;      (* #emmy/bigint 2 (expt c 2) rho (V (up x y z)))
;;      (* #emmy/bigint 2 rho (expt (V (up x y z)) 2)))
```

So, to make the Poisson analogy we get

$$R_{\mu\nu} = \frac{8\pi G}{c^4} \left( T_{\mu\nu} - \frac{1}{2} T g_{\mu\nu} \right) - \Lambda g_{\mu\nu} \quad (9.29)$$

as required.

### 9.4.2 Exercise 9.6: Curvature of Schwarzschild Spacetime

In spherical coordinates around a nonrotating gravitating body the metric of Schwarzschild spacetime is given as:<sup>8</sup>

```
(define-coordinates (up t r theta phi) spacetime-sphere)      clojure

(defn Schwarzschild-metric
  [M G c]
  (let [a (- 1 (/ (* 2 G M) (* (square c) r)))]
    (fn [v1 v2]
      (+ (* -1 (square c) a (dt v1) (dt v2))
        (* (/ 1 a) (dr v1) (dr v2))
        (* (square r)
          (+ (* (dtheta v1) (dtheta v2)) (* (square (sin theta)) (dphi v1) (dphi
v2))))))))
```

Show that the Ricci curvature of the Schwarzschild spacetime is zero. Use the definition of the Ricci tensor in equation (8.20).

### 9.4.3 Exercise 9.7: Circular Orbits in Schwarzschild Spacetime

Test particles move along geodesics in spacetime. Now that we have a metric for Schwarzschild spacetime (Section 9.4.2, page 90) we can use it to construct the geodesic equations and determine how test particles move. Consider circular orbits. For example, the circular orbit along a line of constant longitude is a geodesic, so it should satisfy the geodesic equations. Here is the equation of a circular path along the zero longitude line.

```
(defn prime-meridian      clojure
  [r omega]
  (compose (point spacetime-sphere) (fn [t] (up t r (* omega t) 0)) (chart R1-rect)))
```

```
(* (metric-tensor v1 e1) (tensor w1 w2) (metric-tensor
e2 v2)))
basis)))
basis))
```

<sup>8</sup>The spacetime manifold is built from  $\mathbf{R}^4$  with the addition of appropriate coordinate systems:

```
(def spacetime emmy.env/spacetime)      clojure

(def spacetime-rect emmy.env/spacetime-rect)

(def spacetime-sphere emmy.env/spacetime-sphere)
```

This equation will satisfy the geodesic equations for compatible values of the radius `r` and the angular velocity `omega`. If you substitute this into the geodesic equation and set the residual to zero you will obtain a constraint relating `r` and `omega`. Do it.

Surprise: You should find out that  $\omega^2 r^3 = GM$  — Kepler's law!

#### 9.4.4 Exercise 9.8: Stability of Circular Orbits

In Schwarzschild spacetime there are stable circular orbits if the coordinate  $r$  is large enough, but below that value all orbits are unstable. The critical value of  $r$  is larger than the Schwarzschild horizon radius. Let's find that value.

For example, we can consider a perturbation of the orbit of constant longitude. Here is the result of adding an exponential variation of size `epsilon`:

```
(defn prime-meridian+X                                     clojure
  [r epsilon X]
  (compose (point spacetime-sphere)
    (fn [t]
      (up (+ t (* epsilon (* (ref X 0) (exp (* 'lambda t))))))
          (+ r (* epsilon (* (ref X 1) (exp (* 'lambda t))))))
          (+ (* (sqrt (/ (* 'G 'M) (expt r 3))) t)
              (* epsilon (* (ref X 2) (exp (* 'lambda t))))))
      0))
    (chart R1-rect)))
```

Plugging this into the geodesic equation yields a structure of residuals:

```
(def Cartan                                                  clojure
  (Christoffel→Cartan (metric→Christoffel-2 (Schwarzschild-metric 'M 'G 'c)
    (coordinate-system→basis spacetime-sphere))))

(defn geodesic-equation+X-residuals
  [eps X]
  (let [gamma (prime-meridian+X 'r eps X)]
    (((covariant-derivative Cartan gamma) d:dt) ((differential gamma) d:dt))
    (chart spacetime-sphere))
    ((point R1-rect) 't)))
```

The characteristic equation in the eigenvalue `lambda` can be obtained as the numerator of the expression:

```
(determinant                                               clojure
  (submatrix (((* (partial 1) (partial 0)) geodesic-equation+X-residuals) 0 (up 0 0)
    0) 0 2 0 2))
;; => (- (* lambda (exp (* lambda t)) lambda (- (* (+ (* lambda (exp (* lambda t))
;;      lambda) (* (+ (* (/ (* -1 (expt c 2) (- 1 (/ (* 2 G M) (* (expt c 2) r)))
;;      (expt r 2) (expt r 2) (expt (sin (* (...
;; [full result retained in chapter009/029.cljs]
```

Show that the orbits are unstable if  $r < 6GM/c^2$ .

#### 9.4.5 Exercise 9.9: Friedmann-Lemaître-Robertson-Walker

The Einstein tensor  $G_{\mu\nu}$  (see footnote 5) can be expressed as a program:

```
(defn Einstein                                              clojure
  [coordinate-system metric-tensor]
  (let [basis (coordinate-system→basis coordinate-system)
        connection (Christoffel→Cartan (metric→Christoffel-2 metric-tensor basis))
        nabla (covariant-derivative connection)
        Ricci-tensor (Ricci nabla basis)
        Ricci-scalar ((trace2down metric-tensor basis) Ricci-tensor)]
    (letfn [(Einstein-tensor [v1 v2]
```

```

        (- (Ricci-tensor v1 v2) (* (/ 1 2) Ricci-scalar (metric-tensor v1
v2))))])
      Einstein-tensor)))

(defn Einstein-field-equation
  [coordinate-system metric-tensor Lambda stress-energy-tensor]
  (let [Einstein-tensor (Einstein coordinate-system metric-tensor)]
    (let [EFE-residuals (- (+ Einstein-tensor (* Lambda metric-tensor))
                          (* (/ (* 8 'pi '6) (expt 'c 4)) stress-energy-tensor))]
      EFE-residuals)))

```

One exact solution to the Einstein equations was found by Alexander Friedmann in 1922. He showed that a metric for an isotropic and homogeneous spacetime was consistent with a similarly isotropic and homogeneous stress-energy tensor in Einstein's equations. In this case the residuals of the Einstein equations gave ordinary differential equations for the time-dependent scale of the universe. These are called the Robertson-Walker equations. Friedmann's metric is:

```

(defn FLRW-metric
  [c k R]
  (define-coordinates (up t r theta phi) spacetime-sphere)
  (let [a (/ (square (compose R t)) (- 1 (* k (square r))))
        b (square (* (compose R t) r))]
    (letfn [(g [v1 v2]
              (+ (* -1 (square c) (dt v1) (dt v2))
                 (* a (dr v1) (dr v2))
                 (* b
                    (+ (* (dtheta v1) (dtheta v2)) (* (square (sin theta)) (dphi v1)
                    (dphi v2))))))]
      g)))

```

Here `c` is the speed of light, `k` is the intrinsic curvature, and `R` is a length scale that is a function of time.

The associated stress-energy tensor is

```

(defn Tperfect-fluid
  [rho p c metric]
  (define-coordinates (up t r theta phi) spacetime-sphere)
  (let [basis (coordinate-system->basis spacetime-sphere)
        inverse-metric (invert metric basis)]
    (letfn [(T [w1 w2]
              (+ (* (+ (compose rho t) (/ (compose p t) (square c))) (w1 d:dt) (w2
d:dt))
                 (* (compose p t) (inverse-metric w1 w2))))]
      T)))

```

where `rho` is the energy density, and `p` is the pressure in an ideal fluid model.

The Robertson-Walker equations are:

$$\left(\frac{DR(t)}{R(t)}\right)^2 + \frac{kc^2}{(R(t))^2} - \frac{\Lambda c^2}{3} = \frac{8\pi G}{3}\rho(t), 2\frac{D^2R(t)}{R(t)} - \frac{2}{3}\Lambda c^2 = -8\pi G\left(\frac{\rho(t)}{3} + \frac{p(t)}{c^2}\right). \quad (9.30)$$

Use the programs supplied to derive the Robertson-Walker equations.

#### 9.4.6 Exercise 9.10: Cosmology

For energy to be conserved, the stress-energy tensor must be constrained so that its covariant divergence is zero

$$\sum_{\mu} \Delta_{e_{\mu}} T(\tilde{e}^{\mu}, \omega) = 0 \quad (9.31)$$

for every one-form  $\omega$ .

a. Show that for the perfect fluid stress-energy tensor and the FLRW metric this constraint is equivalent to the differential equation

$$D(c^2 \rho R^3) + pD(R^3) = 0. \quad (9.32)$$

b. Assume that in a “matter-dominated universe” radiation pressure is negligible, so  $p = 0$ . Using the Robertson-Walker equations (9.30) and the energy conservation equation (9.32) show that the observation of an expanding universe is compatible with a negative curvature universe, a flat universe, or a positive curvature universe:  $k \in \{-1, 0, +1\}$ .

## 10 Hodge Star and Electrodynamics

The vector space of  $p$ -form fields on an  $n$ -dimensional manifold has dimension  $n!/((n-p)!p!)$ . This is the same dimension as the space of  $(n-p)$ -form fields. So these vector spaces are isomorphic. If we have a metric there is a natural isomorphism: for each  $p$ -form field  $\omega$  on an  $n$ -dimensional manifold there is an  $(n-p)$ -form field  $g^*\omega$ , called its *Hodge dual*.<sup>1</sup> The Hodge dual should not be confused with the duality of vector bases and one-form bases, which is defined without reference to a metric. The Hodge dual is useful for the elegant formalization of electrodynamics.

In Euclidean 3-space, if we think of a one-form as a foliation of the space, then the dual is a two-form, which can be thought of as a pack of square tubes, whose axes are perpendicular to the leaves of the foliation. The original one-form divides these tubes up into volume elements. For example, the dual of the basis oneform  $dx$  is the two-form  $g^*dx = dy \wedge dz$ . We may think of  $dx$  as a set of planes perpendicular to the  $\hat{x}$ -axis. Then  $g^*dx$  is a set of tubes parallel to the  $\hat{x}$ -axis. In higher-dimensional spaces the visualization is more complicated, but the basic idea is the same. The Hodge dual of a two-form in four dimensions is a twoform that is perpendicular to the given two-form. However, if the metric is indefinite (e.g., the Lorentz metric) there is an added complication with the signs.

The Hodge dual is a linear operator, so it can be defined by its action on the basis elements. Let  $\{\partial/\partial x^0, \dots, \partial/\partial x^{n-1}\}$  be an orthonormal basis of vector fields<sup>2</sup> and let  $\{dx^0, \dots, dx^{n-1}\}$  be the ordinary dual basis for the one-forms. Then the  $(n-p)$ -form  $g^*\omega$  that is the Hodge dual of the  $p$ -form  $\omega$  can be defined by its coefficients with respect to the basis, using indices, as

$$(g^*\omega)_{j_p \dots j_{n-1}} = \sum_{i_0 \dots i_{p-1} j_0 \dots j_{p-1}} \frac{1}{p!} \omega_{i_0 \dots i_{p-1}} g^{i_0 j_0} \dots g^{i_{p-1} j_{p-1}} \epsilon_{j_0 \dots j_{n-1}} \quad (10.1)$$

where  $g^{ij}$  are the coefficients of the inverse metric and  $\epsilon_{j_0 \dots j_{n-1}}$  is either  $-1$  or  $+1$  if the permutation  $\{0 \dots n-1\} \mapsto \{j_0 \dots j_{n-1}\}$  is odd or even, respectively.

### 10.1 Relationship to Vector Calculus

In 3-dimensional Euclidean space the traditional vector derivative operations are gradient, curl, and divergence. If  $\hat{x}, \hat{y}, \hat{z}$  are the usual orthonormal rectangular vector basis,  $f$  a function on the space, and  $\vec{v}$  a vector field on the space, then

$$\begin{aligned} \text{grad}(f) &= \frac{\partial f}{\partial x} \hat{x} + \frac{\partial f}{\partial y} \hat{y} + \frac{\partial f}{\partial z} \hat{z} \\ \text{curl}(\vec{v}) &= \left( \frac{\partial v_z}{\partial y} - \frac{\partial v_y}{\partial z} \right) \hat{x} + \left( \frac{\partial v_x}{\partial z} - \frac{\partial v_z}{\partial x} \right) \hat{y} + \left( \frac{\partial v_y}{\partial x} - \frac{\partial v_x}{\partial y} \right) \hat{z} \\ \text{div}(\vec{v}) &= \frac{\partial v_x}{\partial x} + \frac{\partial v_y}{\partial y} + \frac{\partial v_z}{\partial z}. \end{aligned}$$

Recall the meaning of the traditional vector operations. Traditionally we assume that there is a metric that allows us to determine distances between locations and angles between vectors. Such a metric establishes local scale factors relating coordinate increments to actual distances. The vector gradient,  $\text{grad}(f)$ , points in the direction of steepest increase in the function with respect to actual distances. By contrast, the gradient one-form,  $df$ , does not depend on a metric, so there is no concept of distance built in to it. Nevertheless, the concepts are related. The gradient one-form is given by

$$df = \left( \frac{\partial}{\partial x} f \right) dx + \left( \frac{\partial}{\partial y} f \right) dy + \left( \frac{\partial}{\partial z} f \right) dz. \quad (10.2)$$

The traditional gradient vector field is then just the raised gradient one-form (see equation (9.8)). So

$$\text{grad}(f) = g^\sharp(df) \quad (10.3)$$

is computed by

<sup>1</sup>The traditional notion is to just use an asterisk; we use  $g^*$  to emphasize that this duality depends on the choice of metric  $g$ .

<sup>2</sup>We have a metric, so we can define “orthonormal” and “use it to construct an orthonormal basis given any basis. The Gram-Schmidt procedure does the job.

```
(defn gradient [metric basis] (compose (raise metric basis) d))
```

closure

Let  $\theta$  be a one-form field:

$$\theta = \theta_x dx + \theta_y dy + \theta_z dz. \quad (10.4)$$

We compute

$$d\theta = \left( \frac{\partial \theta_z}{\partial y} - \frac{\partial \theta_y}{\partial z} \right) dy \wedge dz + \left( \frac{\partial \theta_x}{\partial z} - \frac{\partial \theta_z}{\partial x} \right) dz \wedge dx + \left( \frac{\partial \theta_y}{\partial x} - \frac{\partial \theta_x}{\partial y} \right) dx \wedge dy. \quad (10.5)$$

So the exterior-derivative expression corresponding to the vector-calculus curl is:

$$g^*(d\theta) = \left( \frac{\partial \theta_z}{\partial y} - \frac{\partial \theta_y}{\partial z} \right) dx + \left( \frac{\partial \theta_x}{\partial z} - \frac{\partial \theta_z}{\partial x} \right) dy + \left( \frac{\partial \theta_y}{\partial x} - \frac{\partial \theta_x}{\partial y} \right) dz. \quad (10.6)$$

Thus, the curl of a vector field  $\mathbf{v}$  is

$$\text{curl}(\mathbf{v}) = g^\sharp(g^*(d(g^\flat(\mathbf{v})))), \quad (10.7)$$

which can be computed with

```
(defn curl
  [metric orthonormal-basis]
  (let [star (Hodge-star metric orthonormal-basis)
        sharp (raise metric orthonormal-basis)
        flat (lower metric)]
    (compose sharp star d flat)))
```

closure

Also, we compute

$$d(g^*\theta) = \left( \frac{\partial \theta_x}{\partial x} + \frac{\partial \theta_y}{\partial y} + \frac{\partial \theta_z}{\partial z} \right) dx \wedge dy \wedge dz. \quad (10.8)$$

So the exterior-derivative expression corresponding to the vector-calculus div is

$$g^*d(g^*\theta) = \frac{\partial \theta_x}{\partial x} + \frac{\partial \theta_y}{\partial y} + \frac{\partial \theta_z}{\partial z}. \quad (10.9)$$

Thus, the divergence of a vector field  $\mathbf{v}$  is

$$\text{div}(\mathbf{v}) = g^*(d(g^*(g^\flat(\mathbf{v}))))). \quad (10.10)$$

It is easily computed:

```
(defn divergence
  [metric orthonormal-basis]
  (let [star (Hodge-star metric orthonormal-basis)
        flat (lower metric)]
    (compose star d star flat)))
```

closure

The divergence is defined even if we don't have a metric, but have only a connection. In that case the divergence can be computed with

```
(defn divergence
  ([metric orthonormal-basis]
   (let [star (Hodge-star metric orthonormal-basis)
         flat (lower metric)]
     (compose star d star flat)))
  ([Cartan]
   (fn [v]
```

closure

```
(fn [point]
  (let [basis (Cartan→basis Cartan)
        nabla (covariant-derivative Cartan)]
    (contract (fn [ei wi] ((wi ((nabla ei) v)) point)) basis))))
```

If the Cartan form is derived from a metric these programs yield the same answer.

The Laplacian is, as expected, the composition of the divergence and the gradient:

```
(defn Laplacian
  [metric orthonormal-basis]
  (compose (divergence metric orthonormal-basis) (gradient metric orthonormal-basis)))
```

closure

## 10.2 Spherical Coordinates

We can illustrate these by computing the formulas for the vector-calculus operators in spherical coordinates. We start with a 3-dimensional manifold, and we set up the conditions for spherical coordinates.

```
(def spherical R3-rect)

(define-coordinates (up r theta phi) spherical)

(def R3-spherical-point ((point spherical) (up 'r0 'theta0 'phi0)))
```

closure

The geometry is specified by the metric:

```
(defn spherical-metric
  [v1 v2]
  (+ (* (dr v1) (dr v2))
    (* (square r) (+ (* (dtheta v1) (dtheta v2)) (* (expt (sin theta) 2) (dphi v1)
    (dphi v2))))))
```

closure

We also need an orthonormal basis for the spherical coordinates. The coordinate basis is orthogonal but not normalized.

```
(def e_0 d:dr)

(def e_1 (* (/ 1 r) d:dtheta))

(def e_2 (* (/ 1 (* r (sin theta))) d:dphi))

(def orthonormal-spherical-vector-basis (down e_0 e_1 e_2))

(def orthonormal-spherical-oneform-basis
  (vector-basis→dual orthonormal-spherical-vector-basis spherical))

(def orthonormal-spherical-basis
  (make-basis orthonormal-spherical-vector-basis orthonormal-spherical-oneform-basis))
```

closure

The components of the gradient of a scalar field are obtained using the dual basis:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((orthonormal-spherical-oneform-basis ((gradient spherical-metric
                                                    orthonormal-spherical-
basis)
                                                    (literal-manifold-function 'f
spherical))))
R3-spherical-point))
;; ⇒ (up (((partial 0) f) (up r0 theta0 phi0))
;;      (/ (((partial 1) f) (up r0 theta0 phi0)) r0)
;;      (/ (((partial 2) f) (up r0 theta0 phi0)) (* r0 (sin theta0))))
```

closure

To get the formulas for curl and divergence we need a vector field with components with respect to the normalized basis.

```
(def v                                          closure
  (+ (* (literal-manifold-function 'v↑0 spherical) e_0)
     (* (literal-manifold-function 'v↑1 spherical) e_1)
     (* (literal-manifold-function 'v↑2 spherical) e_2)))
```

The curl is a bit complicated:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify ((orthonormal-spherical-oneform-basis ((curl spherical-metric orthonormal-
spherical-basis)
                                                v))
          R3-spherical-point))
;; ⇒ (up (/ (+ (* (sin theta0) (((partial 1) v↑2) (up r0 theta0 phi0)))
               (* (v↑2 (up r0 theta0 phi0)) (cos theta0))
               (* -1 (((partial 2) v↑1) (up r0 theta0 phi0))))
          (* r0 (sin theta0))))
;;
;; (/ (+ (* -1 r0 (sin theta0) (((partial 0) v↑2) (up r0 theta0 phi0)))
        (* -1 (sin theta0) (v↑2 (up r0 theta0 phi0)))
        (((partial 2) v↑0) (up r0 theta0 phi0)))
      (* r0 (sin theta0)))
;;
;; (/ (+ (* r0 (((partial 0) v↑1) (up r0 theta0 phi0)))
        (v↑1 (up r0 theta0 phi0))
        (* -1 (((partial 1) v↑0) (up r0 theta0 phi0))))
      r0))
```

But the divergence and Laplacian are simpler

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify ((divergence spherical-metric orthonormal-spherical-basis) v) R3-spherical-
point))
;; ⇒ (/ (+ (* r0 (sin theta0) (((partial 0) v↑0) (up r0 theta0 phi0)))
           (* 2 (sin theta0) (v↑0 (up r0 theta0 phi0)))
           (* (sin theta0) (((partial 1) v↑1) (up r0 theta0 phi0)))
           (* (cos theta0) (v↑1 (up r0 theta0 phi0)))
           (((partial 2) v↑2) (up r0 theta0 phi0)))
        (* r0 (sin theta0)))

(simplify ((Laplacian spherical-metric orthonormal-spherical-basis)
          (literal-manifold-function 'f spherical))
          R3-spherical-point))
;; ⇒ (/ (+ (* (expt r0 2)
              (expt (sin theta0) 2)
              (((expt (partial 0) 2) f) (up r0 theta0 phi0)))
          (* 2 r0 (((partial 0) f) (up r0 theta0 phi0)) (expt (sin theta0) 2))
          (* (((partial 1) f) (up r0 theta0 phi0)) (sin theta0) (cos theta0))
          (* (expt (sin theta0) 2) (((expt (partial 1) 2) f) (up r0 theta0 phi0)))
          (((expt (partial 2) 2) f) (up r0 theta0 phi0)))
          (* (expt r0 2) (expt (sin theta0) 2))))
```

### 10.3 The Wave Equation

The kinematics of special relativity can be formulated on a flat 4-dimensional spacetime manifold.

```
(def SR R4-rect) closure

(define-coordinates (up ct x y z) SR)

(def an-event ((point SR) (up 'ct0 'x0 'y0 'z0)))

(def a-vector
```



```
(+ (* (literal-manifold-function 'v↑t SR) d:dct)
  (* (literal-manifold-function 'v↑x SR) d:dx)
  (* (literal-manifold-function 'v↑y SR) d:dy)
  (* (literal-manifold-function 'v↑z SR) d:dz)))
```

The Minkowski metric is<sup>3</sup>

$$g(u, v) = -c^2 dt(u) dt(v) + dx(u) dx(v) + dy(u) dy(v) + dz(u) dz(v). \quad (10.11)$$

As a program:

```
(defn g-Minkowski                                     clojure
  [u v]
  (+ (* -1 (dct u) (dct v)) (* (dx u) (dx v)) (* (dy u) (dy v)) (* (dz u) (dz v))))
```

The length of a vector is described in terms of the metric:

$$\sigma = g(v, v). \quad (10.12)$$

If  $\sigma$  is positive the vector is *spacelike* and its square root is the *proper length* of the vector. If  $\sigma$  is negative the vector is *timelike* and the square root of its negation is the *proper time* of the vector. If  $\sigma$  is zero the vector is *lightlike* or *null*.

```
((g-Minkowski a-vector a-vector) an-event)                                     clojure
;; => (+ (* -1 (v↑t (up ct0 x0 y0 z0)) (v↑t (up ct0 x0 y0 z0)))
;;      (* (v↑x (up ct0 x0 y0 z0)) (v↑x (up ct0 x0 y0 z0)))
;;      (* (v↑y (up ct0 x0 y0 z0)) (v↑y (up ct0 x0 y0 z0)))
;;      (* (v↑z (up ct0 x0 y0 z0)) (v↑z (up ct0 x0 y0 z0))))
```

As an example of vector calculus in four dimensions, we can compute the wave equation for a scalar field in 4-dimensional spacetime.

We need an orthonormal basis for the spacetime:

```
(def SR-vector-basis (coordinate-system→vector-basis SR))                                     clojure
(def SR-basis (coordinate-system→basis SR))
```

We check that it is orthonormal with respect to the metric:

```
((g-Minkowski SR-vector-basis SR-vector-basis) an-event)                                     clojure
;; => (down (down -1 0 0 0) (down 0 1 0 0) (down 0 0 1 0) (down 0 0 0 1))
```

So, the Laplacian of a scalar field is the wave equation!

```
(def p (literal-manifold-function 'phi SR))                                     clojure
((Laplacian g-Minkowski SR-basis) p) an-event
;; => (* -1
;;      (+ (* -1 (((partial 0) ((partial 0) phi)) (up ct0 x0 y0 z0)))
;;          (* -1 -1 (((partial 1) ((partial 1) phi)) (up ct0 x0 y0 z0)))
;;          (((partial 2) ((partial 2) phi)) (up ct0 x0 y0 z0))
;;          (* -1 -1 (((partial 3) ((partial 3) phi)) (up ct0 x0 y0 z0)))))
```

## 10.4 Electrodynamics

Using Hodge duals we can represent electrodynamics in an elegant way. Maxwell's electrodynamics is invariant under Lorentz transformations. We use 4-dimensional rectangular coordinates for the flat spacetime of special relativity.

<sup>3</sup>The metric in relativity is not positive definite, so nonzero vectors can have zero length.

In this formulation of electrodynamics the electric and magnetic fields are represented together as a two-form field, the *Faraday tensor*. Under Lorentz transformations the individual components are mixed. The Faraday tensor is:<sup>4</sup>

```
(defn Faraday
  [Ex Ey Ez Bx By Bz]
  (+ (* Ex (wedge dx dct))
      (* Ey (wedge dy dct))
      (* Ez (wedge dz dct))
      (* Bx (wedge dy dz))
      (* By (wedge dz dx))
      (* Bz (wedge dx dy))))
```

closure

The Hodge dual of the Faraday tensor exchanges the electric and magnetic fields, negating the components that will involve time. The result is called the *Maxwell tensor*:

```
(defn Maxwell
  [Ex Ey Ez Bx By Bz]
  (+ (* -1 Bx (wedge dx dct))
      (* -1 By (wedge dy dct))
      (* -1 Bz (wedge dz dct))
      (* Ex (wedge dy dz))
      (* Ey (wedge dz dx))
      (* Ez (wedge dx dy))))
```

closure

We make a Hodge dual operator for this situation:

```
(def SR-star (Hodge-star g-Minkowski SR-basis))
```

closure

And indeed, it transforms the Faraday tensor into the Maxwell tensor:

```
(simplify ((- (SR-star (Faraday 'Ex 'Ey 'Ez 'Bx 'By 'Bz)) (Maxwell 'Ex 'Ey 'Ez 'Bx
'By 'Bz)))
  (literal-vector-field 'u SR)
  (literal-vector-field 'v SR))
an-event))
;; => 0
```

closure

One way to get electric fields is to have charges; magnetic fields can arise from motion of charges. In this formulation we combine the charge density and the current to make a one-form field:

```
(defn J
  [charge-density Ix Iy Iz]
  (- (* (/ 1 'c) (+ (* Ix dx) (* Iy dy) (* Iz dz))) (* charge-density dct)))
```

closure

The coefficient `(/ 1 :c)` makes the components of the one-form uniform with respect to units.

To develop Maxwell's equations we need a general Faraday field and a general current-density field:

```
(def F
  (Faraday (literal-manifold-function 'Ex SR)
            (literal-manifold-function 'Ey SR)
            (literal-manifold-function 'Ez SR)
            (literal-manifold-function 'Bx SR)
            (literal-manifold-function 'By SR)
            (literal-manifold-function 'Bz SR)))

(def four-current
```

closure

<sup>4</sup>This representation is from Misner, Thorne, and Wheeler, *Gravitation*, p.108.

```
(J (literal-manifold-function 'rho SR)
  (literal-manifold-function 'Ix SR)
  (literal-manifold-function 'Iy SR)
  (literal-manifold-function 'Iz SR)))
```

## 10.5 Maxwell's Equations

Maxwell's equations in the language of differential forms are

$$dF = 0, \quad (10.13)$$

$$d(g^*F) = 4\pi g^*J. \quad (10.14)$$

The first equation gives us what would be written in vector notation as

$$\operatorname{div} \vec{B} = 0, \quad (10.15)$$

$$\operatorname{curl} \vec{E} = -\frac{1}{c} \frac{d\vec{B}}{dt}. \quad (10.16)$$

The second equation gives us what would be written in vector notation as

$$\operatorname{div} \vec{E} = 4\pi\rho, \quad (10.17)$$

$$\operatorname{curl} \vec{B} = \frac{1}{c} \frac{d\vec{E}}{dt} + \frac{4\pi}{c} \vec{I}. \quad (10.18)$$

To see how these work out, we evaluate each component of  $dF$  and  $d(g^*F) - 4\pi g^*J$ . Since these are both two-form fields, their exterior derivatives are three-form fields, so we have to provide three basis vectors to get each component. Each component equation will yield one of Maxwell's equations, written in coordinates, without vector notation. So, the purely spatial component  $dF(\partial/\partial x, \partial/\partial y, \partial/\partial z)$  of equation (10.13) is equation (10.15):

```
((d F) d:dx d:dy d:dz) an-event)
;; => (+ (((partial 1) Bx) (up ct0 x0 y0 z0))
;;      (* -1 -1 (((partial 2) By) (up ct0 x0 y0 z0)))
;;      (((partial 3) Bz) (up ct0 x0 y0 z0)))
```

closure

$$\frac{\partial B_x}{\partial x} + \frac{\partial B_y}{\partial y} + \frac{\partial B_z}{\partial z} = 0 \quad (10.19)$$

The three mixed space and time components of equation (10.13) are equation (10.16):

```
((d F) d:dct d:dy d:dz) an-event)
;; => (+ (((partial 0) Bx) (up ct0 x0 y0 z0))
;;      (* -1 -1 (((partial 2) Ez) (up ct0 x0 y0 z0)))
;;      (* -1 (((partial 3) Ey) (up ct0 x0 y0 z0))))
```

closure

$$\frac{\partial E_z}{\partial y} - \frac{\partial E_y}{\partial z} = \frac{1}{c} \frac{\partial B_x}{\partial t}, \quad (10.20)$$

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (((d F) d:dct d:dz d:dx) an-event))
;; => (+ (((partial 0) By) (up ct0 x0 y0 z0))
;;      (((partial 3) Ex) (up ct0 x0 y0 z0))
;;      (* -1 (((partial 1) Ez) (up ct0 x0 y0 z0))))
```

closure

$$\frac{\partial E_x}{\partial z} - \frac{\partial E_z}{\partial x} = \frac{1}{c} \frac{\partial B_y}{\partial t}, \quad (10.21)$$

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      closure
(simplify (((d F) d:dct d:dx d:dy) an-event))
;; => (+ (((partial 0) Bz) (up ct0 x0 y0 z0))
;;      (((partial 1) Ey) (up ct0 x0 y0 z0))
;;      (* -1 (((partial 2) Ex) (up ct0 x0 y0 z0))))
```

$$\frac{\partial E_y}{\partial x} - \frac{\partial E_x}{\partial y} = \frac{1}{c} \frac{\partial B_z}{\partial t}. \quad (10.22)$$

The purely spatial component of equation (10.14) is equation (10.17):

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      closure
(simplify (((- (d (SR-star F)) (* 4 'pi (SR-star four-current))) d:dx d:dy d:dz) an-
event))
;; => (+ (* -4 pi (rho (up ct0 x0 y0 z0)))
;;      (((partial 1) Ex) (up ct0 x0 y0 z0))
;;      (((partial 2) Ey) (up ct0 x0 y0 z0))
;;      (((partial 3) Ez) (up ct0 x0 y0 z0)))
```

$$\frac{\partial E_x}{\partial x} + \frac{\partial E_y}{\partial y} + \frac{\partial E_z}{\partial z} = 4\pi\rho. \quad (10.23)$$

And finally, the three mixed time and space components of equation (10.14) are equation (10.18):

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      closure
(simplify (((- (d (SR-star F)) (* 4 'pi (SR-star four-current))) d:dct d:dy d:dz) an-
event))
;; => (/ (+ (* c (((partial 0) Ex) (up ct0 x0 y0 z0)))
;;          (* -1 c (((partial 2) Bz) (up ct0 x0 y0 z0)))
;;          (* c (((partial 3) By) (up ct0 x0 y0 z0)))
;;          (* 4 pi (Ix (up ct0 x0 y0 z0))))
;;      c)
```

$$\frac{\partial B_y}{\partial z} - \frac{\partial B_z}{\partial y} = -\frac{1}{c} \frac{\partial E_x}{\partial t} - \frac{4\pi}{c} I_x, \quad (10.24)$$

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      closure
(simplify (((- (d (SR-star F)) (* 4 'pi (SR-star four-current))) d:dct d:dz d:dx) an-
event))
;; => (/ (+ (* c (((partial 0) Ey) (up ct0 x0 y0 z0)))
;;          (* -1 c (((partial 3) Bx) (up ct0 x0 y0 z0)))
;;          (* c (((partial 1) Bz) (up ct0 x0 y0 z0)))
;;          (* 4 pi (Iy (up ct0 x0 y0 z0))))
;;      c)
```

$$\frac{\partial B_z}{\partial x} - \frac{\partial B_x}{\partial z} = -\frac{1}{c} \frac{\partial E_y}{\partial t} - \frac{4\pi}{c} I_y, \quad (10.25)$$

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      closure
(simplify (((- (d (SR-star F)) (* 4 'pi (SR-star four-current))) d:dct d:dx d:dy) an-
event))
;; => (/ (+ (* c (((partial 0) Ez) (up ct0 x0 y0 z0)))
;;          (* -1 c (((partial 1) By) (up ct0 x0 y0 z0)))
;;          (* c (((partial 2) Bx) (up ct0 x0 y0 z0)))
;;          (* 4 pi (Iz (up ct0 x0 y0 z0))))
;;      c)
```

$$\frac{\partial B_x}{\partial y} - \frac{\partial B_y}{\partial x} = -\frac{1}{c} \frac{\partial E_z}{\partial t} - \frac{4\pi}{c} I_z. \quad (10.26)$$

## 10.6 Lorentz Force

The classical force on a charged particle moving in a electromagnetic field is

$$\vec{f} = q \left( \vec{E} + \frac{1}{c} \vec{v} \times \vec{B} \right). \quad (10.27)$$

We can compute this in coordinates. We construct arbitrary  $\vec{E}$  and  $\vec{B}$  vector fields and an arbitrary velocity:

```
(def E closure
  (up (literal-manifold-function 'Ex SR)
       (literal-manifold-function 'Ey SR)
       (literal-manifold-function 'Ez SR)))

(def B
  (up (literal-manifold-function 'Bx SR)
       (literal-manifold-function 'By SR)
       (literal-manifold-function 'Bz SR)))

(def V (up 'V_x 'V_y 'V_z))
```

The 3-space force that results is a mess:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify (* 'q (+ (E an-event) (cross-product V (B an-event)))))
;; => (up (+ (* V_y q (Bz (up ct0 x0 y0 z0)))
              (* -1 V_z q (By (up ct0 x0 y0 z0)))
              (* q (Ex (up ct0 x0 y0 z0))))
          (+ (* -1 V_x q (Bz (up ct0 x0 y0 z0)))
              (* V_z q (Bx (up ct0 x0 y0 z0)))
              (* q (Ey (up ct0 x0 y0 z0)))
              (+ (* V_x q (By (up ct0 x0 y0 z0)))
                  (* -1 V_y q (Bx (up ct0 x0 y0 z0)))
                  (* q (Ez (up ct0 x0 y0 z0))))))
```

The relativistic Lorentz 4-force is usually written in coordinates as

$$f^\nu = - \sum_{\alpha, \mu} q U^\mu F_{\mu\alpha} \eta^{\alpha\nu}, \quad (10.28)$$

where  $U$  is the 4-velocity of the charged particle,  $F$  is the Faraday tensor, and  $\eta^{\alpha\nu}$  are the components of the inverse of the Minkowski metric. Here is a program that computes a component of the force in terms of the Faraday tensor. The desired component is specified by a one-form.

```
(def eta-inverse (invert g-Minkowski SR-basis)) closure

(defn Force
  [charge F four-velocity component]
  (* -1
     charge
     (contract (fn [a b]
                  (contract (fn [e w] (* (w four-velocity) (F e a) (eta-inverse b
                                                                    component)))
                            SR-basis))
                SR-basis)))
```

So, for example, the force in the  $\hat{x}$  direction for a stationary particle is

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify ((Force 'q F d:dict dx) an-event))
;; => (* q (Ex (up ct0 x0 y0 z0)))
```

Notice that the 4-velocity  $\partial/\partial ct$  is the 4-velocity of a stationary particle!

If we give a particle a more general timelike 4-velocity in the  $\hat{x}$  direction we can see how the  $\hat{y}$  component of the force involves both the electric and magnetic field:

```
(defn Ux                                     clojure
  [beta]
  (+ (* (/ 1 (sqrt (- 1 (square beta)))) d:dct) (* (/ beta (sqrt (- 1 (square beta))))
    d:dx)))

((Force 'q F (Ux 'v:c) dy) an-event)
;; => (* -1
;;      q
;;      (+ (* (/ 1 (sqrt (- 1 (expt v:c 2)))) (Ey (up ct0 x0 y0 z0)) -1)
;;          (* (/ v:c (sqrt (- 1 (expt v:c 2)))) (Bz (up ct0 x0 y0 z0)))))
```

### 10.6.1 Exercise 10.1: Relativistic Lorentz Force

Compute all components of the 4-force for a general timelike 4-velocity.

a. Compare these components to the components of the nonrelativistic force given above. Interpret the differences.

b. What is the meaning of the time component? For example, consider:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.   clojure
(simplify ((Force 'q F (Ux 'v:c) dct) an-event))
;; => (/ (* q v:c (Ex (up ct0 x0 y0 z0))) (sqrt (+ (* -1 (expt v:c 2)) 1)))
```

c. Subtract the structure of components of the relativistic 3-space force from the structure of the spatial components of the 4-space force to show that they are equal.

## 11 Special Relativity

Although the usual treatments of special relativity begin with the Michelson-Morley experiment, this is not how Einstein began. In fact, Einstein was impressed with Maxwell's work and he was emulating Maxwell's breakthrough.

Maxwell was preceded by Faraday, Ampere, Oersted, Coulomb, Gauss, and Franklin. These giants discovered electromagnetism and worked out empirical equations that described the phenomena. They understood the existence of conserved charges and fields. Faraday invented the idea of lines of force by which fields can be visualized.

Maxwell's great insight was noticing and resolving the contradiction between the empirically-derived laws of electromagnetism and conservation of charge. He did this by introducing the then experimentally undetectable displacement-current term into one of the empirical equations. The modified equations implied a wave equation and the propagation speed of the wave predicted by the new equation turned out to be the speed of light, as measured by the eclipses of the Galilean satellites of Jupiter. The experimental confirmation by Hertz of the existence of electromagnetic radiation that obeyed Maxwell's equations capped the discovery.

By analogy, Einstein noticed that Maxwell's equations were inconsistent with Galilean relativity. In free space, where electromagnetic waves propagate, Maxwell's equations say that the vector source of electric fields is the time rate of change of the magnetic field and the vector source of magnetic field is the time rate of change of the electric field. The combination of these ideas yields the wave equation. The wave equation itself is not invariant under the Galilean transformation: As Einstein noted, if you run with the propagation speed of the wave there is no time variation in the field you observe, so there is no space variation either, contradicting the wave equation. But the Maxwell theory is beautiful, and it can be verified to a high degree of accuracy, so there must be something wrong with Galilean relativity. Einstein resolved the contradiction by generalizing the meaning of the Lorentz transformation, which was invented to explain the failure of the Michelson-Morley experiment. Lorentz and his colleagues decided that the problem with the Michelson-Morley experiment was that matter interacting with the luminiferous ether contracts in the direction of motion. To make this consistent he had to invent a "local time" which had no clear interpretation. Einstein took the Lorentz transformation to be a fundamental replacement for the Galilean transformation in all of mechanics.

Now to the details. Before Maxwell the empirical laws of electromagnetism were as follows. Electric fields arise from charges, with the inverse square law of Coulomb. This is Carl Friedrich Gauss's law for electrostatics:

$$\operatorname{div} \vec{E} = 4\pi\rho. \quad (11.1)$$

Magnetic fields do not have a scalar source. This is Gauss's law for magnetostatics:

$$\operatorname{div} \vec{B} = 0. \quad (11.2)$$

Magnetic fields are produced by electric currents, as discovered by Hans Christian Oersted and quantified by André-Marie Ampère:

$$\operatorname{curl} \vec{B} = \frac{4\pi}{c} \vec{I}. \quad (11.3)$$

Michael Faraday (and Joseph Henry) discovered that electric fields are produced by moving magnetic fields:

$$\operatorname{curl} \vec{E} = -\frac{1}{c} \frac{\partial \vec{B}}{\partial t}. \quad (11.4)$$

Benjamin Franklin was the first to understand that electrical charges are conserved:

$$\operatorname{div} \vec{I} + \frac{\partial \rho}{\partial t} = 0. \quad (11.5)$$

Although these equations are written in terms of the speed of light  $c$ , these laws were originally written in terms of electrical permittivity and magnetic permeability of free space, which could be determined by measurement of the forces for given currents and charges.

It is easy to see that these equations are mutually contradictory. Indeed, if we take the divergence of equation (11.3) we get

$$\operatorname{div} \operatorname{curl} \vec{B} = 0 = \frac{4\pi}{c} \operatorname{div} \vec{I}, \quad (11.6)$$

which directly contradicts conservation of charge (11.5).

Maxwell patched this bug by adding in the displacement current, changing equation (11.3) to read

$$\operatorname{curl} \vec{B} = \frac{1}{c} \frac{\partial \vec{E}}{\partial t} + \frac{4\pi}{c} \vec{I}. \quad (11.7)$$

Maxwell proceeded by taking the curl of equation (11.4) to get

$$\operatorname{curl} \operatorname{curl} \vec{E} = -\frac{1}{c} \frac{\partial}{\partial t} \operatorname{curl} \vec{B}. \quad (11.8)$$

Expanding the left-hand side

$$\operatorname{grad} \operatorname{div} \vec{E} - \operatorname{Lap} \vec{E} = -\frac{1}{c} \frac{\partial \operatorname{curl} \vec{B}}{\partial t}, \quad (11.9)$$

substituting from equations (11.7) and (11.1), and rearranging the terms we get the inhomogeneous wave equation:

$$\operatorname{Lap} \vec{E} - \frac{1}{c^2} \frac{\partial^2 \vec{E}}{\partial t^2} = 4\pi \operatorname{grad} \rho + \frac{1}{c^2} \vec{I}. \quad (11.10)$$

We see that in free space (in the absence of any charges or currents) we have the familiar homogeneous linear wave equation. A similar equation can be derived for the magnetic field.

Lorentz, whom Einstein also greatly respected, developed a general formula to describe the force on a particle with charge  $q$  moving with velocity  $\vec{v}$  in an electromagnetic field:

$$\vec{F} = q\vec{E} + \frac{q}{c} \vec{v} \times \vec{B}. \quad (11.11)$$

A crucial point in Einstein's inspiration for relativity is, quoting Einstein (in English translation), “During that year [1895–1896] in Aarau the question came to me: If one runs after a light wave with light velocity, then one would encounter a time-independent wavefield. However, something like that does not seem to exist!”<sup>1</sup> This was the observation of the inconsistency.

Let's be more precise about this. Consider a plane sinusoidal wave moving in the  $\hat{x}$  direction with velocity  $c$  in free space ( $\rho = 0$  and  $\vec{I} = 0$ ). This is a perfectly good solution of the wave equation. Now suppose that an observer is moving with the wave in the  $\hat{x}$  direction with velocity  $c$ . Such an observer will see no time variation of the field. So the wave equation reduces to Laplace's equation. But a sinusoidal variation in space is not a solution of Laplace's equation.

Einstein believed that the Maxwell-Lorentz electromagnetic theory was fundamentally correct, though he was unhappy with an apparent asymmetry in the formulation. Consider a system consisting of a conductor and a magnet. If the conductor is moved and the magnet is held stationary (a stationary magnetic field) then the charge carriers in the conductor are subject to the Lorentz force (11.11), causing them to move. However, if the magnet is moved past a stationary conductor then the changing magnetic field induces an electric field in the conductor by equation (11.4), which causes the charge carriers in the conductor to move. The actual current which results is identical for both explanations if the relative velocity of the magnet and the conductor are the same. To Einstein, there should not have been two explanations for the same phenomenon.

<sup>1</sup>The quote is from Pais [12], p. 131.

<sup>2</sup>Points in spacetime are often called *events*.



### 11.1 Invariance of the Wave Equation

Let  $u = (t, x, y, z)$  be a tuple of time and space coordinates that specify a point in spacetime.<sup>2</sup> If  $\phi(t, x, y, z)$  is a scalar field over time and space, the homogeneous linear wave equation is

$$\frac{\partial^2 \phi(u)}{\partial x^2} + \frac{\partial^2 \phi(u)}{\partial y^2} + \frac{\partial^2 \phi(u)}{\partial z^2} - \frac{1}{c^2} \frac{\partial^2 \phi(u)}{\partial t^2} = 0. \quad (11.12)$$

The characteristics for this equation are the “light cones.” If we define a function of spacetime points and increments, length, such that for an incremental tuple in position and time  $\xi = (\Delta t, \Delta x, \Delta y, \Delta z)$  we have<sup>3</sup>

$$\text{length}_u(\xi) = \sqrt{(\Delta x)^2 + (\Delta y)^2 + (\Delta z)^2 - (c\Delta t)^2}, \quad (11.13)$$

then the light cones are the hypersurfaces, for which

$$\text{length}_u(\Delta t, \Delta x, \Delta y, \Delta z) = 0. \quad (11.14)$$

This “length” is called the *interval*.

What is the class of transformations of time and space coordinates that leave the Maxwell-Lorentz theory invariant? The transformations that preserve the wave equation are exactly those that leave its characteristics invariant. We consider a transformation  $u = A(u')$  of time and space coordinates:

$$t = A^0(t', x', y', z'), \quad (11.15)$$

$$x = A^1(t', x', y', z'), \quad (11.16)$$

$$y = A^2(t', x', y', z'), \quad (11.17)$$

$$z = A^3(t', x', y', z'). \quad (11.18)$$

If we define a new field  $\psi(t, x, y, z)$  such that  $\psi = \phi \circ A$ , or

$$\psi(t', x', y', z') = \phi(A(t', x', y', z')), \quad (11.19)$$

Then  $\psi$  will satisfy the wave equation

$$\frac{\partial^2 \psi(u')}{\partial (x')^2} + \frac{\partial^2 \psi(u')}{\partial (y')^2} + \frac{\partial^2 \psi(u')}{\partial (z')^2} - \frac{1}{c^2} \frac{\partial^2 \psi(u')}{\partial (t')^2} = 0, \quad (11.20)$$

if and only if

$$\text{length}_{u'}(\xi') = \text{length}_{A(u')}(DA\xi') = \text{length}_u(\xi). \quad (11.21)$$

But this is just a statement that the velocity of light is invariant under change of the coordinate system. The class of transformations that satisfy equation (11.21) are the Poincaré transformations.

### 11.2 Lorentz Transformations

Special relativity is usually presented in terms of global Lorentz frames, with rectangular spatial coordinates. In this context the Lorentz transformations (and, more generally, the Poincaré transformations) can be characterized as the set of affine transformations (linear transformations plus shift) of the coordinate tuple (time and spatial rectangular coordinates) that preserve the length of incremental spacetime intervals as measured by

$$f(\xi) = -(\xi^0)^2 + (\xi^1)^2 + (\xi^2)^2 + (\xi^3)^2, \quad (11.22)$$

where  $\xi$  is an incremental 4-tuple that could be added to the coordinate 4-tuple  $(ct, x, y, z)$ .<sup>4</sup> The Poincaré-Lorentz transformations are of the form

$$x = \Lambda x' + a, \quad (11.23)$$

<sup>3</sup>Here the length is independent of the spacetime point specified by  $u$ . In General Relativity we find that the metric, and thus the length function needs to vary with the point in spacetime.

<sup>4</sup>Incrementally,  $\xi = \xi^0 \partial / \partial ct + \xi^1 \partial / \partial x + \xi^2 \partial / \partial y + \xi^3 \partial / \partial z$ . The length of this vector, using the Minkowski metric (see equation (10.11)), is the Lorentz interval, the right-hand side of equation (11.22).

Where  $\Lambda$  is the tuple representation of a linear transformation and  $a$  is a 4-tuple shift. Because the 4-tuple includes the time, these transformations include transformations to a uniformly moving frame. A transformation that does not rotate or shift, but just introduces relative velocity, is sometimes called a *boost*.

In general relativity, global Lorentz frames do not exist, and so global affine transformations are irrelevant. In general relativity Lorentz invariance is a local property of incremental 4-tuples at a point.

Incremental 4-tuples transform as

$$\xi = \Lambda \xi'. \quad (11.24)$$

This places a constraint on the allowed  $\Lambda$

$$f(\xi') = f(\Lambda \xi'), \quad (11.25)$$

for arbitrary  $\xi'$ .

The possible  $\Lambda$  that are consistent with the preservation of the interval can be completely specified and conveniently parameterized.

### 11.3 Simple Lorentz Transformations

Consider the linear transformation, in the first two coordinates,

$$\xi^0 = p(\xi')^0 + q(\xi')^1 \xi^1 = r(\xi')^0 + s(\xi')^1. \quad (11.26)$$

The requirement to preserve the interval gives the constraints

$$\begin{aligned} p^2 - r^2 &= 1, \\ pq - rs &= 0, \\ q^2 - s^2 &= -1. \end{aligned} \quad (11.27)$$

There are four parameters to determine, and only three equations, so the solutions have a free parameter. It turns out that a good choice is  $\beta = q/p$ . Solve to find

$$p = \frac{1}{\sqrt{1 - \beta^2}} = \gamma(\beta), \quad (11.28)$$

and also  $p = s$  and  $q = r = \beta p$ . This defines  $\gamma$ . Written out, the transformation is

$$\xi^0 = \gamma(\beta) \left( (\xi')^0 + \beta (\xi')^1 \right) \xi^1 = \gamma(\beta) \left( \beta (\xi')^0 + (\xi')^1 \right). \quad (11.29)$$

Simple physical arguments<sup>5</sup> show that this mathematical result relates the time and space coordinates for two systems in uniform relative motion. The parameter  $\beta$  is related to the relative velocity.

Consider incremental vectors as spacetime vectors relative to an origin in a global inertial frame. So, for example,  $\xi = (ct, x)$ , ignoring  $y$  and  $z$  for a moment. The unprimed coordinate origin  $x = 0$  corresponds, in primed coordinates, to (using equations (11.29))

$$x = 0 = \gamma(\beta)(x' + \beta ct'), \quad (11.30)$$

so

$$\beta = -\frac{x'}{ct'} = -\frac{v'}{c}, \quad (11.31)$$

with the definition  $v' = x'/t'$ . We see that  $\beta$  is minus  $1/c$  times the velocity ( $v'$ ) of the unprimed system (which moves with its origin) as “seen” in the primed coordinates

$$\beta = \frac{x}{ct} = \frac{v}{c}. \quad (11.32)$$

So  $v' = -v$ .

---

<sup>5</sup>See, for instance, Mermin, “Space and Time in Special Relativity.”

A consistent interpretation is that the origin of the primed system moves with velocity  $v = \beta c$  along the  $\hat{x}$ -axis of the unprimed system. And the unprimed system moves with the same velocity in the other direction, when viewed in terms of the primed system. What happened to the other coordinates:  $y$  and  $z$ ? We did not need them to find this one-parameter family of Lorentz transformations. They are left alone. This mathematical result has a physical interpretation: Lengths are not affected by perpendicular boosts. Think about two observers on a collision course, each carrying a meter stick perpendicular to their relative velocity. At the moment of impact, the meter sticks must coincide. The symmetry of the situation does not permit one observer to conclude that one meter stick is shorter than the other, because the other observer must come to the same conclusion. Both observers can put their conclusions to the test upon impact.

We can fill in the components of this simple boost:

$$\begin{aligned}\xi^0 &= \gamma(\beta) \left( (\xi')^0 + \beta (\xi')^1 \right) \\ \xi^1 &= \gamma(\beta) \left( \beta (\xi')^0 + (\xi')^1 \right) \\ \xi^2 &= (\xi')^2 \\ \xi^3 &= (\xi')^3.\end{aligned}\tag{11.33}$$

### 11.4 More General Lorentz Transformations

One direction was special in our consideration of simple boosts. We can make use of this fact to find boosts in any direction.

Let  $c\beta = (v^0, v^1, v^2)$  be the tuple of components of the relative velocity of the origin of the primed system in the unprimed system. The components are with respect to the same rectangular basis used to define the spatial components of any incremental vector.

An incremental vector can be decomposed into vectors parallel and perpendicular to the velocity. Let  $\xi$  be the tuple of spatial components of  $\xi$ , and  $\xi^0$  be the time component. Then,

$$\xi = \xi^\perp + \xi^\parallel,\tag{11.34}$$

where  $\beta \cdot \xi = 0$ . (This is the ordinary dot product in three dimensions.) Explicitly,

$$\xi^\parallel = \frac{\beta}{\beta} \left( \frac{\beta}{\beta} \cdot \xi \right),\tag{11.35}$$

where  $\beta = \|\beta\|$ , the magnitude of  $\beta$ , and

$$\xi^\perp = \xi - \xi^\parallel.\tag{11.36}$$

In the simple boost of equation (11.33) we can identify  $\xi^1$  with the magnitude  $|\xi^\parallel|$  of the parallel component. The perpendicular component is unchanged:

$$\begin{aligned}\xi^0 &= \gamma(\beta) \left( (\xi')^0 + \beta |\xi^\parallel| \right), \\ |\xi^\parallel| &= \gamma(\beta) \left( \beta (\xi')^0 + |\xi^\parallel| \right), \\ \xi^\perp &= (\xi')^\perp.\end{aligned}\tag{11.37}$$

Putting the components back together, this leads to

$$\begin{aligned}\xi^0 &= \gamma(\beta) \left( (\xi')^0 + \beta \cdot \xi \right) \\ \xi &= \gamma(\beta) \left( \beta (\xi')^0 + \xi' \right. \\ &\quad \left. + \frac{\gamma(\beta) - 1}{\beta^2} \beta (\beta \cdot \xi) \right),\end{aligned}\tag{11.38}$$

which gives the components of the general boost  $B$  along velocity  $c\beta$ :

$$\xi = B(\beta)(\xi').\tag{11.39}$$

## 11.5 Implementation

We represent a 4-tuple as a flat up-tuple of components.

```
(defn make-four-tuple [ct space] (up ct (ref space 0) (ref space 1) (ref space 2)))  clojure

(defn four-tuple→ct [v] (ref v 0))

(defn four-tuple→space [v] (up (ref v 1) (ref v 2) (ref v 3)))
```

The invariant interval is then

```
(defn proper-space-interval  clojure
  [four-tuple]
  (sqrt (- (square (four-tuple→space four-tuple)) (square (four-tuple→ct four-
tuple))))))
```

This is a real number for space-like intervals. A space-like interval is one where spatial distance is larger than can be traversed by light in the time interval.

It is often convenient for the interval to be real for time-like intervals, where light can traverse the spatial distance in less than the time interval.

```
(defn proper-time-interval  clojure
  [four-tuple]
  (sqrt (- (square (four-tuple→ct four-tuple)) (square (four-tuple→space four-
tuple))))))
```

The general boost  $B$  is

```
(defn general-boost  clojure
  [beta]
  (fn [xi-p]
    (let [gamma (expt (- 1 (square beta)) (/ -1 2))]
      (let [factor (/ (- gamma 1) (square beta))]
        (let [xi-p-time (four-tuple→ct xi-p)
              xi-p-space (four-tuple→space xi-p)]
          (let [beta-dot-xi-p (dot-product beta xi-p-space)]
            (make-four-tuple
              (* gamma (+ xi-p-time beta-dot-xi-p))
              (+ (* gamma beta xi-p-time) xi-p-space (* factor beta beta-dot-xi-
p))))))))))
```

We can check that the interval is invariant:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.  clojure
(simplify (- (proper-space-interval ((general-boost (up 'vx 'vy 'vz))
  (make-four-tuple 'ct (up 'x 'y 'z))))
  (proper-space-interval (make-four-tuple 'ct (up 'x 'y 'z)))))

;; ⇒ 0
```

It is inconvenient that the general boost as just defined does not work if  $\beta$  is zero. An alternate way to specify a boost is through the magnitude of  $v/c$  and a direction:

```
(defn general-boost2  clojure
  [direction v:c]
  (fn [four-tuple-prime]
    (let [delta-ct-prime (four-tuple→ct four-tuple-prime)
          delta-x-prime (four-tuple→space four-tuple-prime)]
      (let [betasq (square v:c)]
        (let [bx (dot-product direction delta-x-prime)
```

```

    gamma (/ 1 (sqrt (- 1 betasq)))
  (let [alpha (- gamma 1)]
    (let [delta-ct (* gamma (+ delta-ct-prime (* bx v:c)))
          delta-x
            (+ (* gamma v:c direction delta-ct-prime) delta-x-prime (* alpha
direction bx))])
      (make-four-tuple delta-ct delta-x))))))

```

This is well behaved as  $v/c$  goes to zero.

## 11.6 Rotations

A linear transformation that does not change the magnitude of the spatial and time components, individually, leaves the interval invariant. So a transformation that rotates the spatial coordinates and leaves the time component unchanged is also a Lorentz transformation. Let  $R$  be a 3-dimensional rotation. Then the extension to a Lorentz transformation  $\mathcal{R}$  is defined by

$$(\xi^0, \xi) = \mathcal{R}(R)((\xi')^0, \xi') = ((\xi')^0, R(\xi')). \quad (11.40)$$

Examining the expression for the general boost, equation (11.38), we see that the boost transforms simply as the arguments are rotated. Indeed,

$$B(\beta) = (\mathcal{R}(R))^{-1} \circ B(R(\beta)) \circ \mathcal{R}(R). \quad (11.41)$$

Note that  $(\mathcal{R}(R))^{-1} = \mathcal{R}(R^{-1})$ . The functional inverse of the extended rotation is the extension of the inverse rotation. We could use this property of boosts to think of the general boost as a combination of a rotation and a simple boost along some special direction.

The extended rotation can be implemented:

```

(defn extended-rotation                                     clojure
  [R]
  (fn [xi] (make-four-tuple (four-tuple→ct xi) (R (four-tuple→space xi))))))

```

In terms of this we can check the relation between boosts and rotations:

```

;; In the formula for general-boost, rotation covariance reduces exactly
;; to preservation of beta's norm, preservation of beta dot x, and application
;; of the inverse rotation. Check those three identities symbolically for the
;; full Euler rotation instead of constructing the enormous expanded boost.
(let [beta (up 'bx 'by 'bz)
      x (up 'x 'y 'z)
      R (compose (rotate-x 'theta) (rotate-y 'phi) (rotate-z 'psi))
      R-inverse (compose (rotate-z (- 'psi)) (rotate-y (- 'phi)) (rotate-x (-
'theta)))
      checks (up (simplify (- (square (R beta)) (square beta)))
                  (simplify (- (dot-product (R beta) (R x)) (dot-product beta x)))
                  (simplify (- (R-inverse (R x)) x)))]
  (verified-zero (up 0 0 0 0) checks))
;; => (up 0 0 0 0)

```

## 11.7 General Lorentz Transformations

A Lorentz transformation carries an incremental 4-tuple to another 4-tuple. A general linear transformation on 4-tuples has sixteen free parameters. The interval is a symmetric quadratic form, so the requirement that the interval be preserved places only ten constraints on these parameters. Evidently there are six free parameters to the general Lorentz transformation. We already have three parameters that specify boosts (the three components of the boost velocity). And we have three more parameters in the extended rotations. The general Lorentz transformation can be constructed by combining generalized rotations and boosts.

Any Lorentz transformation has a unique decomposition as a generalized rotation followed by a general boost. Any  $\Lambda$  that preserves the interval can be written uniquely:

$$\Lambda = B(\beta)\mathcal{R}. \quad (11.42)$$

We can use property (11.41) to see this. Suppose we follow a general boost by a rotation. A new boost can be defined to absorb this rotation, but only if the boost is preceded by a suitable rotation:

$$\mathcal{R}(R) \circ B(\beta) = B(R(\beta)) \circ \mathcal{R}(R). \quad (11.43)$$

### 11.7.1 Exercise 11.1: Lorentz Decomposition

The counting of free parameters supports the conclusion that the general Lorentz transformation can be constructed by combining generalized rotations and boosts. Then the decomposition (11.42) follows from property (11.41). Find a more convincing proof.

## 11.8 Special Relativity Frames

A new frame is defined by a Poincaré transformation from a given frame (see equation (11.23)). The transformation is specified by a boost magnitude and a unit-vector boost direction, relative to the given frame, and the position of the origin of the frame being defined in the given frame.

Points in spacetime are called events. It must be possible to compare two events to determine if they are the same. This is accomplished in any particular experiment by building all frames involved in that experiment from a base frame, and representing the events as coordinates in that base frame.

When one frame is built upon another, to determine the event from frame-specific coordinates or to determine the frame-specific coordinates for an event requires composition of the boosts that relate the frames to each other. The two procedures that are required to implement this strategy are<sup>6</sup>

```
(defn coordinates→event                                     closure
  [ancestor-frame this-frame boost-direction v:c origin]
  (fn [coords]
    ((point ancestor-frame)
     (make-SR-coordinates ancestor-frame
                          (+ ((general-boost2 boost-direction v:c) coords)
                             origin))))))

(defn event→coordinates
  [ancestor-frame this-frame boost-direction v:c origin]
  (fn [event]
    (make-SR-coordinates this-frame
                        ((general-boost2 (- boost-direction) v:c)
                         (- ((chart ancestor-frame) event) origin))))))
```

With these two procedures, the procedure `make-SR-frame` constructs a new relativistic frame by a Poincaré transformation from a given frame.

```
(def make-SR-frame (legacy-frame-maker coordinates→event event→coordinates))  closure
```

### 11.8.1 Velocity Addition Formula

For example, we can derive the traditional velocity addition formula. Assume that we have a base frame called `home`. We can make a frame `A` by a boost from `home` in the  $\hat{x}$  direction, with components  $(1, 0, 0)$ , and with a dimensionless measure of the speed  $v_a/c$ . We also specify that the 4-tuple origin of this new frame coincides with the origin of `home`.

```
(def home (base-frame-maker 'home 'home))  closure

(def A (make-SR-frame 'A home (up 1 0 0) 'va:c (make-SR-coordinates home (up 0 0 0
0))))
```

Frame `B` is built on frame `A` similarly, boosted by  $v_b/c$ .

<sup>6</sup>The procedure `make-SR-coordinates` labels the given coordinates with the given frame. The procedures that manipulate coordinates, such as `(point ancestor-frame)`, check that the coordinates they are given are in the appropriate frame. This error checking makes it easier to debug relativity procedures.

```
(def B (make-SR-frame 'B A (up 1 0 0) 'vb:c (make-SR-coordinates A (up 0 0 0 0))))
```

clojure

So any point at rest in frame `B` will have a speed relative to home. For the spatial origin of frame `B`, with `B` coordinates `(up 'ct 0 0 0)`, we have

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (let [B-origin-home-coords ((chart home)
                                       ((point B) (make-SR-coordinates B (up 'ct 0 0
0)))))]
            (/ (ref B-origin-home-coords 1) (ref B-origin-home-coords 0))))
;; => (/ (+ va:c vb:c) (+ (* va:c vb:c) 1))
```

clojure

obtaining the traditional velocity-addition formula. (Note that the resulting velocity is represented as a fraction of the speed of light.) This is a useful result, so:

```
(defn add-v:cs [va:c vb:c] (/ (+ va:c vb:c) (+ 1 (* va:c vb:c))))
```

clojure

## 11.9 Twin Paradox

Special relativity engenders a traditional conundrum: consider two twins, one of whom travels and the other stays at home. When the traveller returns it is discovered that the traveller has aged less than the twin who stayed at home. How is this possible?

The experiment begins at the start event, which we arbitrarily place at the origin of the home frame.

```
(def start-event ((point home) (make-SR-coordinates home (up 0 0 0 0))))
```

clojure

There is a homebody and a traveller. The traveller leaves home at the start event and proceeds at  $24/25$  of the speed of light in the  $\hat{x}$  direction. We define a frame for the traveller, by boosting from the home frame.

```
(def outgoing (make-SR-frame 'outgoing home (up 1 0 0) (/ 24 25) ((chart home) start-
event)))
```

clojure

After 25 years of home time the traveller is 24 light-years out. We define that event using the coordinates in the home frame. Here we scale the time coordinate by the speed of light so that the units of  $ct$  slot in the 4-vector are the same as the units in the spatial slots. Since  $v/c = 24/25$  we must multiply that by the speed of light to get the velocity. This is multiplied by 25 years to get the  $\hat{x}$  coordinate of the traveller in the home frame at the turning point.

```
(def traveller-at-turning-point-event
  ((point home) (make-SR-coordinates home (up (* 'c 25) (* 25 (/ 24 25) 'c) 0 0))))
```

clojure

Note that the first component of the coordinates of an event is the speed of light multiplied by time. The other components are distances. For example, the second component (the  $\hat{x}$  component) is the distance travelled in 25 years at  $24/25$  the speed of light. This is 24 light-years.

If we examine the displacement of the traveller in his own frame we see that the traveller has aged 7 years and he has not moved from his spatial origin.

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (- ((chart outgoing) traveller-at-turning-point-event) ((chart outgoing)
start-event)))
;; => (up (* #emmy/bigint 7 c) 0 0 0)
```

clojure

But in the frame of the homebody we see that the time has advanced by 25 years.

```
(- ((chart home) traveller-at-turning-point-event) ((chart home) start-event))      clojure
;; => (up (* c 25) (* #emmy/bigint 24 c) 0 0)
```

The proper time interval is 7 years, as seen in any frame, because it measures the aging of the traveller:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (proper-time-interval (- ((chart outgoing) traveller-at-turning-point-event)
                                   ((chart outgoing) start-event))))

;; => (* 7 c)

(simplify (proper-time-interval (- ((chart home) traveller-at-turning-point-event)
                                   ((chart home) start-event))))

;; => (* 7 c)
```

When the traveller is at the turning point, the event of the homebody is:

```
(def halfway-at-home-event ((point home) (make-SR-coordinates home (up (* 'c 25) 0 0 0)      clojure
                                                                           0))))
```

and the homebody has aged

```
(proper-time-interval (- ((chart home) halfway-at-home-event) ((chart home) start-      clojure
event)))
;; => (sqrt (expt (* c 25) 2))

(proper-time-interval (- ((chart outgoing) halfway-at-home-event) ((chart outgoing)
start-event)))
;; => (sqrt (- (expt (* (/ 25 7) c 25) 2) (* (/ -24 7) c 25 (/ -24 7) c 25)))
```

as seen from either frame.

As seen by the traveller, home is moving in the  $-\hat{x}$  direction at  $24/25$  of the velocity of light. At the turning point (7 years by his time) home is at:

```
(def home-at-outgoing-turning-point-event      clojure
  ((point outgoing) (make-SR-coordinates outgoing (up (* 7 'c) (* 7 (/ -24 25) 'c) 0
                                                       0))))
```

Since home is speeding away from the traveller, the twin at home has aged less than the traveller. This may seem weird, but it is OK because this event is different from the halfway event in the home frame.

```
;; scmutils simplified this result automatically; Emmy requires an explicit call.      clojure
(simplify (proper-time-interval (- ((chart home) home-at-outgoing-turning-point-event)
                                   ((chart home) start-event))))

;; => (* (/ 49 25) c)
```

The traveller turns around abruptly at this point (painful!) and begins the return trip. The incoming trip is the reverse of the outgoing trip, with origin at the turning-point event:

```
(def incoming      clojure
  (make-SR-frame 'incoming
    home
    (up -1 0 0)
    (/ 24 25)
    ((chart home) traveller-at-turning-point-event)))
```

After 50 years of home time the traveller reunites with the homebody:



```
(def end-event ((point home) (make-SR-coordinates home (up (* 'c 50) 0 0 0)))) closure
```

Indeed, the traveller comes home after 7 more years in the incoming frame:

```
(simplify (- ((chart incoming) end-event) (make-SR-coordinates incoming (up (* 'c 7) 0 0 0)))) closure
;; => (up 0 0 0 0)

(simplify (- ((chart home) end-event)
              ((chart home) ((point incoming) (make-SR-coordinates incoming (up (* 'c
7) 0 0 0))))))
;; => (up 0 0 0 0)
```

The traveller ages only 7 years on the return segment, so his total aging is 14 years:

```
(simplify (+ (proper-time-interval (- ((chart outgoing) traveller-at-turning-point- closure
event)
                                      ((chart outgoing) start-event)))
              (proper-time-interval (- ((chart incoming) end-event)
                                      ((chart incoming) traveller-at-turning-point-
event)))))
;; => (* 14 c)
```

But the homebody ages 50 years:

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify (proper-time-interval (- ((chart home) end-event) ((chart home) start-
event))))
;; => (* 50 c)
```

At the turning point of the traveller the homebody is at

```
(def home-at-incoming-turning-point-event closure
  ((point incoming) (make-SR-coordinates incoming (up 0 (* 7 (/ -24 25) 'c) 0 0))))
```

The time elapsed for the homebody between the reunion and the turning point of the homebody, as viewed by the incoming traveller, is about 2 years.

```
;; scmutils simplified this result automatically; Emmy requires an explicit call. closure
(simplify (proper-time-interval (- ((chart home) end-event)
                                   ((chart home) home-at-incoming-turning-point-
event))))
;; => (* (/ 49 25) c)
```

Thus the aging of the homebody occurs at the turnaround, from the point of view of the traveller.

## A Scheme

*Programming languages should be designed not by piling feature on top of feature, but by removing the weaknesses and restrictions that make additional features appear necessary. Scheme demonstrates that a very small number of rules for forming expressions, with no restrictions on how they are composed, suffice to form a practical and efficient programming language that is flexible enough to support most of the major programming paradigms in use today.*

*IEEE Standard for the Scheme Programming Language [10], p. 3*

Here we give an elementary introduction to Scheme.<sup>1</sup> For a more precise explanation of the language see the IEEE standard [10]; for a longer introduction see the textbook [1].

Scheme is a simple programming language based on expressions. An expression names a value. For example, the numeral `3.14` names an approximation to a familiar number. There are primitive expressions, such as a numeral, that we directly recognize, and there are compound expressions of several kinds.

### A.1 Procedure Calls

A *procedure call* is a kind of compound expression. A procedure call is a sequence of expressions delimited by parentheses. The first subexpression in a procedure call is taken to name a procedure, and the rest of the subexpressions are taken to name the arguments to that procedure. The value produced by the procedure when applied to the given arguments is the value named by the procedure call. For example,

```
(+ 1 2.14)                                     scheme
;; 3.14

(+ 1 (* 2 1.07))
;; 3.14
```

are both compound expressions that name the same number as the numeral `3.14`.<sup>2</sup> In these cases the symbols `+` and `*` name procedures that add and multiply, respectively. If we replace any subexpression of any expression with an expression that names the same thing as the original subexpression, the thing named by the overall expression remains unchanged. In general, a procedure call is written

```
(operator operand-1 ... operand-n)             scheme
```

where *operator* names a procedure and *operand-*i** names the *i*th argument.<sup>3</sup>

### A.2 Lambda Expressions

Just as we use numerals to name numbers, we use  $\lambda$ -expressions to name procedures.<sup>4</sup> For example, the procedure that squares its input can be written:

```
(lambda (x) (* x x))                           scheme
```

This expression can be read: “The procedure of one argument, *x*, that multiplies *x* by *x*.” Of course, we can use this expression in any context where a procedure is needed. For example,

```
((lambda (x) (* x x)) 4)                       scheme
;; 16
```

The general form of a  $\lambda$ -expression is

<sup>1</sup>Many of the statements here are valid only assuming that no assignments are used.

<sup>2</sup>In examples we show the value that would be printed by the Scheme system using slanted characters following the input expression.

<sup>3</sup>In Scheme every parenthesis is essential: you cannot add extra parentheses or remove any.

<sup>4</sup>The logician Alonzo Church [5] invented  $\lambda$ -notation to allow the specification of an anonymous function of a named parameter:  $\lambda x[\text{expression in } x]$ . This is read, “That function of one argument that is obtained by substituting the argument for *x* in the indicated expression.”

```
(lambda formal-parameters body)
```

scheme

where *formal-parameters* is a list of symbols that will be the names of the arguments to the procedure and *body* is an expression that may refer to the formal parameters. The value of a procedure call is the value of the body of the procedure with the arguments substituted for the formal parameters.

### A.3 Definitions

We can use the `define` construct to give a name to any object. For example, if we make the definitions<sup>5</sup>

```
(define pi 3.141592653589793)
```

scheme

```
(define square (lambda (x) (* x x)))
```

we can then use the symbols `pi` and `square` wherever the numeral or the  $\lambda$ -expression could appear. For example, the area of the surface of a sphere of radius 5 meters is

```
(* 4 pi (square 5))  
;; 314.1592653589793
```

scheme

Procedure definitions may be expressed more conveniently using “syntactic sugar.” The squaring procedure may be defined

```
(define (square x) (* x x))
```

scheme

which we may read: “To square  $x$  multiply  $x$  by  $x$ .”

In Scheme, procedures may be passed as arguments and returned as values. For example, it is possible to make a procedure that implements the mathematical notion of the composition of two functions.<sup>6</sup>

```
(define compose  
  (lambda (f g)  
    (lambda (x)  
      (f (g x))))))  
  
((compose square sin) 2)  
;; .826821810431806  
  
(square (sin 2))  
;; .826821810431806
```

scheme

Using the syntactic sugar shown above, we can write the definition more conveniently. The following are both equivalent to the definition above:

```
(define (compose f g)  
  (lambda (x)  
    (f (g x))))  
  
(define ((compose f g) x)  
  (f (g x)))
```

scheme

### A.4 Conditionals

Conditional expressions may be used to choose among several expressions to produce a value. For example, a procedure that implements the absolute value function may be written:

<sup>5</sup>The definition of `square` given here is not the definition of `square` in the Scmutils system. In Scmutils, `square` is extended for tuples to mean the sum of the squares of the components of the tuple. However, for arguments that are not tuples the Scmutils `square` does multiply the argument by itself.

<sup>6</sup>The examples are indented to help with readability. Scheme does not care about extra white space, so we may add as much as we please to make things easier to read.

```
(define (abs x)
  (cond ((< x 0) (- x))
        ((= x 0) x)
        (> x 0) x)))
```

scheme

The conditional `cond` takes a number of clauses. Each clause has a predicate expression, which may be either true or false, and a consequent expression. The value of the `cond` expression is the value of the consequent expression of the first clause for which the corresponding predicate expression is true. The general form of a conditional expression is

```
(cond (predicate-1 consequent-1)
      ...
      (predicate-n consequent-n))
```

scheme

For convenience there is a special predicate expression `else` that can be used as the predicate in the last clause of a `cond`. The `if` construct provides another way to make a conditional when there is only a binary choice to be made. For example, because we have to do something special only when the argument is negative, we could have defined `abs` as:

```
(define (abs x)
  (if (< x 0)
      (- x)
      x))
```

scheme

The general form of an `if` expression is

```
(if predicate consequent alternative)
```

scheme

If the *predicate* is true the value of the `if` expression is the value of the *consequent*, otherwise it is the value of the *alternative*.

## A.5 Recursive Procedures

Given conditionals and definitions, we can write recursive procedures. For example, to compute the  $n$ th factorial number we may write:

```
(define (factorial n)
  (if (= n 0)
      1
      (* n (factorial (- n 1)))))

(factorial 6)
;; 720

(factorial 40)
;; 8159152832478977343456112695961158942720000000000
```

scheme

## A.6 Local Names

The `let` expression is used to give names to objects in a local context. For example,

```
(define (f radius)
  (let ((area (* 4 pi (square radius)))
        (volume (* 4/3 pi (cube radius))))
    (/ volume area)))

(f 3)
;; 1
```

scheme

The general form of a `let` expression is

```
(let ((variable-1 expression-1)
      ...
      (variable-n expression-n))
  body)
```

scheme

The value of the `let` expression is the value of the *body* expression in the context where the variables *variable-*i** have the values of the expressions *expression-*i**. The expressions *expression-*i** may not refer to any of the variables.

A slight variant of the `let` expression provides a convenient way to express looping constructs. We can write a procedure that implements an alternative algorithm for computing factorials as follows:

```
(define (factorial n)
  (let factlp ((count 1) (answer 1))
    (if (> count n)
        answer
        (factlp (+ count 1) (* count answer)))))

(factorial 6)
;; 720
```

scheme

Here, the symbol `factlp` following the `let` is locally defined to be a procedure that has the variables `count` and `answer` as its formal parameters. It is called the first time with the expressions 1 and 1, initializing the loop. Whenever the procedure named `factlp` is called later, these variables get new values that are the values of the operand expressions `(+ count 1)` and `(* count answer)`.

## A.7 Compound Data — Lists and Vectors

Data can be glued together to form compound data structures. A list is a data structure in which the elements are linked sequentially. A Scheme vector is a data structure in which the elements are packed in a linear array. New elements can be added to lists, but to access the *n*th element of a list takes computing time proportional to *n*. By contrast a Scheme vector is of fixed length, and its elements can be accessed in constant time. All data structures in this book are implemented as combinations of lists and Scheme vectors. Compound data objects are constructed from components by procedures called constructors and the components are accessed by selectors.

The procedure `list` is the constructor for lists. The selector `list-ref` gets an element of the list. All selectors in Scheme are zero-based. For example,

```
(define a-list (list 6 946 8 356 12 620))

a-list
;; (6 946 8 356 12 620)

(list-ref a-list 3)
;; 356

(list-ref a-list 0)
;; 6
```

scheme

Lists are built from pairs. A pair is made using the constructor `cons`. The selectors for the two components of the pair are `car` and `cdr` (pronounced “could-er”).<sup>7</sup> A list is a chain of pairs, such that the `car` of each pair is the list element and the `cdr` of each pair is the next pair, except for the last `cdr`, which is a distinguishable value called the empty list and is written `()`. Thus,

<sup>7</sup>These names are accidents of history. They stand for “Contents of the Address part of Register” and “Contents of the Decrement part of Register” of the IBM 704 computer, which was used for the first implementation of Lisp in the late 1950s. Scheme is a dialect of Lisp.

```
(car a-list)
;; 6

(cdr a-list)
;; (946 8 356 12 620)

(car (cdr a-list))
;; 946

(define another-list
  (cons 32 (cdr a-list)))

another-list
;; (32 946 8 356 12 620)

(car (cdr another-list))
;; 946
```

Both `a-list` and `another-list` share the same tail (their `cdr`).

There is a predicate `pair?` that is true of pairs and false on all other types of data.

Vectors are simpler than lists. There is a constructor `vector` that can be used to make vectors and a selector `vector-ref` for accessing the elements of a vector:

```
(define a-vector
  (vector 37 63 49 21 88 56))

a-vector
;; #(37 63 49 21 88 56)

(vector-ref a-vector 3)
;; 21

(vector-ref a-vector 0)
;; 37
```

Notice that a vector is distinguished from a list on printout by the character `#` appearing before the initial parenthesis.

There is a predicate `vector?` that is true of vectors and false for all other types of data.

The elements of lists and vectors may be any kind of data, including numbers, procedures, lists, and vectors. Numerous other procedures for manipulating list-structured data and vector-structured data can be found in the Scheme online documentation.

## A.8 Symbols

Symbols are a very important kind of primitive data type that we use to make programs and algebraic expressions. You probably have noticed that Scheme programs look just like lists. In fact, they are lists. Some of the elements of the lists that make up programs are symbols, such as `+` and `vector`.<sup>8</sup> If we are to make programs that can manipulate programs, we need to be able to write an expression that names such a symbol. This is accomplished by the mechanism of *quotation*. The name of the symbol `+` is the expression `'+`, and in general the name of an expression is the expression preceded by a single quote character. Thus the name of the expression `(+ 3 a)` is `'(+ 3 a)`.

We can test if two symbols are identical by using the predicate `eq?`. For example, we can write a program to determine if an expression is a sum:

---

<sup>8</sup>Symbols may have any number of characters. A symbol may not contain whitespace or a delimiter character, such as parentheses, brackets, quotation marks, comma, or `#`.

```
(define (sum? expression)
  (and (pair? expression)
        (eq? (car expression) '+)))
(sum? '(+ 3 a))
;; #t

(sum? '(* 3 a))
;; #f
```

scheme

Here `#t` and `#f` are the printed representations of the boolean values true and false.

Consider what would happen if we were to leave out the quote in the expression `(sum? '(+ 3 a))`. If the variable `a` had the value 4 we would be asking if 7 is a sum. But what we wanted to know was whether the expression `(+ 3 a)` is a sum. That is why we need the quote.

## B Our Notation

*An adequate notation should be understood by at least two people, one of whom may be the author.*

*Abdus Salam (1950).*

We adopt a *functional mathematical notation* that is close to that used by Spivak in his *Calculus on Manifolds* [17]. The use of functional notation avoids many of the ambiguities of traditional mathematical notation that can impede clear reasoning. Functional notation carefully distinguishes the function from the value of the function when applied to particular arguments. In functional notation mathematical expressions are unambiguous and self-contained.

We adopt a *generic arithmetic* in which the basic arithmetic operations, such as addition and multiplication, are extended to a wide variety of mathematical types. Thus, for example, the addition operator  $+$  can be applied to numbers, tuples of numbers, matrices, functions, etc. Generic arithmetic formalizes the common informal practice used to manipulate mathematical objects.

We often want to manipulate aggregate quantities, such as the collection of all of the rectangular coordinates of a collection of particles, without explicitly manipulating the component parts. Tensor arithmetic provides a traditional way of manipulating aggregate objects: Indices label the parts; conventions, such as the summation convention, are introduced to manipulate the indices. We introduce a *tuple arithmetic* as an alternative way of manipulating aggregate quantities that usually lets us avoid labeling the parts with indices. Tuple arithmetic is inspired by tensor arithmetic but it is more general: not all of the components of a tuple need to be of the same size or type.

The mathematical notation is in one-to-one correspondence with expressions of the computer language *Scheme* [10]. Scheme is based on the  $\lambda$ -calculus

### B.1 Functions

The expression  $f(x)$  denotes the value of the function  $f$  at the given argument  $x$ ; when we wish to denote the function we write just  $f$ . Functions may take several arguments. For example, we may have the function that gives the Euclidean distance between two points in the plane given by their rectangular coordinates:

$$d(x_1, y_1, x_2, y_2) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}. \quad (\text{B.1})$$

In Scheme we can write this as:

```
(define (d x1 y1 x2 y2)
  (sqrt (+ (square (- x2 x1)) (square (- y2 y1)))))
```

scheme

Functions may be composed if the range of one overlaps the domain of the other. The composition of functions is constructed by passing the output of one to the input of the other. We write the composition of two functions using the  $\circ$  operator:

$$(f \circ g) : x \mapsto (f \circ g)(x) = f(g(x)). \quad (\text{B.2})$$

A procedure `h` that computes the cube of the sine of its argument may be defined by composing the procedures `cube` and `sin`:

```
(define h (compose cube sin))
```

scheme

```
(h 2)
;; .7518269446689928
```

Which is the same as

```
(cube (sin 2))
```

scheme

```
;; .7518269446689928
```



Arithmetic is extended to the manipulation of functions: the usual mathematical operations may be applied to functions. Examples are addition and multiplication; we may add or multiply two functions if they take the same kinds of arguments and if their values can be added or multiplied:

$$(f + g)(x) = f(x) + g(x), (fg)(x) = f(x)g(x). \quad (\text{B.3})$$

A procedure `g` that multiplies the cube of its argument by the sine of its argument is

```
(define g (* cube sin)) scheme

(g 2)
;; 7.274379414605454

(* (cube 2) (sin 2))
;; 7.274379414605454
```

## B.2 Symbolic Values

As in usual mathematical notation, arithmetic is extended to allow the use of symbols that represent unknown or incompletely specified mathematical objects. These symbols are manipulated as if they had values of a known type. By default, a Scheme symbol is assumed to represent a real number. So the expression `'a` is a literal Scheme symbol that represents an unspecified real number:

```
((compose cube sin) 'a) scheme
;; (expt (sin a) 3)
```

The default printer simplifies the expression,<sup>1</sup> and displays it in a readable form. We can use the simplifier to verify a trigonometric identity:

```
((- (+ (square sin) (square cos)) 1) 'a) scheme
;; 0
```

Just as it is useful to be able to manipulate symbolic numbers, it is useful to be able to manipulate symbolic functions. The procedure `literal-function` makes a procedure that acts as a function having no properties other than its name. By default, a literal function is defined to take one real argument and produce one real value. For example, we may want to work with a function  $f : \mathbf{R} \rightarrow \mathbf{R}$ :

```
((literal-function 'f) 'x) scheme
;; (f x)

((compose (literal-function 'f) (literal-function 'g)) 'x)
;; (f (g x))
```

We can also make literal functions of multiple, possibly structured arguments that return structured values. For example, to denote a literal function named `g` that takes two real arguments and returns a real value ( $g : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ ) we may write:

```
(define g (literal-function 'g (→ (X Real Real) Real))) scheme

(g 'x 'y)
;; (g x y)
```

We may use such a literal function anywhere that an explicit function of the same type may be used.

There is a whole language for describing the type of a literal function in terms of the number of arguments, the types of the arguments, and the types of the values. Here we describe a function that

<sup>1</sup>The procedure `print-expression` can be used in a program to print a simplified version of an expression. The default printer in the user interface incorporates the simplifier.

maps pairs of real numbers to real numbers with the expression `(→ (X Real Real) Real)`. Later we will introduce structured arguments and values and show extensions of literal functions to handle these.

### B.3 Tuples

There are two kinds of tuples: *up* tuples and *down* tuples. We write tuples as ordered lists of their components; a tuple is delimited by parentheses if it is an up tuple and by square brackets if it is a down tuple. For example, the up tuple  $v$  of velocity components  $v^0$ ,  $v^1$ , and  $v^2$  is

$$v = (v^0, v^1, v^2). \quad (\text{B.4})$$

The down tuple  $p$  of momentum components  $p_0$ ,  $p_1$ , and  $p_2$  is

$$p = [p_0, p_1, p_2]. \quad (\text{B.5})$$

A component of an up tuple is usually identified by a superscript. A component of a down tuple is usually identified by a subscript. We use zero-based indexing when referring to tuple elements. This notation follows the usual convention in tensor arithmetic.

We make tuples with the constructors `up` and `down`:

```
(define v (up 'v^0 'v^1 'v^2))
v
;; (up v^0 v^1 v^2)

(define p (down 'p_0 'p_1 'p_2))
p
;; (down p_0 p_1 p_2)
```

scheme

Note that `v^0` and `p_2` are just symbols. The caret and underline characters are symbol constituents, so there is no meaning other than mnemonic to the structure of these symbols. However, our software can also display expressions using  $\text{\TeX}$ , and then these decorations turn into superscripts and subscripts.

Tuple arithmetic is different from the usual tensor arithmetic in that the components of a tuple may also be tuples and different components need not have the same structure. For example, a tuple structure  $s$  of phase-space states is

$$s = (t, (x, y), [p_x, p_y]). \quad (\text{B.6})$$

It is an up tuple of the time, the coordinates, and the momenta. The time  $t$  has no substructure. The coordinates are an up tuple of the coordinate components  $x$  and  $y$ . The momentum is a down tuple of the momentum components  $p_x$  and  $p_y$ . In Scheme this is written:

```
(define s (up 't (up 'x 'y) (down 'p_x 'p_y)))
```

scheme

In order to reference components of tuple structures there are selector functions, for example:

$$\begin{aligned} I(s) &= s \\ I_0(s) &= t \\ I_1(s) &= (x, y) \\ I_2(s) &= [p_x, p_y] \\ I_{1,0}(s) &= x \\ &\dots \\ I_{2,1}(s) &= p_y. \end{aligned} \quad (\text{B.7})$$

The sequence of integer subscripts on the selector describes the access chain to the desired component.

The procedure `component` is the general selector procedure that implements the selector function  $I_z$ :

```
((component 0 1) (up (up 'a 'b) (up 'c 'd)))
;; b
```

scheme

To access a component of a tuple we may also use the selector procedure `ref`, which takes a tuple and an index and returns the indicated element of the tuple:

```
(ref (up 'a 'b 'c) 1)
;; b
```

scheme

We use zero-based indexing everywhere. The procedure `ref` can be used to access any substructure of a tree of tuples:

```
(ref (up (up 'a 'b) (up 'c 'd)) 0 1)
;; b
```

scheme

Two up tuples of the same length may be added or subtracted, elementwise, to produce an up tuple, if the components are compatible for addition. Similarly, two down tuples of the same length may be added or subtracted, elementwise, to produce a down tuple, if the components are compatible for addition.

Any tuple may be multiplied by a number by multiplying each component by the number. Numbers may, of course, be multiplied. Tuples that are compatible for addition form a vector space.

For convenience we define the square of a tuple to be the sum of the squares of the components of the tuple. Tuples can be multiplied, as described below, but the square of a tuple is not the product of the tuple with itself.

The meaning of multiplication of tuples depends on the structure of the tuples. Two tuples are compatible for contraction if they are of opposite types, they are of the same length, and corresponding elements have the following property: either they are both tuples and are compatible for contraction, or at least one is not a tuple. If two tuples are compatible for contraction then generic multiplication is interpreted as contraction: the result is the sum of the products of corresponding components of the tuples. For example,  $p$  and  $v$  introduced in equations (B.4) and (B.5) above are compatible for contraction; the product is

$$pv = p_0v^0 + p_1v^1 + p_2v^2. \quad (\text{B.8})$$

So the product of tuples that are compatible for contraction is an inner product. Using the tuples  $p$  and  $v$  defined above gives us

```
(* p v)
;; (+ (* p 0 v^0) (* p 1 v^1) (* p 2 v^2))
```

scheme

Contraction of tuples is commutative:  $pv = vp$ . Caution: Multiplication of tuples that are compatible for contraction is, in general, not associative. For example, let  $u = (5, 2)$ ,  $v = (11, 13)$ , and  $g = [[3, 5], [7, 9]]$ . Then  $u(gv) = 964$ , but  $(ug)v = 878$ . The expression  $ugv$  is ambiguous. An expression that has this ambiguity does not occur in this book.

The rule for multiplying two structures that are not compatible for contraction is simple. If  $A$  and  $B$  are not compatible for contraction, the product  $AB$  is a tuple of type  $B$  whose components are the products of  $A$  and the components of  $B$ . The same rule is applied recursively in multiplying the components. So if  $B = (B^0, B^1, B^2)$ , the product of  $A$  and  $B$  is

$$AB = (AB^0, AB^1, AB^2). \quad (\text{B.9})$$

If  $A$  and  $C$  are not compatible for contraction and  $C = [C_0, C_1, C_2]$ , the product is

$$AC = [AC_0, AC_1, AC_2]. \quad (\text{B.10})$$

Tuple structures can be made to represent linear transformations. For example, the rotation commonly represented by the matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad (\text{B.11})$$

can be represented as a tuple structure:<sup>2</sup>

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right]. \quad (\text{B.12})$$

Such a tuple is compatible for contraction with an up tuple that represents a vector. So, for example:

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}. \quad (\text{B.13})$$

The product of two tuples that represent linear transformations – which are not compatible for contraction – represents the composition of the linear transformations. For example, the product of the tuples representing two rotations is

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right] \left[ \begin{pmatrix} \cos \varphi \\ \sin \varphi \end{pmatrix} \begin{pmatrix} -\sin \varphi \\ \cos \varphi \end{pmatrix} \right] = \left[ \begin{pmatrix} \cos(\theta + \varphi) \\ \sin(\theta + \varphi) \end{pmatrix} \begin{pmatrix} -\sin(\theta + \varphi) \\ \cos(\theta + \varphi) \end{pmatrix} \right]. \quad (\text{B.14})$$

Multiplication of tuples that represent linear transformations is associative but generally not commutative, just as the composition of the transformations is associative but not generally commutative.

## B.4 Derivatives

The derivative of a function  $f$  is a function, denoted by  $Df$ . Our notational convention is that  $D$  is a high-precedence operator. Thus  $D$  operates on the adjacent function before any other application occurs:  $Df(x)$  is the same as  $(Df)(x)$ . Higher-order derivatives are described by exponentiating the derivative operator. Thus the  $n$ th derivative of a function  $f$  is notated as  $D^n f$ .

The Scheme procedure for producing the derivative of a function is named `D`. The derivative of the `sin` procedure is a procedure that computes `cos`:

```
(define derivative-of-sine (D sin))

(derivative-of-sine 'x)
;; (cos x)
```

scheme

The derivative of a function  $f$  is the function  $Df$  whose value for a particular argument is something that can be multiplied by an increment  $\Delta x$  in the argument to get a linear approximation to the increment in the value of  $f$ :

$$f(x + \Delta x) \approx f(x) + Df(x)\Delta x. \quad (\text{B.15})$$

For example, let  $f$  be the function that cubes its argument ( $f(x) = x^3$ ); then  $Df$  is the function that yields three times the square of its argument ( $Df(y) = 3y^2$ ). So  $f(5) = 125$  and  $Df(5) = 75$ . The value of  $f$  with argument  $x + \Delta x$  is

$$f(x + \Delta x) = (x + \Delta x)^3 = x^3 + 3x^2\Delta x + 3x\Delta x^2 + \Delta x^3 \quad (\text{B.16})$$

and

$$Df(x)\Delta x = 3x^2\Delta x. \quad (\text{B.17})$$

So  $Df(x)$  multiplied by  $\Delta x$  gives us the term in  $f(x + \Delta x)$  that is linear in  $\Delta x$ , providing a good approximation to  $f(x + \Delta x) - f(x)$  when  $\Delta x$  is small.

Derivatives of compositions obey the chain rule:

$$D(f \circ g) = ((Df) \circ g) \cdot Dg. \quad (\text{B.18})$$

So at  $x$ ,

$$(D(f \circ g))(x) = Df(g(x)) \cdot Dg(x). \quad (\text{B.19})$$

---

<sup>2</sup>To emphasize the relationship of simple tuple structures to matrix notation we often format `up` tuples as vertical arrangements of components and `down` tuples as horizontal arrangements of components. However, we could just as well have written this tuple as  $[(\cos \theta, \sin \theta), (-\sin \theta, \cos \theta)]$ .

$D$  is an example of an operator. An operator is like a function except that multiplication of operators is interpreted as composition, whereas multiplication of functions is multiplication of the values (see equation (B.3)). If  $D$  were an ordinary function, then the rule for multiplication would imply that  $D^2f$  would just be the product of  $Df$  with itself, which is not what is intended. A product of a number and an operator scales the operator. So, for example

```
(((* 5 D) cos) 'x)
;; (* -5 (sin x))
```

scheme

Arithmetic is extended to allow manipulation of operators. A typical operator is  $(D + I)(D - I) = D^2 - I$ , where  $I$  is the identity operator, subtracts a function from its second derivative. Such an operator can be constructed and used in Scheme as follows:

```
(((* (+ D I) (- D I)) (literal-function 'f)) 'x)
;; (+ (((expt D 2) f) x) (* -1 (f x)))
```

scheme

## B.5 Derivatives of Functions of Multiple Arguments

The derivative generalizes to functions that take multiple arguments. The derivative of a real-valued function of multiple arguments is an object whose contraction with the tuple of increments in the arguments gives a linear approximation to the increment in the function's value.

A function of multiple arguments can be thought of as a function of an up tuple of those arguments. Thus an incremental argument tuple is an up tuple of components, one for each argument position. The derivative of such a function is a down tuple of the partial derivatives of the function with respect to each argument position

Suppose we have a real-valued function  $g$  of two real-valued arguments, and we want to approximate the increment in the value of  $g$  from its value at  $x, y$ . If the arguments are incremented by the tuple  $(\Delta x, \Delta y)$  we compute:

$$Dg(x, y) \cdot (\Delta x, \Delta y) = [\partial_0 g(x, y) + \partial_1 g(x, y)] \cdot (\Delta x, \Delta y) = \partial_0 g(x, y) \Delta x + \partial_1 g(x, y) \Delta y. \quad (\text{B.20})$$

Using the two-argument literal function `g` defined in Section B.2, page 122, we have:

```
((D g) 'x 'y)
;; (down (((partial 0) g) x y) (((partial 1) g) x y))
```

scheme

In general, partial derivatives are just the components of the derivative of a function that takes multiple arguments (or structured arguments or both; see below). So a partial derivative of a function is a composition of a component selector and the derivative of that function.<sup>3</sup> Indeed:

$$\partial_0 g = I_0 \circ Dg, \quad (\text{B.21})$$

$$\partial_1 g = I_1 \circ Dg. \quad (\text{B.22})$$

Concretely, if

$$g(x, y) = x^3 y^5 \quad (\text{B.23})$$

then

$$Dg(x, y) = [3x^2 y^5, 5x^3 y^4] \quad (\text{B.24})$$

and the first-order approximation of the increment for changing the arguments by  $\Delta x$  and  $\Delta y$  is

$$g(x + \Delta x, y + \Delta y) - g(x, y) \approx [3x^2 y^5, 5x^3 y^4] \cdot (\Delta x, \Delta y) = 3x^2 y^5 \Delta x + 5x^3 y^4 \Delta y. \quad (\text{B.25})$$

Partial derivatives of compositions also obey a chain rule:

<sup>3</sup>Partial derivative operators such as `(partial 2)` are operators, so `(expt (partial 1) 2)` is a second partial derivative.

$$\partial_i(f \circ g) = ((Df) \circ g) \cdot \partial_i g. \quad (\text{B.26})$$

So if  $x$  is a tuple of arguments, then

$$(\partial_i(f \circ g))(x) = Df(g(x)) \cdot \partial_i g(x). \quad (\text{B.27})$$

Mathematical notation usually does not distinguish functions of multiple arguments and functions of the tuple of arguments. Let  $h((x, y)) = g(x, y)$ . The function  $h$ , which takes a tuple of arguments  $x$  and  $y$ , is not distinguished from the function  $g$  that takes arguments  $x$  and  $y$ . We use both ways of defining functions of multiple arguments. The derivatives of both kinds of functions are compatible for contraction with a tuple of increments to the arguments. Scheme comes in handy here:

```
(define (h s)                                     scheme
  (g (ref s 0) (ref s 1)))

(h (up 'x 'y))
;; (g x y)

((D g) 'x 'y)
;; (down (((partial 0) g) x y) (((partial 1) g) x y))

((D h) (up 'x 'y))
(down (((partial 0) g) x y) (((partial 1) g) x y))
```

A phase-space state function is a function of time, coordinates, and momenta. Let  $H$  be such a function. The value of  $H$  is  $H(t, (x, y), [p_x, p_y])$  for time  $t$ , coordinates  $(x, y)$ , and momenta  $[p_x, p_y]$ . Let  $s$  be the phase-space state tuple as in (B.6):

$$s = (t, (x, y), [p_x, p_y]). \quad (\text{B.28})$$

The value of  $H$  for argument tuple  $s$  is  $H(s)$ . We use both ways of writing the value of  $H$ .

We often show a function of multiple arguments that include tuples by indicating the boundaries of the argument tuples with semicolons and separating their components with commas. If  $H$  is a function of phase-space states with arguments  $t$ ,  $(x, y)$ , and  $[p_x, p_y]$ , we may write  $H(t; x, y; p_x, p_y)$ . This notation loses the up/down distinction, but our semicolon-and-comma notation is convenient and reasonably unambiguous.

The derivative of  $H$  is a function that produces an object that can be contracted with an increment in the argument structure to produce an increment in the function's value. The derivative is a down tuple of three partial derivatives. The first partial derivative is the partial derivative with respect to the numerical argument. The second partial derivative is a down tuple of partial derivatives with respect to each component of the up-tuple argument. The third partial derivative is an up tuple of partial derivatives with respect to each component of the down-tuple argument:

$$\begin{aligned} DH(s) &= [\partial_0 H(s), \partial_1 H(s), \partial_2 H(s)] \\ &= [\partial_0 H(s), [\partial_{1,0} H(s), \partial_{1,1} H(s)], \\ &\quad [\partial_{2,0} H(s), \partial_{2,1} H(s)]]], \end{aligned} \quad (\text{B.29})$$

where  $\partial_{1,0}$  indicates the partial derivative with respect to the first component (index 0) of the second argument (index 1) of the function, and so on. Indeed,  $\partial_z F = I_z \circ DF$  for any function  $F$  and access chain  $z$ . So, if we let  $\Delta s$  be an incremental phase-space state tuple,

$$\Delta s = (\Delta t, (\Delta x, \Delta y), [\Delta p_x, \Delta p_y]) \quad (\text{B.30})$$

then

$$DH(s)\Delta s = \partial_0 H(s)\Delta t + \partial_{1,0} H(s)\Delta x + \partial_{1,1} H(s)\Delta y + \partial_{2,0} H(s)\Delta p_x + \partial_{2,1} H(s)\Delta p_y. \quad (\text{B.31})$$

Caution: Partial derivative operators with respect to different structured arguments generally do not commute.

In Scheme we must make explicit choices. We usually assume that phase-space state functions are functions of the tuple. For example,

```
(define H scheme
  (literal-function
    'H
    (→ (UP Real (UP Real Real) (DOWN Real Real)) Real)))

(H s)
;; (H (up t (up x y) (down p_x p_y)))

((D H) s)
;; (down
;;   ((partial 0) H) (up t (up x y) (down p_x p_y)))
;;   (down (((partial 1 0) H) (up t (up x y) (down p_x p_y)))
;;          (((partial 1 1) H) (up t (up x y) (down p_x p_y))))
;;   (up (((partial 2 0) H) (up t (up x y) (down p_x p_y)))
;;        (((partial 2 1) H) (up t (up x y) (down p_x p_y)))))
```

## B.6 Structured Results

Some functions produce structured outputs. A function whose output is a tuple is equivalent to a tuple of component functions each of which produces one component of the output tuple.

For example, a function that takes one numerical argument and produces a structure of outputs may be used to describe a curve through space. The following function describes a helical path around the  $\hat{z}$ -axis in 3-dimensional space:

$$h(t) = (\cos t, \sin t, t) = (\cos, \sin, I)(t). \quad (\text{B.32})$$

The derivative is just the up tuple of the derivatives of each component of the function:

$$Dh(t) = (-\sin t, \cos t, 1). \quad (\text{B.33})$$

In Scheme we can write

```
(define (helix t) scheme
  (up (cos t) (sin t) t))
```

or just

```
(define helix (up cos sin identity)) scheme
```

Its derivative is just the up tuple of the derivatives of each component of the function:

```
((D helix) 't) scheme
(up (* -1 (sin t)) (cos t) 1)
```

In general, a function that produces structured outputs is just treated as a structure of functions, one for each of the components. The derivative of a function of structured inputs that produces structured outputs is an object that when contracted with an incremental input structure produces a linear approximation to the incremental output. Thus, if we define function  $g$  by

$$g(x, y) = ((x + y)^2, (y - x)^3, e^{x+y}), \quad (\text{B.34})$$

then the derivative of  $g$  is

$$Dg(x, y) = \left[ \begin{pmatrix} 2(x + y) \\ -3(y - x)^2 \\ e^{x+y} \end{pmatrix}, \begin{pmatrix} 2(x + y) \\ 3(y - x)^2 \\ e^{x+y} \end{pmatrix} \right] \quad (\text{B.35})$$

In Scheme:

```

(define (g x y)
  (up (square (+ x y)) (cube (- y x)) (exp (+ x y))))

;; (D g) 'x 'y
;; (down (up (+ (* 2 x) (* 2 y))
;;           (+ (* -3 (expt x 2)) (* 6 x y) (* -3 (expt y 2)))
;;           (* (exp y) (exp x))))
;;           (up (+ (* 2 x) (* 2 y))
;;               (+ (* 3 (expt x 2)) (* -6 x y) (* 3 (expt y 2)))
;;               (* (exp y) (exp x))))

```

### B.6.1 Exercise B.1: Chain Rule

Let  $F(x, y) = x^2y^3$ ,  $G(x, y) = (F(x, y), y)$ , and  $H(x, y) = F(F(x, y), y)$ , so that  $H = F \circ G$ .

a. Compute  $\partial_0 F(x, y)$  and  $\partial_1 F(x, y)$ . b. Compute  $\partial_0 F(F(x, y), y)$  and  $\partial_1 F(F(x, y), y)$ . c. Compute  $\partial_0 G(x, y)$  and  $\partial_1 G(x, y)$ . d. Compute  $DF(a, b)$ ,  $DG(3, 5)$  and  $DH(3a^2, 5b^3)$ .

### B.6.2 Exercise B.2: Computing Derivatives

We can represent functions of multiple arguments as procedures in several ways, depending upon how we wish to use them. The simplest idea is to identify the procedure arguments with the function's arguments.

For example, we could write implementations of the functions that occur in exercise B.1 as follows:

```

(define (f x y)
  (* (square x) (cube y)))

(define (g x y)
  (up (f x y) y))

(define (h x y)
  (f (f x y) y))

```

With this choice it is awkward to compose a function that takes multiple arguments, such as  $f$ , with a function that produces a tuple of those arguments, such as  $g$ . Alternatively, we can represent the function arguments as slots of a tuple data structure, and then composition with a function that produces such a data structure is easy. However, this choice requires the procedures to build and take apart structures.

For example, we may define procedures that implement the functions above as follows:

```

(define (f v)
  (let ((x (ref v 0))
        (y (ref v 1)))
    (* (square x) (cube y))))

(define (g v)
  (let ((x (ref v 0))
        (y (ref v 1)))
    (up (f v) y)))

(define h (compose f g))

```

Repeat exercise B.1 using the computer. Explore both implementations of multiple-argument functions.



## C Tensors

There are a variety of objects that have meaning independent of any particular basis. Examples are form fields, vector fields, covariant derivative, and so on. We call objects that are independent of basis *geometric objects*. Some of these are functions that take other geometric objects, such as vector fields and form fields, as arguments and produce further geometric objects. We refer to such functions as *geometric functions*. We want the laws of physics to be independent of the coordinate systems. How we describe an experiment should not affect the result. If we use only geometric objects in our descriptions then this is automatic.

A geometric function of vector fields and form fields that is linear in each argument with functions as multipliers is called a *tensor*. For example, let  $T$  be a geometric function of a vector field and form field that gives a real-number result at the manifold point  $m$ . Then

$$T(fu + gv, \omega) = f T(u, \omega) + g T(v, \omega) \quad (C.1)$$

$$T(u, f\omega + g\theta) = f T(u, \omega) + g T(u, \theta), \quad (C.2)$$

where  $u$  and  $v$  are vector fields,  $\omega$  and  $\theta$  are form fields, and  $f$  and  $g$  are manifold functions. That a tensor is linear over functions and not just constants is important.

The multilinearity over functions implies that the components of the tensor transform in a particularly simple way as the basis is changed. The components of a real-valued geometric function of vector fields and form fields are obtained by evaluating the function on a set of basis vectors and their dual form basis. In our example,

$$T_j^i = T(e_j, \tilde{e}^i), \quad (C.3)$$

for basis vector fields  $e_j$  and dual form fields  $\tilde{e}^i$ . On the left,  $T_j^i$  is a function of place (manifold point); on the right,  $T$  is a function of a vector field and a form field that returns a function of place.

Now we consider a change of basis,  $e(f) = e'(f)J$  or

$$e_i(f) = \sum_j e'_j(f) J_i^j, \quad (C.4)$$

Where  $J$  typically depends on place. The corresponding dual basis transforms as

$$\tilde{e}^i(v) = \sum_j K_j^i \tilde{e}'^j(v), \quad (C.5)$$

where  $K = J^{-1}$  or  $\sum_j K_j^i(m) J_k^j(m) = \delta_k^i$ .

Because the tensor is multilinear over functions, we can deduce that the tensor components in the two bases are related by, in our example,

$$T_j^i = \sum_{kl} K_k^i T_l'^k J_j^l, \quad (C.6)$$

or

$$T_j'^i = \sum_{kl} J_k^i T_l^k K_j^l. \quad (C.7)$$

Tensors are a restricted set of mathematical objects that are geometric, so if we restrict our descriptions to tensor expressions they are *prima facie* independent of the coordinates used to represent them. So if we can represent the physical laws in terms of tensors we have built in the coordinate-system independence.

Let's test whether the geometric function  $R$ , which we have called the Riemann tensor (see equation (8.2)), is indeed a tensor field. A real-valued geometric function is a tensor if it is linear (over the functions) in each of its arguments. We can try it for 3-dimensional rectangular coordinates:

```
(let ((cs R3-rect))
  (let ((u (literal-vector-field 'u-coord cs))
        (v (literal-vector-field 'v-coord cs))
```

scheme

```

(w (literal-vector-field 'w-coord cs))
(x (literal-vector-field 'x-coord cs))
(omega (literal-1form-field 'omega-coord cs))
(nu (literal-1form-field 'nu-coord cs))
(f (literal-manifold-function 'f-coord cs))
(g (literal-manifold-function 'g-coord cs))
(nabla (covariant-derivative (literal-Cartan 'G cs)))
(m (typical-point cs)))
(let ((F (Riemann nabla)))
  ((up (- (F (+ (* f omega) (* g nu)) u v w)
            (+ (* f (F omega u v w)) (* g (F nu u v w))))
        (- (F omega (+ (* f u) (* g x)) v w)
            (+ (* f (F omega u v w)) (* g (F omega x v w))))
        (- (F omega v (+ (* f u) (* g x)) w)
            (+ (* f (F omega v u w)) (* g (F omega v x w))))
        (- (F omega v w (+ (* f u) (* g x))
            (+ (* f (F omega v w u)) (* g (F omega v w x))))
        m))))
;; (up 0 0 0 0)

```

Now that we are convinced that the Riemann tensor is indeed a tensor, we know how its components change under a change of basis. Let

$$R_{jkl}^i = R(\tilde{e}^i, e_j, e_k, e_l), \quad (\text{C.8})$$

then

$$R_{jkl}^i = \sum_{mnpq} K_m^i R_{npq}^{\prime m} J_j^n J_k^p J_l^q, \quad (\text{C.9})$$

or

$$R_{jkl}^{\prime i} = \sum_{mnpq} J_m^i R_{npq}^m K_j^n K_k^p K_l^q. \quad (\text{C.10})$$

Whew!

It is easy to generalize these formulas to tensors with general arguments. We have formulated the general tensor test as a program `tensor-test` that takes the procedure `T` to be tested, a list of argument types, and a coordinate system to be used. It tests each argument for linearity (over functions). If the function passed as `T` is a tensor, the result will be a list of zeros.

```

(tensor-test
 (Riemann (covariant-derivative (literal-Cartan 'G R3-rect)))
 '(1form vector vector vector)
 R3-rect)
;; (0 0 0 0)

```

scheme

and so does the torsion (see equation (8.21)):

```

(tensor-test
 (torsion (covariant-derivative (literal-Cartan 'G R3-rect)))
 '(1form vector vector)
 R3-rect)
;; (up 0 0 0 0)

```

scheme

But not all geometric functions are tensors. The covariant derivative is an interesting and important case. The function  $F$ , defined by

$$F(\omega, u, v) = \omega(\nabla_u v), \quad (\text{C.11})$$

is a geometric object, since the result is independent of the coordinate system used to represent the  $\nabla$ . For example:

```

(define ((F nabla) omega u v)
  (omega ((nabla u) v)))
(((- (F (covariant-derivative
  (Christoffel→Cartan
    (metric→Christoffel-2
      (coordinate-system→metric S2-spherical)
      (coordinate-system→basis S2-spherical))))))
  (F (covariant-derivative
    (Christoffel→Cartan
      (metric→Christoffel-2
        (coordinate-system→metric S2-stereographic)
        (coordinate-system→basis S2-stereographic))))))
  (literal-1form-field 'omega S2-spherical)
  (literal-vector-field 'u S2-spherical)
  (literal-vector-field 'v S2-spherical))
  ((point S2-spherical) (up 'theta 'phi)))
;; 0

```

scheme

But it is not a tensor field:

```

(tensor-test
  (F (covariant-derivative (literal-Cartan 'G R3-rect)))
  '(1form vector vector)
  R3-rect)
;; (0 0 MESS)

```

scheme

This result tells us that the function  $F$  is linear in its first two arguments but not in its third argument.

That the covariant derivative is not linear over functions in the second vector argument is easy to understand. The first vector argument takes derivatives of the coefficients of the second vector argument, so multiplying these coefficients by a manifold function changes the derivative.

## D ClojureScript

*Editorial note: This appendix is derived from Appendix A; its examples have been translated to ClojureScript using Emmy, while the original wording has been retained wherever possible.*

*ClojureScript is a dialect of Clojure that compiles to JavaScript and retains Clojure’s emphasis on immutable data and functional programming. Emmy supplies the generic arithmetic and symbolic mathematics used by the examples in this edition.*

Here we give an elementary introduction to the ClojureScript forms used in this book. For a fuller introduction, see the ClojureScript language documentation.

ClojureScript is a programming language based on expressions. An expression names a value. For example, the numeral `3.14` names an approximation to a familiar number. There are primitive expressions, such as a numeral, that we directly recognize, and there are compound expressions of several kinds.

### D.1 Function Calls

A *function call* is a kind of compound expression. A function call is a sequence of expressions delimited by parentheses. The first subexpression in a function call is taken to name a function, and the rest of the subexpressions are taken to name the arguments to that function. The value produced by the function when applied to the given arguments is the value named by the function call. For example,

```
(+ 1 2.14)                                     clojure
;; => 3.14

(+ 1 (* 2 1.07))
;; => 3.14
```

are both compound expressions that name the same number as the numeral `3.14`.<sup>1</sup> In these cases the symbols `+` and `*` name functions that add and multiply, respectively. If we replace any subexpression of any expression with an expression that names the same thing as the original subexpression, the thing named by the overall expression remains unchanged. In general, a function call is written

```
(operator operand-1 ... operand-n)             clojure
```

where *operator* names a function and *operand-*i** names the *i*th argument.<sup>2</sup>

### D.2 Function Expressions

Just as we use numerals to name numbers, we use `fn` expressions to name functions.<sup>3</sup> For example, the function that squares its input can be written:

```
(fn [x] (* x x))                               clojure
;; => <function>
```

This expression can be read: “The function of one argument, *x*, that multiplies *x* by *x*.” Of course, we can use this expression in any context where a function is needed. For example,

```
((fn [x] (* x x)) 4)                           clojure
;; => 16
```

The general form of a `fn` expression is

<sup>1</sup>In examples we show the value that would be printed by the ClojureScript system using slanted characters following the input expression.

<sup>2</sup>In ClojureScript every parenthesis is essential: you cannot add extra parentheses or remove any.

<sup>3</sup>The logician Alonzo Church [5] invented  $\lambda$ -notation to allow the specification of an anonymous function of a named parameter:  $\lambda x[\text{expression in } x]$ . This is read, “That function of one argument that is obtained by substituting the argument for *x* in the indicated expression.”

```
(fn [& formal-parameters] body)
```

clojure

where *formal-parameters* is a vector of symbols naming the function arguments and *body* is an expression that may refer to them. For a variadic function the parameter vector contains `&` before the name that receives the remaining arguments. The value of a function call is the value of the body of the function with the arguments substituted for the formal parameters.

## D.3 Definitions

We can use the `def` construct to give a name to any object. For example, if we make the definitions<sup>4</sup>

```
(def pi 3.141592653589793)
```

clojure

```
(def square (fn [x] (* x x)))
```

we can then use the symbols `pi` and `square` wherever the numeral or the `fn` expression could appear. For example, the area of the surface of a sphere of radius 5 meters is

```
(* 4 pi (square 5))  
;; => 314.1592653589793
```

clojure

### D.3.1 Names, functions, and local bindings

ClojureScript uses several related forms where Scheme uses `define`, `lambda`, and local definitions. Their different scopes are important:

- `def` gives a name to a value in the current namespace. The value may be a number, a function, or any other object. Thus `(def pi 3.14159)` defines a global name.
- `fn` constructs a function value. Its parameters are written in a vector: `(fn [x] (* x x))`. An `fn` may be anonymous, or it may be stored in a name using `def`.
- `defn` is convenient syntax for a `def` whose value is an `fn`. For example, `(defn square [x] (* x x))` is essentially `(def square (fn [x] (* x x)))`. Use `defn` for a named, namespace-level function and `def` for other namespace-level values.
- `let` introduces local names and values. Its binding vector alternates names and expressions: `(let [x 3 y 4] (sqrt (+ (square x) (square y))))`. These names exist only in the body of the `let`.
- `letfn` introduces local function names. Unlike ordinary `let` bindings, the functions may refer to themselves and to one another, so `letfn` is the direct translation for recursive or mutually dependent internal Scheme function definitions.

The names introduced by `def` and `defn` persist in the namespace. The names introduced by `let`, `letfn`, and function parameter vectors are lexical and disappear outside their bodies.

Function definitions may be expressed more conveniently using “syntactic sugar.” The squaring function may be defined

```
(defn square [x] (* x x))
```

clojure

which we may read: “To square  $x$  multiply  $x$  by  $x$ .”

In ClojureScript, functions may be passed as arguments and returned as values. For example, it is possible to make a function that implements the mathematical notion of the composition of two functions:<sup>5</sup>

```
(def compose (fn [f g] (fn [x] (f (g x)))))
```

clojure

<sup>4</sup>The definition of `square` given here is not the definition of `square` in the Emmy system. In Emmy, `square` is extended for tuples to mean the sum of the squares of the components of the tuple. However, for arguments that are not tuples the Emmy `square` does multiply the argument by itself.

<sup>5</sup>The examples are indented to help with readability. ClojureScript does not care about extra white space, so we may add as much as we please to make things easier to read.

```
((compose square sin) 2)
;; => 0.826821810431806

(square (sin 2))
;; => 0.826821810431806
```

Using the syntactic sugar shown above, we can write the definition more conveniently. The following are both equivalent to the definition above:

```
(defn compose [f g] (fn [x] (f (g x))))           clojure

(defn compose [f g] (fn [x] (f (g x))))
```

## D.4 Conditionals

Conditional expressions may be used to choose among several expressions to produce a value. For example, a function that implements the absolute value function may be written:

```
(defn abs                                     clojure
  [x]
  (cond (< x 0) (- x)
        (= x 0) x
        (> x 0) x))
```

The conditional `cond` takes a number of clauses. Each clause has a predicate expression, which may be either true or false, and a consequent expression. The value of the `cond` expression is the value of the consequent expression of the first clause for which the corresponding predicate expression is true. The general form of a conditional expression is

```
(cond predicate-1 consequent-1
      ... predicate-n
      consequent-n)                                clojure
```

For convenience the keyword `:else` can be used as the predicate in the last clause of a `cond`. ClojureScript clauses are written as alternating predicate and consequent expressions, without an extra pair of parentheses around each clause. The `if` construct provides another way to make a conditional when there is only a binary choice to be made. For example, because we have to do something special only when the argument is negative, we could have defined `abs` as:

```
(defn abs [x] (if (< x 0) (- x) x))                clojure
```

The general form of an `if` expression is

```
(if predicate consequent alternative)              clojure
```

If the *predicate* is true the value of the `if` expression is the value of the *consequent*, otherwise it is the value of the *alternative*.

## D.5 Recursive Functions

Given conditionals and definitions, we can write recursive functions. For example, to compute the *n*th factorial number we may write:

```
(defn factorial [n] (if (= n 0) 1 (* (bigint n) (factorial (- n 1)))))  clojure

(factorial 6)
;; => #emmy/bigint 720

;; scmutils simplified this result automatically; Emmy requires an explicit call.
```

```
(simplify (factorial 40))
;; => #emmy/bigint "8159152832478977343456112695961158942720000000000"
```

## D.6 Local Names

The `let` expression is used to give names to objects in a local context. For example,

```
(defn f
  [radius]
  ;; Keep pi symbolic so the common factor cancels before numerical evaluation.
  (let [pi 'pi area (* 4 pi (square radius)) volume (* (/ 4 3) pi (cube radius))] (/
    volume area)))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (f 3))
;; => 1
```

The general form of a `let` expression is

```
(let [variable-1 expression-1 ... variable-n expression-n] body)
```

The value of the `let` expression is the value of the *body* expression in the context where the variables *variable-*i** have the values of the expressions *expression-*i**. The expressions *expression-*i** may not refer to any of the variables.

A slight variant of the `let` expression provides a convenient way to express looping constructs. We can write a function that implements an alternative algorithm for computing factorials as follows:

```
(defn factorial
  [n]
  (letfn [(factlp [count answer] (if (> count n) answer (factlp (+ count 1) (* count
    answer))))]
    (factlp 1 1)))

(factorial 6)
;; => 720
```

Here, the symbol `factlp` following the `let` is locally defined to be a function that has the variables `count` and `answer` as its formal parameters. It is called the first time with the expressions `1` and `1`, initializing the loop. Whenever the function named `factlp` is called later, these variables get new values that are the values of the operand expressions `(+ count 1)` and `(* count answer)`.

## D.7 Compound Data — Lists and Vectors

Data can be glued together to form persistent compound values. ClojureScript lists and other sequences support sequential traversal; vectors support efficient indexed lookup with `nth`. Both are immutable values, and operations that add or replace elements produce new collections. The mathematical `up` and `down` values used throughout this book are Emmy structures, not ordinary ClojureScript vectors: use Emmy's generic arithmetic and `ref` when working with their components.

The function `list` is the constructor for lists. The selector `nth` gets an element of the list. All selectors in ClojureScript are zero-based. For example,

```
(def a-list (list 6 946 8 356 12 620))

a-list
;; => (6 946 8 356 12 620)

(nth a-list 3)
;; => 356
```

```
(nth a-list 0)
;; => 6
```

ClojureScript lists are persistent sequential collections. The functions `first` and `rest` select the first element and the remaining sequence. Thus,

```
(first a-list)
;; => 6

(rest a-list)
;; => (946 8 356 12 620)

(first (rest a-list))
;; => 946

(def another-list (cons 32 (rest a-list)))

another-list
;; => (32 946 8 356 12 620)

(first (rest another-list))
;; => 946
```

Both `a-list` and `another-list` share the same tail (their `rest`).

There is a predicate `pair?` that is true of pairs and false on all other types of data.

Vectors provide convenient literals and indexed access. There is a constructor `vector` that can be used to make vectors and a selector `nth` for accessing the elements of a vector:

```
(def a-vector (vector 37 63 49 21 88 56))

a-vector
;; => [37 63 49 21 88 56]

(nth a-vector 3)
;; => 21

(nth a-vector 0)
;; => 37
```

Notice that a vector is distinguished from a list on printout by the character `#` appearing before the initial parenthesis.

There is a predicate `vector?` that is true of vectors and false for all other types of data.

The elements of lists and vectors may be any kind of data, including numbers, functions, lists, and vectors. Numerous other functions for manipulating list-structured data and vector-structured data can be found in the ClojureScript documentation.

## D.8 Symbols

Symbols are a very important kind of primitive data type that we use to make programs and algebraic expressions. You probably have noticed that ClojureScript programs look just like lists. In fact, they are lists. Some of the elements of the lists that make up programs are symbols, such as `+` and `vector`.<sup>6</sup> If we are to make programs that can manipulate programs, we need to be able to write an expression that names such a symbol. This is accomplished by the mechanism of *quotation*. The name of the symbol `+` is the expression `'+`, and in general the name of an expression is the expression preceded by a single quote character. Thus the name of the expression `(+ 3 a)` is `'(+ 3 a)`.

---

<sup>6</sup>Symbols may have any number of characters. A symbol may not contain whitespace or a delimiter character, such as parentheses, brackets, quotation marks, comma, or `#`.



We can test if two symbols are identical by using the predicate `=`. For example, we can write a program to determine if an expression is a sum:

```
(defn sum? [expression] (and (pair? expression) (= (first expression) '+)))clojure

(sum? '(+ 3 a))
;; => true

(sum? '(* 3 a))
;; => false
```

Here `true` and `false` are the printed representations of the boolean values `true` and `false`.

Consider what would happen if we were to leave out the quote in the expression `(sum? '(+ 3 a))`. If the variable `a` had the value 4 we would be asking if 7 is a sum. But what we wanted to know was whether the expression `(+ 3 a)` is a sum. That is why we need the quote.

## D.9 Scheme–ClojureScript Cheat Sheet

The following correspondences cover the language forms used most often in this book. They describe syntax; Emmy supplies the mathematical operations used by the examples.

| Form                       | Scheme                                                            | ClojureScript                                                                        |
|----------------------------|-------------------------------------------------------------------|--------------------------------------------------------------------------------------|
| Global value               | <code>(define x value)</code>                                     | <code>(def x value)</code>                                                           |
| Global function            | <code>(define (f x) body)</code>                                  | <code>(defn f [x] body)</code>                                                       |
| Anonymous function         | <code>(lambda (x) body)</code>                                    | <code>(fn [x] body)</code>                                                           |
| Function stored as a value | <code>(define f (lambda (x) body))</code>                         | <code>(def f (fn [x] body))</code>                                                   |
| Local values               | <code>(let ((x a) (y b)) body)</code>                             | <code>(let [x a y b] body)</code>                                                    |
| Sequential local values    | <code>(let* ((x 2) (y (+ x 1))) y)</code>                         | <code>(let [x 2 y (+ x 1)] y)</code>                                                 |
| Local functions            | <code>(let () (define (f x) ...) (f 3))</code>                    | <code>(letfn [(f [x] ...)] (f 3))</code>                                             |
| Named local loop           | <code>(let loop ((x 3)) ... (loop ...))</code>                    | <code>(loop [x 3] ... (recur ...))</code>                                            |
| Expression sequence        | <code>(begin (display x) x)</code>                                | <code>(do (println x) x)</code>                                                      |
| Conditional                | <code>(if p a b) (cond (p a) (else b))</code>                     | <code>(if p a b) (cond p a :else b)</code>                                           |
| Boolean literals           | <code>#t</code> and <code>#f</code>                               | <code>true</code> and <code>false</code>                                             |
| Falsey values              | Only <code>#f</code> , as in <code>(if '() 'yes 'no) → yes</code> | <code>false</code> and <code>nil</code> , as in <code>(if nil :yes :no) → :no</code> |
| Exact ratio                | <code>1/2</code>                                                  | <code>(/ 1 2)</code> do not replace exact arithmetic with 0.5                        |
| Vector literal             | <code>#[a b c]</code>                                             | <code>[a b c]</code>                                                                 |
| Empty list                 | <code>'()</code>                                                  | <code>'()</code> or <code>(list)</code>                                              |
| Equality in these examples | <code>(eq? a b)</code>                                            | <code>(= a b)</code>                                                                 |
| Indexed selection          | <code>(list-ref xs 1)</code> or <code>(vector-ref xs 1)</code>    | <code>(nth xs 1)</code> use <code>(ref s 1)</code> for Emmy structures               |
| Sequence selectors         | <code>(car xs)</code> and <code>(cdr xs)</code>                   | <code>(first xs)</code> and <code>(rest xs)</code>                                   |
| Variadic parameters        | <code>(lambda args body)</code> or a dotted parameter list        | <code>(fn [&amp; args] body)</code>                                                  |
| Quotation                  | <code>'(a b)</code>                                               | <code>'(a b)</code> keywords such as <code>:else</code> evaluate to themselves       |
| Function application       | <code>(f x y)</code>                                              | <code>(f x y)</code> parameter and binding lists use vectors                         |

## E Our Notation in Emmy

*Editorial note: This appendix is derived from Appendix B; its examples have been translated to ClojureScript using Emmy, while the original wording has been retained wherever possible.*

*An adequate notation should be understood by at least two people, one of whom may be the author.*

*Abdus Salam (1950).*

We adopt a *functional mathematical notation* that is close to that used by Spivak in his *Calculus on Manifolds* [17]. The use of functional notation avoids many of the ambiguities of traditional mathematical notation that can impede clear reasoning. Functional notation carefully distinguishes the function from the value of the function when applied to particular arguments. In functional notation mathematical expressions are unambiguous and self-contained.

We adopt a *generic arithmetic* in which the basic arithmetic operations, such as addition and multiplication, are extended to a wide variety of mathematical types. Thus, for example, the addition operator  $+$  can be applied to numbers, tuples of numbers, matrices, functions, etc. Generic arithmetic formalizes the common informal practice used to manipulate mathematical objects.

We often want to manipulate aggregate quantities, such as the collection of all of the rectangular coordinates of a collection of particles, without explicitly manipulating the component parts. Tensor arithmetic provides a traditional way of manipulating aggregate objects: Indices label the parts; conventions, such as the summation convention, are introduced to manipulate the indices. We introduce a *tuple arithmetic* as an alternative way of manipulating aggregate quantities that usually lets us avoid labeling the parts with indices. Tuple arithmetic is inspired by tensor arithmetic but it is more general: not all of the components of a tuple need to be of the same size or type.

The mathematical notation is in one-to-one correspondence with expressions of ClojureScript with the Emmy computer algebra library. ClojureScript is a Lisp and is based on the  $\lambda$ -calculus

### E.1 Functions

The expression  $f(x)$  denotes the value of the function  $f$  at the given argument  $x$ ; when we wish to denote the function we write just  $f$ . Functions may take several arguments. For example, we may have the function that gives the Euclidean distance between two points in the plane given by their rectangular coordinates:

$$d(x_1, y_1, x_2, y_2) = \sqrt{(x_2 - x_1)^2 + (y_2 - y_1)^2}. \quad (\text{E.1})$$

In ClojureScript we can write this as:

```
(defn d [x1 y1 x2 y2] (sqrt (+ (square (- x2 x1)) (square (- y2 y1)))))
```

clojure

Functions may be composed if the range of one overlaps the domain of the other. The composition of functions is constructed by passing the output of one to the input of the other. We write the composition of two functions using the  $\circ$  operator:

$$(f \circ g) : x \mapsto (f \circ g)(x) = f(g(x)). \quad (\text{E.2})$$

A function  $h$  that computes the cube of the sine of its argument may be defined by composing the functions `cube` and `sin`:

```
(def h (compose cube sin))

(h 2)
;; => 0.7518269446689928
```

clojure

Which is the same as

```
(cube (sin 2))
;; => 0.7518269446689928
```

closure

Arithmetic is extended to the manipulation of functions: the usual mathematical operations may be applied to functions. Examples are addition and multiplication; we may add or multiply two functions if they take the same kinds of arguments and if their values can be added or multiplied:

$$(f + g)(x) = f(x) + g(x), (fg)(x) = f(x)g(x). \quad (\text{E.3})$$

A function `g` that multiplies the cube of its argument by the sine of its argument is

```
(def g (* cube sin))

(g 2)
;; => 7.274379414605454

(* (cube 2) (sin 2))
;; => 7.274379414605454
```

closure

## E.2 Symbolic Values

As in usual mathematical notation, arithmetic is extended to allow the use of symbols that represent unknown or incompletely specified mathematical objects. These symbols are manipulated as if they had values of a known type. By default, a ClojureScript symbol is assumed to represent a real number. So the expression `'a` is a literal ClojureScript symbol that represents an unspecified real number:

```
((compose cube sin) 'a)
;; => (expt (sin a) 3)
```

closure

The cached runner output records the value returned by each example. Emmy simplifies many symbolic results during generic arithmetic, and examples may call `simplify` explicitly when normalization is required.<sup>1</sup> We can use the simplifier to verify a trigonometric identity:

```
(simplify ((- (+ (square sin) (square cos)) 1) 'a))
;; => 0
```

closure

Just as it is useful to be able to manipulate symbolic numbers, it is useful to be able to manipulate symbolic functions. The function `literal-function` constructs a symbolic function having no properties other than its name and declared signature. By default, a literal function is defined to take one real argument and produce one real value. For example, we may want to work with a function  $f : \mathbf{R} \rightarrow \mathbf{R}$ :

```
((literal-function 'f) 'x)
;; => (f x)

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((compose (literal-function 'f) (literal-function 'g)) 'x))
;; => (f (g x))
```

closure

We can also make literal functions of multiple, possibly structured arguments that return structured values. For example, to denote a literal function named `g` that takes two real arguments and returns a real value ( $g : \mathbf{R} \times \mathbf{R} \rightarrow \mathbf{R}$ ) we may write:

```
(def g (literal-function 'g '(→ (X Real Real) Real)))

(g 'x 'y)
;; => (g x y)
```

closure

<sup>1</sup>Appendix G explains result capture, cached output, and explicit checks. The runner does not rely on scmutils' interactive `print-expression` printer.

We may use such a literal function anywhere that an explicit function of the same type may be used.

There is a whole language for describing the type of a literal function in terms of the number of arguments, the types of the arguments, and the types of the values. Here we describe a function that maps pairs of real numbers to real numbers with the expression `(→ (X Real Real) Real)`. Later we will introduce structured arguments and values and show extensions of literal functions to handle these.

### E.3 Tuples

There are two kinds of tuples: *up* tuples and *down* tuples. We write tuples as ordered lists of their components; a tuple is delimited by parentheses if it is an up tuple and by square brackets if it is a down tuple. For example, the up tuple  $v$  of velocity components  $v^0$ ,  $v^1$ , and  $v^2$  is

$$v = (v^0, v^1, v^2). \quad (\text{E.4})$$

The down tuple  $p$  of momentum components  $p_0$ ,  $p_1$ , and  $p_2$  is

$$p = [p_0, p_1, p_2]. \quad (\text{E.5})$$

A component of an up tuple is usually identified by a superscript. A component of a down tuple is usually identified by a subscript. We use zero-based indexing when referring to tuple elements. This notation follows the usual convention in tensor arithmetic.

We make tuples with the constructors `up` and `down`:

```
(def v (up 'v↑0 'v↑1 'v↑2)) closure

v
;; ⇒ (up v↑0 v↑1 v↑2)

(def p (down 'p_0 'p_1 'p_2))

p
;; ⇒ (down p_0 p_1 p_2)
```

Note that `v↑0` and `p_2` are just symbols. The caret and underline characters are symbol constituents, so there is no meaning other than mnemonic to the structure of these symbols. However, our software can also display expressions using  $\text{\TeX}$ , and then these decorations turn into superscripts and subscripts.

Tuple arithmetic is different from the usual tensor arithmetic in that the components of a tuple may also be tuples and different components need not have the same structure. For example, a tuple structure  $s$  of phase-space states is

$$s = (t, (x, y), [p_x, p_y]). \quad (\text{E.6})$$

It is an up tuple of the time, the coordinates, and the momenta. The time  $t$  has no substructure. The coordinates are an up tuple of the coordinate components  $x$  and  $y$ . The momentum is a down tuple of the momentum components  $p_x$  and  $p_y$ . In ClojureScript this is written:

```
(def s (up 't (up 'x 'y) (down 'p_x 'p_y))) closure
```

In order to reference components of tuple structures there are selector functions, for example:

$$\begin{aligned} I(s) &= s \\ I_0(s) &= t \\ I_1(s) &= (x, y) \\ I_2(s) &= [p_x, p_y] \\ I_{1,0}(s) &= x \\ &\dots \\ I_{2,1}(s) &= p_y. \end{aligned} \quad (\text{E.7})$$

The sequence of integer subscripts on the selector describes the access chain to the desired component.

The function `component` is the general selector that implements the selector function  $I_z$ :

```
((component 0 1) (up (up 'a 'b) (up 'c 'd)))
```

closure

```
;; => b
```

To access a component of a tuple we may also use the selector function `ref`, which takes a tuple and an index and returns the indicated element of the tuple:

```
(ref (up 'a 'b 'c) 1)
```

closure

```
;; => b
```

We use zero-based indexing everywhere. The function `ref` can be used to access any substructure of a tree of tuples:

```
(ref (up (up 'a 'b) (up 'c 'd)) 0 1)
```

closure

```
;; => b
```

Two up tuples of the same length may be added or subtracted, elementwise, to produce an up tuple, if the components are compatible for addition. Similarly, two down tuples of the same length may be added or subtracted, elementwise, to produce a down tuple, if the components are compatible for addition.

Any tuple may be multiplied by a number by multiplying each component by the number. Numbers may, of course, be multiplied. Tuples that are compatible for addition form a vector space.

For convenience we define the square of a tuple to be the sum of the squares of the components of the tuple. Tuples can be multiplied, as described below, but the square of a tuple is not the product of the tuple with itself.

The meaning of multiplication of tuples depends on the structure of the tuples. Two tuples are compatible for contraction if they are of opposite types, they are of the same length, and corresponding elements have the following property: either they are both tuples and are compatible for contraction, or at least one is not a tuple. If two tuples are compatible for contraction then generic multiplication is interpreted as contraction: the result is the sum of the products of corresponding components of the tuples. For example,  $p$  and  $v$  introduced in equations (E.4) and (E.5) above are compatible for contraction; the product is

$$pv = p_0v^0 + p_1v^1 + p_2v^2. \quad (\text{E.8})$$

So the product of tuples that are compatible for contraction is an inner product. Using the tuples  $p$  and  $v$  defined above gives us

```
(* p v)
```

closure

```
;; => (+ (* p_0 v^0) (* p_1 v^1) (* p_2 v^2))
```

Contraction of tuples is commutative:  $pv = vp$ . Caution: Multiplication of tuples that are compatible for contraction is, in general, not associative. For example, let  $u = (5, 2)$ ,  $v = (11, 13)$ , and  $g = [[3, 5], [7, 9]]$ . Then  $u(gv) = 964$ , but  $(ug)v = 878$ . The expression  $ugv$  is ambiguous. An expression that has this ambiguity does not occur in this book.

The rule for multiplying two structures that are not compatible for contraction is simple. If  $A$  and  $B$  are not compatible for contraction, the product  $AB$  is a tuple of type  $B$  whose components are the products of  $A$  and the components of  $B$ . The same rule is applied recursively in multiplying the components. So if  $B = (B^0, B^1, B^2)$ , the product of  $A$  and  $B$  is

$$AB = (AB^0, AB^1, AB^2). \quad (\text{E.9})$$

If  $A$  and  $C$  are not compatible for contraction and  $C = [C_0, C_1, C_2]$ , the product is

$$AC = [AC_0, AC_1, AC_2]. \quad (\text{E.10})$$

Tuple structures can be made to represent linear transformations. For example, the rotation commonly represented by the matrix

$$\begin{bmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{bmatrix} \quad (\text{E.11})$$

can be represented as a tuple structure:<sup>2</sup>

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right]. \quad (\text{E.12})$$

Such a tuple is compatible for contraction with an up tuple that represents a vector. So, for example:

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right] \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} x \cos \theta - y \sin \theta \\ x \sin \theta + y \cos \theta \end{pmatrix}. \quad (\text{E.13})$$

The product of two tuples that represent linear transformations – which are not compatible for contraction – represents the composition of the linear transformations. For example, the product of the tuples representing two rotations is

$$\left[ \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix} \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix} \right] \left[ \begin{pmatrix} \cos \varphi \\ \sin \varphi \end{pmatrix} \begin{pmatrix} -\sin \varphi \\ \cos \varphi \end{pmatrix} \right] = \left[ \begin{pmatrix} \cos(\theta + \varphi) \\ \sin(\theta + \varphi) \end{pmatrix} \begin{pmatrix} -\sin(\theta + \varphi) \\ \cos(\theta + \varphi) \end{pmatrix} \right]. \quad (\text{E.14})$$

Multiplication of tuples that represent linear transformations is associative but generally not commutative, just as the composition of the transformations is associative but not generally commutative.

## E.4 Derivatives

The derivative of a function  $f$  is a function, denoted by  $Df$ . Our notational convention is that  $D$  is a high-precedence operator. Thus  $D$  operates on the adjacent function before any other application occurs:  $Df(x)$  is the same as  $(Df)(x)$ . Higher-order derivatives are described by exponentiating the derivative operator. Thus the  $n$ th derivative of a function  $f$  is notated as  $D^n f$ .

The ClojureScript function for producing the derivative of a function is named `D`. The derivative of the `sin` function is a function that computes `cos`:

```
(def derivative-of-sine (D sin)) clojure

(derivative-of-sine 'x)
;; => (cos x)
```

The derivative of a function  $f$  is the function  $Df$  whose value for a particular argument is something that can be multiplied by an increment  $\Delta x$  in the argument to get a linear approximation to the increment in the value of  $f$ :

$$f(x + \Delta x) \approx f(x) + Df(x)\Delta x. \quad (\text{E.15})$$

For example, let  $f$  be the function that cubes its argument ( $f(x) = x^3$ ); then  $Df$  is the function that yields three times the square of its argument ( $Df(y) = 3y^2$ ). So  $f(5) = 125$  and  $Df(5) = 75$ . The value of  $f$  with argument  $x + \Delta x$  is

$$f(x + \Delta x) = (x + \Delta x)^3 = x^3 + 3x^2\Delta x + 3x\Delta x^2 + \Delta x^3 \quad (\text{E.16})$$

and

$$Df(x)\Delta x = 3x^2\Delta x. \quad (\text{E.17})$$

So  $Df(x)$  multiplied by  $\Delta x$  gives us the term in  $f(x + \Delta x)$  that is linear in  $\Delta x$ , providing a good approximation to  $f(x + \Delta x) - f(x)$  when  $\Delta x$  is small.

Derivatives of compositions obey the chain rule:

---

<sup>2</sup>To emphasize the relationship of simple tuple structures to matrix notation we often format up tuples as vertical arrangements of components and down tuples as horizontal arrangements of components. However, we could just as well have written this tuple as  $[(\cos \theta, \sin \theta), (-\sin \theta, \cos \theta)]$ .

$$D(f \circ g) = ((Df) \circ g) \cdot Dg. \quad (\text{E.18})$$

So at  $x$ ,

$$(D(f \circ g))(x) = Df(g(x)) \cdot Dg(x). \quad (\text{E.19})$$

$D$  is an example of an operator. An operator is like a function except that multiplication of operators is interpreted as composition, whereas multiplication of functions is multiplication of the values (see equation (E.3)). If  $D$  were an ordinary function, then the rule for multiplication would imply that  $D^2 f$  would just be the product of  $Df$  with itself, which is not what is intended. A product of a number and an operator scales the operator. So, for example

```
(((* 5 D) cos) 'x)
;; => (* 5 (- (sin x)))
```

clojure

Arithmetic is extended to allow manipulation of operators. A typical operator is  $(D + I)(D - I) = D^2 - I$ , where  $I$  is the identity operator, subtracts a function from its second derivative. Such an operator can be constructed and used in ClojureScript as follows:

```
(((* (+ D I) (- D I)) (literal-function 'f)) 'x)
;; => (+ (* -1 ((D f) x)) (((expt D 2) f) x) (- ((D f) x) (f x)))
```

clojure

## E.5 Derivatives of Functions of Multiple Arguments

The derivative generalizes to functions that take multiple arguments. The derivative of a real-valued function of multiple arguments is an object whose contraction with the tuple of increments in the arguments gives a linear approximation to the increment in the function's value.

A function of multiple arguments can be thought of as a function of an up tuple of those arguments. Thus an incremental argument tuple is an up tuple of components, one for each argument position. The derivative of such a function is a down tuple of the partial derivatives of the function with respect to each argument position

Suppose we have a real-valued function  $g$  of two real-valued arguments, and we want to approximate the increment in the value of  $g$  from its value at  $x, y$ . If the arguments are incremented by the tuple  $(\Delta x, \Delta y)$  we compute:

$$Dg(x, y) \cdot (\Delta x, \Delta y) = [\partial_0 g(x, y) + \partial_1 g(x, y)] \cdot (\Delta x, \Delta y) = \partial_0 g(x, y) \Delta x + \partial_1 g(x, y) \Delta y. \quad (\text{E.20})$$

Using the two-argument literal function `g` defined in Section E.2, page 141, we have:

```
((D g) 'x 'y)
;; => (down (((partial 0) g) x y) (((partial 1) g) x y))
```

clojure

In general, partial derivatives are just the components of the derivative of a function that takes multiple arguments (or structured arguments or both; see below). So a partial derivative of a function is a composition of a component selector and the derivative of that function.<sup>3</sup> Indeed:

$$\partial_0 g = I_0 \circ Dg, \quad (\text{E.21})$$

$$\partial_1 g = I_1 \circ Dg. \quad (\text{E.22})$$

Concretely, if

$$g(x, y) = x^3 y^5 \quad (\text{E.23})$$

then

$$Dg(x, y) = [3x^2 y^5, 5x^3 y^4] \quad (\text{E.24})$$

<sup>3</sup>Partial derivative operators such as `(partial 2)` are operators, so `(expt (partial 1) 2)` is a second partial derivative.



and the first-order approximation of the increment for changing the arguments by  $\Delta x$  and  $\Delta y$  is

$$g(x + \Delta x, y + \Delta y) - g(x, y) \approx [3x^2y^5, 5x^3y^4] \cdot (\Delta x, \Delta y) = 3x^2y^5\Delta x + 5x^3y^4\Delta y. \quad (\text{E.25})$$

Partial derivatives of compositions also obey a chain rule:

$$\partial_i(f \circ g) = ((Df) \circ g) \cdot \partial_i g. \quad (\text{E.26})$$

So if  $x$  is a tuple of arguments, then

$$(\partial_i(f \circ g))(x) = Df(g(x)) \cdot \partial_i g(x). \quad (\text{E.27})$$

Mathematical notation usually does not distinguish functions of multiple arguments and functions of the tuple of arguments. Let  $h((x, y)) = g(x, y)$ . The function  $h$ , which takes a tuple of arguments  $x$  and  $y$ , is not distinguished from the function  $g$  that takes arguments  $x$  and  $y$ . We use both ways of defining functions of multiple arguments. The derivatives of both kinds of functions are compatible for contraction with a tuple of increments to the arguments. ClojureScript and Emmy make the distinction explicit:

```
(defn h [s] (g (ref s 0) (ref s 1))) clojure

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify (h (up 'x 'y)))
;; => (g x y)

(simplify ((D g) 'x 'y))
;; => (down (((partial 0) g) x y) (((partial 1) g) x y))

(simplify ((D h) (up 'x 'y)))
;; => (down (((partial 0) g) x y) (((partial 1) g) x y))
```

A phase-space state function is a function of time, coordinates, and momenta. Let  $H$  be such a function. The value of  $H$  is  $H(t, (x, y), [p_x, p_y])$  for time  $t$ , coordinates  $(x, y)$ , and momenta  $[p_x, p_y]$ . Let  $s$  be the phase-space state tuple as in (E.6):

$$s = (t, (x, y), [p_x, p_y]). \quad (\text{E.28})$$

The value of  $H$  for argument tuple  $s$  is  $H(s)$ . We use both ways of writing the value of  $H$ .

We often show a function of multiple arguments that include tuples by indicating the boundaries of the argument tuples with semicolons and separating their components with commas. If  $H$  is a function of phase-space states with arguments  $t$ ,  $(x, y)$ , and  $[p_x, p_y]$ , we may write  $H(t; x, y; p_x, p_y)$ . This notation loses the up/down distinction, but our semicolon-and-comma notation is convenient and reasonably unambiguous.

The derivative of  $H$  is a function that produces an object that can be contracted with an increment in the argument structure to produce an increment in the function's value. The derivative is a down tuple of three partial derivatives. The first partial derivative is the partial derivative with respect to the numerical argument. The second partial derivative is a down tuple of partial derivatives with respect to each component of the up-tuple argument. The third partial derivative is an up tuple of partial derivatives with respect to each component of the down-tuple argument:

$$\begin{aligned} DH(s) &= [\partial_0 H(s), \partial_1 H(s), \partial_2 H(s)] \\ &= [\partial_0 H(s), [\partial_{1,0} H(s), \partial_{1,1} H(s)], \\ &\quad [\partial_{2,0} H(s), \partial_{2,1} H(s)]]], \end{aligned} \quad (\text{E.29})$$

where  $\partial_{1,0}$  indicates the partial derivative with respect to the first component (index 0) of the second argument (index 1) of the function, and so on. Indeed,  $\partial_z F = I_z \circ DF$  for any function  $F$  and access chain  $z$ . So, if we let  $\Delta s$  be an incremental phase-space state tuple,

$$\Delta s = (\Delta t, (\Delta x, \Delta y), [\Delta p_x, \Delta p_y]) \quad (\text{E.30})$$

then

$$DH(s)\Delta s = \partial_0 H(s)\Delta t + \partial_{1,0} H(s)\Delta x + \partial_{1,1} H(s)\Delta y + \partial_{2,0} H(s)\Delta p_x + \partial_{2,1} H(s)\Delta p_y. \quad (\text{E.31})$$

Caution: Partial derivative operators with respect to different structured arguments generally do not commute.

In ClojureScript we must make explicit choices. We usually assume that phase-space state functions are functions of the tuple. For example,

```
(def H (literal-function 'H '(→ (UP Real (UP Real Real) (DOWN Real Real)) Real)))  clojure

(H s)
;; ⇒ (H (up t (up x y) (down p_x p_y)))

;; scmutils simplified this result automatically; Emmy requires an explicit call.
(simplify ((D H) s))
;; ⇒ (down (((partial 0) H) (up t (up x y) (down p_x p_y)))
;;       (down (((partial 1 0) H) (up t (up x y) (down p_x p_y)))
;;       (((partial 1 1) H) (up t (up x y) (down p_x p_y)))
;;       (up (((partial 2 0) H) (up t (up x y) (down p_x p_y)))
;;       (((partial 2 1) H) (up t (up x y) (down p_x p_y)))))
```

## E.6 Structured Results

Some functions produce structured outputs. A function whose output is a tuple is equivalent to a tuple of component functions each of which produces one component of the output tuple.

For example, a function that takes one numerical argument and produces a structure of outputs may be used to describe a curve through space. The following function describes a helical path around the  $\hat{z}$ -axis in 3-dimensional space:

$$h(t) = (\cos t, \sin t, t) = (\cos, \sin, I)(t). \quad (\text{E.32})$$

The derivative is just the up tuple of the derivatives of each component of the function:

$$Dh(t) = (-\sin t, \cos t, 1). \quad (\text{E.33})$$

In ClojureScript we can write

```
(defn helix [t] (up (cos t) (sin t) t))  clojure
```

or just

```
(def helix (up cos sin identity))  clojure
```

Its derivative is just the up tuple of the derivatives of each component of the function:

```
((D helix) 't)
;; ⇒ (up (- (sin t)) (cos t) 1)  clojure
```

In general, a function that produces structured outputs is just treated as a structure of functions, one for each of the components. The derivative of a function of structured inputs that produces structured outputs is an object that when contracted with an incremental input structure produces a linear approximation to the incremental output. Thus, if we define function  $g$  by

$$g(x, y) = ((x + y)^2, (y - x)^3, e^{x+y}), \quad (\text{E.34})$$

then the derivative of  $g$  is

$$Dg(x, y) = \left[ \begin{pmatrix} 2(x + y) \\ -3(y - x)^2 \\ e^{x+y} \end{pmatrix}, \begin{pmatrix} 2(x + y) \\ 3(y - x)^2 \\ e^{x+y} \end{pmatrix} \right] \quad (\text{E.35})$$

In ClojureScript:

```
(defn g [x y] (up (square (+ x y)) (cube (- y x)) (exp (+ x y))))
;; ⇒ (down (up (* 2 (+ x y)) (* 3 (expt (- y x) 2) -1) (exp (+ x y))))
;;      (up (* 2 (+ x y)) (* 3 (expt (- y x) 2)) (exp (+ x y))))
```

### E.6.1 Exercise E.1: Chain Rule

Let  $F(x, y) = x^2y^3$ ,  $G(x, y) = (F(x, y), y)$ , and  $H(x, y) = F(F(x, y), y)$ , so that  $H = F \circ G$ .

a. Compute  $\partial_0 F(x, y)$  and  $\partial_1 F(x, y)$ . b. Compute  $\partial_0 F(F(x, y), y)$  and  $\partial_1 F(F(x, y), y)$ . c. Compute  $\partial_0 G(x, y)$  and  $\partial_1 G(x, y)$ . d. Compute  $DF(a, b)$ ,  $DG(3, 5)$  and  $DH(3a^2, 5b^3)$ .

### E.6.2 Exercise E.2: Computing Derivatives

We can represent functions of multiple arguments as ClojureScript functions in several ways, depending upon how we wish to use them. The simplest idea is to identify the ClojureScript function arguments with the function's arguments.

For example, we could write implementations of the functions that occur in exercise E.1 as follows:

```
(defn f [x y] (* (square x) (cube y)))
(defn g [x y] (up (f x y) y))
(defn h [x y] (f (f x y) y))
```

With this choice it is awkward to compose a function that takes multiple arguments, such as  $f$ , with a function that produces a tuple of those arguments, such as  $g$ . Alternatively, we can represent the function arguments as slots of a tuple data structure, and then composition with a function that produces such a data structure is easy. However, this choice requires the functions to build and take apart structures.

For example, we may define functions that implement the functions above as follows:

```
(defn f [v] (let [x (ref v 0) y (ref v 1)] (* (square x) (cube y))))
(defn g [v] (let [x (ref v 0) y (ref v 1)] (up (f v) y)))
(def h (compose f g))
```

Repeat exercise E.1 using the computer. Explore both implementations of multiple-argument functions.

## F Tensors in Emmy

*Editorial note: This appendix is derived from Appendix C; its examples have been translated to ClojureScript using Emmy, while the original wording has been retained wherever possible.*

There are a variety of objects that have meaning independent of any particular basis. Examples are form fields, vector fields, covariant derivative, and so on. We call objects that are independent of basis *geometric objects*. Some of these are functions that take other geometric objects, such as vector fields and form fields, as arguments and produce further geometric objects. We refer to such functions as *geometric functions*. We want the laws of physics to be independent of the coordinate systems. How we describe an experiment should not affect the result. If we use only geometric objects in our descriptions then this is automatic.

A geometric function of vector fields and form fields that is linear in each argument with functions as multipliers is called a *tensor*. For example, let  $T$  be a geometric function of a vector field and form field that gives a real-number result at the manifold point  $m$ . Then

$$T(fu + gv, \omega) = f T(u, \omega) + g T(v, \omega) \quad (\text{F.1})$$

$$T(u, f\omega + g\theta) = f T(u, \omega) + g T(u, \theta), \quad (\text{F.2})$$

where  $u$  and  $v$  are vector fields,  $\omega$  and  $\theta$  are form fields, and  $f$  and  $g$  are manifold functions. That a tensor is linear over functions and not just constants is important.

The multilinearity over functions implies that the components of the tensor transform in a particularly simple way as the basis is changed. The components of a real-valued geometric function of vector fields and form fields are obtained by evaluating the function on a set of basis vectors and their dual form basis. In our example,

$$T_j^i = T(e_j, \tilde{e}^i), \quad (\text{F.3})$$

for basis vector fields  $e_j$  and dual form fields  $\tilde{e}^i$ . On the left,  $T_j^i$  is a function of place (manifold point); on the right,  $T$  is a function of a vector field and a form field that returns a function of place.

Now we consider a change of basis,  $e(f) = e'(f)J$  or

$$e_i(f) = \sum_j e'_j(f) J_i^j, \quad (\text{F.4})$$

Where  $J$  typically depends on place. The corresponding dual basis transforms as

$$\tilde{e}^i(v) = \sum_j K_j^i \tilde{e}'^j(v), \quad (\text{F.5})$$

where  $K = J^{-1}$  or  $\sum_j K_j^i(m) J_k^j(m) = \delta_k^i$ .

Because the tensor is multilinear over functions, we can deduce that the tensor components in the two bases are related by, in our example,

$$T_j^i = \sum_{kl} K_k^i T_l'^k J_j^l, \quad (\text{F.6})$$

or

$$T_j'^i = \sum_{kl} J_k^i T_l^k K_j^l. \quad (\text{F.7})$$

Tensors are a restricted set of mathematical objects that are geometric, so if we restrict our descriptions to tensor expressions they are *prima facie* independent of the coordinates used to represent them. So if we can represent the physical laws in terms of tensors we have built in the coordinate-system independence.

Let's test whether the geometric function  $R$ , which we have called the Riemann tensor (see equation (8.2)), is indeed a tensor field. A real-valued geometric function is a tensor if it is linear (over the functions) in each of its arguments. We can try it for 3-dimensional rectangular coordinates:

```

;; Simplify each tensor-linearity component before assembling the result;
;; simplifying the combined expansion creates a multi-megabyte intermediate.
(clojure
(let [cs R3-rect
      u (literal-vector-field 'u-coord cs)
      v (literal-vector-field 'v-coord cs)
      w (literal-vector-field 'w-coord cs)
      x (literal-vector-field 'x-coord cs)
      omega (literal-oneform-field 'omega-coord cs)
      nu (literal-oneform-field 'nu-coord cs)
      f (literal-manifold-function 'f-coord cs)
      g (literal-manifold-function 'g-coord cs)
      nabla (covariant-derivative (literal-Cartan 'G cs))
      m (typical-point cs)
      F (Riemann nabla)]
  (mapr (fn [component] (freeze (simplify (component m))))
    (up (- (F (+ (* f omega) (* g nu)) u v w) (+ (* f (F omega u v w)) (* g (F nu
u v w))))
        (- (F omega (+ (* f u) (* g x)) v w) (+ (* f (F omega u v w)) (* g (F
omega x v w))))
        (- (F omega v (+ (* f u) (* g x)) w) (+ (* f (F omega v u w)) (* g (F
omega v x w))))
        (- (F omega v w (+ (* f u) (* g x)) (+ (* f (F omega v w u)) (* g (F
omega v w x))))))
    ;; => (up 0 0 0 0)

```

Now that we are convinced that the Riemann tensor is indeed a tensor, we know how its components change under a change of basis. Let

$$R_{jkl}^i = R(\tilde{e}^i, e_j, e_k, e_l), \quad (\text{F.8})$$

then

$$R_{jkl}^i = \sum_{mnpq} K_m^i R_{npq}^m J_j^n J_k^p J_l^q, \quad (\text{F.9})$$

or

$$R_{jkl}^i = \sum_{mnpq} J_m^i R_{npq}^m K_j^n K_k^p K_l^q. \quad (\text{F.10})$$

Whew!

It is easy to generalize these formulas to tensors with general arguments. We have formulated the general tensor test as a program `tensor-test` that takes the Emmy/ClojureScript function `T` to be tested, a list of argument types, and a coordinate system to be used. It tests each argument for linearity (over functions). If the function passed as `T` is a tensor, the result will be a list of zeros.

```

(tensor-test (Riemann (covariant-derivative (literal-Cartan 'G R3-rect)))
  '(oneform vector vector vector)
  R3-rect)
;; => [0 0 0 0]
(clojure

```

and so does the torsion (see equation (8.21)):

```

(tensor-test (torsion (covariant-derivative (literal-Cartan 'G R3-rect)))
  '(oneform vector vector)
  R3-rect)
;; => [0 0 0]
(clojure

```

But not all geometric functions are tensors. The covariant derivative is an interesting and important case. The function  $F$ , defined by

$$F(\omega, u, v) = \omega(\nabla_u v), \quad (\text{F.11})$$

is a geometric object, since the result is independent of the coordinate system used to represent the  $\nabla$ . For example:

```
(defn F [nabla] (fn [omega u v] (omega ((nabla u) v)))) closure

;; Emmy's generic symbolic chart conversion introduces inverse trigonometric
;; functions and square roots before simplifying them away. Verify the same
;; statement exactly, component by component. Everything below remains visible
;; in the book except the repeated 2-by-2 transformation contraction, which is
;; compiled so SCI does not wrap its differentiable function as a MetaFn.
;;
;; Exact compiled helper source used below:
;; (ns fdg.slow-checks
;;   (:refer-clojure :exclude [+ - * /])
;;   (:require [emmy.env :as e]))
;;
;; (defn- sum2 [f] (e/+ (f 0) (f 1)))
;;
;; (defn spherical→stereographic
;;   "Exact north-pole stereographic coordinates in a native Emmy function."
;;   [[theta phi]]
;;   (let [scale (e// (e/sin theta) (e/- 1 (e/cos theta)))]
;;     (e/up (e/* scale (e/cos phi))
;;           (e/* scale (e/sin phi)))))
;;
;; (defn- old-term
;;   [old-symbols J i b c]
;;   (e/simplify
;;     (sum2
;;       (fn [j]
;;         (e/simplify
;;           (sum2
;;             (fn [k]
;;               (e/simplify
;;                 (e/* (get-in old-symbols [j k i])
;;                     (get-in J [b j])
;;                     (get-in J [c k])))))))))
;;
;; (defn- transformed-coefficient
;;   [old-symbols J J-inverse dJ b c a]
;;   (e/simplify
;;     (sum2
;;       (fn [i]
;;         (e/simplify
;;           (e/* (get-in J-inverse [i a])
;;               (e/simplify
;;                 (e/+ (get-in dJ [b c i])
;;                     (old-term old-symbols J i b c)))))))
;;
;; (defn- transform-symbols
;;   [old-symbols q]
;;   (let [J (e/mapr e/simplify ((e/D spherical→stereographic) q))
;;         J-inverse (e/mapr e/simplify (e/invert J))
;;         dJ (e/mapr e/simplify ((e/D (e/D spherical→stereographic)) q))
;;         gamma (fn [b c a]
;;                   (transformed-coefficient
;;                     old-symbols J J-inverse dJ b c a))]
;;     (e/down
;;       (e/down
;;         (e/up (gamma 0 0 0) (gamma 0 0 1))
;;         (e/up (gamma 0 1 0) (gamma 0 1 1)))
;;       (e/down
;;         (e/up (gamma 1 0 0) (gamma 1 0 1))
;;         (e/up (gamma 1 1 0) (gamma 1 1 1))))))
```

```

;; (defn transform-stereographic-Christoffel-to-spherical
;;   "Apply the exact 2D Christoffel coordinate-transformation law, simplifying
;;   each contraction before assembling the next one."
;;   [stereographic-symbols spherical-coordinates]
;;   (transform-symbols stereographic-symbols spherical-coordinates))
(let [q (up 'theta 'phi)
      spherical-basis (coordinate-system→basis S2-spherical)
      stereographic-basis (coordinate-system→basis S2-stereographic)
      Gamma-for (fn [coordinate-system basis coordinates]
                   (let [metric (coordinate-system→metric coordinate-system)
                         christoffel (metric→Christoffel-2 metric basis)
                         symbols (Christoffel→symbols christoffel)]
                     (mapr simplify (symbols ((point coordinate-system)
                                              coordinates))))))
      Gamma-sphere (Gamma-for S2-spherical spherical-basis q)
      Gamma-stereo (fn [[x y]]
                     (let [d (+ 1 (square x) (square y))
                           minus-x (/ (* -2 x) d)
                           plus-x (/ (* 2 x) d)
                           minus-y (/ (* -2 y) d)
                           plus-y (/ (* 2 y) d)]
                         (down (down (up minus-x plus-y) (up minus-y minus-x))
                                (down (up minus-y minus-x) (up plus-x minus-y))))
                     Gamma-derived (Gamma-for S2-stereographic stereographic-basis (up 'x 'y))
                     residuals (fn [actual expected] (mapr (fn [a e] (freeze (simplify (- a e))))
                                                                actual expected))
                     derivation (residuals Gamma-derived (Gamma-stereo (up 'x 'y)))
                     scale (/ (sin 'theta) (- 1 (cos 'theta)))
                     xy (up (* scale (cos 'phi)) (* scale (sin 'phi)))
                     Gamma-transformed (transform-stereographic-Christoffel-to-spherical (Gamma-stereo xy) q)
                     transformation (residuals Gamma-transformed Gamma-sphere)]
                       (verified-zero 0 (up derivation transformation)))
      ;; ⇒ 0

```

But it is not a tensor field:

```

(tensor-test (F (covariant-derivative (literal-Cartan 'G R3-rect)))
              '(oneform vector vector)
              R3-rect)
;; ⇒ [0 0
;;      (+ (* (omega41535_0 (up x041872 x141873 x241874))
;;             (v41537↑0 (up x041872 x141873 x241874))
;;             ((partial 0) g41538) (up x041872 x141873 x241874))
;;          (v41536↑0 (up x041872 x141873 x241874)))
;;      (* (omega41535_0 (up x041872 x141873 x241874))
;;          (v41537↑0 (up x041872 x141873 x241874))
;;          ((partial 1) g41538) (up x041872 x141873 x241874))
;;          (v41536↑1 (up x041872 x141873 x241874)))
;;      (* (omega41535_0 (up x041872 x141873 x241874))
;;          (v41537↑0 (up x041872 x141873 x241874))
;;          ((partial 2) g41538) (up x041872 x141873 x241874))
;;          (v41536↑2 (up x041872 x141873 x241874)))
;;      (* (((partial 0) g41538) (up x041872 x141873 x241874))
;;          (v41536↑0 (up x041872 x141873 x241874))
;;          (v41537↑1 (up x041872 x141873 x241874))
;;          (omega41535_1 (up x041872 x141873 x241874))))
;;      (* (((partial 0) g41538) (up x041872 x141873 x241874))
;;          (v41536↑0 (up x041872 x141873 x241874))
;;          (v41537↑2 (up x041872 x141873 x241874))
;;          (omega41535_2 (up x041872 x141873 x241874))))
;;      (* (((partial 1) g41538) (up x041872 x141873 x241874))
;;          (v41536↑0 (up x041872 x141873 x241874))
;;          (v41537↑1 (up x041872 x141873 x241874))
;;          (v41536↑1 (up x041872 x141873 x241874))
;;          (v41537↑1 (up x041872 x141873 x241874)))

```

```

;;      (omega41535_1 (up x041872 x141873 x241874)))
;;      (* (((partial 1) g41538) (up x041872 x141873 x241874))
;;      (v41536↑1 (up x041872 x141873 x241874))
;;      (v41537↑2 (up x041872 x141873 x241874))
;;      (omega41535_2 (up x041872 x141873 x241874)))
;;      (* (((partial 2) g41538) (up x041872 x141873 x241874))
;;      (v41536↑2 (up x041872 x141873 x241874))
;;      (v41537↑1 (up x041872 x141873 x241874))
;;      (omega41535_1 (up x041872 x141873 x241874)))
;;      (* (((partial 2) g41538) (up x041872 x141873 x241874))
;;      (v41536↑2 (up x041872 x141873 x241874))
;;      (v41537↑2 (up x041872 x141873 x241874))
;;      (omega41535_2 (up x041872 x141873 x241874))))])

```

This result tells us that the function  $F$  is linear in its first two arguments but not in its third argument.

That the covariant derivative is not linear over functions in the second vector argument is easy to understand. The first vector argument takes derivatives of the coefficients of the second vector argument, so multiplying these coefficients by a manifold function changes the derivative.



## G Running the Emmy Examples

The ClojureScript listings in this edition are executable programs using [Emmy](#), a computer algebra and mathematical physics library in the `scmutils` tradition. This appendix describes the runner supplied with this book, the intended workflow for checking and modifying examples, and the smaller setup needed to run an example in a clean ClojureScript project.

### G.1 What the Book Runner Does

The runner has two related execution paths. The browser interface uses `emmy.sci` and the Small Clojure Interpreter (SCI) to evaluate source text interactively. The smoke runner is compiled by `shadow-cljs` as a Node.js program and uses the same SCI environment to check every runnable block without manual interaction. In both cases Emmy's public environment is installed in a session namespace together with the small adapters in `fdg.compat`.

The converter writes each example to a stable pair of files:

```
codeblocks/chapter006/003.scm      text
codeblocks/chapter006/003.cljs
```

The `.scm` file is the clean Scheme snapshot extracted from the generated book. The neighboring `.cljs` file is the ClojureScript/Emmy port. The files are the durable conversion state; the browser editor is a temporary scratch editor and does not save changes.

The converter also creates `emmy-runner/public/generated/blocks.json`. For each block the manifest records its chapter, order, heading, source location, definitions, top-level forms, and whether it is an executable example. The runner uses this information to reproduce the order in which a reader encounters definitions.

Examples are grouped into chapter sessions. A block may use names introduced by an earlier block in the same chapter, so evaluating an arbitrary listing in isolation is not generally equivalent to following the book. Each chapter starts in a fresh context. Before it runs a block, the runner handles definitions that intentionally replace names supplied by `emmy.env` and declares mutually dependent definitions from the same listing.

### G.2 Preparing and Starting the Runner

From the root of this repository, generate and check the ClojureScript blocks with

```
make emmy-blocks      sh
```

This converts the blocks, compiles the Node smoke program, executes the runnable manifest, captures results, formats the generated ClojureScript, and runs conversion assertions. To start the browser interface, use

```
make emmy-runner      sh
```

Then open <http://localhost:8080>. The first compilation can take longer because Clojure and JavaScript dependencies must be downloaded. Leave the command running while using the page; `shadow-cljs` watches the source and reloads the application after changes.

The browser has three actions:

- **Run through this block** resets the context and executes the runnable blocks from the beginning of the selected chapter through the selected block. This is the normal and most reliable way to check a book example.
- **Run editor only** evaluates only the text currently in the editor, preserving the existing session. Use this for a small experiment after running through the required setup. It is not evidence that the block works from a clean chapter context.
- **Reset context** discards all definitions made during the current session and reinstalls Emmy and `fdg.compat`.

A setup listing that replaces the historical Scheme `load` form is installed by the runner and is not evaluated as an ordinary example. If a successful expression returns `nil`, the interface says that no output was produced; otherwise it prints the returned value. Errors identify the block at which sequential execution stopped. Compare that block's `.cljs` file with its `.scm` neighbor and with the surrounding text.

For a lasting correction, edit the file under `codeblocks/` or, preferably, implement a recurring translation issue in the appropriate converter or compatibility layer. Rerunning the conversion copies the durable block into the browser's generated directory. Do not edit `emmy-runner/public/generated/` directly.

### G.3 Automated Runs and Captured Results

The compiled smoke program can be invoked directly:

```
cd emmy-runner
clojure -M:shadow-cljs compile smoke
node target/smoke.js
```

sh

Useful options are

```
node target/smoke.js --verbose
node target/smoke.js --chapter=chapter008
node target/smoke.js --through=chapter008-024
node target/smoke.js --capture-results
```

sh

`--chapter` limits execution to one chapter, `--through` stops at a manifest identifier, and `--verbose` prints each block as it begins. Capture mode writes returned values immediately after the corresponding top-level expressions as `;; =>` comments. Very large results are represented by a short prefix, an omission marker, and their full character count; inspect such a value interactively when the complete expression is needed.

The smoke runner continues after an individual block failure so that one invocation reports all known failures. It prints a consolidated summary and returns a nonzero exit status. The book build is configured to continue far enough to produce PDFs even when the Emmy stage reports problems, but a successful PDF build does not mean the ClojureScript checks passed: the Emmy summary must also be clean.

Selected results are additionally simplified, frozen into stable data, and compared with corresponding cases from Emmy's FDG test suite. These oracle checks detect mathematical regressions; ordinary successful execution only establishes that a block is accepted and runs to completion.

#### G.3.1 Exact checks for expensive symbolic blocks

Some `scmutils` calculations are mathematically modest but create very large intermediate expressions in Emmy. The runner does not replace these calculations with floating-point samples. Instead, the converter rewrites the affected listing into smaller exact identities whose conjunction proves the original result. In the checks that call `verified-zero`, a nonzero component fails the block. The tensor-linearity listing instead returns its simplified residuals; its captured result records that all four were zero in the most recent run.

Four listings currently use this treatment:

- In Chapter 7 (page 67), Emmy's generic Legendre transform is evaluated at a fully symbolic phase-space state and checked against the closed-form Hamiltonian. The following listing differentiates that exact closed form. This avoids repeatedly differentiating through the generic transform while still checking that the transform produced the function being differentiated.
- In Chapter 11 (page 110), boost covariance is reduced directly from the definition of `general-boost` to the three rotation facts it uses: preservation of the boost vector's squared norm, preservation of the relevant dot product, and application of the inverse rotation. All three are checked for the complete symbolic Euler rotation.
- In Appendix F's tensor-linearity check (page 150), the four residuals are simplified independently before being assembled. Its cached `;; => (up 0 0 0 0)` comment records the zero result returned by the most recent captured run, although this listing does not call `verified-zero`.
- In Appendix F's coordinate-change check (page 151), both Christoffel tables are derived symbolically from the exact spherical and stereographic metric-component matrices. Each derived table is checked against its closed form, after which the stereographic table is transformed component by component

and compared with the spherical table. Simplifying each two-dimensional contraction before assembling the next prevents the generic arbitrary-field calculation from producing a multi-megabyte intermediate expression.

The displayed Appendix F coordinate-change listing contains the coordinate bases, both Emmy-derived Christoffel tables, the closed stereographic table, the explicit stereographic coordinate formula, and both sets of exact residuals. Its only compiled operation is `transform-stereographic-Christoffel-to-spherical`, defined in `fdg.slow-checks` and re-exported by `fdg.compat`. That helper differentiates the same factored coordinate formula to obtain its Jacobian and Hessian and performs the repeated two-dimensional contractions.

Compilation is faster here because SCI interprets each nested generic arithmetic operation and represents functions defined in evaluated source as `MetaFn` wrappers. Emmy’s differentiation and structure dispatch operate directly on ordinary ClojureScript functions in the compiled helper, and the JavaScript engine can optimize the repeated small contractions. The helper still uses Emmy’s `D`, generic arithmetic, and `simplify` it changes the execution boundary, not the algebra. Keeping only that contraction kernel compiled makes the proof readable in the book while avoiding the pathological arbitrary-field expansion. For auditability, the converter copies the exact compiled `fdg.slow-checks` source into the displayed `.cljs` listing as comments immediately before its use.

When another block exceeds its chapter watchdog, first factor the mathematical claim into exact intermediate identities and simplify each component at the earliest useful point. Keep the original generic operation in at least one symbolic comparison, as in Chapter 7, or derive both sides from Emmy’s native objects, as in Appendix F. A numerical sample is useful for diagnosis but is not a substitute for the checked book result.

## G.4 Running a Repository Block Manually

For most investigations, use the browser’s **Run through this block** action. If a normal compiled ClojureScript program is preferable, start from the relevant `.cljs` file and include every definition on which it depends. These dependencies are usually the preceding listings in the same chapter. A block that uses names from `fdg.compat` also needs that namespace from this repository.

Captured `:: =>` lines are comments and may remain in the copied source. They document previous output but are not assertions. Print a value explicitly with `println` and usually pass symbolic results through `simplify` and `freeze` before printing:

```
(println (pr-str (freeze (simplify expression))))
```

clojure

Do not replace exact ratios such as `(/ 1 2)` with JavaScript decimals unless a numerical approximation is intended. Emmy’s generic arithmetic sees the explicit division and can retain exact symbolic structure.

## G.5 A Clean ClojureScript and Emmy Project

The reproducible route is to use the versions pinned by this repository. The following minimal Node-target project uses the same Emmy commit, ClojureScript compiler, shadow-cljs version, and JavaScript numerical dependencies as the book runner.

Install a Java JDK, the Clojure CLI, and Node.js with npm. The current official Clojure instructions are at [clojure.org/guides/install\\_clojure](https://clojure.org/guides/install_clojure). On macOS, the documented Homebrew command is

```
brew install clojure/tools/clojure
```

sh

On Linux and other systems, follow the official installer instructions rather than copying a versioned installer command from this book. Confirm the tools before continuing:

```
java --version
clojure --version
node --version
npm --version
```

sh

Create a project:

```
mkdir -p fdg-emmy-demo/src/demo
cd fdg-emmy-demo
```

sh

Create `deps.edn` with

```
{:paths ["src"]
 :deps {io.github.mentat-collective/emmy
        {:git/sha "5a2e1470087d41ec43e73d661bfceb941a6e8870"}}
 :aliases
 {:shadow-cljs
  {:extra-deps
   {thheller/shadow-cljs {:mvn/version "2.20.14"}
    org.clojure/clojurescript {:mvn/version "1.11.60"}}
   :main-opts ["-m" "shadow.cljs.devtools.cli"]}}}
```

clojure

Create `package.json` with the JavaScript dependencies required by this Emmy revision:

```
{
  "private": true,
  "dependencies": {
    "fraction.js": "4.2.1",
    "odex": "3.0.0-rc.4"
  }
}
```

json

Create `shadow-cljs.edn`:

```
{:deps true
 :builds
 {:app
  {:target :node-script
   :output-to "target/main.js"
   :main demo.main/main}}}
```

clojure

Finally, create `src/demo/main.cljs`:

```
(ns demo.main
  (:refer-clojure :exclude [*])
  (:require [emmy.env :refer [* D freeze simplify]]))

(defn cube [x]
  (* x x x))

(defn main []
  (println
   (pr-str
    (freeze
     (simplify
      ((D cube) 'x)))))))
```

clojure

Install the npm dependencies, compile, and run:

```
npm install
clojure -M:shadow-cljs compile app
node target/main.js
```

sh

The program differentiates the function `cube` symbolically and prints its simplified result. Replace the body of `main` with a self-contained book example, then copy any earlier definitions it needs. ClojureScript

does not support `:refer :all` in an `ns` declaration, so explicitly add every Emmy name used by the copied example to the `:refer` vector. If an Emmy generic operation has the same name as an operation in `cljs.core`, add that name to `:refer-clojure :exclude` as shown for `*`. This ensures that symbolic values reach Emmy's generic operation rather than the narrower JavaScript operation.

For an independently maintained project, consult the installation instructions at [emmy.mentat.org](http://emmy.mentat.org) and the [shadow-cljs User's Guide](#), then choose and record current dependency versions. For reproducing this book, retain the pinned files above and the repository's `package-lock.json`.

## H Edition-specific Prose Record

This table is generated from the converter's edition-aware prose replacements. Each row identifies text for which the ClojureScript edition or the combined edition prints a variant of the Scheme wording. The page links are resolved from labels placed directly at the modified text, so they follow pagination changes automatically.

| Location  | Page                      | Edition-specific change                  |
|-----------|---------------------------|------------------------------------------|
| Prologue  | <a href="#">page xiii</a> | functional-derivative appendix reference |
| Prologue  | <a href="#">page xiii</a> | language footnote                        |
| Prologue  | <a href="#">page xv</a>   | closing appendix references              |
| Prologue  | <a href="#">page xiv</a>  | Lagrange-equations function terminology  |
| Prologue  | <a href="#">page xiv</a>  | Lagrange-equations result terminology    |
| Chapter 1 | <a href="#">page 2</a>    | functional-notation appendix reference   |
| Chapter 1 | <a href="#">page 2</a>    | Lfree language and appendix footnote     |
| Chapter 1 | <a href="#">page 2</a>    | sphere->R3 terminology                   |
| Chapter 1 | <a href="#">page 2</a>    | F->C terminology                         |
| Chapter 1 | <a href="#">page 4</a>    | literal metric footnote                  |
| Chapter 1 | <a href="#">page 5</a>    | real-line coordinate setup               |
| Chapter 2 | <a href="#">page 9</a>    | structured derivative appendix reference |
| Chapter 6 | <a href="#">page 44</a>   | real-line coordinate footnote            |
| Chapter 7 | <a href="#">page 50</a>   | tensor footnote appendix reference       |
| Chapter 7 | <a href="#">page 59</a>   | tensor transformation appendix reference |
| Chapter 7 | <a href="#">page 60</a>   | recursive map description                |
| Chapter 7 | <a href="#">page 66</a>   | F->C footnote                            |
| Chapter 8 | <a href="#">page 79</a>   | cyclic-sum terminology                   |
| Chapter 8 | <a href="#">page 79</a>   | Bianchi footnote terminology             |
| Chapter 8 | <a href="#">page 80</a>   | tensor definition appendix reference     |
| Chapter 9 | <a href="#">page 88</a>   | Einstein tensor appendix reference       |
| Errata    | <a href="#">page 171</a>  | rules.scm Emmy note                      |
| Errata    | <a href="#">page 172</a>  | expensive simplification Emmy wording    |

## I Source and Edition Change Log

This appendix is the complete maintained ledger of durable differences between the pinned Org snapshot, the published PDF, and the Typst editions in this repository. Every entry has a stable identifier. A change is recorded here only when it is implemented in the converter, an authored Typst source file, an authored ClojureScript/Emmy source file, or a documented build policy. The vendored Org snapshot remains unchanged; source corrections are applied to a temporary conversion copy.

The ledger is exhaustive for the maintained change surface. The audit’s explicitly unchecked comparison work is not silently converted into a claimed correction: it remains an open audit item in `typ/audit.typ`.

### I.1 Conversion contract

- **I-001 — Pinned-source protection.** `scripts/convert-org-to-typst.mjs` verifies checksums for every vendored Org file before conversion and fails if the upstream snapshot has been edited.
- **I-002 — Temporary source copy.** All normalization and fidelity repairs operate on a temporary source copy; no file under `fdg-book/scheme/org/` is rewritten.
- **I-003 — Raw evaluator results.** Org `:results` value raw output is attached to the preceding Scheme listing as result comments before Pandoc conversion, so cached values are not mistaken for a second executable listing.
- **I-004 — Bare Scheme forms.** Bare parenthesized Scheme listings and slash-delimited displayed results are wrapped as explicit Scheme source blocks; slash-delimited results become comments.
- **I-005 — Stable code records.** Each extracted listing receives a stable chapter/ordinal identifier and is emitted to the generated codeblock records used by the book and runner.
- **I-006 — Subheadings.** Org headings below the level that Pandoc preserves are marked before conversion and restored as visible Typst subheadings.
- **I-007 — Display-math normalization.** Imported `equation`, `align`, `aligned`, and `split` environments have wrapper commands, labels, duplicate rows, and unsupported alignment artifacts normalized before Typst rendering.
- **I-008 — Footnote display boundary.** Main-text displays receive equation labels; displays in the Org footnote section do not consume the main equation counter unless the published source explicitly labels them.
- **I-009 — Published unnumbered displays.** Chapter 7’s published unnumbered derivation rows, code-result arrays, and intermediate displays are excluded from the generated equation sequence.
- **I-010 — Citation mapping.** Numeric Org citations are mapped to the seeded bibliography keys, preserving the published citation order used by the generated references.
- **I-011 — Equation and property references.** Numeric equation, property, section, appendix, range, and list references are rewritten to Typst labels and references; generic parenthesized equation references become one Typst reference wrapper.
- **I-012 — Page-reference mapping.** Stable section and content labels replace hard-coded internal page references so references follow the A4 reflow and edition-specific pagination.
- **I-013 — Figure insertion.** The converter inserts the eight imported published figure PDFs at their corresponding prose anchors and keeps their captions adjacent to the assets.
- **I-014 — Typst math cleanup.** The final math pass normalizes inverse powers, operators, vector notation, wedge notation, spacing, delimiters, and generated line-break markers to the Typst math vocabulary used by this project.

### I.2 Source/PDF fidelity corrections

- **I-015 — Chapter 1 transition.** The sentence introducing the coordinate-writing listing is restored after the cached Cartan result, matching the published prose sequence.
- **I-016 — Chapter 2 heading.** The manifold-functions heading is rendered with the published spelling **Independent**.
- **I-017 — Appendix B, equation B.7.** The selector result is  $I_0(s)=t$ , matching the published equation.
- **I-018 — Appendix B, equation B.10.** The left side is  $AC$ , matching the published tensor name.
- **I-019 — Appendix B phase-space result.** The cached phase-space tuple uses `(down p_x p_y)` rather than two unlabelled momentum symbols.
- **I-020 — Appendix B tuple syntax.** The malformed tuple form `(up_x y)` is rendered as `(up x y)`.



- **I-021 — Appendix C, equation C.1.** The second term uses the second vector argument  $\mathbf{v}$  in  $\mathbf{g} \cdot \mathbf{T}(\mathbf{v}, \boldsymbol{\omega})$ .
- **I-022 — Appendix C, equation C.10.** The primed Riemann component is placed on the left side of the transformation equation.
- **I-023 — Chapter 7 state advancer.** The extra grouping around `sphere-Cartan` is removed from the cached state-advancer listing.
- **I-024 — Chapter 9, equation 9.24.** The missing metric display is restored with the Newtonian-limit expression shown in the published book.
- **I-025 — Chapter 9, equation 9.25.** The missing second-order position equation is restored after the Newtonian-limit prose.
- **I-026 — Chapter 9, equation 9.26.** The missing Einstein equation is restored before the explanation of  $R_{(\mu \nu)}$ .
- **I-027 — Chapter 10, equation 10.3.** The gradient equation is rendered as  $\text{grad}(\mathbf{f}) = \mathbf{g}^{\text{sharp}}(d\mathbf{f})$  without the trailing duplicate differential fragment.
- **I-028 — Chapter 10, equation 10.6.** The second curl component is attached to  $d\mathbf{y}$ , matching the published coordinate basis.
- **I-029 — Chapter 11 code closure.** The unmatched closing parenthesis in the `make-4tuple` listing is removed.
- **I-030 — Chapter 11, equations 11.1–11.5.** The five introductory Maxwell and charge-conservation equations are restored from the published PDF.
- **I-031 — Chapter 11, equations 11.6–11.10.** The divergence, corrected Ampère, wave, and inhomogeneous wave equations are restored from the published PDF.
- **I-032 — Chapter 11, equations 11.11–11.14.** The Lorentz-force, scalar wave, interval, and light-cone equations are restored from the published PDF.
- **I-033 — Chapter 11, equations 11.20–11.21.** The transformed wave equation and invariant-length condition are restored from the published PDF.
- **I-034 — Chapter 11 Lorentz-coordinate display.** The two-coordinate transformation is separated into the two published rows with the correct punctuation.
- **I-035 — Chapter 11 Lorentz constraints.** The three constraints on  $\mathbf{p}$ ,  $\mathbf{q}$ ,  $\mathbf{r}$ , and  $\mathbf{s}$  are separated and punctuated as displayed in the published PDF.
- **I-036 — Chapter 11 boost display.** The four-coordinate boost transformation is separated into the four published rows.
- **I-037 — Chapter 11 vector decomposition.** The parallel/perpendicular decomposition is rendered with norm notation and separate rows.
- **I-038 — Chapter 11 general boost.** The general vector boost is rendered as a two-row display with its dot product and inverse-rotation terms intact.
- **I-039 — Chapter 1 internal references.** References that formerly named Chapter 7 directly target Section 7.10, preserving the current section granularity.
- **I-040 — Prologue citation.** The Papert reference is attached to the bibliography key used by the generated bibliography instead of remaining a broken numeric prose citation.

### I.3 Source text, code, and math normalization

The following entries enumerate every current literal repair in the shared Org normalization layer. They are applied to the generated copy and are not general claims about the upstream project.

- **I-041 — Psi subscript.** The malformed `psi_a` command is restored.
- **I-042 — Indexed  $\mathbb{M}$ .** The extra closing brace in  $\mathbb{M}_i$  is removed.
- **I-043 — Real-vector space exponent.** The extra closing brace in  $\text{mathsf{R}}^n$  is removed.
- **I-044 —  $\mathfrak{m}$  math delimiters.** Two malformed `mathsf{m}` forms are converted to valid math markup.
- **I-045 — Scalar  $n$  math.** The escaped line-break form is reduced to the scalar  $n$ .
- **I-046 — French title accent.** The title *Les Misérables* receives its correct accent.
- **I-047 — Euler–Lagrange spelling.** The joined word is rendered as **Euler-Lagrange**.
- **I-048 — Manifold  $\mathbb{R}^5$  prose.** The  $\mathbb{R}^5$  expression is placed inside one valid math span.
- **I-049 — Lagrangian capitalization.** The sentence beginning “This Lagrangian” uses the published capitalization.
- **I-050 — Longitude spacing.** The first longitude prose uses  $90^\circ \text{ E}$  spacing.



- **I-051 — Hard-hyphen removal for direction.** The line-break hyphen in **direction** is removed.
- **I-052 — Hard-hyphen removal for derivatives.** Both one-line and split-line forms of **derivatives** are joined.
- **I-053 — Hard-hyphen removal for components.** Both one-line and split-line forms of **components** are joined.
- **I-054 — Coordinate-system identifier.** The Chapter 6 identifier is `coordinate-system-at`.
- **I-055 — Pullback identifier.** Whitespace inside `pullback` is removed wherever the Org import split the identifier.
- **I-056 — Reference-coordinate phrase.** The damaged coordinate phrase is normalized to `ref coords`.
- **I-057 — Sine-longitude phrase.** The damaged trigonometric identifier is normalized to `sin long`.
- **I-058 — Longitude-momentum identifier.** The split identifier is normalized to `longp`.
- **I-059 — Pullback parameter case.** The pullback parameter is `f-on-M`.
- **I-060 — Assume spelling.** The prose uses **assume**.
- **I-061 — Comma spacing.** The prose uses `However, if`.
- **I-062 — Symmetric connection.** The lower Christoffel indices on the right side are swapped to express symmetry.
- **I-063 — Function-map prose.** The sentence uses “functions that map”.
- **I-064 — Stray sentence punctuation.** The period before the coordinate-functions sentence is removed.
- **I-065 — Function/math spacing.** The phrase “function `f` at” has a valid prose/math boundary.
- **I-066 — Mapping sentence punctuation.** The period after `m = mu(n)` is restored before the following sentence.
- **I-067 — `mathsf{N}` delimiters.** The malformed math span is repaired.
- **I-068 — Primed-coordinate line.** The `x'` and `chi'` expression is kept on one math line with its comma.
- **I-069 — Basis coefficient index.** The basis formula uses `X_k(f)` rather than an unindexed `X(f)`.
- **I-070 — First inverse-map derivative.** The derivative of `f` is written with `(chi')^(-1)`.
- **I-071 — Coordinate-change inverse.** The two Jacobian factors use inverse maps with exponent `-1`.
- **I-072 — Vector-component transformation.** The transformed component formula uses the inverse Jacobian.
- **I-073 — Vector-field display.** The two-component vector-field formula is compacted into one valid math display.
- **I-074 — Area-form display.** `A = dx wedge dy` is compacted into one valid math display.
- **I-075 — Euler operator display.** The malformed `E[L]` expression is rendered as `\mathbf{E}[L]`.
- **I-076 — Differential delimiters.** The `dx` and `dy` phrase uses complete math spans.
- **I-077 — Metric punctuation.** The period after `g` is outside the math command.
- **I-078 — `u` delimiter.** The malformed closing bracket after `u` is replaced by a brace.
- **I-079 — `w` delimiter.** The malformed closing bracket after `w` is replaced by a brace.
- **I-080 — Basis-set braces.** The set of basis vectors uses explicit left and right braces.
- **I-081 — Metric-index brace.** The `g^(ij)` exponent is closed correctly.
- **I-082 — Epsilon subscript.** The epsilon subscript receives its missing closing brace and math delimiter.
- **I-083 — Prime command.** The misspelled prime command is restored.
- **I-084 — Einstein spelling.** The author’s name is rendered as **Einstein**.
- **I-085 — TeX command.** The TeX name is rendered with `\mathrm` rather than an unsupported macro.
- **I-086 — Ellipsis command.** Imported `\ldots` is normalized to the supported centered ellipsis.
- **I-087 — Half factor.** The imported fraction is normalized to `\frac{1}{2}`.
- **I-088 — Negative thin-space commands.** Unsupported `! spacing` commands are removed.
- **I-089 — Curve derivative delimiters.** The `D(chi compose gamma)(t)` expression has balanced math delimiters.
- **I-090 — Summation braces.** The `d_l^i` and `d_k^i` summands lose imported grouping braces that break Typst parsing.
- **I-091 — Summation continuation.** The `d_k^i(m) c_j^k(m)` row is made into one valid summand.
- **I-092 — Lorentz quadratic form.** The duplicated `xi^2` term is corrected to `xi^1`.
- **I-093 — Parallel-vector notation.** The imported vertical-bar notation is rendered as bold `xi` parallel notation.

- **I-094 — Inverse curvature argument.** The malformed `R^(=1)` is rendered as `R^(-1)`.
- **I-095 — Four-tuple identifier.** The four-tuple constructor is normalized to `make-4tuple`.
- **I-096 — Curvature equality.** The stray backslash before the equality sign is removed.
- **I-097 — Curvature-vector equality.** The missing equality sign before the omega vector is restored.
- **I-098 — Exponential delimiter.** The extra brace in the exponential identity is removed.

## I.4 Imported setup and executable-claim repairs

- **I-099 — Division spacing.** A digit immediately following a Scheme division operator receives the required separator.
- **I-100 — Fake vector field.** The Chapter 6 call uses `make-fake-vector-field`.
- **I-101 — R2 point closure.** The `R2-rect-point` definition has one balanced closing parenthesis.
- **I-102 — SR basis setup.** The Chapter 10 setup defines `SR-basis` from `SR`.
- **I-103 — Spherical basis setup.** The spherical coordinate declaration is emitted before `S2-basis` is used.
- **I-104 — Spherical one-form setup.** The Chapter 8 setup defines the omega one-form field and the manifold function used by the torsion listing.
- **I-105 — Force setup.** `eta-inverse` is defined before the `Force` listing uses it.
- **I-106 — Pullback function closure.** The imported `coordinatize` listing closes its inner expression at the correct level.
- **I-107 — Euler-angle setup.** The `SO3` listing receives the Euler coordinate system, coordinates, point, scalar function, and three frame-vector definitions it uses.
- **I-108 — Differential dual result.** The cached result of `dx(d/dx)` is corrected to `1`.
- **I-109 — Spelling and operator repairs.** `illustrate`, `basis→1form-basis`, `metric basis`, and the `Tij` tensor identifier are normalized.
- **I-110 — Newton metric setup.** The spacetime coordinates and `spacetime-rect-basis` are defined before the Newton metric.
- **I-111 — Schwarzschild connection setup.** The Schwarzschild Cartan form is built before the geodesic residual function.
- **I-112 — Newton potential binding.** `v` is bound once and passed to the Newton connection.
- **I-113 — Base-frame setup.** The `home` frame is defined before the special-relativistic frame `A`.
- **I-114 — Errata energy result.** The complete three-term cached value is retained instead of only its leading term.
- **I-115 — Spherical manifold family.** Generated code uses the valid `S^2-type` manifold family.
- **I-116 — Polar-Cartan operator.** The polar-Cartan listing calls `circular`, the operator defined by the surrounding source.
- **I-117 — Errata code names.** Generated errata annotations retain the historical source forms while the corrected generated code uses the repaired names.

## I.5 Equation layout and counter changes

- **I-118 — Chapter 2 equation split.** Equations 2.6 and 2.7 are separated into two displays instead of being merged with prose.
- **I-119 — Appendix A general forms.** The operator, `lambda`, `cond`, `if`, and `let` templates are rendered as Scheme code blocks rather than malformed centered math.
- **I-120 — Chapter 3 footnote equation.** The 3.3 footnote is prose plus an inline equation, not a nested code block with a second equation label.
- **I-121 — Chapter 4 equation 4.1.** The short two-equality display is kept on one line.
- **I-122 — Chapter 4 equation 4.6.** The `cjk(m)` product is kept with the equality row.
- **I-123 — Chapter 4 equation 4.9.** The second equality is not forced onto a new source line.
- **I-124 — Chapter 4 equation 4.19.** The Jacobian equality is compacted.
- **I-125 — Chapter 4 equation 4.29.** The spherical-basis equality is compacted.
- **I-126 — Chapter 4 equation 4.34.** The long bracket calculation uses an aligned continuation marker rather than a crowded plus row.
- **I-127 — Chapter 4 equation 4.35.** The coefficient identity is compacted and its minus term remains on the same equality.
- **I-128 — Chapter 4 equation 4.38.** The closing parenthesis is kept inside the displayed summand.

- **I-129 — Chapter 4 exponential identity.** The product of exponentials is compacted without changing its commutator term.
- **I-130 — Chapter 4 equations 4.40–4.42.** The short flow identities are compacted into the available width.
- **I-131 — Chapter 5 equation 5.24.** The commutator is `[v_i, v_j]`, the first summation stays on the equality row, and continuation rows use `&quad`.
- **I-132 — Chapter 5 equation 5.34.** Boundary-integral terms use explicit `+` and `-` continuation markers.
- **I-133 — Chapter 7 equation 7.41.** The display's unnecessary leading break and oversized delimiter wrapper are removed.
- **I-134 — Chapter 7 equation 7.59.** Repeated summand parentheses are retained where required by the displayed algebra, and the first equality remains aligned.
- **I-135 — Chapter 7 equation labels 7.77–7.79.** The geodesic displays receive the published labels after the preceding unnumbered display is suppressed.
- **I-136 — Chapter 8 equation 8.2.** The short curvature definition is compacted.
- **I-137 — Chapter 8 equation 8.16.** The covariant-derivative display begins with its equality and uses aligned continuation rows.
- **I-138 — Chapter 8 equation 8.18.** The short torsion display is compacted.
- **I-139 — Chapter 10 compact display set.** Displays 10.1, 10.2, 10.5–10.11, and 10.19–10.26 use the available width and avoid unnecessary source breaks.
- **I-140 — Chapter 10 operator alignment.** Displays 10.3, 10.7, and 10.10 keep their left-hand operator and first equality on the same aligned row.
- **I-141 — Equation counter reset.** Each chapter and numbered appendix resets the equation counter at its wrapper.
- **I-142 — Footnote counter reset.** Each chapter and numbered appendix resets the footnote counter at its wrapper.
- **I-143 — Figure counter reset.** Each chapter and numbered appendix resets the figure counter at its wrapper.
- **I-144 — Footnote-only labels.** The generated sequence excludes the footnote-only labels 3.58, 3.59, 5.40, 7.83, 7.84, 8.33, and 8.34.
- **I-145 — Generated equation sequences.** Regression locks the contiguous sequences 1.1–1.6, 2.1–2.7, 3.1–3.57, 4.1–4.44, 5.1–5.39, 6.1–6.28, 7.1–7.79, 8.1–8.32, 9.1–9.32, 10.1–10.28, 11.1–11.43, B.1–B.35, and C.1–C.11.
- **I-146 — Chapter footnote counts.** Regression locks the generated counts for the Prologue, Chapters 1–11, Appendices A–C, and the project-authored appendices.

## I.6 ClojureScript and Emmy edition transformations

- **I-147 — Appendix D source.** Appendix D is derived from Appendix A with ClojureScript language names, ClojureScript code syntax, Emmy terminology, and D-section labels.
- **I-148 — Appendix E source.** Appendix E is derived from Appendix B with ClojureScript function terminology, Emmy notation, and E-section labels.
- **I-149 — Appendix F source.** Appendix F is derived from Appendix C with Emmy tensor terminology and F-section labels.
- **I-150 — ClojureScript renderer.** Derived appendices use the ClojureScript code renderer and `lang: "clojure"` blocks.
- **I-151 — Editorial derivation notes.** Derived appendices receive a visible note identifying their Scheme source appendix and Emmy translation.
- **I-152 — Appendix-wide references.** Source appendix labels, equation labels, appendix references, and cross-references are changed from A/B/C to D/E/F as appropriate.
- **I-153 — Scheme implementation vocabulary.** `Scmutils` and its lower-case form become `Emmy` in the translated appendices.
- **I-154 — Procedure vocabulary.** Scheme procedure and program wording becomes ClojureScript function and program wording.
- **I-155 — Scheme system vocabulary.** Scheme system, symbol, vector, and selector wording becomes ClojureScript wording.

- **I-156 — Scheme language description.** The Scheme language paragraph becomes a ClojureScript/Emmy description with the JavaScript compilation context.
- **I-157 — Appendix D language introduction.** The elementary Scheme introduction is replaced by an introduction to the ClojureScript forms used by this edition.
- **I-158 — Appendix D function terminology.** Procedure calls, definitions, lambda expressions, recursive procedures, and the `define` construct become function calls, definitions, function expressions, recursive functions, and `def`.
- **I-159 — Appendix D collections.** The pair/list discussion is replaced by persistent ClojureScript sequences and Emmy up/down structures.
- **I-160 — Appendix D selectors.** `list-ref` and `vector-ref` become `nth`; `car` and `cdr` become `first` and `rest`; Emmy structures use `ref`.
- **I-161 — Appendix D equality.** The equality form becomes `=`.
- **I-162 — Appendix D booleans.** `#t` and `#f` become `true` and `false`.
- **I-163 — Appendix D authored names section.** The appendix adds the scopes and roles of `def`, `fn`, `defn`, `let`, and `letfn`.
- **I-164 — Appendix D parameter vectors.** Function parameters are documented as vectors, including the variadic `&` marker.
- **I-165 — Appendix D conditional syntax.** The final `cond` predicate is documented as the `:else` keyword, with ClojureScript's alternating predicate/consequent syntax.
- **I-166 — Appendix D collection semantics.** The appendix states that ClojureScript collections are immutable and that Emmy structures are not ordinary ClojureScript vectors.
- **I-167 — Appendix D cheat-sheet global values.** The cheat sheet records `(define x value)` to `(def x value)`.
- **I-168 — Appendix D cheat-sheet global functions.** The cheat sheet records `(define (f x) body)` to `(defn f [x] body)`.
- **I-169 — Appendix D cheat-sheet anonymous functions.** The cheat sheet records `(lambda (x) body)` to `(fn [x] body)`.
- **I-170 — Appendix D cheat-sheet function values.** The cheat sheet records a named lambda value to a named `fn` value.
- **I-171 — Appendix D cheat-sheet local values.** The cheat sheet records Scheme `let` bindings to ClojureScript binding vectors.
- **I-172 — Appendix D cheat-sheet sequential local values.** The cheat sheet records `let*` sequencing through ordered ClojureScript `let` bindings.
- **I-173 — Appendix D cheat-sheet local functions.** The cheat sheet records local Scheme definitions to `letfn`.
- **I-174 — Appendix D cheat-sheet named loops.** The cheat sheet records named Scheme loops to `loop` and `recur`.
- **I-175 — Appendix D cheat-sheet sequencing.** The cheat sheet records `begin` to `do`.
- **I-176 — Appendix D cheat-sheet conditionals.** The cheat sheet records Scheme `if/cond` clauses to ClojureScript `if/else` forms.
- **I-177 — Appendix D cheat-sheet booleans.** The cheat sheet records Scheme boolean literals to ClojureScript boolean literals.
- **I-178 — Appendix D cheat-sheet falsey values.** The cheat sheet records Scheme's false-only truth rule and ClojureScript's `false/nil` falsey rule.
- **I-179 — Appendix D cheat-sheet exact ratios.** The cheat sheet records `1/2` to `(/ 1 2)` and preserves exact arithmetic.
- **I-180 — Appendix D cheat-sheet vectors.** The cheat sheet records Scheme vector literals to ClojureScript vector literals.
- **I-181 — Appendix D cheat-sheet empty lists.** The cheat sheet records the empty-list form and ClojureScript's list constructor alternative.
- **I-182 — Appendix D cheat-sheet equality.** The cheat sheet records `eq?` to `=`.
- **I-183 — Appendix D cheat-sheet indexed selection.** The cheat sheet records list/vector selection to `nth` and Emmy `ref`.
- **I-184 — Appendix D cheat-sheet sequence selectors.** The cheat sheet records `car/cdr` to `first/rest`.

- **I-185 — Appendix D cheat-sheet variadic parameters.** The cheat sheet records dotted Scheme parameters to a ClojureScript parameter vector containing `&`.
- **I-186 — Appendix D cheat-sheet quotation.** The cheat sheet records quotation and keyword self-evaluation.
- **I-187 — Appendix D cheat-sheet function application.** The cheat sheet records identical call syntax and notes vector-based parameter and binding lists.
- **I-188 — Appendix E composition wording.** Function composition is described with ClojureScript functions rather than Scheme procedures.
- **I-189 — Appendix E arithmetic wording.** The `g` construction is described as a ClojureScript function.
- **I-190 — Appendix E cached simplification.** The interactive printer description is replaced by cached runner output, explicit `simplify`, and Appendix G guidance.
- **I-191 — Appendix E literal functions.** `literal-function` is described as constructing a symbolic function with a declared signature.
- **I-192 — Appendix E multiple arguments.** The Scheme-specific paragraph becomes a ClojureScript/Emmy distinction.
- **I-193 — Appendix E selector vocabulary.** `component` and `ref` are described as functions.
- **I-194 — Appendix E derivative vocabulary.** Derivatives of functions are described with ClojureScript function terminology.
- **I-195 — Appendix E exercise labels.** Exercises B.1 and B.2 become E.1 and E.2.
- **I-196 — Appendix F tensor test.** The tensor test is described as taking an Emmy/ClojureScript function.

## I.7 Project-authored material and execution records

- **I-197 — ClojureScript preface.** The ClojureScript editions add an authored preface describing provenance, repository variants, the runner, translation confidence, artwork, and update date.
- **I-198 — Appendix D.** The project adds the ClojureScript language appendix.
- **I-199 — Appendix E.** The project adds the Emmy notation appendix.
- **I-200 — Appendix F.** The project adds the Emmy tensors appendix.
- **I-201 — Appendix G.** The project adds the runner, generated block records, sequential chapter sessions, browser controls, smoke checks, captured results, exact symbolic checks, and clean-project instructions.
- **I-202 — Appendix H.** The project adds the generated edition-specific prose table; its 23 records are listed individually in Appendix H and repeated by location below.
- **I-203 — Appendix I.** The project adds this source-and-edition ledger.
- **I-204 — Annotated errata.** The project adds the upstream errata as end matter and marks the fourteen repairs listed in the annotated-errata section.
- **I-205 — Generated code artifacts.** Each Scheme block has a durable `.scm` record; each translated block has a durable `.cljs` record; the runner manifest records order, heading, source location, definitions, top-level forms, and run status.
- **I-206 — Browser and smoke execution.** The project adds an SCI browser runner, a compiled Node smoke runner, chapter-session execution, result capture, and regression assertions.
- **I-207 — Expensive symbolic checks.** The project adds exact intermediate-identity checks for the Chapter 7 Legendre transform, Chapter 11 boost covariance, Appendix F tensor linearity, and Appendix F coordinate-change calculation.
- **I-208 — Edition gating.** Appendices D–I and the ClojureScript preface appear only in the ClojureScript and combined editions.
- **I-209 — Manifest registration.** The generated source manifest includes Appendices D–I with their display titles.

## I.8 Annotated errata corrections

These are the individual errata markers currently annotated as corrected in the generated errata section.

- **I-210 — Chapter 1 Cartan order.** The generated Chapter 1 setup places the Cartan definition before the geodesic residual listing that uses it.
- **I-211 — Chapter 6 manifold family.** The generated `S2` definition uses `S^2`-type.
- **I-212 — Chapter 7 circular operator.** The generated polar-Cartan listing uses `circular`.



- **I-213 — Chapter 7 S2-Christoffel coordinates.** The generated setup installs `S2-spherical` before `S2-Christoffel` is used.
- **I-214 — Chapter 7 sphere manifold family.** The generated `sphere` definition uses `S^2-type`.
- **I-215 — Chapter 8 spherical coordinates.** The generated setup installs `S2-spherical`.
- **I-216 — Chapter 8 omega field.** The generated setup defines the omega one-form field.
- **I-217 — Chapter 8 torsion function.** The generated setup defines the `f` manifold function used by the torsion listing.
- **I-218 — Chapter 9 spacetime basis.** The generated setup defines `spacetime-rect` and `spacetime-rect-basis`.
- **I-219 — Chapter 9 Newton potential.** The generated setup binds `v` before passing it to `Newton-metric`.
- **I-220 — Chapter 9 returned value.** The generated result preserves all terms reported by the upstream erratum.
- **I-221 — Chapter 10 SR basis.** The generated setup defines `SR-basis` before the last section listing uses it.
- **I-222 — Chapter 10 eta inverse.** The generated setup defines `eta-inverse` before `Force` uses it.
- **I-223 — Chapter 10 base frame.** The generated setup includes the `home` frame definition before constructing frame `A`.

## I.9 Edition-specific prose records

Appendix H is the generated page-linked record of every edition-aware prose replacement. The following list is the complete replacement set, in converter order.

- **I-224 — Prologue functional-derivative reference.** Scheme points to Appendix B; ClojureScript points to Appendix E; the combined edition names both.
- **I-225 — Prologue language footnote.** Scheme describes Scheme and Scmutils; ClojureScript describes ClojureScript and Emmy; the combined edition describes both pairs.
- **I-226 — Prologue closing references.** Scheme points to Appendices A/B; ClojureScript points to D/E; the combined edition names both pairs.
- **I-227 — Prologue Lagrange-equations terminology.** Scheme uses procedure terminology; ClojureScript uses function terminology; the combined edition names both function forms.
- **I-228 — Prologue Lagrange-equations result.** The result description is rewritten for function-valued ClojureScript code in the translated edition.
- **I-229 — Chapter 1 functional-notation reference.** The appendix reference changes from B to E for ClojureScript and names B/E in the combined edition.
- **I-230 — Chapter 1 Lfree footnote.** The procedure/function wording and language appendix reference change by edition.
- **I-231 — Chapter 1 sphere→R3.** Procedure becomes function in ClojureScript and both names appear in the combined edition.
- **I-232 — Chapter 1 F→C terminology.** Procedure becomes function in ClojureScript and both names appear in the combined edition.
- **I-233 — Chapter 1 literal metric footnote.** `literal-metric`, quoted symbol wording, and notation appendix references change by edition.
- **I-234 — Chapter 1 real-line coordinate setup.** Scheme uses `d/dt`; ClojureScript uses `d:dt`; the combined edition names both.
- **I-235 — Chapter 2 structured derivative reference.** The tuple appendix reference changes from B to E by edition.
- **I-236 — Chapter 6 real-line coordinate footnote.** The coordinate-vector-field name changes from `d/dt` to `d:dt` in ClojureScript.
- **I-237 — Chapter 7 tensor footnote reference.** The tensor appendix reference changes from C to F by edition.
- **I-238 — Chapter 7 tensor-transformation reference.** The tensor appendix reference changes from C to F by edition.
- **I-239 — Chapter 7 recursive-map description.** Scheme describes `s:map/r`; ClojureScript describes Emmy `mapr`; the combined edition names both.

- **I-240 — Chapter 7  $F \rightarrow C$  footnote.** Procedure becomes function in ClojureScript and both names appear in the combined edition.
- **I-241 — Chapter 8 cyclic-sum terminology.** Procedure becomes function in ClojureScript and generic function wording is used in the combined edition.
- **I-242 — Chapter 8 Bianchi terminology.** Scheme programs become ClojureScript programs or executable programs by edition.
- **I-243 — Chapter 8 tensor-definition reference.** The tensor appendix reference changes from C to F by edition.
- **I-244 — Chapter 9 Einstein-tensor reference.** The tensor appendix reference changes from C to F by edition.
- **I-245 — Errata rules.scm note.** The translated editions explain Emmy’s replacement role and point to Appendix G; the Scheme edition retains the Scheme note.
- **I-246 — Errata expensive simplification note.** The translated editions identify the cached Emmy check; the Scheme edition retains the scmutils performance wording.

## I.10 Presentation and policy changes

- **I-247 — Paper size.** The Typst editions use A4 page geometry, with repository margins and regenerated pagination; they do not imitate the published PDF’s trim size.
- **I-248 — Paragraph indentation.** Paragraphs after the first paragraph of a section use a 1.5em first-line indent; the first paragraph remains unindented.
- **I-249 — Typography.** The Typst editions use New Computer Modern, project-defined leading, justified paragraphs, and tracked justification limits.
- **I-250 — Running heads.** Odd and even pages use mirrored running heads with the current section title and page number.
- **I-251 — Code presentation.** Scheme and ClojureScript listings use project syntax highlighting, language tags, stable anchors, and edition-aware rendering.
- **I-252 — Cover.** The project uses an authored plotted cover rather than the published cover artwork.
- **I-253 — Figures.** The project uses imported figure PDFs at the eight preserved figure anchors and maintains a separate CeTZ redraw collection for selected figures.
- **I-254 — Front matter.** The project-authored title material includes the current license, sales contact, CIP spacing, print-line spacing, dc23 punctuation, and right-aligned control number treatment.
- **I-255 — Contents and index.** Contents, cross-references, page references, and index locators are regenerated from Typst labels and the current edition’s pagination.
- **I-256 — End-matter order.** The current book places errata, references, and the generated index after the main text and appendices.
- **I-257 — Combined edition.** The combined edition retains the Scheme material and adds the translated ClojureScript material and edition-specific prose branches.
- **I-258 — Source-of-truth policy.** Durable changes belong in the converter, authored Typst library, authored translation, or runner source; generated `typ/content/` files are refreshable outputs.
- **I-259 — Regression policy.** Source/PDF repairs, edition gates, counter behavior, generated labels, and required appendix records are checked by the repository regression scripts.

## I.11 Maintenance rule

When a new Org/PDF disagreement or source-derived project change is found, add the next identifier here with its source location, generated location, current form, target form, and reason. Add or update a regression assertion when the change is mechanically testable. If the discrepancy is still under comparison, record it in the unchecked inventory in `typ/audit.typ` instead of presenting it as a completed correction.

## Errata for FDG

This section is an annotated version of the errata maintained by [mentat-collective/fdg-book](https://github.com/mentat-collective/fdg-book). Notes marked “Corrected in this edition” identify issues already repaired in the generated text or code.

### Chapter 1

Page 9: `Cartan` is used in `geodesic-equation-residuals` before it is defined just after. Be careful to flip evaluation order of these two listings. *Status: Corrected in this edition.*

### Chapter 3

Page 24: the definition of `v` assumes that `R2→R` from page 16 is still defined. This might be nice to add to the `scmutils` library so that this code will work if users attempt chapter 3 independently.

The listing at the bottom of the page assumes `R2-rect-point` is still defined.

Page 38: Note that `omega` was already defined on page 35.

### Chapter 4

Page 44: Note that you need to install the `R2-rect` coordinate system:

```
(define-coordinates (up x y) R2-rect) closure
```

### Chapter 5

Page 60: This page restates:

```
(define-coordinates (up x y z) R3-rect) closure
```

But assumes that the `R3-rect-point` binding still exists from the previous chapter:

```
(def R3-rect-point ((point R3-rect) (up 'x0 'y0 'z0))) closure
```

Page 60: I was concerned that

```
(define-coordinates (up r theta z) R3-cyl) closure
```

installs `z`, `dz` and `d/dz` from cylindrical coordinates over top of the rectangular coordinates. I know that these are equivalent for all intents and purposes... just noting that maybe

```
(define-coordinates (up r theta z-cyl) R3-cyl) closure
```

would be more pedantic, and, maybe, more correct? Or maybe this affects nothing.

Page 67: the definitions of `alpha` and `beta` assume that `R2→R` from page 16 is still defined.

Page 67 uses `R2-rect-point` without ever defining it. The definition is, of course,

```
(def R2-rect-point ((point R2-rect) (up 'x0 'y0))) closure
```

### Chapter 6

Page 75: the definition of `S2` references the nonexistent manifold family `S^2` instead of `S^2-type`: *Status: Corrected in this edition.*

```
(def S2 (make-manifold S2-type 2 3)) closure
```

### Chapter 7

On page 92, the result at the top of the page is not correct, and `scmutils` won't produce it for the supplied code... I don't know enough to see how the book result comes about, so I submit it to you for examination! The code on the page is written for `R3` but the result is for `R2`. If you run the code in the book, you get:



```
(((partial 1) f-rect) (up 1 0 0))
(* a (((partial 0) f-rect) (up 1 0 0)))
(* (/ -1 2) (expt a 2) (((partial 1) f-rect) (up 1 0 0)))
(* (/ -1 6) (expt a 3) (((partial 0) f-rect) (up 1 0 0)))
(* (/ 1 24) (expt a 4) (((partial 1) f-rect) (up 1 0 0)))
```

closure

While the book shows:

```
(((partial 0) f-rect) (up 1 0))
(* -1 a (((partial 1) f-rect) (up 1 0)))
(* (/ -1 2) (expt a 2) (((partial 1) f-rect) (up 1 0)))
(* (/ 1 6) (expt a 3) (((partial 0) f-rect) (up 1 0)))
(* (/ 1 24) (expt a 4) (((partial 1) f-rect) (up 1 0)))
```

closure

The `(partial 0)` and `(partial 1)` are switched, the point is in R2 vs R3 and the negative signs are distributed differently. Strange!

Page 103: I believe the code listing at the end of the page subs in `J` where `circular` belongs. *Status: Corrected in this edition.* The code shown is:

```
(((covariant-derivative R2-polar-Cartan) d:dx) J) f) R2-rect-point)
```

closure

The correct version is:

```
(((covariant-derivative R2-polar-Cartan) d:dx) circular) f) R2-rect-point)
```

closure

Page 107: the definition of `S2-Christoffel` will not work without `S2-spherical` coordinates installed. *Status: Corrected in this edition.*

```
(define-coordinates (up theta phi) S2-spherical)
```

closure

Page 107: The definition of `sphere` references the nonexistent `S^2` manifold family instead of the correct `S^2-type`. *Status: Corrected in this edition.*

## Chapter 8

Page 116: The code beginning here requires the `S2-spherical` coordinate system: *Status: Corrected in this edition.*

```
(define-coordinates (up theta phi) S2-spherical)
```

closure

Page 127 states “Where `omega` is an arbitrary one-form field.” It would be nice to add this definition to the setup in footnote 8: *Status: Corrected in this edition.*

```
(def omega (literal-oneform-field 'omega S2-spherical))
```

closure

The torsion example on page 127 uses an `f` that has not yet been defined:

```
(let [X (literal-vector-field 'X-sphere S2-spherical)
      Y (literal-vector-field 'Y-sphere S2-spherical)]
  (((torsion-vector nabla) X Y) f) m))
```

closure

This can be fixed by adding the following to footnote 8's setup instructions: *Status: Corrected in this edition.*

```
(def f (literal-manifold-function 'f S2-spherical))
```

closure

## Chapter 9

Page 135: I'm not sure if it causes any problems, but `raise` as defined in the book does not wrap its return procedure in `procedure→vector-field`, like the `scmutils` version does. Other functions in the book are careful to show this, so it might be worth a note.

Page 136: The code beginning here requires the `S2-spherical` coordinate system and `S2-basis`:

```
(define-coordinates (up theta phi) S2-spherical) closure

(def S2-basis (coordinate-system→basis S2-spherical))
```

Page 141: The simplifier in the current build of `scmutils` can't simplify the denominators to the book's terms with the  $3/2$  power. If this was hand-simplified, great! Otherwise, maybe this is a regression in the simplifier. I can't see a setting in `rules.scm` that would allow this, but I haven't looked at the full set of rules in a while... Emmy has no corresponding `rules.scm` file; translated examples use Emmy's simplifier and explicit cached checks as described in Appendix G.

Page 146: The code in section 9.3 requires the `spacetime-rect` coordinate system to be installed. `spacetime-rect-basis` is also used in the first code block on this page without definition: *Status: Corrected in this edition.*

```
(define-coordinates (up t x y z) spacetime-rect) closure

(def spacetime-rect-basis (coordinate-system→basis spacetime-rect))
```

Page 147: `V` is passed as an argument to `Newton-metric` without first being defined. `V` was declared inline above in the definition of `nabla`, and should be explicitly defined like so: *Status: Corrected in this edition.*

```
(def V (literal-function 'V '(→ (UP Real Real Real) Real))) closure
```

Page 147: the expression after “If we evaluate the right-hand side expression we obtain” actually returns

```
(+ (* (/ 1 2) (expt 'c 4) rho) closure
  (* 2 (expt 'c 2) rho (V (up x y z)))
  (* 2 rho (expt (V (up x y z)) 2)))
```

instead of the stated return value:

```
(* (/ 1 2) (expt 'c 4) rho) closure
```

Maybe the shown value is meant to be just the leading term, but this is worth explaining. *Status: Corrected in this edition.*

## Chapter 10

Page 159: the setup block should define `SR-basis`, as it is used in the last example of the section, on page 160: *Status: Corrected in this edition.*

```
(def SR-basis (coordinate-system→basis SR)) closure
```

If the setup block defined `SR-basis` then the `SR-vector-basis` definition on page 160 could become:

```
(def SR-vector-basis (basis→vector-basis SR-basis)) closure
```

Page 159: the Minkowski metric is written with a  $c^2$  term, but all of the following functions, results and discussion seem to assume that `c` is normalized to 1. I would recommend a note on page 159 to this effect!

Including the  $c^2$  term explicitly would require, I believe, the following substitutions (along with result substitutions for the forms below):

```
;; page 159
(defn g-Minkowski
  [u v]
  (+ (* -1 (square 'c) (dct u) (dct v)) (* (dx u) (dx v)) (* (dy u) (dy v)) (* (dz u)
    (dz v))))

(def SR-vector-basis (down (* (/ 1 'c) d:dct) d:dx d:dy d:dz))

(def SR-oneform-basis (up (* 'c dct) dx dy dz))

(def SR-basis (make-basis SR-vector-basis SR-oneform-basis))

(defn Faraday
  [Ex Ey Ez Bx By Bz]
  (+ (* Ex 'c (wedge dx dct))
    (* Ey 'c (wedge dy dct))
    (* Ez 'c (wedge dz dct))
    (* Bx (wedge dy dz))
    (* By (wedge dz dx))
    (* Bz (wedge dx dy))))

(defn Maxwell
  [Ex Ey Ez Bx By Bz]
  (+ (* -1 'c Bx (wedge dx dct))
    (* -1 'c By (wedge dy dct))
    (* -1 'c Bz (wedge dz dct))
    (* Ex (wedge dy dz))
    (* Ey (wedge dz dx))
    (* Ez (wedge dx dy))))

(defn J
  [charge-density Ix Iy Iz]
  (- (* (/ 1 'c) (+ (* Ix dx) (* Iy dy) (* Iz dz))) (* charge-density 'c dct)))

(((d F) (* (/ 1 'c) d:dct) d:dy d:dz) an-event)

(((d F) (* (/ 1 'c) d:dct) d:dz d:dx) an-event)

(((d F) (* (/ 1 'c) d:dct) d:dx d:dy) an-event)

(((d (- (d (SR-star F)) (* 4 'pi (SR-star four-current))) (* (/ 1 'c) d:dct) d:dy d:dz)
  an-event)

(((d (- (d (SR-star F)) (* 4 'pi (SR-star four-current))) (* (/ 1 'c) d:dct) d:dz d:dx)
  an-event)

(((d (- (d (SR-star F)) (* 4 'pi (SR-star four-current))) (* (/ 1 'c) d:dct) d:dx d:dy)
  an-event)
```

Page 165: In the definition of `Force`, `eta-inverse` is not defined, so the following two code examples (and, presumably, exercise 10.1b) will not run! *Status: Corrected in this edition.*

## Chapter 11

Page 178: This symbolic simplification is also an expensive check in Emmy. Appendix G identifies the cached check and its result.

Page 180 states “Assume that we have a base frame called `home`.” The base frame defined in the library is `the-ether`; I would recommend including one of the following definitions as setup: *Status: Corrected in this edition.*

```
(def home (base-frame-maker 'home 'home))  
  
(def home the-ether)
```

closure

## References

- [1] H. Abelson, G. J. Sussman, and J. Sussman, *Structure and Interpretation of Computer Programs*. Cambridge, MA: MIT Press, 1996.
- [2] H. Abelson and A. deSessa, *Turtle Geometry*. Cambridge, MA: MIT Press, 1980.
- [3] R. L. Bishop and S. I. Goldberg, *Tensor Analysis on Manifolds*. New York: MacMillan, 1968.
- [4] S. Carroll, *Spacetime and Geometry: An Introduction to General Relativity*. Benjamin Cummings, 2003.
- [5] A. Church, *The Calculi of Lambda-Conversion*. Princeton University Press, 1941.
- [6] H. Flanders, *Differential Forms with Applications to the Physical Sciences*. New York: Academic Press, 1963.
- [7] T. Frankel, *The Geometry of Physics*. Cambridge University Press, 1997.
- [8] G. Galilei, *Il Saggiatore (The Assayer)*. 1623.
- [9] S. W. Hawking and G. F. R. Ellis, *The Large Scale Structure of Space-Time*. Cambridge University Press, 1973.
- [10] “IEEE Standard for the Scheme Programming Language.” 1991.
- [11] C. W. Misner, K. S. Thorne, and J. A. Wheeler, *Gravitation*. San Francisco: W. H. Freeman and Company, 1973.
- [12] A. Pais, *Subtle is the Lord: The Science and the Life of Albert Einstein*. Oxford, UK: Oxford University Press, 1982.
- [13] S. A. Papert, *Mindstorms: Children, Computers, and Powerful Ideas*. Basic Books, 1980.
- [14] B. Schutz, *A First Course in General Relativity*. Cambridge University Press, 1985.
- [15] I. M. Singer and J. A. Thorpe, *Lecture Notes on Elementary Topology and Geometry*. Glenview, Illinois: Scott, Foresman and Company, 1967.
- [16] M. Spivak, *A Comprehensive Introduction to Differential Geometry*. Houston, Texas: Publish or Perish, 1970.
- [17] M. Spivak, *Calculus on Manifolds*. New York, NY: W. A. Benjamin, 1965.
- [18] G. J. Sussman and J. Wisdom, “The Role of Programming in the Formulation of Ideas,” technical report AIM-2002-18, Nov. 2002.
- [19] G. J. Sussman, J. Wisdom, and M. E. Mayer, *Structure and Interpretation of Classical Mechanics*. Cambridge, MA: MIT Press, 2001.
- [20] R. M. Wald, *General Relativity*. University of Chicago Press, 1984.
- [21] “Free software.” [Online]. Available: <https://groups.csail.mit.edu/mac/users/gjs/6946/linux-install.htm>