The van der Corput trick, and equidistribution on nilmanifolds

Terence Tao · 14 June 2008

https://terrytao.wordpress.com/2008/06/14/the-van-der-corputs-trick-and-equidistribution-on-nilmanifolds/

tao2typ · converted to Typst 2026-10-14 (UTC)

This week I was in Columbus, Ohio, attending a conference on equidistribution on manifolds. I talked about my recent paper with Ben Green on the quantitative behaviour of polynomial sequences in nilmanifolds, which I have blogged about previously. During my talk (and inspired by the immediately preceding talk of Vitaly Bergelson), I stated explicitly for the first time a generalisation of the van der Corput trick which morally underlies our paper, though it is somewhat buried there as we specialised it to our application at hand (and also had to deal with various quantitative issues that made the presentation more complicated). After the talk, several people asked me for a more precise statement of this trick, so I am presenting it here, and as an application reproving an old theorem of Leon Green that gives a necessary and sufficient condition as to whether a linear sequence $(g^n x)_{n=1}^\infty$ on a nilmanifold G/Γ is equidistributed, which generalises the famous theorem of Weyl on equidistribution of polynomials.

UPDATE, Feb 2013: It has been pointed out to me by Pavel Zorin that this argument does not fully recover the theorem of Leon Green; to cover all cases, one needs the more complicated van der Corput argument in our paper.

— The classical van der Corput trick —

The classical van der Corput trick (first used implicitly by Weyl) gives a means to establish the equidistribution of a sequence $(x_n)_{n=1}^\infty$ in a torus $\mathbb{T}^d$ (e.g. a sequence $(P(n) \bmod 1)_{n=1}^\infty$ in the unit circle $\mathbb{T} = \mathbb{R}/\mathbb{Z}$ for some function P, such as a polynomial.) Recall that such a sequence is said to be *equidistributed* if one has

$$\frac{1}{N}\sum_{n=1}^N f(x_n) \to \int_{\mathbb{T}^d} f \tag{1}$$

as $N \to \infty$ for every continuous function $f : \mathbb{T}^d \to \mathbb{C}$; an equivalent formulation of equidistribution is that

$$\frac{1}{N}|\{1 \leq n \leq N : x_n \in B\}| \to \mathrm{vol}(B)$$

for every box B in the torus $\mathbb{T}^d$. (The equivalence can be deduced easily from Urysohn's lemma.) Equidistribution is an important phenomenon to study in ergodic theory and number theory, but also arises in applications such as Monte Carlo integration and pseudorandom number generation.

The fundamental equidistribution theorem of Weyl states that a sequence is equidistributed if and only if the exponential sums

$$\frac{1}{N}\sum_{n=1}^N e^{2\pi i \chi(x_n)} \tag{2}$$

converge to zero for every non-trivial character $\chi : \mathbb{T}^d \to \mathbb{T}$, i.e. a non-zero continuous homomorphism to the unit cicle. Indeed, it is clear that (2) is a special case of (1), and conversely the general case of (1) can be deduced from (2) and either the Weierstrass approximation theorem or basic Fourier analysis.

The significance of the equidistribution theorem is that it reduces the study of equidistribution to the question of estimating exponential sums, which is a problem in analysis and number theory. For

instance, from the equidistribution theorem and the geometric series formula we immediately obtain the following result (stronger than Kronecker's approximation theorem):

> **Corollary 1.** (Equidistribution of linear sequences in torii) Let $\alpha \in \mathbb{T}^d$. Then the sequence $(\alpha n)_{n=1}^\infty$ is equidistributed in $\mathbb{T}^d$ if and only if α is *totally irrational*, which means that $\chi(\alpha) \neq 0$ for all non-zero characters χ.

For instance, the linear sequence $(\sqrt{2}n \mod 1, \sqrt{3}n \mod 1)$ is equidistributed in the two-torus $\mathbb{T}^2$, since $(\sqrt{2}, \sqrt{3})$ is totally irrational, but the linear sequence $(\sqrt{2}n \mod 1, \sqrt{8}n \mod 1)$ is not (the character $\chi : (x,y) \mapsto y - 2x$ annihilates $(\sqrt{2}, \sqrt{8})$ and thus obstructs equidistribution). [Of course, in the latter case, the orbit is still equidistributed in a smaller torus, namely the kernel of the character χ mentioned above; this is an extremely simple case of a much more general result known as Ratner's theorem, which I will not talk further about here.]

One elementary but very useful tool for estimating exponential sums is *Weyl's differencing trick*, that ultimately rests on the humble Cauchy-Schwarz inequality. One formulation of this trick can be phrased as the following inequality:

> **Lemma 1.** (van der Corput inequality) Let $a_1, a_2, \ldots$ be a sequence of complex numbers bounded in magnitude by 1. Then for any $1 \leq H \leq N$ we have
>
> $$\left|\frac{1}{N}\sum_{n=1}^N a_n\right| \ll (\frac{1}{H}\sum_{h=0}^{H-1}\left|\frac{1}{N}\sum_{n=1}^N a_{n+h}\overline{a_n}\right|)^{1/2} + O(\frac{H}{N}). \tag{3}$$

Proof. Observe that

$$\frac{1}{N}\sum_{n=1}^N a_n = \frac{1}{N}\sum_{n=1}^N a_{n+h} + O(\frac{H}{N})$$

for every $0 \leq h \leq H-1$. Averaging this in h we obtain

$$\frac{1}{N}\sum_{n=1}^N a_n = \frac{1}{N}\sum_{n=1}^N \frac{1}{H}\sum_{h=0}^{H-1} a_{n+h} + O(\frac{H}{N})$$

and hence by the Cauchy-Schwarz inequality

$$\left|\frac{1}{N}\sum_{n=1}^N a_n\right| \leq (\frac{1}{N}\sum_{n=1}^N \left|\frac{1}{H}\sum_{h=0}^{H-1} a_{n+h}\right|^2)^{1/2} + O(\frac{H}{N}).$$

Expanding out the square and rearranging a bit, we soon obtain the upper bound (3) (in fact one can sharpen the constants slightly here, though this will not be important for this discussion). □

The significance of this inequality is that it replaces the task of bounding a sum of coefficients a_n by that of bounding a sum of "differentiated" coefficients $a_{n+h}\overline{a_n}$. This trick is thus useful in "polynomial" type situations when the differentiated coefficients are often simpler than the original coefficients. One particularly clean application of this inequality is as follows:

> **Corollary 2.** (Van der Corput's difference theorem) Let $(x_n)_{n=1}^\infty$ be a sequence in a torus $\mathbb{T}^d$ such that the difference sequences $(x_{n+h} - x_n)_{n=1}^\infty$ are equidistributed for every non-zero h. Then $(x_n)_{n=1}^\infty$ is itself equidistributed.

Proof. By Weyl's equidistribution theorem, it suffices to show that (2) holds for every non-trivial character χ. But by Lemma 1, we can bound the magnitude of the left-hand side of (2) by

$$\ll (\frac{1}{H}\sum_{h=0}^{H-1}\left|\frac{1}{N}\sum_{n=1}^N e^{2\pi i\chi(x_{n+h})}e^{-2\pi i\chi(x_n)}\right|)^{1/2} + O(\frac{H}{N}) \tag{4}$$

for any fixed H.

Now we use the fact that χ is a character to simplify $e^{2\pi i\chi(x_{n+h})}e^{-2\pi i\chi(x_n)}$ as $e^{2\pi i\chi(x_{n+h}-x_n)}$. By hypothesis and the equidistribution theorem, the inner sum $\frac{1}{N}\sum_{n=1}^{N} e^{2\pi i\chi(x_{n+h})}e^{-2\pi i\chi(x_n)}$ goes to zero as $N \to \infty$ for any fixed non-zero h; when instead h is zero, this sum is of course just 1. We conclude that for fixed H, the expression (4) is bounded by O(1/H) in the limit $N \to \infty$. Thus the limit (or limit superior) of the magnitude of (2) is bounded in magnitude by O(1/H) for every H, and is thus zero. The claim follows. □

By iterating this theorem, and using the observation that the difference sequence $(P(n+h) - P(n))_{n=1}^{\infty}$ of a polynomial sequence $(P(n))_{n=1}^{\infty}$ of degree d becomes a polynomial sequence of degree d-1 for any non-zero h, we can conclude by induction the following famous result of Weyl, generalising Corollary 1:

> **Theorem 1.** (Equidistribution of polynomial sequences in torii) Let $P : \mathbb{Z} \to \mathbb{T}^d$ be a polynomial sequence taking values in a torus. Then the sequence $(P(n))_{n=1}^{\infty}$ is equidistributed in $\mathbb{T}^d$ if and only if $\chi(P(\cdot))$ is non-constant for all non-zero characters χ.

In the one-dimensional case d=1, this theorem asserts that a polynomial $P : \mathbb{Z} \to \mathbb{R}$ with real coefficients is equidistributed modulo one if and only if it has at least one irrational non-constant coefficient; thus for instance the sequence $(\pi n^3 + \sqrt{2}n^2 + \frac{1}{4}n \bmod 1)_{n=1}^{\infty}$ is equidistributed.

— A variant of the trick —

It turns out that van der Corput's difference theorem (Corollary 2) can be generalised to deal not just on torii, but on more general measure spaces with a torus action. Given a topological probability space (X, μ) (which one should probably take to be a Polish space to avoid various technicalities) and a sequence $(x_n)_{n=1}^{\infty}$ in X, we say that such a sequence is equidistributed with respect to μ if we have

$$\frac{1}{N}\sum_{n=1}^{N} f(x_n) \to \int_X f \, d\mu \tag{5}$$

for all continuous compactly supported functions $f : X \to \mathbb{C}$. This clearly generalises the previous notion of equidistribution, in which X was a torus and μ was uniform probability measure.

To motivate our generalised version of Corollary 2, we observe that the hypothesis "the sequence $(x_{n+h} - x_n)_{n=1}^{\infty}$ is equidistributed in $\mathbb{T}^d$" can be phrased in a more dynamical fashion (eliminating the subtraction operation, which is algebraic) as the equivalent assertion that the sequence of pairs $((x_{n+h}, x_n))_{n=1}^{\infty}$ in $\mathbb{T}^d \times \mathbb{T}^d$, after quotienting out by the action of the diagonal subgroup $(\mathbb{T})^d)^{\Delta} := \{(y, y) : y \in \mathbb{T}^d\}$, becomes equidistributed on the quotient space $\mathbb{T}^d \times \mathbb{T}^d/(\mathbb{T})^d)^{\Delta}$. This convoluted reformulation is necessary for generalisations, in which we do not have a good notion of subtraction, but we still have a good notion of group action and quotient spaces.

We can now prove

> **Proposition 1.** (Generalised van der Corput difference theorem) Let (X, μ) be a (Polish) probability space with a continuous (right-)action of a torus $\mathbb{T}^d$, and let $\pi : X \to X/\mathbb{T}^d$ be the projection map onto the quotient space (which then has the pushforward measure $\pi_*\mu$. Let $(x_n)_{n=1}^{\infty}$ be a sequence in X obeying the following properties:
>
> 1. (Horizontal equidistribution) The projected sequence $(\pi(x_n))_{n=1}^{\infty}$ in $X/\mathbb{T}^d$ is equidistributed with respect to $\pi_*\mu$.
> 2. (Vertical differenced equidistribution) For every non-zero h, the sequence $((x_{n+h}, x_n)(\mathbb{T}^d)^{\Delta})_{n=1}^{\infty}$ in the quotiented product space $(X \times X)/(\mathbb{T}^d)^{\Delta}$ is equidistributed with respect to some measure ν_h which is invariant under the action of the torus $\mathbb{T}^d \times \mathbb{T}^d/(\mathbb{T}^d)^{\Delta}$.
>
> Then $(x_n)_{n=1}^{\infty}$ is equidistributed with respect to μ.

Note that Corollary 2 is the special case of Proposition 1 in which X is itself the torus $\mathbb{T}^d$ with the usual translation action and uniform measure (so that the quotient space is a point).

Proof. We need to verify the property (1). If the function f was invariant under the action of the torus $\mathbb{T}^d$, then we could push it down to the quotient space $X/\mathbb{T}^d$ and the claim would follow from hypothesis 1. We may therefore subtract off the invariant component $\int_{\mathbb{T}^d} f(\cdot\, y)\, dy$ from our function and assume instead that f has zero vertical mean in the sense that $\int_{\mathbb{T}^d} f(xy)\, dy = 0$ for all x. A Fourier expansion in the vertical variable (or the Weierstrass approximation theorem) then allows us to reduce to the case when f has a *vertical frequency* given by some non-zero character $\chi : \mathbb{T}^d \to \mathbb{T}$ of the torus, in the sense that $f(xy) = f(x)e^{2\pi i\chi(y)}$ for all $x \in X$ and $y \in \mathbb{T}^d$.

Now we apply van der Corput's inequality as in the proof of Corollary 2. Using these arguments, we find that it suffices to show that

$$\frac{1}{N}\sum_{n=1}^{N} f(x_{n+h})\overline{f(x_n)} \to 0$$

for each non-zero h. But the summand here is just the tensor product function $f \otimes \overline{f} : X \times X \to \mathbb{C}$ applied to the pair (x_{n+h}, x_n). The fact that f has a vertical frequency implies that $f \otimes \overline{f}$ is invariant with respect to the diagonal action $(\mathbb{T}^d)^\Delta$, and thus this function descends to the quotient space $(X \times X)/(\mathbb{T}^d)^\Delta$. On the other hand, as the vertical frequency is non-trivial, the latter function also has zero mean on every orbit of $\mathbb{T}^d \times \mathbb{T}^d/(\mathbb{T}^d)^\Delta$ and thus vanishes when integrated against ν_h. The claim then follows from hypothesis 2. □

As an application, let us prove the following result, first established by (Leon) Green:

> **Theorem 2.** (Equidistribution of linear sequences in nilmanifolds) Let G/Γ be a nilmanifold (where we take the nilpotent group G to be connected for simplicity, although this is not strictly necessary), and let $g \in G$ and $x \in G/\Gamma$. Then $(g^n x)_{n=1}^\infty$ is equidistributed with respect to Haar measure on G/Γ if and only if $\chi(g^n x)$ is non-constant in n for every non-trivial *horizontal character* $\chi : G/\Gamma \to \mathbb{T}$, where a horizontal character is any continuous homomorphism $\chi : G \to \mathbb{T}$ that vanishes on Γ (and thus descends to G/Γ).

This statement happens to contain Weyl's result (Theorem 1) as a special case, because polynomial sequences can be encoded as linear sequences in nilmanifolds; but it is actually stronger, allowing extensions to generalised polynomials that involve the floor function $\lfloor\cdot\rfloor$ or the fractional part function $\{\}$. For instance, if we take

$$G := \begin{pmatrix} 1 & \mathbb{R} & \mathbb{R} \\ 0 & 1 & \mathbb{R} \\ 0 & 0 & 1 \end{pmatrix}; \Gamma := \begin{pmatrix} 1 & \mathbb{Z} & \mathbb{Z} \\ 0 & 1 & \mathbb{Z} \\ 0 & 0 & 1 \end{pmatrix}$$

and

$$g := \begin{pmatrix} 1 & \alpha & \beta \\ 0 & 1 & \gamma \\ 0 & 0 & 1 \end{pmatrix}; x = \Gamma$$

for some real numbers α, β, γ then a computation shows that

$$g^n x = \begin{pmatrix} 1 & \{\alpha n\} & \{\beta n + \alpha\frac{n(n-1)}{2} - \{\alpha n\}\lfloor \gamma n\rfloor\} \\ 0 & 1 & \{\gamma n\} \\ 0 & 0 & 1 \end{pmatrix}\Gamma$$

and then Green's theorem asserts that the triple

$$(\{\alpha n\}, \{\beta n + \alpha\frac{n(n-1)}{2} - \{\alpha n\}\lfloor \gamma n\rfloor\}, \{\gamma n\})$$

is equidistributed in the unit cube $[0,1]^3$ if and only if the pair (α, γ) is totally irrational (the rationality of β turns out to be irrelevant). Even for concrete values such as $\alpha = \sqrt{2}, \beta = 0, \gamma = \sqrt{3}$, it is not obvious how to establish this fact directly; for instance a direct application of Corollary 2 does not obviously simplify the situation.

Proof of Theorem 2. (Sketch) It is clear that if $\chi(g^n x)$ is constant for some non-trivial character, then the orbit $g^n x$ is trapped on a level set of χ and thus cannot equidistribute. Conversely, suppose that $\chi(g^n x)$ is never constant. We induct on the step s of the nilmanifold. The case s=0 is trivial, and the case s=1 follows from Corollary 1, so suppose inductively that $s \geq 2$ and that the claim has already been proven for smaller s. We then look at the vertical torus $G_s/(\Gamma \cap G_s) \equiv \mathbb{T}^d$, where G_s is the last non-trivial group in the lower central series (and thus central). The quotient of the nilmanifold G/Γ by this torus action turns out to be a nilmanifold of one lower step (in which G is replaced by G/G_s) and so the projection of the orbit $(g^n x)_{n=1}^{\infty}$ is then equidistributed by induction hypothesis. Applying Proposition 1, it thus suffices to check that for each non-zero h, the sequence of pairs $(g^{n+h}x, g^n x)$ in $G/\Gamma \times G/\Gamma$, after quotienting out by the diagonal action of the torus, is equidistributed with respect to some measure which is invariant under the residual torus $\mathbb{T}^d \times \mathbb{T}^d/(\mathbb{T}^d)^{\Delta}$.

We first pass to the abelianisation (or *horizontal torus*) $G/G_2\Gamma$ of the nilmanifold, and observe that the projections $\pi(g^{n+h}x), \pi(g^n x)$ of the coefficients of the pair $(g^{n+h}x, g^n x)$ to this torus only differ by a constant $\pi(g^h)$. Thus the pair $(g^{n+h}x, g^n x)$ does not range freely in $G/\Gamma \times G/\Gamma$, but is instead constrained to a translate of a smaller nilmanifold $G/\Gamma \times_{\pi} G/\Gamma$, defined as the space of pairs (x,y) with $\pi(x) = \pi(y)$. After quotienting out also by the diagonal vertical torus, we obtain a nilmanifold coming from the group $(G \times_{G_2} G)/G_s^{\Delta}$, where $G \times_{G_2} G$ is the space of pairs (g,h) of group elements $g, h \in G$ whose projections to the abelianisation G/G_2 agree, and $G_s^{\Delta} := \{(g_s, g_s) : g_s \in G_s\}$ is the vertical diagonal group. But a short computation shows that this new group is at most s-1 step nilpotent. One can then apply the induction hypothesis to show the required equidistribution properties of (x_{n+h}, x_n), thus closing the induction by Proposition 1. [UPDATE, Feb 2013: This doesn't work in all cases, because sometimes the orbit $(g^{n+h}x, g^n x)$ is not equidistributed in the abelianisation of this nilmanifold.] □

There are many further generalisations of these results, including a polynomial version of Theorem 2 due to Leibman (that also permits G to be disconnected), and quantitative versions of all of these results in my paper with Ben Green that I discuss in my earlier blog post.

Comments (25)

 **A** · 16 June 2008

It's a nitpick, but the statement of corollary 2 isn't parsing correctly.

 **Terence Tao** · 16 June 2008

Thanks for the correction!

 **Marker lectures II, "Linear equations in primes" « What's new** · 18 November 2008

[...] turns out to be somewhat unpleasant; the standard technology of Weyl differencing and the van der Corput lemma eventually works (see this paper of Ben Green and myself), but does not scale well to bracket [...]

ERT2: Polynomial Von Neumann's Theorem « Disquisitiones Mathematicae · 25 October 2009

[...] true, by the induction hypothesis). This is done with the use of Van der Corput's Trick (see this lecture of Terry Tao for a broader discussion on this trick). Theorem 3 (Van der Corput Trick) If is a bounded sequence [...]

ERT9: Weak Mixing implies Weak Mixing of all orders « Disquisitiones Mathematicae · 2 March 2010

[...] van der Corput trick allows an inductive [...]

 **Nilotpal Sinha** · 24 October 2010

A remark on Corollary 1.
Reading this post reminded me of an old observation of mine of which I thought would be fit to share in this post on equidistribution.
Let u(n) be any sequence such that for every in (0,1), u(nt) ~ t u(n). If α is any real such that αu(n) and n are linearly independent then the sequence αu(n) is equidistributed mod 1.
Applications:
Eg 1. Take u(n) = n and α irrational. Then the above remark implies that the sequence αn is equidistributed mod 1. This was the original equidistribution theorem of Weyl in 1909.
Eg 2. Take u(n) = p(n) where p(n) is the n-th prime. Then our remark implies that the sequence α p(n) is equidistributed mod 1. This was the result of I.M. Vinogradov, 1935.
Taking advantage of the liner independence condition in the remark, we have have similar results for rational numbers too.
Eg 3. Taking α = 1 and u(n) ~ 2Πn/log(n) we see that the imaginary parts of the non-trivial zeta zeros are equidistributed mod 1. This was the result of Hlawka.
In short the remark looks very powerful and if applied in the right places, it can yield many interesting results.

 **Walsh's ergodic theorem, metastability, and external Cauchy convergence « What's new** · 25 October 2012

[...] for . (We will see why these particular functions arise in the argument shortly.) The key step in proving Theorem 8 is then the following result, reminiscent of the van der Corput lemma in ergodic theory (see e.g. this blog post). [...]

 **Sean Eberhard** · 4 October 2013

Dear Prof Tao,
I'm trying to understand the proof of Theorem 2, whether the sketch you give here or a proof in the literature somewhere. I thought I understood the proof until I read your update from Feb 2013. It suffices to check that $(g^n x, g^n x)$ equidistributes in the nilmanifold coming from $(G \times_{G_2} G)/G_s^{\Delta}$: it would then follow that $(g^{n+h} x, g^n x)$ is equidistributed in a translate of this nilmanifold. Since this

nilmanifold has step $s-1$, if this fails it follows by induction that there is a nontrivial horizontal character χ of $(G \times_{G_2} G)/G_s^\Delta$ such that $\chi(g^n x, g^n x)$ is constant. But then by precomposing with the diagonal embedding $\Delta : G \to G \times_{G_2} G$ we get a nontrivial character $\chi \circ \Delta$ of G such that $\chi(\Delta(g^n x))$ is constant, contradicting our hypothesis. Doesn't this work?



Terence Tao · 4 October 2013

Unfortunately, the non-triviality of χ does not imply the non-triviality of $\chi \circ \Delta$,and in any event $(g^n x, g^n x)$ is clearly trapped in the diagonal $\Delta(G)/G_s^\Delta$ of $G \times_{G_2} G/G_s^\Delta$ and so cannot equidistribute. (A similar problem was pointed out to me by Pavel Zorin in https://terrytao.wordpress.com/2010/05/29/254b-notes-6-the-inverse-conjecture-for-the-gowers-norm-ii-the-integer-case/#comment-217771 , which prompted the above correction.)



Sean Eberhard · 4 October 2013

I see. The point that was getting me is that $G \times_{G_2} G/G_s^\Delta$ typically has abelianisation larger than just G/G_2. Where is the best place to find the missing details?



Terence Tao · 4 October 2013

Unfortunately I do not know of a quick way to make the strategy outlined in this blog post to work in full generality; one either has to work in a fully quantitative setting (as in my paper with Ben, http://arxiv.org/abs/0709.3562) or else work in an ergodic theory setting (taking full advantage of the ergodic theorem) as in the original paper of Leibman, http://www.ams.org/mathscinet-getitem?mr=2122919 .



Cyril · 1 July 2014

Hello. I am a beginner in the theory you are dealing with. However I think there is a link with an old problem I have for a long time:
Is the sequence $\left(\sum_{k=0}^{n} \cos(k^2)\right)_n$ bounded ?



Terence Tao · 1 July 2014

No, it is unbounded. If it were bounded, then the sequence $n \mapsto \left|\sum_{k=0}^{K} \cos((n+k)^2)\right|^2$ would be bounded uniformly in K, but from Weyl's theorem and the irrationality of π, the asymptotic mean of this sequence is comparable to K, a contradiction.



Cyril · 1 July 2014

Thank you very much but I don't really understand. Can you give me a link to the Weyl's theorem you are talking about ?
When you say "bounded uniformly in K" you mean that it exists a constant C such that for all K, for all n, the sequence is bounded by C ?



Terence Tao · 2 July 2014

Yes.
Weyl's theorem can be found for instance in Corollary 6 of my notes https://terrytao.wordpress.com/2010/03/28/254b-notes-1-equidistribution-of-polynomial-sequences-in-torii/ . The point is that when one expands out the square in the indicated sequence in terms of complex exponentials, one gets terms such as $e(\frac{1}{2\pi}((n+k)^2 - (n+k')^2))$ for various k, k'. For $k \neq k'$, the phase here contains an irrational nonconstant coefficient and so has asymptotic mean zero in n. If one worked with $\cos(k)$ instead of $\cos(k^2)$ then $e(\frac{1}{2\pi}((n+k) - (n+k')))$ would instead be constant in n and Weyl's theorem would not apply.



Cyril · 2 July 2014

Ouawww I got it !!!! Thank you very much ! I love this trick.



Eytan Paldi · 2 July 2014

Thanks! I expected this question to be much more difficult – because it seems related to the complicated behavior of the (theta) function
$\varphi(q) := 1 + 2\sum_{k\geq 1} q^{k^2}$, $|q| < 1$
near the irrational point e^i on the unit circle – its natural boundary!
The function $\varphi(q)$ (defined e.g. in page 36, Entry 22(i) in Berndt's book "Ramanujan Notebooks" Part III, 1991)
From the proof of Entry 23(i) (page 38 in this book) it follows that
$\varphi(-q) = \prod_{k\geq 1} \frac{1-q^k}{1+q^k}$
$= \frac{F(q^2)}{F(q)^2}$
Where $F(q) := \prod_{k\geq 1}(1-q^k)^{-1}$ is the generating function for the partition function $p(n)$.
Therefore, $\varphi(q)$ behavior near the unit circle may be studied via the well known complicated behavior of $F(q)$ there (in particular, it follows that $\varphi(q)$ has the unit circle as its natural boundary!)



Eytan Paldi · 2 July 2014

It is not clear why this argument is working for $\cos(k^2)$ but not for $\cos k$.



Cyril · 2 July 2014

That's why I'm asking the reference of the theorem.



Anonymous · 3 July 2014

In your paper, I think there is a typo in line 2 in the introduction. Shouldn't it be "role" rather than "rôle"? (English is not my first language.)



Anonymous · 3 July 2014

I see that I'm wrong. (Please delete this and the previous comment.)

Frogs and Lily Pads and Discrepancy | Gödel's Lost Letter and P=NP · 24 September 2015

[...] The connection with products of the form is analyzed by an old trick which Tao once discussed here. There is finally an important use of Andrew Granville's notion of pretending, which we once [...]



Anonymous · 14 September 2017

Dear Prof. Tao,
I am trying to understand the proof of Proposition 1. I got stuck in the first part of the proof, where you claim that we can assume the continuous function f has the property that $f(xy) = f(x)e^{2\pi i\chi(y)}$ for all $x \in X$ and $y \in \mathbb{T}^d$, where χ is some non-zero character of the torus. I assume this is true because functions of this kind generate an algebra that separates points and hence, by Stone-Weierstrass (like you suggested in the proof), this algebra is dense in $C(X)$. However, I am stuck on proving that this family actually separates points. Can you please clarify the argument that I am missing here. Thank you very much!



Terence Tao · 14 September 2017

If x_1, x_2 do not lie in the same $\mathbb{T}^d$-orbit, one can take a bump function ϕ supported in a small neighbourhood of x_1 and average it along the action to obtain a suitable function $\int_{\mathbb{T}^d} \phi(xy)\ dy$ to separate x_1, x_2. If instead x_2 lies in the orbit of x_1, then the set $\{y \in \mathbb{T}^d : x_2 = x_1 y\}$ is a non-trivial coset $y_0 + H$ of some closed subgroup H of $\mathbb{T}^d$. One can then locate a frequency $k \in \mathbb{Z}^d$ which annihilates H but not y_0 (because the characters on $\mathbb{T}^d/H$ separate points, by Plancherel's theorem), and then the modulated average $\int_{\mathbb{T}^d} \phi(xy) e(k \cdot y)\ dy$ will work if the support of ϕ is small enough.



Anonymous · 6 January 2025

For a particular problem coming from information theory I was asked to check your paper 0709.3562 for "quantitative (or effective) equidistribution," (in particular Theorem 1.16 (Quantitative Leibman theorem)). What background do I need to build to this level? My background is in Electrical Engineering and taken engineering versions of Fourier transforms. Essentially I was told to convert 1.16 to statement about sequences being dense in the torus. Thank you.