A small remark on the Elliott conjecture

Terence Tao · 30 October 2015

https://terrytao.wordpress.com/2015/10/30/a-small-remark-on-the-elliott-conjecture/

tao2typ · converted to Typst 2026-10-14 (UTC)

The Chowla conjecture asserts, among other things, that one has the asymptotic

$$\frac{1}{X}\sum_{n\leq X}\lambda(n+h_1)\ldots\lambda(n+h_k) = o(1)$$

as $X\to\infty$ for any distinct integers $h_1,\ldots,h_k$, where λ is the Liouville function. (The usual formulation of the conjecture also allows one to consider more general linear forms $a_i n + b_i$ than the shifts $n+h_i$, but for sake of discussion let us focus on the shift case.) This conjecture remains open for $k\geq 2$, though there are now some partial results when one averages either in x or in the $h_1,\ldots,h_k$, as discussed in this recent post.

A natural generalisation of the Chowla conjecture is the Elliott conjecture. Its original formulation was basically as follows: one had

$$\frac{1}{X}\sum_{n\leq X} g_1(n+h_1)\ldots g_k(n+h_k) = o(1) \tag{1}$$

whenever $g_1,\ldots,g_k$ were bounded completely multiplicative functions and $h_1,\ldots,h_k$ were distinct integers, and one of the g_i was "non-pretentious" in the sense that

$$\sum_p \frac{1-\mathrm{Re}(g_i(p)\overline{\chi(p)}p^{-it})}{p} = +\infty \tag{2}$$

for all Dirichlet characters χ and real numbers t. It is easy to see that some condition like (2) is necessary; for instance if $g(n) := \chi(n)n^{it}$ and χ has period q then $\frac{1}{X}\sum_{n\leq X} g(n+q)\overline{g(n)}$ can be verified to be bounded away from zero as $X\to\infty$.

In a previous paper with Matomaki and Radziwill, we provided a counterexample to the original formulation of the Elliott conjecture, and proposed that (2) be replaced with the stronger condition

$$\inf_{|t|\leq X}\sum_{p\leq X}\frac{1-\mathrm{Re}(g_i(p)\overline{\chi(p)}p^{-it})}{p} \to +\infty \tag{3}$$

as $X\to\infty$ for any Dirichlet character χ. To support this conjecture, we proved an averaged and non-asymptotic version of this conjecture which roughly speaking showed a bound of the form

$$\frac{1}{H^k}\sum_{h_1,\ldots,h_k\leq H}\left|\frac{1}{X}\sum_{n\leq X} g_1(n+h_1)\ldots g_k(n+h_k)\right| \leq \varepsilon$$

whenever H was an arbitrarily slowly growing function of X, X was sufficiently large (depending on ε, k and the rate at which H grows), and one of the g_i obeyed the condition

$$\inf_{|t|\leq AX}\sum_{p\leq X}\frac{1-\mathrm{Re}(g_i(p)\overline{\chi(p)}p^{-it})}{p} \geq A \tag{4}$$

for some A that was sufficiently large depending on k,ε, and all Dirichlet characters χ of period at most A. As further support of this conjecture, I recently established the bound

$$\frac{1}{\log \omega} \left| \sum_{X/\omega \leq n \leq X} \frac{g_1(n+h_1) g_2(n+h_2)}{n} \right| \leq \varepsilon$$

under the same hypotheses, where ω is an arbitrarily slowly growing function of X.

In view of these results, it is tempting to conjecture that the condition (4) for one of the g_i should be sufficient to obtain the bound

$$\left| \frac{1}{X} \sum_{n \leq X} g_1(n+h_1) \dots g_k(n+h_k) \right| \leq \varepsilon$$

when A is large enough depending on k, ε. This may well be the case for $k = 2$. However, the purpose of this blog post is to record a simple counterexample for $k > 2$. Let's take $k = 3$ for simplicity. Let t_0 be a quantity much larger than X but much smaller than X^2 (e.g. $t = X^{3/2}$), and set

$$g_1(n) := n^{it_0}; \quad g_2(n) := n^{-2it_0}; \quad g_3(n) := n^{it_0}.$$

For $X/2 \leq n \leq X$, Taylor expansion gives

$$(n+1)^{it} = n^{it_0} \exp(it_0/n) + o(1)$$

and

$$(n+2)^{it} = n^{it_0} \exp(2it_0/n) + o(1)$$

and hence

$$g_1(n) g_2(n+1) g_3(n+2) = 1 + o(1)$$

and hence

$$\left| \frac{1}{X} \sum_{X/2 \leq n \leq X} g_1(n) g_2(n+1) g_3(n+2) \right| \gg 1.$$

On the other hand one can easily verify that all of the g_1, g_2, g_3 obey (4) (the restriction $|t| \leq AX$ there prevents t from getting anywhere close to t_0). So it seems the correct non-asymptotic version of the Elliott conjecture is the following:

> **Conjecture 1 (Non-asymptotic Elliott conjecture)** Let k be a natural number, and let $h_1, \dots, h_k$ be integers. Let $\varepsilon > 0$, let A be sufficiently large depending on $k, \varepsilon, h_1, \dots, h_k$, and let X be sufficiently large depending on $k, \varepsilon, h_1, \dots, h_k, A$. Let $g_1, \dots, g_k$ be bounded multiplicative functions such that for some $1 \leq i \leq k$, one has
>
> $$\inf_{|t| \leq AX^{k-1}} \sum_{p \leq X} \frac{1 - \mathrm{Re}(g_i(p) \overline{\chi(p)} p^{-it})}{p} \geq A$$
>
> for all Dirichlet characters χ of conductor at most A. Then
>
> $$\left| \frac{1}{X} \sum_{n \leq X} g_1(n+h_1) \dots g_k(n+h_k) \right| \leq \varepsilon.$$

The $k = 1$ case of this conjecture follows from the work of Halasz; in my recent paper a logarithmically averaged version of the $k = 2$ case of this conjecture is established. The requirement to take t to be as large as AX^{k-1} does not emerge in the averaged Elliott conjecture in my previous paper with Matomaki and Radziwill; it thus seems that this averaging has concealed some of the subtler features of the Elliott conjecture. (However, this subtlety does not seem to affect the asymptotic version of the conjecture formulated in that paper, in which the hypothesis is of the form (3), and the conclusion is of the form (1).)

A similar subtlety arises when trying to control the maximal integral

$$\frac{1}{X}\int_X^{2X} \sup_\alpha \frac{1}{H} |\sum_{x\leq n\leq x+H} g(n)e(\alpha n)|\ dx. \tag{5}$$

In my previous paper with Matomaki and Radziwill, we could show that easier expression

$$\frac{1}{X}\sup_\alpha \int_X^{2X} \frac{1}{H} |\sum_{x\leq n\leq x+H} g(n)e(\alpha n)|\ dx. \tag{6}$$

was small (for H a slowly growing function of X) if g was bounded and completely multiplicative, and one had a condition of the form

$$\inf_{|t|\leq AX} \sum_{p\leq X} \frac{1-\mathrm{Re}(g(p)\overline{\chi(p)}p^{-it})}{p} \geq A \tag{7}$$

for some large A. However, to obtain an analogous bound for (5) it now appears that one needs to strengthen the above condition to

$$\inf_{|t|\leq AX^2} \sum_{p\leq X} \frac{1-\mathrm{Re}(g(p)\overline{\chi(p)}p^{-it})}{p} \geq A$$

in order to address the counterexample in which $g(n) = n^{it_0}$ for some t_0 between X and X^2. This seems to suggest that proving (5) (which is closely related to the $k=3$ case of the Chowla conjecture) could in fact be rather difficult; the estimation of (6) relied primarily of prior work of Matomaki and Radziwill which used the hypothesis (7), but as this hypothesis is not sufficient to conclude (5), some additional input must also be used.

Comments (4)

 **Anonymous** · 31 October 2015

It seems that the generalized non pretentious restriction in conjecture 1, was constructed to avoid the above counterexample (for which $g_1 \cdots g_k = 1$). Is it possible to relax this restriction by adding an appropriate restriction on the product $g_1 \cdots g_k$ (e.g. to be $\neq 1$) ?

 **Terence Tao** · 1 November 2015

At a bare minimum, one will need $g_1 ... g_k$ to be non-pretentious. For instance suppose that $g_1 = ... = g_1 = \chi$ is a Dirichlet character. Then the function $n \mapsto g_1(n+h_1)...g_k(n+h_k)$ will be periodic and will typically not have mean zero (one expects a square root cancellation, but not perfect cancellation). In this case $g_1 ... g_k$ will be a Dirichlet character which will usually not be principal. One can also multiply one or more of the g_i by $n \mapsto n^{it}$ for some small t and still have some non-cancellation in the sum $\sum_{n \leq x} g_1(n+h_1)...g_k(n+h_k)$.

In principle, once one is in the "structured" case where each of the g_i pretends to be a modulated Dirichlet character $n \mapsto \chi_i(n) n^{it_i}$, one can write g_i as a Dirichlet convolution of this modulated character with another arithmetic function that pretends to be 1, and one can evaluate the sum $\sum_{n \leq x} g_1(n+h_1)...g_k(n+h_k)$ more or less explicitly (though to deal with some error terms one may need something like a generalised Elliott-Halberstam conjecture, or else do some smoothing in the x variable). This should in principle let one refine the condition yielding cancellation in the sum $\sum_{n \leq x} g_1(n+h_1)...g_k(n+h_k)$ to obtain a near-optimal criterion for this cancellation. (This is analogous to the situation involving finding linear patterns in dense sets, where there is a crude "Cauchy-Schwarz complexity" criterion that handles all the cases in which a certain high order Gowers norm is small, but then there is a more refined "true complexity" criterion that uses some more advanced additive combinatorics (such as the equidistribution theory of nilsequences) to find a more optimal criterion to get equidistribution.)

 **sylvainjulien** · 31 October 2015

Am I wrong, or is a closing parenthesis missing in the expression of the "non-pretentious" quantity?

 **sylvainjulien** · 31 October 2015

Ok, it's been corrected, my apologies.