What's new

A counterexample to a strong polynomial Freiman-Ruzsa conjecture

Terence Tao · 9 November 2008

https://terrytao.wordpress.com/2008/11/09/a-counterexample-to-a-strong-polynomial-freiman-ruzsa-conjecture/

tao2typ · converted to Typst 2026-10-14 (UTC)

One of my favourite open problems in additive combinatorics is the polynomial Freiman-Ruzsa conjecture, which Ben Green guest blogged about here some time ago. It has many equivalent formulations (which is always a healthy sign when considering a conjecture), but here is one involving "approximate homomorphisms":

> **Polynomial Freiman-Ruzsa conjecture.** Let $f : F_2^n \to F_2^m$ be a function which is an approximate homomorphism in the sense that $f(x+y) - f(x) - f(y) \in S$ for all $x, y \in F_2^n$ and some set $S \subset F_2^m$. Then there exists a genuine homomorphism $g : F_2^n \to F_2^m$ such that $f - g$ takes at most $O(|S|^{O(1)})$ values.

Remark 1. The key point here is that the bound on the range of $f - g$ is at most polynomial in |S|. An exponential bound of $2^{|S|}$ can be trivially established by splitting F_2^m into the subspace spanned by S (which has size at most $2^{|S|}$) and some complementary subspace, and then letting g be the projection of f to that complementary subspace. ◇

Recently, Ben Green and I have shown that this conjecture is equivalent to a certain polynomially quantitative strengthening of the inverse conjecture for the Gowers norm $U^3(F_2^n)$; I hope to talk about this in a future post. For this (somewhat technical) post, I want to comment on a possible further strengthening of this conjecture, namely

> **Strong Polynomial Freiman-Ruzsa conjecture.** Let $f : F_2^n \to F_2^m$ be a function which is an approximate homomorphism in the sense that $f(x+y) - f(x) - f(y) \in S$ for all $x, y \in F_2^n$ and some set $S \subset F_2^m$. Then there exists a genuine homomorphism $g : F_2^n \to F_2^m$ such that $f - g$ takes values in the sumset $CS := S + \ldots + S$ for some fixed $C = O(1)$.

This conjecture is known to be true for certain types of set S (e.g. for Hamming balls, this is a result of Farah). Unfortunately, it is false in general; the purpose of this post is to describe one counterexample (related to the failure of the inverse conjecture for the Gowers norm for $U^4(F_2^n)$ for classical polynomials; in particular, the arguments here have several features in common with those in the papers of Lovett-Meshulam-Samorodnitsky and Green-Tao). [A somewhat different counterexample also appears in the paper of Farah.] The verification of the counterexample is surprisingly involved, ultimately relying on the multidimensional Szemerédi theorem of Furstenberg and Katznelson.

(The results here are derived from forthcoming joint work with Ben Green.)

— Description of counterexample —

We let n be a large number, and replace F_2^m by the $\frac{n(n+1)}{2}$-dimensional vector space V of quadratic forms $Q : F_2^n \to F_2$ (with a basis given by the monomials $x_i x_j$ with $1 \leq i \leq j \leq n$). We let $f : F_2^n \to V$ be defined by the formula

$$f(h_1, ..., h_n)(x_1, ..., x_n) := \sum_{1 \leq i < j \leq n} h_i x_i h_j x_j.$$

A brief computation shows that for any $h, k \in F_2^n$, the quadratic form $f(h+k) - f(h) - f(k)$ is of rank at most three, by which we mean that it is a function of at most three linear forms. More specifically, we have

$$f(h+k) - f(h) - f(k) = a_{h,k} b_{h,k} + b_{h,k} c_{h,k} + c_{h,k} a_{h,k} \tag{1}$$

where

$$a_{h,k}(x) = \sum_{i=1}^{n} h_i (1 - k_i) x_i$$

$$b_{h,k}(x) = \sum_{i=1}^{n} (1 - h_i) k_i x_i$$

$$c_{h,k}(x) = \sum_{i=1}^{n} h_i k_i x_i$$

Thus, if we let S be the space of quadratic forms of rank at most 3, the hypotheses of the polynomial Freiman-Ruzsa conjecture hold.

— Verification of counterexample —

To establish the counterexample, we assume for contradiction that there exists a linear function $g : F_2^n \to V$ such that $f(h) - g(h)$ has bounded rank for all h, and deduce a contradiction (for n sufficiently large).

By hypothesis, we have linear forms $L_{h,1}, ..., L_{h,d}$ for all $h \in F_2^n$ and some $d = O(1)$ and coefficients $c_{h,i,j} \in F_2$ for all $1 \leq i \leq j \leq d$ such that

$$f(h) - g(h) = \sum_{1 \leq i \leq j \leq n} c_{h,i,j} L_{h,i} L_{h,j}$$

and in particular (by (1) and linearity of g)

$$a_{h,k} b_{h,k} + b_{h,k} c_{h,k} + c_{h,k} a_{h,k}$$

$$= \sum_{1 \leq i \leq j \leq n} c_{h+k,i,j} L_{h+k,i} L_{h+k,j} - c_{h,i,j} L_{h,i} L_{h,j} - c_{k,i,j} L_{k,i} L_{k,j}. \tag{2}$$

The key point is that the linear forms $a_{h,k}, b_{h,k}, c_{h,k}$ are usually "independent" of the linear forms $L_{h,i}, L_{k,i}, L_{h+k,i}$. The key lemma in this regard is

> **Lemma 1.** If h, k are selected uniformly and independently at random, then with probability 1-o(1), $a_{h,k}$ is not a linear combination of the $L_{h,i}, L_{k,i}, L_{h+k,i}$. Similarly for $b_{h,k}, c_{h,k}$.

Proof. By cyclically permuting h,k,h+k it suffices to show this for $c_{h,k}$. Since there are at most O(1) possible linear combinations amongst the $L_{h,i}, L_{k,i}, L_{h+k,i}$, it suffices to show that for any given assignments $h \mapsto L'_h, k \mapsto L''_k, h+k \mapsto L'''_{h+k}$ of linear forms, that the probability of the event

$$c_{h,k} = L'_h + L''_k + L'''_{h+k} \tag{3}$$

is o(1). Suppose for contradiction that the event (3) holds for a set E of pairs (h,k) in $F_2^n \times F_2^n$ of positive density. Applying the Furstenberg-Katznelson multidimensional Szemerédi theorem) we can find (for n large enough) a square $(h,k), (h+r,k), (h,k+r), (h+r,k+r)$ in E with r non-zero. Applying (3) for all four pairs and summing, we obtain

$$c_{h,k} + c_{h+r,k} + c_{h,k+r} + c_{h+r,k+r} = 0$$

(recall we are in characteristic 2). But the left-hand side is equal to the linear form $\sum_i r_i x_i$, which is non-zero, a contradiction. □

Now we can obtain the desired contradiction. For a generic choice of h,k, we now know that none of the $a_{h,k}, b_{h,k}, c_{h,k}$ are linear combinations of the $L_{h,i}, L_{k,i}, L_{h+k,i}$. Thus, on a given level set of the $L_{h,i}, L_{k,i}, L_{h+k,i}$ (which form a subspace of F_2^n), the linear functions $a_{h,k}, b_{h,k}, c_{h,k}$ are non-constant, and so the range of the triplet $(a_{h,k}, b_{h,k}, c_{h,k})$ must be an affine subspace of F_2^3 which is not contained in any plane in F_2^3 that is parallel to the two of the coordinate axes (i.e. of the form $a = const$, $b = const$, or $c = const$. This forces the subspace to be parallel to (1,1,1). But this implies that $(a,b,c) \mapsto ab + bc + ca$ is non-constant on this space, contradicting (2).

Remark 2. The function f appearing in the above example is closely related to the symmetric polynomial

$$S_4(x) := \sum_{1 \leq i < j < k < l \leq n} x_i x_j x_k x_l.$$

Indeed, one can show that the derivative $S_4(x+h) - S_4(x)$ of S_4 is equal to $f(h)$, plus some additional terms which involve only a finite number of linear forms, and the quadratic polynomial $S_2(x) := \sum_{1 \leq i < j \leq n} x_i x_j$. If it was the case that f could be approximated by a linear map g modulo low rank errors, then it one could use this to eventually show that S_4 correlated with a cubic polynomial; but it is known (from the papers of Lovett-Meshulam-Samorodnitsky and Green-Tao) that this is not the case. Thus there is an alternate way to verify that the above example is indeed a counterexample to the strong polynomial Freiman-Ruzsa conjecture. ◇

Comments (7)



bengreen · 11 November 2008

Terry,

I know that your blog has a wide readership and so it would be irresponsible of me to endorse gambling in any way. However I should perhaps mention the informal bet I had with Jean Bourgain, whereby he wins 100 dollars from me if a counterexample to PFR is found and I win the same if a proof of the conjecture is found within 5 years.

This counterexample to the `strong' form of the conjecture (which I knew about before making the bet....) does suggest that a proof of the statement might have to be a little weird.

I'd like to take this rare opportunity to use the word `cohomology' on your website (or indeed at all). Emmanuel Breuillard pointed out to me that there is this thing called bounded cohomology, which roughly speaking measures the obstruction to maps $f : G \to \mathbb{R}$ with $|f(xy) - f(x)f(y)| \leq C$ being a bounded distance from genuine homomorphisms. Apparently quite a lot of the theory is developed in a paper of Gromov which I have yet to obtain. See also an article by Kotschick in Notices of the AMS, vol 51. no .2 called "What is a quasimorphism"?

Perhaps `Ruzsa cohomology' can resolve these issues but suddenly 5 years looks like a very short period of time :-)

Ben

 **Harald** · 9 March 2009

Hi –

I think there are a couple of mistakes. The definitions of a_{h,k}, b_{h,k} and c_{h,k} after equation (1) can't be right. What are the right polynomials? In the section right thereafter, all occurences of L_{h,d}, L_{k,d}, L_{h+k,d} should be
L_{h,j},L_{k,j}, L_{h+k,j} instead.

 **Terence Tao** · 9 March 2009

Thanks for the correction! The $b_{h,k}$ was missing a k_i in their definition; I've fixed it now.

 **Harald** · 9 March 2009

Also – what does "any coordinate plane" mean? Does that mean the xy, yz, xz planes? If so, how does the fact that the range of (a_{h,k},b_{h,k},c_{h,k}) is not contained in one of those imply that it is at least two-dimensional?

Harald

 **Terence Tao** · 10 March 2009

Dear Harald,

By coordinate plane I meant the planes in F_2^3 in which x, y, or z is held constant. The only one-dimensional spaces which are not in coordinate planes are those lines in the direction (1,1,1), but ab+bc+ca is not constant on those lines. I'll reword the argument there to make it a bit clearer.

 **An equivalence between inverse sumset theorems and inverse conjectures for the U^3 norm « What's new** · 20 June 2009

[...] Math. Proc. Camb. Phil. Soc.. The main result of this short paper (which was briefly announced in this earlier post) is a connection between two types of inverse theorems in additive combinatorics, namely the [...]



Metric entropy analogues of sum set theory | What's new · 19 March 2014

[…] due to the failure of a certain strong version of Freiman's theorem, as discussed in this previous post); nevertheless, the proofs of the discrete arguments are elementary enough that they can be […]