Covering a non-closed interval by disjoint closed intervals

Terence Tao · 4 October 2010

https://terrytao.wordpress.com/2010/10/04/covering-a-non-closed-interval-by-disjoint-closed-intervals/

tao2typ · converted to Typst 2026-10-14 (UTC)

The following question came up in my 245A class today:

> Is it possible to express a non-closed interval in the real line, such as [0,1), as a countable union of disjoint closed intervals?

I was not able to answer the question immediately, but by the end of the class some of the students had come up with an answer. It is actually a nice little test of one's basic knowledge of real analysis, so I am posing it here as well for anyone else who is interested. Below the fold is the answer to the question (whited out; one has to highlight the text in order to read it).

The answer to the question is… no.

First of all, it suffices to prove the claim for an open interval such as (0,1). For, if one partitions a half-open interval such as [0,1) into closed intervals, then after selecting any one of these intervals [a,b], the open interval (b,1) must then also be partitioned into countably many disjoint closed intervals.

By mapping (0,1) homeomorphically to the real line, it suffices to show that the real line R cannot be partitioned into closed intervals.

As each interval is bounded, we need an infinite number of intervals $I_n = [a_n, b_n]$ to cover the real line. Now consider the set

$$E := \bigcup_{n=1}^{\infty} \{a_n, b_n\}$$

consisting of the endpoints of the intervals I_n. Clearly, E is countably infinite. Also, as the I_n form a disjoint cover of E, E is the complement of the open set $\bigcup_{n=1}^{\infty} (a_n, b_n)$ and is hence closed. Finally, we claim that E is perfect: that not only is E closed, but every point in E is a limit point in E. Indeed, if x lies in E, then x is either the left or right endpoint of an interval I_n, but it is not both. If it is, say, the right-endpoint of an interval, then by approaching x from the right we see that x is the limit of the left endpoint of intervals to the right of x.

Now we appeal to a general theorem (a special case of the even more general Baire category theorem) that asserts that a perfect subset of a complete metric space cannot be countably infinite. A proof is as follows. Suppose for contradiction that $E = \{x_1, x_2, x_3, ...\}$ is a countably infinite perfect set. Let B_1 be any closed ball of positive radius whose centre lies in E (e.g. one can take the closed ball of radius 1 centered at x_1). Using the fact that x_1 is a limit point, one can then find a closed ball B_2 inside B_1 whose centre still lies in E, but which is disjoint from

x_1, and whose radius is at most half that of B_1. (Indeed, if x_1 does not already lie in the interior of B_1, one simply has to shrink B_1 by a little bit and recenter to another element of E to move x_1 off the boundary of B_1; otherwise, one uses the limit point property to find another point in the interior of B_1 other than x_1, and makes a small closed ball around that.) Then one can find a closed ball B_3 inside B_2 of radius at most half that of B_2 whose centre still lies in E, but is now also disjoint from x_2 in addition to x_1. Continuing in this fashion we find a sequence $B_1, B_2, B_3, \ldots$ of nested closed balls with radii going to zero, each of which has centre in E. The centres of these balls then form a Cauchy sequence, which by completeness, must converge to a limit in E that also lies in $\bigcap_{n=1}^{\infty} B_n$. But by construction, $\bigcap_{n=1}^{\infty} B_n$ lies outside each element x_n of E, and is thus disjoint from E, a contradiction.

Several of the students also came up with a proof based on the fact that the set E had the topological structure of a Cantor set, which is necessarily uncountable. This is close to the proof given above, though the above proof has the advantage that it is more readily generalisable to higher dimensions, showing for instance that a non-closed box cannot be expressed as the countable union of closed boxes. (For the generalisation, one has to use the full strength of the Baire category theorem, rather than just relying on the theory of perfect sets.) UPDATE: as pointed out in comments, the higher-dimensional result can also be easily deduced from the one-dimensional result by looking at a slice.

Note that if one replaces the reals in the question by the rationals, then the answer to the question becomes "yes" (this is easiest to see if one uses closed intervals of zero length, though one can also concoct examples using positive length intervals also). This shows that any proof must use a property of the real line, such as completeness, that is not shared by the rationals, and so it is unlikely that there is any elementary proof that is significantly shorter than the one given above.

Comments (59)



Anonymous · 4 October 2010

"If it is, say, the __right__-endpoint of an interval, then by approaching x from the right we see that x is the limit of the left endpoint of intervals to the right of x." *[Corrected, thanks – T.]*



porton · 4 October 2010

The formulas in this post are completely messed up.



porton · 4 October 2010

Oh, no. It is not messed formulas. It is (in some mysterious reason) invisible text on the page (white on white).



porton · 4 October 2010

Oh, sorry, Terry, for this stupid comment. Now I see that you intentionally whited up the text.



Anonymous · 28 September 2019

oh I am still confused about the text

Student · 4 October 2010

Another question that I still dont know how to prove is:

Thm:Show that a union of open intervals can be written as a disjoint union of open intervals.

Induction isn't enough because induction just works for a finite number of arbitrary open intervals.

I know how where to find the answer, but I want to solve it by myself. I am more than one week trying to solve it. :o)

Best

> **Anonymous** · 3 January 2011
>
> I do not have access to Follands book right now, but nevertheless, I tried to proof Students question.
>
> Theorem: Any open subset U of R may be expressed as a disjoint union of open intervals. (this implies his question as any union of open intervals is an open subset of R) [R being the real set]
> Proof: Consider the intersection V of U and Q [Q being the rational numbers]. Note that V is countably infinite.
> For each q in V, define
> A_q = inf{a in U | (a,q) is a subset of U}
> B_q = sup{b in U | (q, b) is a subset of U}
> Both values exist in R because U is open.
> Now, I_q = (A_q, B_q) is an open interval fully contained in U. Because if x is in I_q but not in U, it would contradict our definition of either A_q or B_q.
> We obtain a countably infinite amount of intervals I_q, and for q1 and q2 in V we either have
> I_q1 = I_q2, or
> I_q1 and I_q2 are disjoint. (because if two intervals intersect in any point, they have to be the same by our construction)
> Also, because Q is a dense subset of R and U is open, any x in U is an element of I_q for some q in V (consider an open ball around x, and a rational number q within that ball), so we conclude that the union of all I_q equals U.
>
> This may seem sketchy, but being a foreign undergraduate is leaving me with some problems to express myself precisely.
>
> > **Anonymous** · 28 September 2019
> >
> > Any open set in $\mathbb{R}^n$ (and in other topological spaces) is a disjoint union of its (open) connected components.

Gautam · 2 June 2013

"A Course on Borel Sets" by S M Srivastava. pg.51.

Anonymous · 11 August 2016

The intersection of open intervals is open. Deleting the intersection from one of the offending pairs of open intervals gives two open intervals with equal union.

Anonymous · 11 August 2016

*one from

Anonymous · 11 August 2016

Intersection of open intervals is not necessarily open (e.g. the intersection of all open intervals containing a closed interval).



ugroh · 4 October 2010

Remark to the Student-question: A proof can be found e.g. in Folland, Real Analysis, Prop. 0.21 .

Ulrich

David Milovich · 4 October 2010

Cantor's original proof of the reals' uncountability
can be adapted to get a shorter elementary proof.

Start with [0,1) and a sequence (C_n:n=0,1,2,...)
of disjoint closed subintervals of [0,1).
Choose i_0 and j_0 such that C_(i_0)<C_(j_0).
Given C_(i_n)<C_(j_n),
choose the least i_(n+1) such that
C_(i_n)<C_(i_(n+1))<C_(j_n).
Then choose some j_(n+1) such that
C_(i_(n+1))<C_(j_(n+1))<C_(j_n).

Let x be the supremum of the union
of all sets of the form C_(i_n).
Given any n, since C_(i_n)<x<C_(j_n) and i_(n+1) is
minimal with respect to C_(i_n)<C_(i_(n+1))<C_(j_n),
x cannot be in C_k for any k<i_(n+1).
Thus, for all n, x is not in C_k for any k<i_(n+1).
But C_(i_n)<C_(i_(n+1)), so the sequence (i_n:n=0,1,2,...)
never repeats, which implies that the sequence is
unbounded, which implies that x is not in any C_k.

Marton Eekes · 5 October 2010

A slightly harder nice exercise is (I only modified the very last word):

Is it possible to express a non-closed interval in the real line, such as [0,1), as a countable union of disjoint closed sets?

 **pavel zorin** · 5 October 2010

Using the hint on the Baire category theorem one could also infer that a complete connected metric space is not a countable union of disjoint closed subsets. I am not sure whether this result generalizes to Baire spaces, though.

best regards, pavel

 **pavel zorin** · 5 October 2010

I have noticed an error in my reasoning: the space should also be locally connected. Nevertheless, I do not have a counterexample if local connectedness does not hold.

 **pavel zorin** · 7 October 2010

Here is a counterexample, i.e. a connected complete metric space which is a countable union of closed sets:

Let {e_{i,j}; i>0, j≥0} be orthonormal vectors in a Hilbert space.

Let S_i consist of the closed interval [e_{i,0}, 2 e_{i,0}] and the polylines from 2 e_{i,0} to 2 e_{i,0}+e_{i,j} to e_{j,0}+e_{i,j} to e_{j,0}+e_{i,j}/i.

Then S_i are closed and their union is complete and connected. This set is best visualized as countably many intervals, each of which has antennae in the direction of an endpoint of each other set.

Note that the boundary of each S_i is just {e_{i,0}}, hence the union of the boundaries is a discrete set (as opposed to a perfect set in the case of intervals on the real line)

 **domotorp** · 5 October 2010

I remembered that we used to call such sets F_σ^* on the problem solving seminar.

 **Anonymous** · 9 October 2010

I am ignorant of the difference of a closed set and a closed interval of the real line. Can't any closed set ve decomposed into closed intervals?

 **mircea petrache** · 5 October 2010

Take a V-shaped function u:[0,1]->[0,1] such that u(0)=u(1)=1 and u(1/2)=0, and such that u is piecewise affine on [0,1/2], [1/2,1].

Take now a family of intervals which would falsify the theorem. On one such interval [a,b] put v(x)=u((x-a)/(b-a)), then define v(1)=v(0)=1. So from the set of intervals of the text, we got a continuous function v:[0,1]->[0,1], which is =1 just on the union of extrema of the segments.

v is uniformly continuous, but this contradicts the fact that the lenghts of intervals can be arbitrarily small (thus the slope of v is unbounded).

does this make sense?

 **Tobias Hagge** · 5 October 2010

Thank you for your wonderful blog. Here is an argument that uses the compactness of closed intervals in $\mathbb{R}$.

Given a sequence $([a_i, b_i])_{i \in \mathbb{N}^+}$ of disjoint closed intervals in $(0,1)$, let $C_n = \cup_{i=1}^n [a_i, b_i]$. We'll construct a nested sequence of nonempty closed intervals S_i such that C_i and S_i are disjoint. The intersection of such is nonempty and does not intersect any $[a_i, b_i]$.

Choose $S_0 = [c_0, d_0]$ such that $c_0 < a_1$ and $a_1 < d_0 < b_1$. If $[a_i, b_i]$ is disjoint from S_{i-1}, let $S_i = S_{i-1}$. Otherwise, suppose that $S_{i-1} - C_i = [c, d)$. This lies in some component (e, d) of the complement of C_i in $(0,1)$. Set $S_i = [c_i, d_i]$, where $c < c_i < d_i < d$, chosen so that the least-indexed $[a_i,b_i]$ which lies in (e, d) contains c_i but not d_i. The case $S_{i-1} - C_i = (c, d]$ is similar.

 **Danny Calegari** · 5 October 2010

I think this is a theorem of Sierpinski; the reference I have in mind is "Un theoreme sur les continus" (Tohoku Math 13 (1918) 300-303) but I don't have the reference handy.

Top Posts — WordPress.com · 5 October 2010

[...] Covering a non-closed interval by disjoint closed intervals The following question came up in my 245A class today: Is it possible to express a non-closed interval in the real [...] [...]

 **Tobias Hagge** · 5 October 2010

One may give a purely topological proof by constructing a nested sequence of closed intervals S_n, the intersection of which is disjoint from the union of the I_n. Suppose S_n is chosen so that for the least k such that I_k is not disjoint from S_n, S_n – I_k is a half-open interval. Inside of this half-open interval is a closed subinterval S_{n+1} with the same property and larger least k.

This gives the result for any dense total order on which closed intervals are compact. However, I'm having a little trouble comparing the generality of that argument to the argument in the post. Are such spaces always metrizable?

 **Tobias Hagge** · 7 October 2010

Why the down vote?

 **Tobias Hagge** · 11 October 2010

To answer my own question, no. Counterexample: the hyperreals. More general counterexample: the Dedekind-MacNeille completion of any non-metrizable total order. By the least upper bound property, closed intervals are compact. The completion is not metrizable since it has a non-metrizable subspace.

 **gowers** · 6 October 2010

In Cambridge this is a well-known "exercise for the enthusiast" that has been tackled by several generations of the brighter undergraduates.

As a little experiment, I just typed "Can [0,1] be covered by countably many disjoint closed intervals?" into Google (with no inverted commas) and this post came up first. So now there will be an easy way of solving the problem. I wonder if eventually this method will work

for all exercises. (I've just tried it for one or two other questions and it didn't work, so we haven't reached this state yet.)

> **anonymous** · 6 October 2010
>
> No, in fact we have the opposite problem: it's often inconveniently difficult to find the answers to basic mathematical questions by searching online.
>
> (I had a rare success with this the other day when looking for an example of a differentiable function that isn't continuously differentiable. After a surprisingly long few minutes I found what I was looking for, but in general this is the type of thing that is difficult to find: things that "everyone knows" but that don't have standard names.)

Xueping · 6 October 2010

I have a different proof which I can't judge whether it's available.
Observation: Let $ (a_n, b_n)$ be a strictly decreasing sequence of open intervals. If $a_n < a_{n+1}< b_{n+1}<b_{n}$ is always true, then their intersection is not empty.

Suppose that $(0,1)= \cup [c_n , d_n]$, where $ [c_n , d_n]$ are disjoint. Take $A_0 =(0,1)$.
Consider the complement of $[c_1 ,d_1]$, $(0, c_1) \cup (c_2, 1)$, $[c_2, d_2]$ must lie in one of the two intervals, say $(0, c_1) $ for example.
Then take $A_1 =(d_2 , c_1)$. Wait until the next $[c_i, d_i]$ lies in A_1 and another $[c_k, d_k]$ divides an interval of the complement of $[c_i, d_i]$ in A_1. Then in the resulting two intervals, choose one that has no common endpoints with A_1 to be A_2. Inductively we can get a strictly decreasing sequence of open intervals A_n, which satisfies that the endpoints never coincide. As a consequence, the intersection of them is nonempty.
From the construction, we can see that this intersection is a subset of the complement $\cup [c_n , d_n]$, which is empty. A contradiction.

huangxueping · 6 October 2010

I have a different proof which I can't judge whether it's available.
Observation: Let (a_n, b_n) be a strictly decreasing sequence of open intervals. If $a_n < a_{n+1} < b_{n+1} < b_n$ is always true, then their intersection is not empty.

Suppose that $(0, 1) = \cup [c_n, d_n]$, where $[c_n, d_n]$ are disjoint. Take $A_0 = (0, 1)$.
Consider the complement of $[c_1, d_1]$, $(0, c_1) \cup (c_2, 1)$, $[c_2, d_2]$ must lie in one of the two intervals, say $(0, c_1)$ for example.
Then take $A_1 = (d_2, c_1)$. Wait until the next $[c_i, d_i]$ lies in A_1 and another $[c_k, d_k]$ divides one interval of the complement of $[c_i, d_i]$ in A_1 into two intervals . Then in the resulting two intervals, choose one that has no common endpoints with A_1 to be A_2. Inductively we can get a strictly decreasing sequence of open intervals A_n, which satisfies that their endpoints never coincide. As a consequence, the intersection of them is nonempty.
From the construction, we can see that this intersection is a subset of the complement $\cup [c_n, d_n]$, which is empty. A contradiction.

> **Leena** · 9 October 2010
>
> :huangxueping I am sorry but I feel this does not complete the proof... cuz a closed set in (0,1) need not necessarily be in the form [c,d] it could be something like {a,b,c,d} etc...

The above proof proves that (0,1) can not be written as a union of disjoint connected and closed subsets of (0,1). I guess We need a more general prof for the above thm.

Fan · 29 July 2013

I think this proof also generalizes to the case of closed sets. Consider the complement of each closed set, which is open and can be written as a countable union of open intervals. It is easy to see that there cannot be only one open interval (since the complement is not connected), so you still have the freedom of choice to make a nested sequence of open intervals whose intersection is not empty.

Fan · 29 July 2013

Is this proof morally the same as Hagge's?

J Balachandran · 6 October 2010

I might be misunderstanding the question. If I'm correct a finite union of closed sets in a closed set. However the given set is not closed since the limit point 1 does not belong to the set. So there is no way we can have a closed set to represent this non-closed set. Is this reasoning correct? If wrong can someone highlight me where the mistake is

Anonymous · 6 October 2010

Not quite. The question is whether the open set can be expressed as a *countable* union of closed sets (ie infinite unions are allowed). Your explanation only holds if we are constrained to a *finite* union of closed sets.

J Balachandran · 7 October 2010

Oh ok. sorry. I got the mistake

Jonathan Weinstein · 6 October 2010

J Balachandran: It's a countable union, not finite, so not necessarily closed.

Huang Xueping: Very nice. You ruined my day though, I solved it your way and scrolled through the first 15 comments without seeing this...oh well.

Leena · 9 October 2010

but the sets are disjoint... so even if it is a countable union and not finite ... it is still a closed set..... if we take its closure ... we realize its a closed set.

Will · 12 October 2010

What about intervals of the form
[1/(2^(2n+1)),1/(2^(2n)]
Take the union (for natural numbers n) of all such intervals and 0 is a limit point.
Thus, a countable union of disjoint closed intervals need not be a closed set.

Apoorva Khare · 6 October 2010

First, Huang Xueping's proof can basically be modified to work for the more general question of covering (or not) by closed SETS (posed by Marton Eekes).

Second, to generalize this to higher dimensions (can the open unit ball be covered by countably many disjoint closed SETS in Euclidean space?), is it not enough to intersect with

the "X-axis"? For if there were such a cover, then the intersection with the X-axis would give a countable cover of (-1,1) by disjoint closed sets, which is false.

[Good point; I've added a mention of this in the post. -T.]

 **Apoorva Khare** · 6 October 2010

Curiously, I first saw the (more general, ref. Eekes) problem eight years ago in graduate school (and I solved it essentially by Huang Xueping's method too), as well as the "higher dimensions" variant. Just recently, I asked it of my students here at Yale, some three weeks ago! Then I posed it to a friend in Ann Arbor, and he asked it of his students 1-2 weeks ago. Curiously, the same problem showed up in Prof. Tao's class in UCLA this week.

 **Koen van Woerden** · 7 October 2010

Suppose a countable cover I_n of closed disjoint sets of the interval $(0,1)$ exists.

Define the function $f : (0,1) \to \mathbf{R}$ as follows: each x in $(0,1)$ is covered by precisely one I_n. Send x to n. Then f is continuous, and since the image of f is contained in the natural numbers, by the intermediate value theorem f is constant. Hence $(0,1)$ is covered by a single closed interval. Hence $(0,1)$ is closed. Contradiction.

 **Koen van Woerden** · 7 October 2010

Never mind, this is not a good proof. Sorry for that.

 **Liu Shiping** · 7 October 2010

Also, as the I_n form a disjoint partition of E,

I guess it should be a disjoint partition of the real line.

[Corrected, thanks – T.]

 **some graduate student** · 7 October 2010

This confused me too. Usually a "partition" is a collection of subsets of the set satisfying certain conditions — but the I_n aren't subsets of E.

 **mircea petrache** · 8 October 2010

I try to make a proof like Koen van Woerden's, i.e. using functions and continuity. It looks elementary just because the key points are left to the reader.

Take a square [0,1]x[-1,0], and consider the inscribed semicircle with diameter the "upper" side [0,1]x{0}. Now suppose a family as in the theorem exists. Then for each segment I in the family, put a shrinked copy of the above square. Then the union of the "shrinked semicircles" will describe the graph of a continuous function (since at the end points of any segment, shorter segments accumulate from one side).

But this also would define a continuous function on [0,1] (if you extend to zero at the extrema). Thus it should also be uniformly continuous. Still, the fact that the length of segments is arbitrarily close to 0 (and the fact that the tangent to the semicircles is vertical), contradicts any choice of modulus of continuity.

 **mircea petrache** · 8 October 2010

again the proof was wrong, maybe stretching the vertical direction of the squares to square root of I makes it work

 **Vladimir Slepnev** · 8 October 2010

I got the answer “no” by the following argument that sounds very natural to me: 1) reduce the problem to the open interval like you did, 2) removing a closed interval turns an open interval into two open intervals, 3) repeating the process countably many times leaves us with a binary tree of intervals, 4) if some interval never gets split then we have our answer, 5) if all intervals get split then alternating left-right-left-right-… gives us an intersection point. Completeness is used in the last step.

 **Arindam Bandyopadhyay** · 8 October 2010

Dear Sir,
Please discuss about the following problem:
Show that $\{\sum\limits_{k=0}^{n}\frac{1}{k^{2}}\}$ is a Cauchy sequence by using the following definition of Cauchy sequence:
A sequence $\{x_{n}\}$ is said to be a Cauchy sequence if for any $\epsilon >0$, there exists a $m\in\mathbb{N}$ such that $|x_{n+p}-x_{n}|<\epsilon$ for all $n\geq m$ and all $p\in\mathbb{N}$.

 **Leena** · 8 October 2010

let (0,1) = union of countable union of I(n) where each I (n) is closed and all of them are disjoint. Then denote (0,1) by set A and the countable union of I(n) as B. We claim sets A and B are equal. then Closures of both of them also must be equal.. closure of (0,1) is [0,1]. Now taking the closure of the countable union of I(n) …. all the sets are disjoint and hence closure of union of I (n) is equal to union of closures of each of I (n). but all of I (n) are closed. So its the same set B again. Hence the closure of A is not equal to closure of B . Which is a contradiction. so (0,1) can not be written as a union of countable union of disjoint closed sets. Its applicable to all open intervals too…

 Lewis Bowen · 10 October 2010

I think I found a different proof. It begins the same way; it suffices to show that the real line cannot be covered by countably many disjoint intervals. Suppose for a contradiction that it can. Let C_1, C_2,… be such a covering. Let O_i be the union of the complement of C_i with the interior of C_i (so O_i is the whole line minus the two endpoints of C_i). So O_i is a dense open set. Accordingly the intersection of all the O_i’s is a dense G_delta. But it must be empty if the C_i’s cover and are disjoint. This contradiction proves it.

 **Local Connectedness and Some Weird Counterexamples « Gracious Living and the Two Meat Meal** · 11 November 2010

[…] only way to do this is by letting itself be the only one in the union; proofs are, for example, here (read the comments for other viewpoints/try it yourself first!). So the only paths in the set are […]

Manuel · 17 March 2011

To prove that [0,1) can't be expressed as a union of countable many disjoint closed intervals, can't we just say de following?

Suppose we can. We have then a countable family F of closed intervals, which can be ordered in the sense reals are (?). We order them creating the sequence {[a_n, b_n]}, where b_1<a_2, b_2<a *[text lost here due to unintended HTML processing of the < and > signs – T.]* >b_k (since disjoint). So we can find some c in (b_k, a_(k+1)) opened set which hasn't been covered by the family F.

Maybe there's an error when saying we can order that family F, because if not this would sound good to me...

Sergei Ovchinnikov · 22 September 2011

See American Mathematical Monthly, 84 (1977),827-828.

Luqing Ye · 29 July 2013

I feel frustrated to see that the answer is "no" because this means that my intuition is not reliable when faced with infinity...

Which means I need to adjust my intuition.

Now I give a proof to show why it is impossible to cover a non-closed interval by countable disjoint closed intervals.

In order to prove that it is impossible to cover [0,1) by countable disjoint closed intervals,it is clear that we just need to prove that it is impossible to cover [0,1] by countable disjoint closed intervals.

We prove by contradiction.

Otherwise,

There must exists a sequence of real numbers $a_1, \cdots, a_n$ in [0,1] which is convergent to a real number in [0,1],and every item in this sequence is exactly covered by one non-degenerate closed interval(we call the interval of the shape [a,a] as the degenerate closed interval),and any of the two non-degenerate closed intervals are disjoint.

This is impossible,because at the very end of this sequence,two items will be very near that they must be in one non-degenerate intervals.

Luqing Ye · 15 August 2013

Dear Prof.Tao,

I don't know whether you regard [1,1] as a closed interval or not.

Anonymous · 9 November 2021

Since it is singleton it is close

myc10003 · 29 April 2015

I have a different proof. Consider an arbitrary covering of (0, 1) by pairwise disjoint, closed intervals I_x, each I_x is in (0, 1). Denote the covering set by L. One can show that L is a linear continuum and hence uncountable.

Since the closed intervals are pairwise disjoint, there is a natural total order amongst them. Since L covers (0, 1), for any I_x < I_y in L, there exists an I_z in L such that I_x < I_z < I_y.

It remains to show that L has the least upper bound property. Suppose a subset of L, S, is bounded above by an I_x. There must exist a least upper bound of the right endpoints of the intervals in this subset. Call it a. Obviously a is either to the left of I_x or in I_x. Hence a is in (0, 1). If a is the right endpoint of an interval in S, that interval is the least upper bound of S. Otherwise, there exists an I_y in L − S that covers a. Since a is the least upper bound of the right ends of intervals in S, a must be the left endpoint of I_y. Hence I_y is the least upper bound. Therefore L has the least upper bound property.