Higher uniformity of arithmetic functions in short intervals II. Almost all intervals

Terence Tao · 11 November 2024

https://terrytao.wordpress.com/2024/11/11/higher-uniformity-of-arithmetic-functions-in-short-intervals-ii-almost-all-intervals/

tao2typ · converted to Typst 2026-10-14 (UTC)

Kaisa Matomäki, Maksym Radziwill, Fernando Xuancheng Shao, Joni Teräväinen, and myself have (finally) uploaded to the arXiv our paper "Higher uniformity of arithmetic functions in short intervals II. Almost all intervals". This is a sequel to our previous paper from 2022. In that paper, discorrelation estimates such as

$$\sum_{x \leq n \leq x+H} (\Lambda(n) - \Lambda^\sharp(n)) \bar{F}(g(n)\Gamma) = o(H)$$

were established, where Λ is the von Mangoldt function, $\Lambda^\sharp$ was some suitable approximant to that function, $F(g(n)\Gamma)$ was a nilsequence, and $[x, x+H]$ was a reasonably short interval in the sense that $H \sim x^{\theta+\varepsilon}$ for some $0 < \theta < 1$ and some small $\varepsilon > 0$. In that paper, we were able to obtain non-trivial estimates for θ as small as $5/8$, and for some other functions such as divisor functions d_k for small values of k, we could lower θ somewhat to values such as $3/5$, $5/9$, $1/3$ of θ. This had a number of analytic number theory consequences, for instance in obtaining asymptotics for additive patterns in primes in such intervals. However, there were multiple obstructions to lowering θ much further. Even for the model problem when $F(g(n)\Gamma) = 1$, that is to say the study of primes in short intervals, until recently the best value of θ available was $7/12$, although this was very recently improved to $17/30$ by Guth and Maynard.

However, the situation is better when one is willing to consider estimates that are valid for *almost all* intervals, rather than all intervals, so that one now studies local higher order uniformity estimates of the form

$$\int_X^{2X} \sup_{F,g} \left| \sum_{x \leq n \leq x+H} (\Lambda(n) - \Lambda^\sharp(n)) \bar{F}(g(n)\Gamma) \right| dx = o(XH)$$

where $H = X^{\theta+\varepsilon}$ and the supremum is over all nilsequences of a certain Lipschitz constant on a fixed nilmanifold G/Γ. This generalizes local Fourier uniformity estimates of the form

$$\int_X^{2X} \sup_{\alpha} \left| \sum_{x \leq n \leq x+H} (\Lambda(n) - \Lambda^\sharp(n)) e(-\alpha n) \right| dx = o(XH).$$

There is particular interest in such estimates in the case of the Möbius function $\mu(n)$ (where, as per the Möbius pseudorandomness conjecture, the approximant $\mu^\sharp$ should be taken to be zero, at least in the absence of a Siegel zero). This is because if one could get estimates of this form for any H that grows sufficiently slowly in X (in particular $H = \log^{o(1)} X$), this would imply the (logarithmically averaged) Chowla conjecture, as I showed in a previous paper.

While one can lower θ somewhat, there are still barriers. For instance, in the model case $F \equiv 1$, that is to say prime number theorems in almost all short intervals, until very recently the best value of θ was $1/6$, recently lowered to $2/15$ by Guth and Maynard (and can be lowered all the way to zero on the Density Hypothesis). Nevertheless, we are able to get some improvements at higher orders:

- For the von Mangoldt function, we can get θ as low as $1/3$, with an arbitrary logarithmic saving $\log^{-A} X$ in the error terms; for divisor functions, one can even get power savings in this regime.

- For the Möbius function, we can get $\theta = 0$, recovering our previous result with Tamar Ziegler, but now with $\log^{-A} X$ type savings in the exceptional set (though not in the pointwise bound outside of the set).
- We can now also get comparable results for the divisor function.

As sample applications, we can obtain Hardy-Littlewood conjecture asymptotics for arithmetic progressions of almost all given steps $h \sim X^{1/3+\varepsilon}$, and divisor correlation estimates on arithmetic progressions for almost all $h \sim X^{\varepsilon}$.

Our proofs are rather long, but broadly follow the "contagion" strategy of Walsh, generalized from the Fourier setting to the higher order setting. Firstly, by standard Heath–Brown type decompositions, and previous results, it suffices to control "Type II" discorrelations such as

$$\sup_{F,g} \left| \sum_{x \leq n \leq x+H} \alpha^* \beta(n) \bar{F}(g(n)\Gamma) \right|$$

for almost all x, and some suitable functions α, β supported on medium scales. So the bad case is when for most x, one has a discorrelation

$$\left| \sum_{x \leq n \leq x+H} \alpha^* \beta(n) \bar{F}_x(g_x(n)\Gamma) \right| \gg H$$

for some nilsequence $F_x(g_x(n)\Gamma)$ that depends on x.

The main issue is the dependency of the polynomial g_x on x. By using a "nilsequence large sieve" introduced in our previous paper, and removing degenerate cases, we can show a functional relationship amongst the g_x that is very roughly of the form

$$g_x(an) \approx g_{x'}(a'n)$$

whenever $n \sim x/a \sim x'/a'$ (and I am being extremely vague as to what the relation "$\approx$" means here). By a higher order (and quantitatively stronger) version of Walsh's contagion analysis (which is ultimately to do with separation properties of Farey sequences), we can show that this implies that these polynomials $g_x(n)$ (which exert influence over intervals $[x, x+H]$) can "infect" longer intervals $[x', x'+Ha]$ with some new polynomials $\tilde{g}_{x'}(n)$ and various $x' \sim Xa$, which are related to many of the previous polynomials by a relationship that looks very roughly like

$$g_x(n) \approx \tilde{g}_{ax}(an).$$

This can be viewed as a rather complicated generalization of the following vaguely "cohomological"-looking observation: if one has some real numbers α_i and some primes p_i with $p_j \alpha_i \approx p_i \alpha_j$ for all i, j, then one should have $\alpha_i \approx p_i \alpha$ for some α, where I am being vague here about what $\approx$ means (and why it might be useful to have primes). By iterating this sort of contagion relationship, one can eventually get the $g_x(n)$ to behave like an Archimedean character n^{iT} for some T that is not too large (polynomial size in X), and then one can use relatively standard (but technically a bit lengthy) "major arc" techiques based on various integral estimates for zeta and L functions to conclude.

Comments (7)



Anonymous · 11 November 2024

Possible typo in reference [4]: Should "Ueber" be "Über"?



Terence Tao · 11 November 2024

Strangely enough, this was Hilbert's actual title in his 1892 paper. It is possible that German spelling standards have changed slightly over the last century and a quarter. (The title could also have been an artefact of the typesetting technology of the era.)

Anonymous · 14 November 2024

From wikipedia (https://de.wikipedia.org/wiki/%C3%9C)
In the umlaut meaning, the Ü was first written as UE. This later evolved into a U with a small E above it (Ů /ů). Over time, this E was stylized into two dots.

Budha Gautama sidarta · 12 November 2024

Is mathematics discovered or created?Platonism (discovery) argues that mathematical entities exist independently of human minds, in an abstract "Platonic realm." Mathematicians, therefore, discover these concepts, rather than invent them. G.H. Hardy supported this view, noting that the beauty of certain formulas suggests a preexisting mathematical reality.Formalism and Constructivism (creation), on the other hand, claim that mathematics is a human invention. Concepts are created, with rules and axioms established to form a logical system. David Hilbert saw mathematics as a symbolic system without inherent external reality, while Constructivism adds that only concepts we can construct and verify have true validity.

goingtoinfinity · 1 December 2024

I am very curious about your estimation of how difficult it would be to formalize works in this specific area of research. I read that the formalization of PFR was achieved so quickly because the field of combinatorics is well-developed in Lean. How does this compare to analytic number theory? How reliant is this field on other deep results in the literature? As far as I understand, PFR was largely self-contained. Is this also true for this result, or would it require reliance on many other complex results from the literature? Thank you very much!



Terence Tao · 1 December 2024

Right now we have a project to formalize the prime number theorem (PNT) in Lean, as well as generalizations such as the prime number theorem in arithmetic progressions and the Chebotarev density theorem. We did manage to formalize the PNT, and are well on our way to formalizing the PNT in arithmetic progressions, but one stumbling block has been the ability to manipulate asymptotic estimates in the way that analysts are used to doing (e.g., combining together multiple O() or o() error terms into a single term, etc.). We can do it with the current Mathlib framework, but it is quite tedious; in particular, the "easy" parts of the PNT in which one converts from, say, control of the von Mangoldt function to control of the prime counting function, has turned out to be one of the more difficult steps in the argument to formalize. This is an issue that remains something of an open problem to resolve in the future. One possibility is to eschew asymptotic notation altogether and work with completely explicit estimates, which are avoided in human-readable proofs due to their fragility to arithmetic errors, but in the fully verified framework of Lean, this is not an issue.

Sumit shingare · 29 December 2024

Dear Professor Tao,
In a recent lecture, you mentioned that we might be just one breakthrough away from resolving all the logarithmically averaged versions of the Chowla and Sarnak conjectures. If such a breakthrough happens, how much closer do you think it would bring us to proving the full Chowla conjecture or even the Twin Prime Conjecture?
Additionally, do you believe the tools and techniques you and your collaborators have been developing will eventually play a role in tackling these conjectures directly, or do you think completely different methods will be required?
Your decades-long dedication to the study of primes is truly inspiring. I deeply appreciate your work and wish you all the best as you continue your groundbreaking research.

Sincerely,
Sumit