275A, Notes 4: The central limit theorem

Terence Tao · 2 November 2015

https://terrytao.wordpress.com/2015/11/02/275a-notes-4-the-central-limit-theorem/

tao2typ · converted to Typst 2026-11-02 (UTC)

Let $X_1, X_2, \ldots$ be iid copies of an absolutely integrable real scalar random variable X, and form the partial sums $S_n := X_1 + \ldots + X_n$. As we saw in the last set of notes, the law of large numbers ensures that the empirical averages S_n/n converge (both in probability and almost surely) to a deterministic limit, namely the mean $\mu = \mathbf{E}X$ of the reference variable X. Furthermore, under some additional moment hypotheses on the underlying variable X, we can obtain *square root cancellation* for the fluctuation $\frac{S_n}{n} - \mu$ of the empirical average from the mean. To simplify the calculations, let us first restrict to the case $\mu = 0, \sigma^2 = 1$ of mean zero and variance one, thus

$$\mathbf{E}X = 0$$

and

$$\mathbf{Var}(X) = \mathbf{E}X^2 = 1.$$

Then, as computed in previous notes, the normalised fluctuation $S_n/\sqrt{n}$ also has mean zero and variance one:

$$\mathbf{E}\frac{S_n}{\sqrt{n}} = 0$$
$$\mathbf{Var}(\frac{S_n}{\sqrt{n}}) = \mathbf{E}(\frac{S_n}{\sqrt{n}})^2 = 1.$$

This and Chebyshev's inequality already indicates that the "typical" size of S_n is $O(\sqrt{n})$, thus for instance $\frac{S_n}{\sqrt{n}\omega(n)}$ goes to zero in probability for any $\omega(n)$ that goes to infinity as $n \to \infty$. If we also have a finite fourth moment $\mathbf{E}|X|^4 < \infty$, then the calculations of the previous notes also give a fourth moment estimate

$$\mathbf{E}(\frac{S_n}{\sqrt{n}})^4 = 3 + O(\frac{\mathbf{E}|X|^4}{n}).$$

From this and the Paley-Zygmund inequality (Exercise 44 of Notes 1) we also get some lower bound for $\frac{S_n}{\sqrt{n}}$ of the form

$$\mathbf{P}(\left|\frac{S_n}{\sqrt{n}}\right| \geq \varepsilon) \geq \varepsilon$$

for some absolute constant $\varepsilon > 0$ and for n sufficiently large; this indicates in particular that $\frac{S_n\omega(n)}{\sqrt{n}}$ does not converge in any reasonable sense to something finite for any $\omega(n)$ that goes to infinity.

The question remains as to what happens to the ratio $S_n/\sqrt{n}$ itself, without multiplying or dividing by any factor $\omega(n)$. A first guess would be that these ratios converge in probability or almost surely, but this is unfortunately not the case:

Proposition 1 Let $X_1, X_2, \ldots$ be iid copies of an absolutely integrable real scalar random variable X with mean zero, variance one, and finite fourth moment, and write $S_n := X_1 + \ldots + X_n$. Then the random variables $S_n/\sqrt{n}$ do not converge in probability or almost surely to any limit, and neither does any subsequence of these random variables.

Proof: Suppose for contradiction that some sequence $S_{n_j}/\sqrt{n_j}$ converged in probability or almost surely to a limit Y. By passing to a further subsequence we may assume that the convergence is in the almost sure sense. Since all of the $S_{n_j}/\sqrt{n_j}$ have mean zero, variance one, and bounded fourth moment, Theorem 25 of Notes 1 implies that the limit Y also has mean zero and variance one. On the other hand, Y is a tail random variable and is thus almost surely constant by the Kolmogorov zero-one law from Notes 3. Since constants have variance zero, we obtain the required contradiction. □

Nevertheless there is an important limit for the ratio $S_n/\sqrt{n}$, which requires one to replace the notions of convergence in probability or almost sure convergence by the weaker concept of convergence in distribution.

Definition 2 (Vague convergence and convergence in distribution) Let R be a locally compact Hausdorff topological space with the Borel σ-algebra. A sequence of finite measures μ_n on R is said to converge vaguely to another finite measure μ if one has

$$\int_R G(x)\ d\mu_n(x) \to \int_R G(x)\ d\mu(x)$$

as $n \to \infty$ for all continuous compactly supported functions $G : R \to \mathbf{R}$. (Vague convergence is also known as *weak convergence*, although strictly speaking the terminology weak-* convergence would be more accurate.) A sequence of random variables X_n taking values in R is said to *converge in distribution* (or *converge weakly* or *converge in law*) to another random variable X if the distributions μ_{X_n} converge vaguely to the distribution μ_X, or equivalently if

$$\mathbf{E}G(X_n) \to \mathbf{E}G(X)$$

as $n \to \infty$ for all continuous compactly supported functions $G : R \to \mathbf{R}$.

One could in principle try to extend this definition beyond the locally compact Hausdorff setting, but certain pathologies can occur when doing so (e.g. failure of the Riesz representation theorem), and we will never need to consider vague convergence in spaces that are not locally compact Hausdorff, so we restrict to this setting for simplicity.

Note that the notion of convergence in distribution depends only on the distribution of the random variables involved. One consequence of this is that convergence in distribution does not produce unique limits: if X_n converges in distribution to X, and Y has the same distribution as X, then X_n also converges in distribution to Y. However, limits are unique up to equivalence in distribution (this is a consequence of the Riesz representation theorem, discussed for instance in this blog post). As a consequence of the insensitivity of convergence in distribution to equivalence in distribution, we may also legitimately talk about convergence of distribution of a sequence of random variables X_n to another random variable X even when all the random variables $X_1, X_2, \ldots$ and X involved are being modeled by different probability

spaces (e.g. each X_n is modeled by Ω_n, and X is modeled by Ω, with no coupling presumed between these spaces). This is in contrast to the stronger notions of convergence in probability or almost sure convergence, which require all the random variables to be modeled by a common probability space. Also, by an abuse of notation, we can say that a sequence X_n of random variables converges in distribution to a probability measure μ, when μ_{X_n} converges vaguely to μ. Thus we can talk about a sequence of random variables converging in distribution to a uniform distribution, a gaussian distribution, etc..

From the dominated convergence theorem (available for both convergence in probability and almost sure convergence) we see that convergence in probability or almost sure convergence implies convergence in distribution. The converse is not true, due to the insensitivity of convergence in distribution to equivalence in distribution; for instance, if $X_1, X_2, \ldots$ are iid copies of a non-deterministic scalar random variable X, then the X_n trivially converge in distribution to X, but will not converge in probability or almost surely (as one can see from the zero-one law). However, there are some partial converses that relate convergence in distribution to convergence in probability; see Exercise 10 below.

> **Remark 3** The notion of convergence in distribution is somewhat similar to the notion of convergence in the sense of distributions that arises in distribution theory (discussed for instance in this previous blog post), however strictly speaking the two notions of convergence are distinct and should not be confused with each other, despite the very similar names.

The notion of convergence in distribution simplifies in the case of real scalar random variables:

> **Proposition 4** Let $X_1, X_2, \ldots$ be a sequence of scalar random variables, and let X be another scalar random variable. Then the following are equivalent:
>
> - (i) X_n converges in distribution to X.
> - (ii) $F_{X_n}(t)$ converges to $F_X(t)$ for each continuity point t of F_X (i.e. for all real numbers $t \in \mathbf{R}$ at which F_X is continuous). Here $F_X(t) := \mathbf{P}(X \leq t)$ is the cumulative distribution function of X.

Proof: First suppose that X_n converges in distribution to X, and F_X is continuous at t. For any $\varepsilon > 0$, one can find a δ such that

$$F_X(t) - \varepsilon \leq F_X(t') \leq F_X(t) + \varepsilon$$

for every $t' \in [t-\delta, t+\delta]$. One can also find an N larger than $|t| + \delta$ such that $F_X(-N) \leq \varepsilon$ and $F_X(N) \geq 1 - \varepsilon$. Thus

$$\mathbf{P}(|X| \geq N) = O(\varepsilon)$$

and

$$\mathbf{P}(|X - t| \leq \delta) = O(\varepsilon).$$

Let $G : \mathbf{R} \to [0,1]$ be a continuous function supported on $[-2N, t]$ that equals 1 on $[-N, t-\delta]$. Then by the above discussion we have

$$\mathbf{E}G(X) = F_X(t) + O(\varepsilon)$$

and hence

$$\mathbf{E}G(X_n) = F_X(t) + O(\varepsilon)$$

for large enough n. In particular

$$\mathbf{P}(X_n \leq t) \geq F_X(t) - O(\varepsilon).$$

A similar argument, replacing G with a continuous function supported on $[t, 2N]$ that equals 1 on $[t+\delta, N]$ gives

$$\mathbf{P}(X_n > t) \geq 1 - F_X(t) - O(\varepsilon)$$

for n large enough. Putting the two estimates together gives

$$F_{X_n}(t) = F_X(t) + O(\varepsilon)$$

for n large enough; sending $\varepsilon \to 0$, we obtain the claim.

Conversely, suppose that $F_{X_n}(t)$ converges to $F_X(t)$ at every continuity point t of F_X. Let $G : \mathbf{R} \to \mathbf{R}$ be a continuous compactly supported function, then it is uniformly continuous. As F_X is monotone increasing, it can only have countably many points of discontinuity. From these two facts one can find, for any $\varepsilon > 0$, a simple function $G_\varepsilon(t) = \sum_{i=1}^n c_i 1_{(t_i, t_{i+1}]}(t)$ for some $t_1 < \ldots < t_n$ that are points of continuity of F_X, and real numbers c_i, such that $|G(t) - G_\varepsilon(t)| \leq \varepsilon$ for all t. Thus

$$\mathbf{E}G(X_n) = \mathbf{E}G_\varepsilon(X_n) + O(\varepsilon)$$
$$= \sum_{i=1}^n c_i (F_{X_n}(t_{i+1}) - F_{X_n}(t_i)) + O(\varepsilon).$$

Similarly for X_n replaced by X. Subtracting and taking limit superior, we conclude that

$$\limsup_{n\to\infty} |\mathbf{E}G(X_n) - \mathbf{E}G(X)| = O(\varepsilon),$$

and on sending $\varepsilon \to 0$, we obtain that X_n converges in distribution to X as claimed. □

The restriction to continuity points of t is necessary. Consider for instance the deterministic random variables $X_n = 1/n$, then X_n converges almost surely (and hence in distribution) to 0, but $F_{X_n}(0) = 0$ does not converge to $F_X(0) = 1$.

> **Example 5** For any natural number n, let X_n be a discrete random variable drawn uniformly from the finite set $\{0/n, 1/n, \ldots, (n-1)/n\}$, and let X be the continuous random variable drawn uniformly from $[0, 1]$. Then X_n converges in distribution to X. Thus we see that a continuous random variable can emerge as the limit of discrete random variables.

> **Example 6** For any natural number n, let X_n be a continuous random variable drawn uniformly from $[0, 1/n]$, then X_n converges in distribution to the deterministic real number 0. Thus we see that discrete (or even deterministic) random variables can emerge as the limit of continuous random variables.

> **Exercise 7 (Portmanteau theorem)** Show that the properties (i) and (ii) in Proposition 4 are also equivalent to the following three statements:
>
> - (iii) One has $\limsup_{n\to\infty} \mathbf{P}(X_n \in K) \leq \mathbf{P}(X \in K)$ for all closed sets $K \subset \mathbf{R}$.

- (iv) One has $\liminf_{n\to\infty} \mathbf{P}(X_n \in U) \geq \mathbf{P}(X \in U)$ for all open sets $U \subset \mathbf{R}$.
- (v) For any Borel set $E \subset \mathbf{R}$ whose topological boundary ∂E is such that $\mathbf{P}(X \in \partial E) = 0$, one has $\lim_{n\to\infty} \mathbf{P}(X_n \in E) = \mathbf{P}(X \in E)$.

(Note: to prove this theorem, you may wish to invoke Urysohn's lemma. To deduce (iii) from (i), you may wish to start with the case of compact K.)

We can now state the famous central limit theorem:

Theorem 8 (Central limit theorem) Let $X_1, X_2, \ldots$ be iid copies of a scalar random variable X of finite mean $\mu := \mathbf{E}X$ and finite non-zero variance $\sigma^2 := \mathbf{Var}(X)$. Let $S_n := X_1 + \ldots + X_n$. Then the random variables $\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu)$ converges in distribution to a random variable with the standard normal distribution $N(0,1)$ (that is to say, a random variable with probability density function $x \mapsto \frac{1}{\sqrt{2\pi}} e^{-x^2/2}$). Thus, by abuse of notation

$$\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu) \to N(0,1).$$

In the normalised case $\mu = 0, \sigma^2 = 1$ when X has mean zero and unit variance, this simplifies to

$$\frac{S_n}{\sqrt{n}} \to N(0,1).$$

Using Proposition 4 (and the fact that the cumulative distribution function associated to $N(0,1)$ is continuous, the central limit theorem is equivalent to asserting that

$$\mathbf{P}(\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu) \leq t) \to \frac{1}{\sqrt{2\pi}} \int_{-\infty}^{t} e^{-x^2/2}\ dx$$

as $n \to \infty$ for any $t \in \mathbf{R}$, or equivalently that

$$\mathbf{P}(a \leq \frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu) \leq b) \to \frac{1}{\sqrt{2\pi}} \int_{a}^{b} e^{-x^2/2}\ dx.$$

Informally, one can think of the central limit theorem as asserting that S_n approximately behaves like it has distribution $N(n\mu, n\sigma^2)$ for large n, where $N(\mu, \sigma^2)$ is the normal distribution with mean μ and variance σ^2, that is to say the distribution with probability density function $x \mapsto \frac{1}{\sqrt{2\pi}\sigma} e^{-(x-\mu)^2/2\sigma^2}$. The integrals $\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{t} e^{-x^2/2}\ dx$ can be written in terms of the error function erf as $\frac{1}{2} + \frac{1}{2}\ \mathrm{erf}(t/\sqrt{2})$.

The central limit theorem is a basic example of the *universality phenomenon* in probability – many statistics involving a large system of many independent (or weakly dependent) variables (such as the normalised sums $\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu)$) end up having a universal asymptotic limit (in this case, the normal distribution), regardless of the precise makeup of the underlying random variable X that comprised that system. Indeed, the universality of the normal distribution is such that it arises in many other contexts than the fluctuation of iid random variables; the central limit theorem is merely the first place in probability theory where it makes a prominent appearance.

We will give several proofs of the central limit theorem in these notes; each of these proofs has their advantages and disadvantages, and can each extend to prove many further results beyond the central limit theorem. We first give Lindeberg's proof of the central limit theorem, based on

exchanging (or swapping) each component $X_1, \ldots, X_n$ of the sum S_n in turn. This proof gives an accessible explanation as to why there should be a universal limit for the central limit theorem; one then computes directly with gaussians to verify that it is the normal distribution which is the universal limit. Our second proof is the most popular one taught in probability texts, namely the Fourier-analytic proof based around the concept of the characteristic function $t \mapsto \mathbf{E}e^{itX}$ of a real random variable X. Thanks to the powerful identities and other results of Fourier analysis, this gives a quite short and direct proof of the central limit theorem, although the arguments may seem rather magical to readers who are not already familiar with Fourier methods. Finally, we give a proof based on the moment method, in the spirit of the arguments in the previous notes; this argument is more combinatorial, but is straightforward and is particularly robust, in particular being well equipped to handle some dependencies between components; we will illustrate this by proving the Erdos-Kac law in number theory by this method. Some further discussion of the central limit theorem (including some further proofs, such as one based on Stein's method) can be found in this blog post. Some further variants of the central limit theorem, such as local limit theorems, stable laws, and large deviation inequalities, will be discussed in the next (and final) set of notes.

The following exercise illustrates the power of the central limit theorem, by establishing combinatorial estimates which would otherwise require the use of Stirling's formula to establish.

Exercise 9 (De Moivre-Laplace theorem) Let X be a Bernoulli random variable, taking values in $\{0,1\}$ with $\mathbf{P}(X=0) = \mathbf{P}(X=1) = 1/2$, thus X has mean $1/2$ and variance $1/4$. Let $X_1, X_2, \ldots$ be iid copies of X, and write $S_n := X_1 + \ldots + X_n$.

- (i) Show that S_n takes values in $\{0, \ldots, n\}$ with $\mathbf{P}(S_n = i) = \frac{1}{2^n}\binom{n}{i}$. (This is an example of a binomial distribution.)
- (ii) Assume Stirling's formula

$$n! = (1+o(1))\sqrt{2\pi n} n^n e^{-n} \tag{1}$$

where $o(1)$ is a function of n that goes to zero as $n \to \infty$. (A proof of this formula may be found in this previous blog post.) Using this formula, and without using the central limit theorem, show that

$$\mathbf{P}(a \leq 2\sqrt{n}(\frac{S_n}{n} - \frac{1}{2}) \leq b) \to \frac{1}{\sqrt{2\pi}} \int_a^b e^{-x^2/2}\ dx$$

as $n \to \infty$ for any fixed real numbers $a < b$.

The above special case of the central limit theorem was first established by de Moivre and Laplace.

We close this section with some basic facts about convergence of distribution that will be useful in the sequel.

Exercise 10 Let $X_1, X_2, \ldots, Y_1, Y_2, \ldots$ be sequences of real random variables, and let X, Y be further real random variables.

- (i) If X is deterministic, show that X_n converges in distribution to X if and only if X_n converges in probability to X.
- (ii) Suppose that X_n is independent of Y_n for each n, and X independent of Y. Show that $X_n + iY_n$ converges in distribution to $X + iY$ if and only if X_n converges in distribution to X and Y_n converges in distribution to Y. (The shortest way to prove this is by invoking the Stone-Weierstrass theorem, but one can also proceed by proving some version of Proposition 4.) What happens if the independence hypothesis is dropped?
- (iii) If X_n converges in distribution to X, show that for every $\varepsilon > 0$ there exists $K > 0$ such that $\mathbf{P}(|X_n| \geq K) < \varepsilon$ for all sufficiently large n. (That is to say, X_n is a tight sequence of random variables.)
- (iv) Show that X_n converges in distribution to X if and only if, after extending the probability space model if necessary, one can find copies $Z_1, Z_2, \ldots$ and Z of $X_1, X_2, \ldots$ and X respectively such that Z_n converges almost surely to Z. (*Hint:* use the Skorohod representation, Exercise 29 of Notes 0.)
- (v) If $X_1, X_2, \ldots$ converges in distribution to X, and $F : \mathbf{R} \to \mathbf{R}$ is continuous, show that $F(X_1), F(X_2), \ldots$ converges in distribution to $F(X)$. Generalise this claim to the case when X takes values in an arbitrary locally compact Hausdorff space.
- (vi) (Slutsky's theorem) If X_n converges in distribution to X, and Y_n converges in probability to a *deterministic* limit Y, show that $X_n + Y_n$ converges in distribution to $X + Y$, and $X_n Y_n$ converges in distribution to XY. (Hint: either use (iv), or else use (iii) to control some error terms.) This statement combines particularly well with (i). What happens if Y is not assumed to be deterministic?
- (vii) (Fatou lemma) If $G : \mathbf{R} \to [0, +\infty)$ is continuous, and X_n converges in distribution to X, show that $\liminf_{n\to\infty} \mathbf{E}G(X_n) \geq \mathbf{E}G(X)$.
- (viii) (Bounded convergence) If $G : \mathbf{R} \to \mathbf{R}$ is continuous and bounded, and X_n converges in distribution to X, show that $\lim_{n\to\infty} \mathbf{E}G(X_n) = \mathbf{E}G(X)$.
- (ix) (Dominated convergence) If X_n converges in distribution to X, and there is an absolutely integrable Y such that $|X_n| \leq Y$ almost surely for all n, show that $\lim_{n\to\infty} \mathbf{E}X_n = \mathbf{E}X$.

For future reference we also mention (but will not prove) Prokhorov's theorem that gives a partial converse to part (iii) of the above exercise:

Theorem 11 (Prokhorov's theorem) Let $X_1, X_2, \ldots$ be a sequence of real random variables which is tight (that is, for every $\varepsilon > 0$ there exists $K > 0$ such that $\mathbf{P}(|X_n| \geq K) < \varepsilon$ for all sufficiently large n). Then there exists a subsequence X_{n_j} which converges in distribution to some random variable X (which may possibly be modeled by a different probability space model than the $X_1, X_2, \ldots$.)

The proof of this theorem relies on the Riesz representation theorem, and is beyond the scope of this course; but see for instance Exercise 29 of this previous blog post. (See also the closely related Helly selection theorem, covered in Exercise 30 of the same post.)

— 1. The Lindeberg approach to the central limit theorem —

We now give the Lindeberg argument establishing the central limit theorem. The proof splits into two unrelated components. The first component is to establish the central limit theorem for a *single* choice of underlying random variable X. The second component is to show that the limiting distribution of $\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu)$ is *universal* in the sense that it does not depend on the choice of underlying random variable. Putting the two components together gives Theorem 8.

We begin with the first component of the argument. One could use the Bernoulli distribution from Exercise 9 as the choice of underlying random variable, but a simpler choice of distribution (in the sense that no appeal to Stirling's formula is required) is the normal distribution $N(\mu, \sigma^2)$ itself. The key computation is:

> **Lemma 12 (Sum of independent Gaussians)** Let X_1, X_2 be independent real random variables with normal distributions $N(\mu_1, \sigma_1^2)$, $N(\mu_2, \sigma_2^2)$ respectively for some $\mu_1, \mu_2 \in \mathbf{R}$ and $\sigma_1, \sigma_2 > 0$. Then $X_1 + X_2$ has the normal distribution $N(\mu_1 + \mu_2, \sigma_1^2 + \sigma_2^2)$.

This is of course consistent with the additivity of mean and variance for independent random variables, given that random variables with the distribution $N(\mu, \sigma^2)$ have mean μ and variance σ^2.

Proof: By subtracting μ_1 and μ_2 from X_1, X_2 respectively, we may normalise $\mu_1 = \mu_2 = 0$, by dividing through by $\sqrt{\sigma_1^2 + \sigma_2^2}$ we may also normalise $\sigma_1^2 + \sigma_2^2 = 1$. Thus

$$\mathbf{P}(X_1 \in E_1) = \frac{1}{\sqrt{2\pi}\sigma_1} \int_{E_1} e^{-x_1^2/2\sigma_1^2}\, dx_1$$

and

$$\mathbf{P}(X_2 \in E_2) = \frac{1}{\sqrt{2\pi}\sigma_2} \int_{E_2} e^{-x_2^2/2\sigma_2^2}\, dx_2$$

for any Borel sets $E_1, E_2 \subset \mathbf{R}$. As X_1, X_2 are independent, this implies that

$$\mathbf{P}((X_1, X_2) \in E) = \frac{1}{2\pi\sigma_1\sigma_2} \int_{E} e^{-x_1^2/2\sigma_1^2 - x_2^2/2\sigma_2^2}\, dx_1 dx_2$$

for any Borel set $E \subset \mathbf{R}^2$ (this follows from the uniqueness of product measure, or equivalently one can use the monotone class lemma starting from the case when E is a finite boolean combination of product sets $E_1 \times E_2$). In particular, we have

$$\mathbf{P}(X_1 + X_2 \leq t) = \frac{1}{2\pi\sigma_1\sigma_2} \int_{\mathbf{R}^2} 1_{x_1+x_2 \leq t} e^{-x_1^2/2\sigma_1^2 - x_2^2/2\sigma_2^2}\, dx_1 dx_2$$

for any $t \in \mathbf{R}$. Making the change of variables $x = x_1 + x_2$ (and using the Fubini-Tonelli theorem as necessary) we can write the right-hand side as

$$\frac{1}{2\pi\sigma_1\sigma_2} \int_{-\infty}^{t} \int_{\mathbf{R}} e^{-x_1^2/2\sigma_1^2 - (x-x_1)^2/2\sigma_2^2}\, dx_1 dx. \quad (2)$$

We can complete the square using $\sigma_1^2 + \sigma_2^2 = 1$ to write (after some routine algebra)

$$e^{-x_1^2/2\sigma_1^2 - (x-x_1)/2\sigma_2^2} = e^{-x^2/2} e^{-(x_1 - x\sigma_1^2)^2/2\sigma_1^2\sigma_2^2}$$

so on using the identity $\frac{1}{\sqrt{2\pi}\sigma} \int_{\mathbf{R}} e^{-(x_1-\mu)^2/2\sigma^2}\ dx_1 = 1$ for any $\mu \in \mathbf{R}$ and $\sigma > 0$, we can write (2) as

$$\frac{1}{\sqrt{2\pi}} \int_{-\infty}^{t} e^{-x^2/2}\ dx$$

and so $X_1 + X_2$ has the cumulative distribution function of $N(0,1)$, giving the claim. □

In the next section we give an alternate proof of the above lemma using the machinery of characteristic functions. A more geometric argument can be given as follows. With the same normalisations as in the above proof, we can write $\sigma_1 = \cos\theta$ and $\sigma_2 = \sin\theta$ for some $0 < \theta < \pi/2$. Then we can write $X_1 = \cos\theta Y_1$ and $X_2 = \sin\theta Y_2$ where Y_1, Y_2 are iid copies of $N(0,1)$. But the joint probability density function $(x_1, x_2) \mapsto \frac{1}{2\pi} e^{-(x_1^2+x_2^2)/2}$ of (Y_1, Y_2) is rotation invariant, so $X_1 + X_2 = \cos\theta Y_1 + \sin\theta Y_2$ has the same distribution as Y_1, and the claim follows.

From the above lemma we see that if $X_1, X_2, \ldots$ are iid copies of a normal distribution $N(\mu, \sigma^2)$ of mean μ and variance σ^2, then $S_n = X_1 + \ldots X_n$ has distribution $N(n\mu, n\sigma^2)$, and hence $\frac{\sqrt{n}}{\sigma}(\frac{S_n}{n} - \mu)$ has distribution exactly equal to $N(0,1)$. Thus the central limit theorem is clearly true in this case.

> **Exercise 13 (Probabilistic interpretation of convolution)** Let $f, g : \mathbf{R} \to [0, +\infty]$ be measurable functions with $\int_{\mathbf{R}} f(x)\ dx = \int_{\mathbf{R}} g(x)\ dx = 1$. Define the convolution $f*g$ of f and g to be
>
> $$f*g(x) := \int_{\mathbf{R}} f(y) g(x-y)\ dy.$$
>
> Show that if X, Y are independent real random variables with probability density functions f, g respectively, then $X + Y$ has probability density function $f*g$.

Now we turn to the general case of the central limit theorem. By subtracting μ from X (and from each of the X_i) we may normalise $\mu = 0$; by dividing X (and each of the X_i) by σ we may also normalise $\sigma = 1$. Thus $\mathbf{E}X = 0, \mathbf{E}X^2 = 1$, and our task is to show that

$$\mathbf{E}G(\frac{X_1 + \ldots + X_n}{\sqrt{n}}) \to \mathbf{E}G(N) \qquad (3)$$

as $n \to \infty$, for any continuous compactly supported functions $G : \mathbf{R} \to \mathbf{R}$, where N is a random variable distributed according to $N(0,1)$ (possibly modeled by a different probability space than the original $X_1, X_2, \ldots$). Since any continuous compactly supported function can be approximated uniformly by smooth compactly supported functions (as can be seen from the Weierstrass or Stone-Weierstrass theorems), it suffices to show this for smooth compactly supported G.

Let $Y_1, Y_2, \ldots$ be iid copies of N; by extending the probability space used to model $X_1, X_2, \ldots$ (using Proposition 26 from Notes 2), we can model the $Y_1, Y_2, \ldots$ and $X_1, X_2, \ldots$ by a common model, in such a way that the combined collection $(X_n, Y_n)_{n=1}^{\infty}$ of random variables are jointly independent. As we have already proved the central limit theorem in the normally distributed case, we already have

$$\mathbf{E}G(\frac{Y_1+...+Y_n}{\sqrt{n}}) \to \mathbf{E}G(N) \quad (4)$$

as $n \to \infty$ (indeed we even have equality here). So it suffices to show that

$$\mathbf{E}G(\frac{X_1+...+X_n}{\sqrt{n}}) - \mathbf{E}G(\frac{Y_1+...+Y_n}{\sqrt{n}}) \to 0 \quad (5)$$

as $n \to \infty$.

We first establish this claim under the additional simplifying assumption of a finite third moment: $\mathbf{E}|X|^3 < \infty$. Rather than swap all of the $X_1, ..., X_n$ with all of the $Y_1, ..., Y_n$, let us just swap the final X_n to a Y_n, that is to say let us consider the expression

$$\mathbf{E}G(\frac{X_1+...+X_n}{\sqrt{n}}) - \mathbf{E}G(\frac{X_1+...+X_{n-1}+Y_n}{\sqrt{n}}).$$

Writing $Z := \frac{X_1+...+X_{n-1}}{\sqrt{n}}$, we can write this as

$$\mathbf{E}G(Z + \frac{1}{\sqrt{n}}X_n) - \mathbf{E}G(Z + \frac{1}{\sqrt{n}}Y_n).$$

To compute this expression we use Taylor expansion. As G is smooth and compactly supported, the first three derivatives of G are bounded, leading to the Taylor approximation

$$G(Z + \frac{1}{\sqrt{n}}X_n) = G(Z) + \frac{1}{\sqrt{n}}G'(Z)X_n + \frac{1}{2n}G''(Z)X_n^2 + O(\frac{1}{n^{3/2}}|X_n|^3)$$

where the implied constant depends on G. Taking expectations, we conclude that

$$\mathbf{E}G(Z + \frac{1}{\sqrt{n}}X_n) = \mathbf{E}G(Z) + \frac{1}{\sqrt{n}}\mathbf{E}G'(Z)X_n + \frac{1}{2n}\mathbf{E}G''(Z)X_n^2$$
$$+O(\frac{1}{n^{3/2}}\mathbf{E}|X|^3).$$

Now for a key point: as the random variable Z only depends on $X_1, ..., X_{n-1}$, it is independent of X_n, and so we can decouple the expectations to obtain

$$\mathbf{E}G(Z + \frac{1}{\sqrt{n}}X_n) = \mathbf{E}G(Z) + \frac{1}{\sqrt{n}}\mathbf{E}G'(Z)\mathbf{E}X_n + \frac{1}{2n}\mathbf{E}G''(Z)\mathbf{E}X_n^2$$
$$+O(\frac{1}{n^{3/2}}\mathbf{E}|X|^3).$$

The same considerations apply after swapping X_n with Y_n (which also has a bounded third moment):

$$\mathbf{E}G(Z + \frac{1}{\sqrt{n}}Y_n) = \mathbf{E}G(Z) + \frac{1}{\sqrt{n}}\mathbf{E}G'(Z)\mathbf{E}Y_n + \frac{1}{2n}\mathbf{E}G''(Z)\mathbf{E}Y_n^2$$
$$+O(\frac{1}{n^{3/2}}).$$

But by hypothesis, X_n and Y_n have matching moments to second order: $\mathbf{E}X_n = \mathbf{E}Y_n = 0$ and $\mathbf{E}X_n^2 = \mathbf{E}Y_n^2 = 1$. Thus on subtraction we have

$$\mathbf{E}G(\frac{X_1+...+X_n}{\sqrt{n}}) - \mathbf{E}G(\frac{X_1+...+X_{n-1}+Y_n}{\sqrt{n}})$$
$$= O(\frac{1}{n^{3/2}}(1 + \mathbf{E}|X|^3)).$$

A similar argument (permuting the indices, and replacing some of the X_i with Y_i) gives

$$\mathbf{E}G(\frac{X_1+...+X_i+Y_{i+1}+...+Y_n}{\sqrt{n}})$$
$$-\mathbf{E}G(\frac{X_1+...+X_{i-1}+Y_i+...+Y_n}{\sqrt{n}})$$
$$= O(\frac{1}{n^{3/2}}(1+\mathbf{E}|X|^3))$$

for all $1 \leq i \leq n$. Summing the telescoping series, we conclude that

$$\mathbf{E}G(\frac{X_1+...+X_n}{\sqrt{n}}) - \mathbf{E}G(\frac{Y_1+...+Y_n}{\sqrt{n}}) = O(\frac{1}{n^{1/2}}(1+\mathbf{E}|X|^3))$$

which gives (5). Note how it was important to Taylor expand to at least third order to obtain a total error bound that went to zero, which explains why it is the first two moments $\mathbf{E}X, \mathbf{E}X^2$ of X (or equivalently, the mean and variance) that play such a decisive role in the central limit theorem.

Now we remove the hypothesis of finite third moment. As in the previous set of notes, we use the truncation method, taking advantage of the "room" inherent in the $\frac{1}{n^{1/2}}$ factor in the error term of the above analysis. For technical reasons we have to modify the usual truncation slightly to preserve the mean zero condition. Let $\varepsilon > 0$. We split $X_i = X_{i,\leq} + X_{i,>}$, where $X_{i,\leq} := X_i 1_{|X_i| \leq \varepsilon\sqrt{n}} - \mu_n$ and $X_{i,>} := X_i 1_{|X_i| > \varepsilon\sqrt{n}} + \mu_n$ with $\mu_n := \mathbf{E}X1_{|X|\leq\varepsilon\sqrt{n}}$; we split $X = X_\leq + X_>$ respectively (we are suppressing the dependence of $X_{i,\leq}, X_{i,>}, X_\leq, X_>$ on ε and n to simplify the notation). The random variables $X_\leq$ (and $X_{i,\leq}$) are chosen to have mean zero, but their variance is not quite one. However, from dominated convergence, the quantity $\mu_n := \mathbf{E}X1_{|X|\leq\varepsilon\sqrt{n}}$ converges to 0, and the variance $\sigma_n^2 := \mathbf{Var}(X_\leq) = \mathbf{E}|X|^2 1_{|X|\leq\varepsilon\sqrt{n}} - \mu_n^2$ converges to 1, as $n \to \infty$.

Let $Y_1', ..., Y_n'$ be iid copies of $N(0, \sigma_n^2)$. The previous arguments then give

$$\mathbf{E}G(\frac{X_{1,\leq}+...+X_{n,\leq}}{\sqrt{n}}) - \mathbf{E}G(\frac{Y_1'+...+Y_n'}{\sqrt{n}}) = O(\frac{1}{n^{1/2}}(1+\mathbf{E}\left|X_\leq\right|^3))$$

for n large enough. We can bound

$$\mathbf{E}\left|X_\leq\right|^3 \leq \mathbf{E}\varepsilon n^{1/2}|X|^2 = \varepsilon n^{1/2}$$

and thus

$$\mathbf{E}G(\frac{X_{1,\leq}+...+X_{n,\leq}}{\sqrt{n}}) - \mathbf{E}G(\frac{Y_1'+...+Y_n'}{\sqrt{n}}) = O(\varepsilon)$$

for large enough n. By Lemma 12, $\frac{Y_1'+...+Y_n'}{\sqrt{n}}$ is distributed as $N(0, \sigma_n)$, and hence

$$\mathbf{E}G(\frac{Y_1'+...+Y_n'}{\sqrt{n}}) \to \mathbf{E}G(N)$$

as $n \to \infty$. We conclude that

$$\mathbf{E}G(\frac{X_{1,\leq}+...+X_{n,\leq}}{\sqrt{n}}) = \mathbf{E}G(N) + O(\varepsilon).$$

Next, we consider the error term

$$\frac{X_{1,>}+...+X_{n,>}}{\sqrt{n}}.$$

The variable $X_>$ has mean zero, and by dominated convergence, the variance of $X_>$ goes to zero as $n \to \infty$. The above error term is also mean zero and has the same variance as $X_>$. In particular, from Cauchy-Schwarz we have

$$\mathbf{E}\left|\frac{X_{1,>} + \ldots + X_{n,>}}{\sqrt{n}}\right| \to 0.$$

As G is smooth and compactly supported, it is Lipschitz continuous, and hence

$$G(\frac{X_1 + \ldots + X_n}{\sqrt{n}}) = G(\frac{X_{1,\leq} + \ldots + X_{n,\leq}}{\sqrt{n}}) + O(\left|\frac{X_{1,>} + \ldots + X_{n,>}}{\sqrt{n}}\right|).$$

Taking expectations, and then combining all these estimates, we conclude that

$$\mathbf{E}G(\frac{X_1 + \ldots + X_n}{\sqrt{n}}) = \mathbf{E}G(N) + O(\varepsilon)$$

for n sufficiently large; letting n go to infinity we obtain (3) as required. This concludes the proof of Theorem 8.

The above argument can be generalised to a central limit theorem for certain triangular arrays, known as the Lindeberg central limit theorem:

Exercise 14 (Lindeberg central limit theorem) Let k_n be a sequence of natural numbers going to infinity in n. For each natural number n, let $X_{1,n}, \ldots, X_{k_n,n}$ be jointly independent real random variables of mean zero and finite variance. (We do not require the random variables $(X_{1,n}, \ldots, X_{k_n,n})$ to be jointly independent in n, or even to be modeled by a common probability space.) Let σ_n be defined by

$$\sigma_n^2 := \sum_{i=1}^{k_n} \mathbf{Var}(X_{i,n})$$

and assume that $\sigma_n > 0$ for all n.

- (i) If one assumes the Lindeberg condition that

$$\frac{1}{\sigma_n^2} \sum_{i=1}^{k_n} \mathbf{E}(|X_{i,n}|^2 1_{|X_{i,n}| > \varepsilon \sigma_n}) \to 0$$

as $n \to \infty$ for any $\varepsilon > 0$, then show that the random variables $\frac{X_{1,n} + \ldots + X_{k_n,n}}{\sigma_n}$ converge in distribution to a random variable with the normal distribution $N(0,1)$.

- (ii) Show that the Lindeberg condition implies the *Feller condition*

$$\frac{1}{\sigma_n^2} \max_{1 \leq i \leq k_n} \mathbf{E}|X_{i,n}|^2 \to 0$$

as $n \to \infty$.

Note that Theorem 8 (after normalising to the mean zero case $\mu = 0$) corresponds to the special case $k_n = n$ and $X_{i,n} := X_i$ (or, if one wishes, $X_{i,n} := X_i/\sqrt{n}$) of the Lindeberg central limit theorem. It was shown by Feller that if the situation as in the above exercise and the Feller

condition holds, then the Lindeberg condition is necessary as well as sufficient for $X_{1,n} + ... + X_{k_n,n}$ to converge in distribution to a random variable with normal distribution $N(0,\sigma^2)$; the combined result is sometimes known as the *Lindeberg-Feller theorem*.

Exercise 15 (Weak Berry-Esséen theorem) Let $X_1, ..., X_n$ be iid copies of a real random variable X of mean zero, unit variance, and finite third moment.

- (i) Show that

$$\mathbf{E}G(\frac{X_1 + ... + X_n}{\sqrt{n}}) = \mathbf{E}G(N) + O(n^{-1/2}(\sup_{x \in \mathbf{R}} |G'''(x)|)\mathbf{E}|X|^3)$$

  whenever G is three times continuously differentiable and compactly supported, with N distributed as $N(0,1)$ and the implied constant in the $O()$ notation absolute.
- (ii) Show that

$$\mathbf{P}(\frac{X_1 + ... + X_n}{\sqrt{n}} \leq t) = \mathbf{P}(N \leq t) + O(n^{-1/2}\mathbf{E}|X|^3)^{1/4}$$

  for any $t \in \mathbf{R}$, with the implied constant absolute.

We will strengthen the conclusion of this theorem in Theorem 37 below.

Remark 16 The Lindeberg exchange method explains why the limiting distribution of statistics such as $\frac{X_1+...+X_n}{\sqrt{n}}$ depend primarily on the first two moments of the component random variables X_i, if there is a suitable amount of independence between the X_i. It turns out that there is an analogous application of the Lindeberg method in random matrix theory, which (very roughly speaking) asserts that appropriate statistics of random matrices such as $(X_{ij})_{1 \leq i,j \leq n}$ depend primarily on the first four moments of the matrix components X_{ij}, if there is a suitable amount of independence between the X_{ij}. See for instance this survey article of Van Vu and myself for more discussion of this. The Lindeberg method also suggests that the more moments of X one assumes to match with the Gaussian variable N, the faster the rate of convergence (because one can use higher order Taylor expansions).

We now use the Lindeberg central limit theorem to obtain the converse direction of the Kolmogorov three-series theorem (Exercise 29 of Notes 3).

Exercise 17 (Kolmogorov three-series theorem, converse direction) Let $X_1, X_2, ...$ be a sequence of jointly independent real random variables, with the property that the series $\sum_{n=1}^\infty X_n$ is almost surely convergent (i.e., the partial sums are almost surely convergent), and let $A > 0$.

- (i) Show that $\sum_{n=1}^\infty \mathbf{P}(|X_n| > A)$ is finite. (*Hint:* argue by contradiction and use the second Borel-Cantelli lemma.)
- (ii) Show that $\sum_{n=1}^\infty \mathbf{Var}(X_n 1_{|X_n| \leq A})$ is finite. (*Hint:* first use (i) and the Borel-Cantelli lemma to reduce to the case where $|X_n| \leq A$ almost surely. If $\sum_{n=1}^\infty \mathbf{Var}(X_n)$

is infinite, use Exercise 14 to show that $\sum_{n=1}^N (X_n - \mathbf{E}X_n)/\sqrt{\sum_{n=1}^N \mathbf{Var}(X_n)}$ converges in distribution to a standard normal distribution, and use this to contradict the almost sure convergence of $\sum_{n=1}^\infty X_n$.)

- (iii) Show that the series $\sum_{n=1}^\infty \mathbf{E}X_n 1_{|X_n| \leq A}$ is convergent. (*Hint:* reduce as before to the case where $|X_n| \leq A$ almost surely, and apply the forward direction of the three-series theorem to $\sum_n X_n - \mathbf{E}X_n$.)

— 2. The Fourier-analytic approach to the central limit theorem —

Let us now give the standard Fourier-analytic proof of the central limit theorem. Given any real random variable X, we introduce the characteristic function $\varphi_X : \mathbf{R} \to \mathbf{C}$, defined by the formula

$$\varphi_X(t) := \mathbf{E}e^{itX}. \quad (6)$$

Equivalently, φ_X is the Fourier transform of the probability measure μ_X. One should caution that the term "characteristic function" has several other unrelated meanings in mathematics; particularly confusingly, in real analysis "characteristic function" is used to denote what in probability one would call an "indicator function". Note that no moment hypotheses are required to define the characteristic function, because the random variable e^{itX} is bounded even when X is not absolutely integrable.

Example 18 The signed Bernoulli distribution, which takes the values $+1$ and -1 with probabilities of $1/2$ each, has characteristic function $\phi(t) = \cos(t)$.

Most of the standard random variables in probability have characteristic functions that are quite simple and explicit. For the purposes of proving the central limit theorem, the most important such explicit form of the characteristic function is of the normal distribution:

Exercise 19 Show that the normal distribution $N(\mu, \sigma^2)$ has characteristic function $\phi(t) = e^{it\mu}e^{-\sigma^2 t^2/2}$.

We record the explicit characteristic functions of some other standard distributions:

Exercise 20 Let $\lambda > 0$, and let X be a Poisson random variable with intensity λ, thus X takes values in the non-negative integers with $\mathbf{P}(X = k) = e^{-\lambda}\frac{\lambda^k}{k!}$. Show that $\varphi_X(t) = \exp(\lambda(e^{it} - 1))$ for all $t \in \mathbf{R}$.

Exercise 21 Let X be uniformly distributed in some interval $[a, b]$. Show that $\varphi_X(t) = \frac{e^{itb} - e^{ita}}{it(b-a)}$ for all non-zero t.

Exercise 22 Let $x_0 \in \mathbf{R}$ and $\gamma > 0$, and let X be a Cauchy random variable with parameters x_0, γ, which means that X is a real random variable with probability density function $\frac{\gamma}{\pi((x-x_0)^2+\gamma^2)}$. Show that $\varphi_X(t) = e^{ix_0 t}e^{-\gamma|t|}$ for all $t \in \mathbf{R}$.

The characteristic function is clearly bounded in magnitude by 1, and equals 1 at the origin. By the dominated convergence theorem, φ_X is continuous in t.

Exercise 23 (Riemann-Lebesgue lemma) Show that if X is a real random variable that has an absolutely integrable probability density function f, then $\varphi_X(t) \to 0$ as $t \to \infty$. (*Hint:* first show the claim when f is a finite linear combination of intervals, then for the general case show that f can be approximated in L^1 norm by such finite linear combinations.) Note from Example 18 that the claim can fail if X does not have a probability density function. (In Fourier analysis, this fact is known as the Riemann-Lebesgue lemma.)

Exercise 24 Show that the characteristic function φ_X of a real random variable X is in fact uniformly continuous on its domain.

Let X be a real random variable. If we Taylor expand e^{itX} and formally interchange the series and expectation, we arrive at the heuristic identity

$$\varphi_X(t) = \sum_{k=0}^{\infty} \frac{(it)^k}{k!} \mathbf{E}X^k \quad (7)$$

which thus interprets the characteristic function of a real random variable X as a kind of generating function for the moments. One rigorous version of this identity is as follows.

Exercise 25 (Taylor expansion of characteristic function) Let X be a real random variable with finite k^{th} moment for some $k \geq 1$. Show that φ_X is k times continuously differentiable, with

$$\frac{d^j}{dt^j}\varphi_X(t) = i^j \mathbf{E}X^j e^{itX}$$

for all $0 \leq j \leq k$. Conclude in particular the partial Taylor expansion

$$\varphi_X(t) = \sum_{j=0}^{k} \frac{(it)^j}{j!} \mathbf{E}X^j + o(|t|^k)$$

where $o(|t|^k)$ is a quantity that goes to zero as $t \to 0$, times $|t|^k$.

Exercise 26 Let X be a real random variable, and assume that it is *subgaussian* in the sense that there exist constants $C, c > 0$ such that

$$\mathbf{P}(|X| \geq t) \leq Ce^{-ct^2}$$

for all $t \in \mathbf{R}$. (Thus for instance a bounded random variable is subgaussian, as is any gaussian random variable.) Rigorously establish (7) in this case, and show that the series converges locally uniformly in t.

Note that the characteristic function depends only on the distribution of X: if X and Y are equal in distribution, then $\varphi_X = \varphi_Y$. The converse statement is true also: if $\varphi_X = \varphi_Y$, then X and Y are equal in distribution. This follows from a more general (and useful) fact, known as Lévy's continuity theorem.

Theorem 27 (Lévy continuity theorem, special case) Let X_n be a sequence of real random variables, and let X be an additional real random variable. Then the following statements are equivalent:

- (i) φ_{X_n} converges pointwise to φ_X.
- (ii) X_n converges in distribution to X.

Proof: The implication of (i) from (ii) is immediate from (6) and Exercise 10(viii).

Now suppose that (i) holds, and we wish to show that (ii) holds. We need to show that

$$\mathbf{E}G(X_n) \to \mathbf{E}G(X)$$

whenever $G : \mathbf{R} \to \mathbf{R}$ is a continuous, compactly supported function. As in the Lindeberg argument, it suffices to prove this when G is smooth and compactly supported, in particular G is a Schwartz function (infinitely differentiable, with all derivatives rapidly decreasing). But then we have the Fourier inversion formula

$$G(X_n) = \int_{\mathbf{R}} \hat{G}(t) e^{it \cdot X_n}\ dt$$

where

$$\hat{G}(t) := \frac{1}{2\pi} \int_{\mathbf{R}} G(x) e^{-it \cdot x}\ dx$$

is Schwartz, and is in particular absolutely integrable (see e.g. these lecture notes of mine). From the Fubini-Tonelli theorem, we thus have

$$\mathbf{E}G(X_n) = \int_{\mathbf{R}} \hat{G}(t) \varphi_{X_n}(t)\ dt \qquad (8)$$

and similarly for X. The claim now follows from the Lebesgue dominated convergence theorem. □

Remark 28 Setting $X_n := Y$ for all n, we see in particular the previous claim that $\varphi_X = \varphi_Y$ if and only if X, Y have the same distribution. It is instructive to use the above proof as a guide to prove this claim directly.

There is one subtlety with the Lévy continuity theorem: it is possible for a sequence φ_{X_n} of characteristic functions to converge pointwise, but for the limit to not be the characteristic function of any random variable, in which case X_n will not converge in distribution. For instance, if $X_n \equiv N(0,n)$, then $\varphi_{X_n}(t) = e^{-t^2/2n}$ converges pointwise to $1_{t=0}$ for any t, but this is clearly not the characteristic function of any random variable (as characteristic functions are continuous). However, this lack of continuity is the only obstruction:

Exercise 29 (Lévy's continuity theorem, full version) Let X_n be a sequence of real valued random variables. Suppose that φ_{X_n} converges pointwise to a limit φ. Show that the following are equivalent:

- (i) φ is continuous at 0.
- (ii) X_n is a tight sequence (as in Exercise 10(iii)).
- (iii) φ is the characteristic function of a real valued random variable X (possibly after extending the sample space).
- (iv) X_n converges in distribution to some real valued random variable X (possibly after extending the sample space).

Hint: To get from (ii) to the other conclusions, use Theorem 11 and Theorem 27. To get back to (ii) from (i), use (8) for a suitable Schwartz function G. The other implications are easy once Theorem 27 is in hand.

Remark 30 Lévy's continuity theorem is very similar in spirit to Weyl's criterion in equidistribution theory.

Exercise 31 (Esséen concentration inequality) Let X be a random variable taking values in $\mathbf{R}$. Then for any $r > 0$, $\varepsilon > 0$, show that

$$\sup_{x_0 \in \mathbf{R}} \mathbf{P}(|X - x_0| \leq r) \leq C_\varepsilon r \int_{t \in \mathbf{R}: |t| \leq \varepsilon/r} |\varphi_X(t)|\, dt \qquad (9)$$

for some constant C_ε depending only on ε. (*Hint:* Use (8) for a suitable Schwartz function G.) The left-hand side of (9) (as well as higher dimensional analogues of this quantity) is known as the *small ball probability* of X at radius r.

In Fourier analysis, we learn that the Fourier transform is a particularly well-suited tool for studying convolutions. The probability theory analogue of this fact is that characteristic functions are a particularly well-suited tool for studying sums of independent random variables. More precisely, we have

Exercise 32 (Fourier identities) Let X, Y be independent real random variables. Then

$$\varphi_{X+Y}(t) = \varphi_X(t)\varphi_Y(t) \qquad (10)$$

for all $t \in V$. Also, for any scalar c, one has

$$\varphi_{cX}(t) = \varphi_X(\overline{c}t)$$

and more generally, for any linear transformation $T : V \to V$, one has

$$\varphi_{TX}(t) = \varphi_X(T^* t).$$

Remark 33 Note that this identity (10), combined with Exercise 19 and Remark 28, gives a quick alternate proof of Lemma 12.

In particular, we have the simple relationship

$$\varphi_{S_n/\sqrt{n}}(t) = \varphi_X(t/\sqrt{n})^n \qquad (11)$$

that describes the characteristic function of $S_n/\sqrt{n}$ in terms of that of X.

We now have enough machinery to give a quick proof of the central limit theorem:

Proof: (Fourier proof of Theorem 8) We may normalise X to have mean zero and variance 1. By Exercise 25, we thus have

$$\varphi_X(t) = 1 - t^2/2 + o(|t|^2)$$

for sufficiently small t, or equivalently

$$\varphi_X(t) = \exp(-t^2/2 + o(|t|^2))$$

for sufficiently small t. Applying (11), we conclude that

$$\varphi_{S_n/\sqrt{n}}(t) \to \exp(-t^2/2)$$

as $n \to \infty$ for any fixed t. But by Exercise 19, $\exp(-t^2/2)$ is the characteristic function of the normal distribution $N(0,1)$. The claim now follows from the Lévy continuity theorem. □

The above machinery extends without difficulty to vector-valued random variables $\vec{X} = (X_1, ..., X_d)$ taking values in $\mathbf{R}^d$. The analogue of the characteristic function is then the function $\varphi_{\vec{X}} : \mathbf{R}^d \to \mathbf{C}$ defined by

$$\varphi_{\vec{X}}(t_1, ..., t_d) := \mathbf{E} e^{i(t_1 X_1 + ... + t_d X_d)}.$$

We leave the routine extension of the above results and proofs to the higher dimensional case to the reader. Most interesting is what happens to the central limit theorem:

Exercise 34 (Vector-valued central limit theorem) Let $\vec{X} = (X_1, ..., X_d)$ be a random variable taking values in $\mathbf{R}^d$ with finite second moment. Define the covariance matrix $\Sigma(\vec{X})$ to be the $d \times d$ matrix Σ whose ij^{th} entry is the covariance $\mathbf{E}(X_i - \mathbf{E}(X_i))(X_j - \mathbf{E}(X_j))$.

- Show that the covariance matrix is positive semi-definite real symmetric.
- Conversely, given any positive definite real symmetric $d \times d$ matrix Σ and $\mu \in \mathbf{R}^d$, show that the multivariate normal distribution $N(\mu, \Sigma)_{\mathbf{R}^d}$, given by the absolutely continuous measure

$$\frac{1}{((2\pi)^d \det \Sigma)^{1/2}} e^{-(x-\mu)\cdot\Sigma^{-1}(x-\mu)/2}\, dx,$$

  has mean μ and covariance matrix Σ, and has a characteristic function given by

$$\varphi_{N(\mu,\Sigma)_{\mathbf{R}^d}}(t) = e^{i\mu\cdot t} e^{-t\cdot\Sigma t/2}.$$

  How would one define the normal distribution $N(\mu, \Sigma)_{\mathbf{R}^d}$ if Σ degenerated to be merely positive semi-definite instead of positive definite?
- If $\vec{S}_n := \vec{X}_1 + ... + \vec{X}_n$ is the sum of n iid copies of $\vec{X}$, show that $\frac{\vec{S}_n - n\mu}{\sqrt{n}}$ converges in distribution to $N(0, \Sigma(X))_{\mathbf{R}^d}$. (For this exercise, you may assume without proof that the Lévy continuity theorem extends to $\mathbf{R}^d$.)

Exercise 35 (Complex central limit theorem) Let X be a complex random variable of mean $\mu \in \mathbf{C}$, whose real and imaginary parts have variance $\sigma^2/2$ and covariance 0. Let $X_1, ..., X_n \equiv X$ be iid copies of X. Show that as $n \to \infty$, the normalised sums $\frac{\sqrt{n}}{\sigma}(\frac{X_1 + ... + X_n}{n} - \mu)$ converge in distribution to the standard complex gaussian $N(0,1)_{\mathbf{C}}$, defined as the measure μ on $\mathbf{C}$ with

$$\mu(S) := \frac{1}{\pi} \int_S e^{-|z|^2}\, dz$$

for Borel $S \in \mathbf{C}$, where dz is Lebesgue measure on $\mathbf{C}$ (identified with $\mathbf{R}^2$ in the usual fashion).

Exercise 36 Use characteristic functions and the truncation argument to give an alternate proof of the Lindeberg central limit theorem (Theorem 14).

A more sophisticated version of the Fourier-analytic method gives a more quantitative form of the central limit theorem, namely the Berry-Esséen theorem.

Theorem 37 (Berry-Esséen theorem) Let X have mean zero and unit variance. Let $S_n := X_1 + ... + X_n$, where $X_1, ..., X_n$ are iid copies of X. Then we have

$$\mathbf{P}(\frac{S_n}{\sqrt{n}} < a) = \mathbf{P}(N < a) + O(\frac{1}{\sqrt{n}}(\mathbf{E}|X|^3)) \quad (12)$$

uniformly for all $a \in \mathbf{R}$, where N has the distribution of $N(0,1)$, and the implied constant is absolute.

Proof: (Optional) Write $\varepsilon := \mathbf{E}|X|^3/\sqrt{n}$; our task is to show that

$$\mathbf{P}(\frac{S_n}{\sqrt{n}} < a) = \mathbf{P}(N < a) + O(\varepsilon) \quad (13)$$

for all a. We may of course assume that $\varepsilon < 1$, as the claim is trivial otherwise. (In particular, X has finite third moment.)

Let $c > 0$ be a small absolute constant to be chosen later. Let $\eta : \mathbf{R} \to \mathbf{R}$ be an non-negative Schwartz function with total mass 1 whose Fourier transform is supported in $[-c, c]$; such an η can be constructed by taking the inverse Fourier transform of a smooth function supported on $[-c/2, c/2]$ and equal to one at the origin, and then multiplying that transform by its complex conjugate to make it non-negative.

Let $\psi : \mathbf{R} \to \mathbf{R}$ be the smoothed out version of $1_{(-\infty,0]}$, defined as

$$\psi(x) := \int_{\mathbf{R}} 1_{(-\infty,0]}(x - \varepsilon y)\eta(y)\, dy$$
$$= \int_{x/\varepsilon}^{\infty} \eta(y)\, dy.$$

Observe that ψ is decreasing from 1 to 0. As η is rapidly decreasing and has mean one, we also have the bound

$$\psi(x) = 1_{(-\infty,0]}(x) + O_\eta((1 + |x|/\varepsilon)^{-100}) \quad (14)$$

(say) for any x, where subscript indicates that the implied constant depends on η.

We claim that it suffices to show that

$$\mathbf{E}\psi(\frac{S_n}{\sqrt{n}} - a) = \mathbf{E}\psi(N - a) + O_\eta(\varepsilon) \quad (15)$$

for every a, where the subscript means that the implied constant depends on η. Indeed, suppose

that (15) held. Replacing a by $a + C\varepsilon$ and $a - C\varepsilon$ for some large absolute constant C and subtracting, we have

$$\mathbf{E}\psi(\frac{S_n}{\sqrt{n}} - a - C\varepsilon) - \psi(\frac{S_n}{\sqrt{n}} - a + C\varepsilon) = \mathbf{E}\psi(N - a - C\varepsilon) - \mathbf{E}\psi(N - a + C\varepsilon)$$
$$+O_\eta(\varepsilon)$$

for any a. From (14) we see that the function $\psi(t - a - C\varepsilon) - \psi(t - a + C\varepsilon)$ is bounded by $O_{\eta,C}(1 + |x - a|/\varepsilon)^{-100})$, and hence by the bounded probability density of N

$$\mathbf{E}\psi(N - a - C\varepsilon) - \mathbf{E}\psi(N - a + C\varepsilon) = O_{\eta,C}(\varepsilon).$$

Also, $\psi(t - a - C\varepsilon) - \psi(t - a + C\varepsilon)$ is non-negative, and for C large enough, it is bounded from below by (say) $1/2$ on $[a, a + \varepsilon]$. We conclude (after choosing C appropriately) that

$$\mathbf{E}(\frac{S_n}{\sqrt{n}} \in [a, a + \varepsilon]) = O_\eta(\varepsilon) \quad (16)$$

for all real a. This implies that

$$\mathbf{E}(1 + \left|\frac{S_n}{\sqrt{n}} - a\right|/\varepsilon)^{-100} = O_\eta(\varepsilon),$$

as can be seen by covering the real line by intervals $[a + j\varepsilon, a + (j + 1)\varepsilon)]$ and applying (16) to each such interval. From (14) we conclude that

$$\mathbf{E}\psi(\frac{S_n}{\sqrt{n}} - a) = \mathbf{E}1(-\infty, 0](\frac{S_n}{\sqrt{n}} - a) + O_\eta(\varepsilon).$$

A similar argument gives

$$\mathbf{E}\psi(N - a) = \mathbf{E}1(-\infty, 0](N - a) + O_\eta(\varepsilon),$$

and (13) then follows from (15).

It remains to establish (15). We write

$$\psi(x) = \lim_{M\to\infty} \int_{x/\varepsilon}^{x/\varepsilon+M} \eta(y)\, dy$$

(with the expression in the limit being uniformly bounded) and

$$\eta(y) = \int_{\mathbf{R}} \hat{\eta}(t) e^{ity}\, dt$$

to conclude (after applying Fubini's theorem) that

$$\psi(x) = \lim_{M\to\infty} \int_{-c}^{c} \hat{\eta}(t) e^{itx/\varepsilon} \frac{e^{itM} - 1}{it}\, dt$$

using the compact support of $\hat{\eta}$. Hence

$$\psi(\frac{S_n}{\sqrt{n}} - a) - \psi(N - a) = \lim_{M\to\infty} \int_{-c}^{c} \hat{\eta}(t)(e^{it\frac{S_n}{\sqrt{n}}/\varepsilon} - e^{itN/\varepsilon}) \frac{e^{itM} - 1}{it} e^{-ita/\varepsilon}\, dt.$$

By the fundamental theorem of calculus we have $e^{it\frac{S_n}{\sqrt{n}}/\varepsilon} - e^{itN/\varepsilon} = O(|t|(\left|\frac{S_n}{\sqrt{n}}\right| + |N|)/\varepsilon)$. This factor of $|t|$ cancels a similar factor on the denominator to make the expression inside the limit dominated by an absolutely integrable random variable. Thus by the dominated convergence theorem

$$\mathbf{E}\psi(\frac{S_n}{\sqrt{n}}) - \mathbf{E}\psi(N) = \lim_{M\to\infty} \int_{-c}^{c} \hat{\eta}(t)(\varphi_{S_n/\sqrt{n}}(t/\varepsilon) - \varphi_N(t/\varepsilon))\frac{e^{itM}-1}{it} e^{-ita/\varepsilon}\ dt$$

and so it suffices to show that

$$\int_{-c}^{c} \hat{\eta}(t)(\varphi_{S_n/\sqrt{n}}(t/\varepsilon) - \varphi_N(t/\varepsilon))\frac{e^{itM}-1}{it} e^{-ita/\varepsilon}\ dt = O_\eta(\varepsilon)$$

uniformly in M. Applying the triangle inequality and the compact support of $\hat{\eta}$, it suffices to show that

$$\int_{|t|\leq c} |\ \varphi_{S_n/\sqrt{n}}(t/\varepsilon) - \varphi_N(t/\varepsilon)|\ \frac{dt}{|t|} = O(\varepsilon). \quad (17)$$

From Taylor expansion we have

$$e^{itX} = 1 + itX - \frac{t^2}{2}X^2 + O(|t|^3|X|^3)$$

for any t; taking expectations and using the definition of ε we have

$$\varphi_X(t) = 1 - t^2/2 + O(\varepsilon\sqrt{n}|t|^3)$$

and in particular

$$\varphi_X(t) = \exp(-t^2/2 + O(\varepsilon\sqrt{n}|t|^3))$$

if $|t| \leq c/\varepsilon\sqrt{n}$ and c is small enough. Applying (11), we conclude that

$$\varphi_{S_n/\sqrt{n}}(t) = \exp(-t^2/2 + O(\varepsilon|t|^3))$$

if $|t| \leq c/\varepsilon$. Meanwhile, from Exercise 19 we have $\varphi_N(t) = \exp(-t^2/2)$. Elementary calculus then gives us

$$\left|\varphi_{S_n/\sqrt{n}}(t) - \varphi_N(t)\right| \leq O(\varepsilon|t|^3 \exp(-t^2/4))$$

(say) if c is small enough. Inserting this bound into (17) we obtain the claim. □

> **Exercise 38** Show that the error terms in Theorem 37 are sharp (up to constants) when X is a signed Bernoulli random variable, or more precisely when X takes values in $\{-1, +1\}$ with probability $1/2$ for each.

> **Exercise 39** Let X_n be a sequence of real random variables which converge in distribution to a real random variable X, and let Y_n be a sequence of real random variables which converge in distribution to a real random variable Y. Suppose that, for each n, X_n and Y_n are independent, and suppose also that X and Y are independent. Show that $X_n + Y_n$ converges in distribution to $X + Y$. (*Hint:* use the Lévy continuity theorem.)

The following exercise shows that the central limit theorem fails when the variance is infinite.

> **Exercise 40** Let $X_1, X_2, \ldots$ be iid copies of an absolutely integrable random variable X of mean zero.
>
> - (i) In this part we assume that X is *symmetric*, which means that X and $-X$ have the same distribution. Show that for any $t > 0$ and $M > 0$,

$$\mathbf{P}(X_1 + ... + X_n \geq t)$$

$$\geq \frac{1}{2}\mathbf{P}(X_1 1_{|X_1| \leq M} + ... + X_n 1_{|X_n| \leq M} \geq t).$$

(*Hint:* relate both sides of this inequality to the probability of the event that $X_1 1_{|X_1| \leq M} + ... + X_n 1_{|X_n| \leq M} \geq t$ and $X_1 1_{|X_1| > M} + ... + X_n 1_{|X_n| > M} \geq 0$, using the symmetry of the situation.)

- (ii) If X is symmetric and $\frac{X_1+...+X_n}{\sqrt{n}}$ converges in distribution to a real random variable Y, show that X has finite variance. (*Hint:* if this is not the case, then $X1_{|X| \leq M}$ will have arbitrarily large variance as M increases. On the other hand, $\mathbf{P}(Y \geq u)$ can be made arbitrarily small by taking u large enough. For a large threshold M, apply (i) (with $t = u\sqrt{n}$) to obtain a contradiction.
- (iii) Generalise (ii) by removing the hypothesis that X is symmetric. (*Hint:* apply the *symmetrisation* trick of replacing X by $X - X'$, where X' is an independent copy of X, and use the previous exercise. One may need to utilise some truncation arguments to show that $X - X'$ has infinite variance whenever X has infinite variance.)

Exercise 41

- (i) If X is a real random variable of mean zero and variance σ^2, and t is a real number, show that

$$\varphi_X(t) = 1 + O(\sigma^2 t^2)$$

and that

$$\varphi_X(t) = 1 - \frac{1}{2}\sigma^2 t^2 + O(t^2 \mathbf{E} \min(|X|^2, t|X|^3)).$$

(*Hint:* first establish the Taylor bounds $e^{itX} = 1 + itX + O(t^2|X|^2)$ and $e^{itX} = 1 + itX + \frac{1}{2}t^2|X|^2 + O(t^3|X|^3)$.)

- (ii) Establish the pointwise inequality

$$|z_1 ... z_n - z'_1 ... z'_n| \leq \sum_{i=1}^n |z_i - z'_i|$$

whenever $z_1, ..., z_n, z'_1, ..., z'_n$ are complex numbers in the disk $\{z \in \mathbf{C} : |z| \leq 1\}$.

- (iii) Suppose that for each n, $X_{1,n}, ..., X_{n,n}$ are jointly independent real random variables of mean zero and finite variance, obeying the uniform bound

$$|X_{i,n}| \leq \varepsilon_n \sqrt{n}$$

for all $i = 1, \ldots, n$ and some ε_n going to zero as $n \to \infty$, and also obeying the variance bound

$$\sum_{i=1}^{n} \mathbf{Var}(X_{i,n}) \to \sigma^2$$

as $n \to \infty$ for some $0 < \sigma < \infty$. If $S_n := X_{1,n} + \ldots + X_{n,n}$, use (i) and (ii) to show that

$$\varphi_{S_n/\sqrt{n}}(t) \to e^{-\sigma^2 t^2/2}$$

as $n \to \infty$ for any given real t.

- (iv) Use (iii) and a truncation argument to give an alternate proof of the Lindeberg central limit theorem (Theorem 14). (Note: one has to address the issue that truncating a random variable may alter its mean slightly.)

— 3. The moment method —

The above Fourier-analytic proof of the central limit theorem is one of the quickest (and slickest) proofs available for this theorem, and is accordingly the "standard" proof given in probability textbooks. However, it relies quite heavily on the Fourier-analytic identities in Exercise 32, which in turn are extremely dependent on both the commutative nature of the situation (as it uses the identity $e^{A+B} = e^A e^B$) and on the independence of the situation (as it uses identities of the form $\mathbf{E}(e^A e^B) = (\mathbf{E}e^A)(\mathbf{E}e^B)$). As one or both of these factors can be absent when trying to generalise this theorem, it is also important to look for non-Fourier based methods to prove results such as the central limit theorem. These methods often lead to proofs that are lengthier and more technical than the Fourier proofs, but also tend to be more robust.

The most elementary (but still remarkably effective) method available in this regard is the *moment method*, which we have already used in previous notes. In principle, this method is equivalent to the Fourier method, through the identity (7); but in practice, the moment method proofs tend to look somewhat different than the Fourier-analytic ones, and it is often more apparent how to modify them to non-independent or non-commutative settings.

We first need an analogue of the Lévy continuity theorem. Here we encounter a technical issue: whereas the Fourier phases $x \mapsto e^{itx}$ were bounded, the moment functions $x \mapsto x^k$ become unbounded at infinity. However, one can deal with this issue as long as one has sufficient decay:

Exercise 42 (Subgaussian random variables) Let X be a real random variable. Show that the following statements are equivalent:

- (i) There exist $C, c > 0$ such that $\mathbf{P}(|X| \geq t) \leq Ce^{-ct^2}$ for all $t > 0$.
- (ii) There exist $C' > 0$ such that $\mathbf{E}|X|^k \leq (C'k)^{k/2}$ for all $k \geq 1$.
- (iii) There exist $C'', c'' > 0$ such that $\mathbf{E}e^{tX} \leq C''e^{c''t^2}$ for all $t \in \mathbf{R}$.

Furthermore, show that if (i) holds for some C, c, then (ii) holds for C' depending only on C, c, and similarly for any of the other implications. Variables obeying (i), (ii), or (iii) are called *subgaussian*. The function $t \mapsto \mathbf{E}e^{tX}$ is known as the moment generating function of X; it is of course closely related to the characteristic function of X.

Exercise 43 Use the truncation method to show that in order to prove the central limit theorem (Theorem 8), it suffices to do so in the case when the underlying random variable X is bounded (and in particular subgaussian).

Once we restrict to the subgaussian case, we have an analogue of the Lévy continuity theorem:

Theorem 44 (Moment continuity theorem) Let X_n be a sequence of real random variables, and let X be a subgaussian random variable. Suppose that for every $k = 0, 1, 2, \ldots$, $\mathbf{E}X_n^k$ converges pointwise to $\mathbf{E}X^k$. Then X_n converges in distribution to X.

Proof: Let $k = 0, 1, 2, \ldots$, then by the preceding exercise we have

$$\mathbf{E}|X|^{k+1} \leq (C(k+1))^{(k+1)/2}$$

for some C independent of k. In particular,

$$\mathbf{E}|X_n|^{k+1} \leq 2(C(k+1))^{(k+1)/2}$$

when n is sufficiently large depending on k. From Taylor's theorem with remainder (and Stirling's formula (1)) we conclude

$$\varphi_{X_n}(t) = \sum_{j=0}^{k} \frac{(it)^j}{j!}\mathbf{E}X_n^j + O((C(k+1))^{-(k+1)/2}|t|^{k+1})$$

uniformly in t, for n sufficiently large. Similarly for X. Taking limits using (i) we see that

$$\limsup_{n\to\infty} \left|\varphi_{X_n}(t) - \varphi_X(t)\right| = O((C(k+1))^{-(k+1)/2}|t|^{k+1}).$$

Then letting $k \to \infty$, keeping t fixed, we see that $\varphi_{X_n}(t)$ converges pointwise to $\varphi_X(t)$ for each t, and the claim now follows from the Lévy continuity theorem. □

We remark that in place of Stirling's formula, one can use the cruder lower bound $n! \geq n^n/e^n$, which is immediate from the observation that $n^n/n!$ occurs in the Taylor expansion of e^n.

Remark 45 One corollary of Theorem 44 is that the distribution of a subgaussian random variable is uniquely determined by its moments (actually, this could already be deduced from Exercise 26 and Remark 28). The situation can fail for distributions with slower tails, for much the same reason that a smooth function is not determined by its derivatives at one point if that function is not analytic.

The Fourier inversion formula provides an easy way to recover the distribution from the

characteristic function. Recovering a distribution from its moments is more difficult, and sometimes requires tools such as analytic continuation; this problem is known as the *inverse moment problem* and will not be discussed here.

Exercise 46 (Converse direction of moment continuity theorem) Let X_n be a sequence of uniformly subgaussian random variables (thus there exist $C, c > 0$ such that $\mathbf{P}(|X_n| \geq t) \leq Ce^{-ct^2}$ for all $t > 0$ and all n), and suppose X_n converges in distribution to a limit X. Show that for any $k = 0, 1, 2, \ldots$, $\mathbf{E}X_n^k$ converges pointwise to $\mathbf{E}X^k$.

We now give the moment method proof of the central limit theorem. As discussed above we may assume without loss of generality that X is bounded (and in particular subgaussian); we may also normalise X to have mean zero and unit variance. By Theorem 44, it suffices to show that

$$\mathbf{E}S_n^k/n^{k/2} \to \mathbf{E}N^k$$

for all $k = 0, 1, 2, \ldots$, where $N \equiv N(0,1)_{\mathbf{R}}$ is a standard gaussian variable.

The moments $\mathbf{E}N^k$ were already computed in Exercise 37 of Notes 1. So now we need to compute $\mathbf{E}S_n^k/n^{k/2}$. Using linearity of expectation, we can expand this as

$$\mathbf{E}S_n^k/n^{k/2} = n^{-k/2} \sum_{1 \leq i_1, \ldots, i_k \leq n} \mathbf{E}X_{i_1} \ldots X_{i_k}.$$

To understand this expression, let us first look at some small values of k.

- For $k = 0$, this expression is trivially 1.
- For $k = 1$, this expression is trivially 0, thanks to the mean zero hypothesis on X.
- For $k = 2$, we can split this expression into the diagonal and off-diagonal components:

$$n^{-1} \sum_{1 \leq i \leq n} \mathbf{E}X_i^2 + n^{-1} \sum_{1 \leq i < j \leq n} \mathbf{E}2X_iX_j.$$

  Each summand in the first sum is 1, as X has unit variance. Each summand in the second sum is 0, as the X_i have mean zero and are independent. So the second moment $\mathbf{E}Z_n^2$ is 1.
- For $k = 3$, we have a similar expansion

$$n^{-3/2} \sum_{1 \leq i \leq n} \mathbf{E}X_i^3 + n^{-3/2} \sum_{1 \leq i < j \leq n} \mathbf{E}3X_i^2X_j + 3X_iX_j^2$$

$$+n^{-3/2} \sum_{1 \leq i < j < k \leq n} \mathbf{E}6X_iX_jX_k.$$

  The summands in the latter two sums vanish because of the (joint) independence and mean zero hypotheses. The summands in the first sum need not vanish, but are $O(1)$, so the first term is $O(n^{-1/2})$, which is asymptotically negligible, so the third moment $\mathbf{E}Z_n^3$ goes to 0.

- For $k = 4$, the expansion becomes quite complicated:

$$n^{-2} \sum_{1 \leq i \leq n} \mathbf{E} X_i^4 + n^{-2} \sum_{1 \leq i < j \leq n} \mathbf{E} 4X_i^3 X_j + 6X_i^2 X_j^2 + 4X_i X_j^3$$

$$+n^{-2} \sum_{1 \leq i < j < k \leq n} \mathbf{E} 12X_i^2 X_j X_k + 12X_i X_j^2 X_k + 12X_i X_j X_k^2$$

$$+n^{-2} \sum_{1 \leq i < j < k < l \leq n} \mathbf{E} 24X_i X_j X_k X_l.$$

  Again, most terms vanish, except for the first sum, which is $O(n^{-1})$ and is asymptotically negligible, and the sum $n^{-2} \sum_{1 \leq i < j \leq n} \mathbf{E} 6X_i^2 X_j^2$, which by the independence and unit variance assumptions works out to $n^{-2} 6 \binom{n}{2} = 3 + o(1)$. Thus the fourth moment $\mathbf{E} Z_n^4$ goes to 3 (as it should).

Now we tackle the general case. Ordering the indices $i_1, \ldots, i_k$ as $j_1 < \ldots < j_m$ for some $1 \leq m \leq k$, with each j_r occuring with multiplicity $a_r \geq 1$ and using elementary enumerative combinatorics, we see that $\mathbf{E} S_n^k / n^{k/2}$ is the sum of all terms of the form

$$n^{-k/2} \sum_{1 \leq j_1 < \ldots < j_m \leq n} c_{k,a_1,\ldots,a_m} \mathbf{E} X_{j_1}^{a_1} \ldots X_{j_m}^{a_m} \quad (18)$$

where $1 \leq m \leq k$, $a_1, \ldots, a_m$ are positive integers adding up to k, and $c_{k,a_1,\ldots,a_m}$ is the multinomial coefficient

$$c_{k,a_1,\ldots,a_m} := \frac{k!}{a_{(1)!} \ldots a_{(m)!}}.$$

The total number of such terms depends only on k (in fact, it is 2^{k-1} (exercise!), though we will not need this fact).

As we already saw from the small k examples, most of the terms vanish, and many of the other terms are negligible in the limit $n \to \infty$. Indeed, if any of the a_r are equal to 1, then every summand in (18) vanishes, by joint independence and the mean zero hypothesis. Thus, we may restrict attention to those expressions (18) for which all the a_r are at least 2. Since the a_r sum up to k, we conclude that m is at most $k/2$.

On the other hand, the total number of summands in (18) is clearly at most n^m (in fact it is $\binom{n}{m}$), and the summands are bounded (for fixed k) since X is bounded. Thus, if m is *strictly* less than $k/2$, then the expression in (18) is $O(n^{m-k/2})$ and goes to zero as $n \to \infty$. So, asymptotically, the only terms (18) which are still relevant are those for which m is *equal* to $k/2$. This already shows that $\mathbf{E} Z_n^k$ goes to zero when k is odd. When k is even, the only surviving term in the limit is now when $m = k/2$ and $a_1 = \ldots = a_m = 2$. But then by independence and unit variance, the expectation in (18) is 1, and so this term is equal to

$$n^{-k/2}\binom{n}{m} c_{k,2,\dots,2} = \frac{1}{(k/2)!}\frac{k!}{2^{k/2}} + o(1),$$

and the main term is happily equal to the moment $\mathbf{E}N^k$ as computed in Exercise 37 of Notes 1. (One could also appeal to Lemma 12 here, specialising to the case when X is normally distributed, to explain this coincidence.) This concludes the proof of the central limit theorem.

Exercise 47 (Chernoff bound) Let $X_1, \dots, X_n$ be iid copies of a real random variable X of mean zero and unit variance, which is subgaussian in the sense of Exercise 42. Write $S_n := X_1 + \dots + X_n$.

- (i) Show that there exist $c'' > 0$ such that $\mathbf{E}e^{tX} \leq e^{c''t^2}$ for all $t \in \mathbf{R}$. Conclude that $\mathbf{E}e^{tS_n/\sqrt{n}} \leq e^{c''t^2}$ for all $t \in \mathbf{R}$. (*Hint:* the first claim follows directly from Exercise 42 when $|t| \geq 1$; for $|t| < 1$, use the Taylor approximation $e^{tX} = 1 + tX + O(t^2(e^{2X} + e^{-2X}))$.)
- (ii) Conclude the Chernoff bound

$$\mathbf{P}(\left|\frac{S_n}{\sqrt{n}}\right| \geq \lambda) \leq Ce^{-c\lambda^2}$$

for some $C, c > 0$, all $\lambda > 0$, and all $n \geq 1$.

Exercise 48 (Erdös-Kac theorem) For any natural number $x \geq 100$, let n be a natural number drawn uniformly at random from the natural numbers $\{1, \dots, x\}$, and let $\omega(n)$ denote the number of distinct prime factors of n.

- (i) Show that for any $k = 0, 1, 2, \dots$, one has

$$\mathbf{E}(\frac{\omega(n) - \log\log x}{\sqrt{\log\log x}})^k \to 0$$

as $x \to \infty$ if k is odd, and

$$\mathbf{E}(\frac{\omega(n) - \log\log x}{\sqrt{\log\log x}})^k \to \frac{k!}{2^{k/2}(k/2)!}$$

as $x \to \infty$ if k is even. (*Hint:* adapt the arguments in Exercise 16 of Notes 3, estimating $\omega(n) - \log\log x$ by $\sum_{p \leq x^{1/10k}} 1_{p \mid n} - \frac{1}{p}$, using Mertens' theorem and induction on k to deal with lower order errors, and treating the random variables $1_{p \mid n} - \frac{1}{p}$ as being approximately independent and approximately of mean zero.)
- (ii) Establish the Erdös-Kac theorem

$$\frac{1}{x}\left|\{n \leq x : a \leq \frac{\omega(n) - \log\log x}{\sqrt{\log\log x}} \leq b\}\right| \to \frac{1}{\sqrt{2\pi}}\int_a^b e^{-x^2/2}\,dx$$

> as $x \to \infty$ for any fixed $a < b$.
>
> Informally, the Erdös-Kac theorem asserts that $\omega(n)$ behaves like $N(\log\log n, \log\log n)$ for "random" n. Note that this refines the Hardy-Ramanujan theorem (Exercise 16 of Notes 3).

Comments (127)

 · 2 November 2015

may want to add "275A – probability theory" in the tags

[Added, thanks – T.]

 · 3 November 2015

In theorem 36, the assumption $E(|X|^3) < \infty$ is not really needed (since otherwise (11) holds trivially!)

[Assumption removed, thanks – T.]

 · 3 November 2015

You should use … instead of ... and $\cdots$, resp.

 · 3 November 2015

In exercise 13(ii), "sup" is in fact "max", and the expectation should be dependent on the index i.

[Corrected, thanks – T.]

 · 4 November 2015

According to the Wikipedia articles on the CLT and the Berry-Esseen theorem:

1. The term "central limit theorem" (in German) was first used by Polya (in 1920) where the word "central" is due to its central importance in probability theory. The later interpretation of "central" in the sense that the CLT describes the behaviour of the distribution near its center is (according to Le Cam) due to the French school on probability.

2. The best implied constant in the error term in 11, is lower-bounded by 0.4097 (Esseen, 1956) and upper-bounded by 0.4748 (Shevtsova, 2011).

 · 4 November 2015

In the proof of Theorem 26 you briefly switch to R^d

[Corrected, thanks – T.]

 · 4 November 2015

A useful lemma is that if X_n converge in distribution they can be coupled to converge almost surely. This is also an instructive alternate approach to the second direction in Proposition 4.

[This was the intention of Exercise 10(iv), though I had previously only asked for convergence in probability, and have now strengthened the conclusion. -T.]

 · 4 November 2015

It seems that exercise 10(vi) generalizes Slutsky's theorem (which deals with a degenerate(!) limit Y), but in the Wikipedia article on Slutsky's theorem (according to the second note) it is claimed that the requirement that Y is degenerate(!) is important (the theorem is not (!) valid in general for a non-degenerate limit Y) – is this claim true?

[Oops, this was an error – the exercise has now been corrected. -T.]

 · 5 November 2015

Exercise 9, definition of distribution at the beginning has typo.

[Corrected, thanks – T.]

 **John Mangual** · 5 November 2015

The Lindberg approach is related to the "renormalization group approach", as I learned from Yakov Sinai.

However, I couldn't not reconcile the discussions of RG in hep-th and the discussion in the probability textbook.

 **Joop de Maat** · 5 November 2015

A typo ("equvialence" before Remark 3) and a question: How exactly are the two notions of distributions from Remark 3 similar ? You only indicated in what the where different, not in what they are alike.

 **Terence Tao** · 5 November 2015

Thanks for the correction! The similarity is both notions of convergence require testing against "nice" functions (continuous compactly supported functions in the case of convergence in distribution, and smooth compactly supported functions in the case of convergence in the sense of distributions). In particular, it is true that a sequence of random variables X_n converge in distribution to a random variable X if and only if the probability measures μ_{X_n} converge in the sense of distributions to μ_X. (But once one leaves the setting of probability measures, one can have more exotic distributions as the distributional limit, e.g. derivatives of delta functions.)

 **Venky** · 14 November 2015

In the Lindberg approach, I was trying to see where the Gaussian nature of the pdf of Y_i is important other than (3). Can one say that an implication of CLT is that the Gaussian pdf is the only pdf that preserves its functional form over countable addition of independent variables? Great notes, thanks.

 **Terence Tao** · 14 November 2015

Yes, the gaussian is the only finite-variance stable law, and the Lindeberg argument explains why there can be at most one such law. (But there are other stable laws in the infinite variance case, which lie outside of the reach of the CLT: for instance, the Cauchy distribution is also stable.)

· 14 November 2015

I have a question on the Example 5, we have known that Xn converges to X in distribution, where Xn is uniform distributed on {1/n, 2/n, ..., n/n} and X is uniform[0,1]. How can we check that if Xn converges to X in prob?

Terence Tao · 14 November 2015

This is not possible from the information given, because we do not know the joint distribution of (X, X_n), only the individual distributions of X and X_n; the joint distribution is irrelevant for the question of whether there is convergence in distribution, but is very relevant for convergence in probability For instance, X_n could be X rounded up to the next multiple of $1/n$, in which case X_n converges to X in probability (and almost surely). Or, X_n could be independent of X, in which case it will not converge in probability or almost surely to X.

Terence Tao · 18 November 2015

There appears to be a strange issue with one of the wordpress LaTeX renderers right now, as some simple expressions such as $\varphi = 0$ are not rendering properly while others such as φ are fine. For those in the class following the notes online, a temporary workaround is to hover one's mouse over the improperly rendered images to see the uncompiled LaTeX. I do not seem to be able to fix the issue on my end, but I'm hoping that the problem will resolve itself shortly (at which point I will remove this comment).

Terence Tao · 19 November 2015

OK, the problem seems to be that as of about 24 hours ago, wordpress has become unable to render any latex fragment that it has not previously rendered, presumably due to the machine that does all the new renderings being somehow offline. I'd be interested to see if other wordpress.com sites are having the same issue, in which case it would make sense to report it. In the meantime, I've rolled back these notes to an earlier version for now to avoid the rendering errors. *[Update: it now appears that the issue is affecting all wordpress.com blogs, and has been reported. -T.]*

ianmarqz · 19 November 2015

I have the same problem of well rendering : (sample: https://juanmarqz.wordpress.com/about/sum-of-reciprocals/

275A, Notes 5: Variants of the central limit theorem | What's new · 19 November 2015

[...] the previous set of notes we established the central limit theorem, which we formulate here as [...]

Dillon · 23 November 2015

The characteristic function for a uniform distribution on an interval (Exercise 21) is missing an i in the denominator. Also some of the preceding exercises still say F for the characteristic function where φ would be more consistent.

[Corrected, thanks – T.]

· 24 November 2015

In exercise 10 (vii), should it be liminf EG(X_n)>= EG(X)?

[Corrected, thanks – T.]

· 30 November 2015

'non-engative' should be 'non-negative' (search and replace time)

[Corrected, thanks – T.]

· 30 November 2015

'Levi continuity theorem' should be 'Lévy continuity theorem' (search and replace again)

[Corrected, thanks – T.]

jura05 · 2 December 2015

thank you for these lectures!
some typos:
– formula after (14) – missing term O(.)
– before (15): instead of "below by 1/2 on [0,eps]" should be "below by 1/2 on [a,a+eps]"
– proof of (14) goes for the special case a=0; although there is no difference.. maybe it is convenient to write a general bound
$|E\psi(X) - E\psi(Y)| \leq \int_{-c}^{c} |\varphi_X(t/\varepsilon) - \varphi_Y(t/\varepsilon)|\, dt/t$ and apply it to X=S_n/sqrt{n}-a, Y=N-a

[Corrected, thanks – T.]

· 30 May 2016

Thanks for the great notes. Small typos:

1. inequality (2) and the following equation are missing a square in the $(x-x_1)^2$ term.

2. Closed bracket missing in Ex 17 (ii)

3. Ex 17 (iii), the indicator function doesn't appear correctly.

[Corrected, thanks – T.]

· 30 May 2016

What is known about the convergence rate in Erdos-Kac theorem?

Terence Tao · 30 May 2016

I think this paper by Harper represents the current state of the art: http://www.ams.org/mathscinet-getitem?mr=2507311

Anonymous · 2 October 2016

In Exercise 14, for each n, $(X_{1,n}, ..., X_{k_n,n})$ are jointly independent. What do you mean by "We do not require the random variables $(X_{1,n}, ..., X_{k_n,n})$ to be jointly independent in n"? Would you illustrate by giving an example that what could be dependent here?

[For instance, we do not require the tuple $(X_{1,1}, ..., X_{k_1,1})$ to be independent of $(X_{1,2}, ..., X_{k_2,2})$. -T.]

Anonymous · 4 October 2016

In Exercise 25, can one avoid calculating the j-th derivative of $\varphi_X(t)$ by just directly integrating the Taylor expansion of e^{it}?

[This should work also, as long as one is careful with estimating error terms – T.]

Anonymous · 4 October 2016

For Exercise 19, all the textbooks I have seen use the contour integrals. Can one do it with other methods than the one in complex analysis?

[One such method begins by observing that the pdf of the normal distribution obeys an ODE which then translates to an ODE for the characteristic function (cf. Stein's method). -T]

Sam · 13 November 2024

Dear professor Tao:

Could you elaborate a bit more on this method by translating an ODE for the pdf of the normal distribution to an ODE for the characteristic function?

Terence Tao · 14 November 2024

For instance, the pdf $f(x) = \frac{1}{2\pi} e^{-x^2/2}$ obeys the ODE $f'(x) = -xf(x)$. I invite you to take Fourier transforms of both sides and derive a corresponding ODE for the characteristic function $\Phi(t)$ of this pdf.

Sam · 16 November 2024

Thank you professor. If one takes the derivative of the characteristic function inside the integral using dominated convergence, does one still need to justify manipulating complex valued functions?

Anonymous · 4 October 2016

In the Fourier proof of Theorem 8, would you elaborate the "equivalently" part? Is the parenthesis a typo there?

[We are using the Taylor approximation $1 - x = \exp(-x + o(x))$ for small x (specifically, $x = t^2/2n - o(|t|^2/n)$). -T.]

Anonymous · 4 October 2016

By direct substitution, I got
$\varphi_X(t/\sqrt{n})^n = \exp(-t^2 + n \cdot o(|t|^2/n)) = \exp(-t^2 + \cdot o(|t|^2))$
For fixed \$t\$, when taking the limit, how do you get rid of the little o term?

Terence Tao · 4 October 2016

The asymptotic notation $o()$ here is with respect to the limit $n \to \infty$, rather than $t \to 0$,so the $o(|t|^2)$ term goes to zero in the limit when t is fixed and n goes to infinity. (By the way, I corrected a typo in my previous answer, in which I had omitted a factor of $1/n$ in the choice of x.)

Anonymous · 5 October 2016

In Exercise 14, what is the intuition of the Lindeberg condition?

$$\frac{1}{\sigma_n^2}\sum_{i=1}^{k_n}\mathbf{E}(|X_{i,n}|^2 1_{|X_{i,n}|>\varepsilon\sigma_n}) \to 0$$

Why would one expect such a condition implies the desired result?

> **Terence Tao** · 6 October 2016
>
> The central limit theorem can fail if there is a significant probability that one of the summands $X_{i,n}$ is so large that it is comparable to the total standard deviation σ_n. The Lindeberg condition is basically an assertion that these sort of outlier events are vanishingly rare (and their contribution to the sum has vanishingly small variance) in the limit $n \to \infty$.

Anonymous · 6 October 2016

I saw Billingsley in his Probability and Measure proves Remark 28 by proving the inversion formula:

If the probability measure μ has characteristic function φ and $\mu\{a\}=\mu\{b\}=0$, then

$$\mu(a,b] = \lim_{T\to\infty}\frac{1}{2\pi}\int_{-T}^{T}\frac{e^{-ita}-e^{-itb}}{it}\,dt.$$

This approach looks rather different from the one in the proof of Theorem 27, which uses the Schwartz functions and is much shorter. What makes the Schwartz function-approach faster and is it hinted somewhere that these two approaches are essentially the same?

> **Anonymous** · 8 October 2016
>
> Ah, the inversion formula should be
>
> $$\mu(a,b] = \lim_{T\to\infty}\frac{1}{2\pi}\int_{-T}^{T}\frac{e^{-ita}-e^{-itb}}{it}\varphi(t)dt.$$
>
> Actually, the proof of the inversion formula is not that long. The key step is just exchanging the integral using Fubini and taking the limit. Still it looks rather different from the Schwartz-function-approach.
>
> > **Terence Tao** · 8 October 2016
> >
> > Basically the difference lies in the use of the smoothing sums technique: replacing a rough cutoff such as $1_{[a,b]}$ with a smoother cutoff in order to enjoy better decay of the Fourier transform. The price one pays for this is that the formulae become a little less explicit, possibly leading to worse quantitative bounds in some cases, but in many applications this sort of loss is acceptable.

Anonymous · 12 October 2016

In theorem 11, can we replace random variables with random vectors?

[Yes; see for instance the linked exercise. -T.]

Anonymous · 13 October 2016

Also for Proposition 4 and Exercise 7 one can have random vectors instead? I'm wondering if i missed reading some of the notes about the random vector part or it is just trivial generalization.

Terence Tao · 14 October 2016

If one deletes (ii) then the remaining claims are equivalent in the vector-valued case. But it is not clear to me whether there is an analogue for (ii) in the vector-valued case (one has to define the cumulative distribution function properly, and specify exactly what a "point of continuity" would mean for that function); the proof given is restricted to the scalar case.

Anonymous · 14 October 2016

https://en.wikipedia.org/wiki/Cumulative_distribution_function#Multivariate_case

Anonymous · 13 October 2016

In Exercise 7, do you have a hint for (v) —>(i)? I don't have any idea why the boundary of E would have anything to do with continuity of a function.

[Consider sets E of the form $E = (-\infty, t]$. -T.]

Anonymous · 13 October 2016

In the proof of Proposition 4, is $PG(X_n)$ a typo? Likewise for several others?

[Corrected, thanks – T.]

Quantitative continuity estimates | What's new · 22 May 2017

[...] starting point for the Lindeberg exchange method in probability theory, discussed for instance in this previous post. The identity in (ii) can also be used in the Lindeberg exchange method; the terms in the [...]

Discotech · 25 June 2017

In exercise 35, should the exponent have a π? I think maybe it should not. Unless I'm misunderstanding the meaning of the Lebesgue integral over the complex plane, the measure defined isn't normalized to 1.

In the hint on exercise 40, do you mean to refer to the probability $P(X_1 1_{|X_1|>M} + \dots + X_n 1_{|X_d|>M} \geq 0)$ (greater than signs instead of less than signs)?

[Corrected, thanks – T.]

· 26 October 2018

In Slutsky's theorem, can one relax the condition regarding the convergence of Y_n to be convergence in distribution?

Terence Tao · 27 October 2018

When the limit Y is deterministic, convergence in distribution is the same as convergence in probability.

haonanz · 20 February 2019

Hello Prof. Tao, I have a question regard Exercise 7 prove ii) => v). I am thinking if I could use sigma algebra like proof. I believe the difficulty comes from the fact we have only convergence at continuity point. If we assume it's true for all t, then I think I can show that the set of E satisfy this property is closed under countable union and countable closure), then use the fact Borel sets are the smallest sigma algebra generated by the half-intervals to complete the proof. I wonder the fact we know those discontinuity is at most countably many points can somehow be used to show this generated sigma algebra still contains the half-intervals thus the borel set.

Terence Tao · 21 February 2019

I'm not sure a sigma algebra approach will be easy to execute, because the set of Borel sets B with ∂B of measure zero is not closed under countable union or countable intersection. (The strategy I had in mind instead is to approximate B by its interior and its exterior and use (iii), (iv).)

haonanz · 23 February 2019

A follow up question regard proving i)=>iii), I can prove the claim for the compact case using upper approximation of a CTS and compact supported function. Do you mind providing hint on how can I extend this to the closed case? Thanks again!

Terence Tao · 23 February 2019

Show that for large R, the probability $\mathbf{P}(|X| \leq R)$ is close to 1, then show that the probability $\mathbf{P}(|X_n| \leq 2R)$ is close to 1 for large n, thus the probability $\mathbf{P}(|X_n| > 2R)$ is small.

· 11 August 2024

Dear professor Tao:

As of the direction (i) implies (iii) in Exercise 7, I think the compact case follows directly from using the function 1_K in the definition of weak convergence. Are we supposed to use inner regularity of the probability measure to generalize to the closed set setting? If possible I would like to hear a little more on this earlier comment of yours on showing that the probability of $|X_n|$ being large is small, thank you.

[As already stated in the exercise, the hint is to use Urysohn's lemma. -T]

anonymous · 6 November 2019

There should be a k_n instead of n in point (ii) of exercise 14.

[Corrected, thanks – T]

· 18 April 2020

Let $X_1, X_2, \ldots$ be iid Bernoulli(p) random variables, that is, $P(X_i = 1) = p$ and $P(X_i = 0) = 1 - p$.

Let

$$D_n = \sum_{i=1}^{n} Y_i$$

where $Y_i = I_{(X_i=1, X_{i+1}=1)}$.

Show that $\sqrt{n}\left(\frac{D_n}{n} - p^2\right) \xrightarrow{D} N(0, \sigma^2)$ and determine σ^2.

How can we prove this by using some elementary tools?

 **Aditya Guha Roy** · 24 December 2020

You may try to solve it using the central limit theorem and Slutsky's theorem. I have not done this very exercise, but it seems that this can be done by using these tools. Try it and let us know here.

 **N is a number** · 1 January 2021

Yes that would work. I just checked it.

 **Aditya Guha Roy** · 23 December 2020

On line number 22 of these notes, you have written about applying Paley Zygmund inequality to get the mentioned lower bound.
I think there is a typo in labelling it as Exercise 42 of Notes 1 ; it should be Exercise 44 of Notes 1.

[Corrected, thanks – T.]

 **N is a number** · 1 January 2021

In Exercise 19, we see that from the definition of characteristic function it suffices to show that the result holds when $\mu = 0$ and $\sigma^2 = 1$. Now that calls for showing $\mathbf{E}[e^{itX}] = e^{-t^2/2}$ for all complex numbers t. Now rewriting the LHS as an integral one has a holomorphic function on the LHS and the RHS is also a holomorphic function. Now, one can prove this identity for the case when t is purely imaginary so that it is purely real by a direct evaluation of the integral which appears on the LHS.
Now, using the fact that non-trivial holomorphic maps have isolated zeroes, and identifying that the holomorphic map LHS – RHS is zero identically on the imaginary axis, one concludes that the identity holds for all complex numbers t.

Is that correct ?

[Sure, this works. One can also compute the integral for arbitrary t by completing the square and contour shifting instead of relying on analytic continuation. -T]

 **Aditya Guha Roy** · 1 January 2021

I think to show that $\mathbf{E}[e^{itX}]$ is holomorphic over $\mathbf{C}$, an easy way is to notice that it is the derivative of the holomorphic map $t \mapsto \mathbf{E}[Xe^{itX}]$ and hence the former must also be holomorphic.

 **Aditya Guha Roy** · 1 January 2021

Filling a gap in my previous comment: we need to notice that we can interchange the derivative and expectation, thanks to Lebesgue's dominated convergence theorem.

N is a number · 1 January 2021

Prof. Tao in lines 4-5 of the proof of Proposition 1 you have cited Theorem 24 from Notes 1. I think you wanted to cite Theorem 25 from Notes 1 (since here we have the premises satisfied for $\varepsilon = 2$).

[Corrected, thanks – T.]

Orestis Plevrakis · 17 March 2021

I tried to prove Levy's continuity theorem without using Prokhorov's theorem, but I failed. However, I cannot see why one must pass through such an argument to get Levy's theorem. Can you give some intuition?

Anonymous · 27 July 2021

In the proof of the moment method (after exercise 46) there are two mentions of exercise 36 in Notes 1, but it should be exercise 37.

Further, a question about theorem 44: in the final equation, the RHS seems to be tending to ∞ for fixed t. How can we argue that $\phi_{X_n}(t) \to \phi_X(t)$ point-wise for t?

[Corrected, thanks. There was a negative sign that was omitted by mistake in the exponent of the RHS that has now been restored – T.]

Aditya Guha Roy · 26 March 2022

In Exercise 15 (Weak Berry-Esseen bounds) part (i) is it the right way to do as follows:

We would do everything verbatim as in Lindeberg's proof except now we expand the Taylor series till the third derivative, and in the first step instead of specializing to finite third moment we consider the special case of finite fourth moment; then for the general case we use the same truncation method.

> **Aditya Guha Roy** · 26 March 2022
>
> More precisely:
> first of all we consider the case when the fourth moment of X is finite, then notice that we can just imitate the construction (X_n, Y_n) as done in Lindeberg's proof of the central limit theorem, with the only difference being that now we are expanding the Taylor series of $G(Z + \frac{1}{\sqrt{n}}X_n)$ upto third order derivatives to get
> $G(Z + \frac{1}{\sqrt{n}}X_n) = G(Z) + \frac{1}{\sqrt{n}}G'(Z)X_n + \frac{1}{2n}G''(Z)X_n^2 +$
> $\frac{1}{6\sqrt{n^3}}G'''(Z)X_n^3 + O(n^{-2}|X_n|^4)$;
> now we just imitate all the other steps of Lindeberg's proof (for the case of finite third moments) the only distinction being that now we keep carrying the third derivative term throughout,
> thus we would obtain at the final step
> $\mathbf{E}[G(\frac{X_1+.....+X_n}{\sqrt{n}})] - \frac{E}{[}G(N)] = c \cdot \sup_x|G'''(x)|\mathbf{E}[|X|^3]n^{-1/2} + o(n^{-1/2})$
> where c is an absolute constant (the role of G got nullified by taking the supremum).
>
> Now we relax the condition of finite fourth moment in the exact same manner by constructing a truncation and then bounding the remaining portion of the original series to show that $\mathbf{E}[\frac{|X_{1,>}+....+X_{n,>}|}{\sqrt{n}}] \to 0$ as $n \to \infty$ and then using the Lipschitz property of

G which is guaranteed as G is three times differentiable and hence at least continuously differentiable, and also compactly supported (thanks to the mean value theorem).
Now about the truncated portion in a similar way we show that it satisfies the conclusions we had for the finite fourth moment case (the same arguments as in Lindeberg's proof would go through).

Finally putting all these together would give us the weak Berry-Esseen bounds.

 **Aditya Guha Roy** · 26 March 2022

I see some typos: that should be $o(n^{-1/2})$, and $\mathbf{E}[G(N)]$ respectively in the 13th line of display containing the sup term.

 **Terence Tao** · 26 March 2022

This will not work because the implied constant in the error term depends on bounds on the fourth derivative of G, which will diverge in the limit. A Taylor expansion to third order will suffice here, with no truncation required.

 **Aditya Guha Roy** · 26 March 2022

Yes sir, I realize this now. Thank you for pointing it out to me.

 **Aditya Guha Roy** · 26 March 2022

Sir, towards the end of Lindeberg's proof you have said that this proves (3).
I think you meant to label the equation in the 5th line of display after Exercise 13 involving X_i s as (3) instead of the one with Y_i s.

[Corrected, thanks – T.]

 **Aditya Guha Roy** · 31 March 2022

Prof Tao in the statement of Berry Esseen theorem (Theorem 37) I think you may like to add the condition that X needs to have a finite third moment.
(Though technically speaking what you have written is still quite correct.)

Convergence of mean from almost sure convergence in a super-linear bounded mean setting | Aditya Guha Roy's weblog · 3 December 2023

[...] (together with Kolmogorov's 0-1 law) is demonstrated in the proof of the opening proposition here. In a later blogpost I shall discuss consequences of Proposition 1 which are of a similar [...]

 · 9 August 2024

Dear professor Tao:

To clarify, in the proof of Proposition 4, the fact that F_X has at most countable number of discontinuities is used to ensure the partition by the t_i exists, is this correct?

[Yes – T.]

 · 10 August 2024

Dear professor Tao;

Possible minor typos in the proof of Proposition 1:

t should be changed to t_i in the expectation of $G_\varepsilon(X_n)$, and the remark should be "the restriction to continuity points of F_X is necessary..."

[Corrected, thanks – T.]

· 16 August 2024

Dear professor Tao:

To deduce (i) from (iii) in Exercise 7, I can see the Urysohn's lemma is invoked to establish the compact case. From the compact case, it seems that one can get the the limit superior of the probability of X_n in K bounded by the probability of X in K plus the limit superior of the probability of $|X_n| > R$. Hence my question concerns only the factor 2 in your earlier comment. Thank you.

Terence Tao · 16 August 2024

Apply Urysohn's lemma to obtain a non-negative continuous function supported on $B(0, 2R)$ and equal to one on $B(0, R)$. Apply this function to both X_n and X to be able to compare the indicated probabilities $\mathbf{P}(|X| \leq R)$ and $\mathbf{P}(|X_n| \leq 2R)$.

· 17 August 2024

Dear professor Tao:

Thank you so much for the feedback. For a compact K, if we set U_i to be the set of distance smaller than $1/i$ to K, and apply Urysohn's lemma to find a continuous function f_i that equals 1 on K and 0 on the complement of U_i, then continuity from above establishes the compact case.

By (i) the probability of $X_nleq + / - R$ converges to the probability of $Xleq + / - R$, isn't it follows then that the probability of $|X_n|leqR$ converges to the probability of $|X|leqR$?

Terence Tao · 18 August 2024

Part (i) only gives one-sided bounds; it upper bounds the lim sup, but does not lower bound the lim inf.

· 20 August 2024

Dear professor Tao:

In the proof of De-Moivre Laplace theorem, is it advisable to split into the case of odd/even i as in Durrett's text?

· 20 August 2024

And if possible, can you provide a bit more hint on how to proceed here? thank you

Terence Tao · 20 August 2024

There are multiple ways to proceed here, which have varying degrees of efficacy. In general, I recommend you actually *try* multiple approaches, before asking for advice whether it is going to be work or not – these sorts of "failures" are in fact extremely instructive and crucial in building one's one mathematical confidence; they are part of why these problems are assigned as exercises in the first place. Nobody is going to

administer any sort of punishment to you for selecting an initial approach that turns out not to be fruitful.

See also my advice post "Ask yourself dumb questions – and answer them!", as well as this blog post of Tim Gowers warning about being stymied by "fake difficulties".

· 20 August 2024

Thank you so much for the encouragement professor Tao.

· 31 August 2024

Dear professor Tao:

In Exercise 10(iv) (Skorokhod's representation theorem), if one chooses the unit interval $[0,1]$ with Borel measure as the probability space model for the $Z_n := X_n^+$ (defined as in Exercise 29 of note 0) and $Z := X^+$, can you give some directions as to what the null set might be outside of which one has convergence almost surely?

· 2 September 2024

Dear professor Tao:

For any ω in $(0,1)$, I worked out that we can use $X^+(\omega)$ to bound both the limsup of $X_n^+(\omega)$ from above, and the liminf of $X_n^+(\omega)$ from below, therefore get convergence almost surely. May I ask if 0 and 1 capture the entire null set on which the convergence might fail?

· 8 September 2024

Dear professor Tao:

In the geometric argument for Lemma 12, is the fact that the joint density of independent Gaussians the density of a multivariate Gaussian variable assumed to be known, as the result of some Exercises in Note 2?

Terence Tao · 8 September 2024

The joint density of two independent random variables is always the tensor product of the individual densities (this is an easy consequence of Tonelli's theorem).

· 18 September 2024

Dear professor Tao:

To derive the error term in the Taylor approximation in the Lindeberg approach, one needs the sum of $G^{(3+i)}(Z)/(3+i)!$ times $(X_n/\sqrt{n})^i$ to be $O(1)$, where the implied constant depends on G. I'd like to ask for some little clarification on this particular step. Thank you.

· 18 September 2024

Dear professor Tao:

I apologize for the bothering. It is the Taylor Theorem with remainder.

· 20 September 2024

Dear Professor Tao:

In the Lindeberg approach to the CLT, the error term in the display right below the statement "The same considerations apply after swapping X_n and Y_n..." should be the same as above.

 · 20 September 2024

My mistake, Y_n is a copy of N.

 · 20 September 2024

Dear Professor Tao:

In the Lindeberg approach to the CLT, I doubt the expression for variance of $X_{\leq}$ involves $+\mu_n^2$ rather than $-\mu_n^2$.

 · 29 September 2024

Dear professor Tao:

To generalize the Lindeberg swapping trick to triangular arrays, I attempted extending the probability space for each row n to get iid normal variables $Y_{i,n}$ for i between 1 and n, with mean 0 and variance that equals the variance of $1/\sigma_n X_{k_n,n}$. Yet it's not clear that on subtraction the error term goes to zero with this construction. Can you provide some hint on how the Lindeberg condition is used in this setting? Thank you.

 · 30 September 2024

Dear professor Tao:

I see it now that one should extend the probability spaces to get a triangular array of normal variables such that each row have mean zero and variances of the corresponding row of the original variables. Yet it is still a bit unclear as to how the Lindeberg condition is used to control the error term. A little suggestion will be greatly appreciated, thank you.

 · 7 October 2024

Dear professor Tao:

For the Lindeberg CLT, denoting the variance of $X_{i,n}$ by $alpha_{i,n}^2$, and assuming finite third moment for the entire triangular array, one can get the difference between the expectation of the scaled row sums to be such that $bfEG(fracX_{1,n} + dots + X_{k_n,n}sigma_n) - bfEG(fracY_{1,n} + dots + Y_{k_n,n}sigma_n) = O(frack_n sigma_n^3(alpha_{k_n,n}^3 + bfE\left|X_{k_n,n}\right|^3))$. But then it's unclear how the Lindeberg condition implies that the RHS tends to 0. May I ask if there is something missing?

 · 9 October 2024

Dear professor Tao:

I apologize for the previous poorly-complied comment. To prove the Lindeberg CLT, assuming finite third moment for the entire triangular array, I got the expectation difference of G(row sum of $X_{i,n}$) scaled by $1/\sigma_n$ and G(row sum of $Y_{i,n}$) scaled by $1/\sigma_n$, to be the big-O of k_n(the standard deviation of the last element in row n cube + the third moment of the last element in row n), over σ_n cube, which I find hard to control by the given condition. Can you provide some hint on the correct way to generalize the original swapping argument?

· 26 October 2024

Dear professor Tao:

In the hint for part (ii) of Exercise 17, shouldn't the index in the denominator for the sum of variances ranges from 1 to N instead of from 1 to infinity according to the Lindeberg CLT?

[Corrected, thanks – T.]

Sam · 29 October 2024

As for part (iii) of the converse direction of Kolmogorov's three-series theorem, can you give some hint on how to show that the variance of the normalized sum is absolutely convergent?



Terence Tao · 29 October 2024

Use part (ii).

Sam · 31 October 2024

Dear professor Tao

How much of complex/Fourier analysis do you suggest to be a reasonable prerequisite to the study of the Fourier approach of CLT, including the optional proof of Berry-Essen theorem?

Sam · 21 November 2024

Dear professor Tao:

The characteristic function of the Cauchy random variable boils down to calculating an integral with integrand of the form cos(aw)/(1+w^2), is there a way to get the result of Exercise 22 without resorting to complex integration?

Sam · 22 December 2024

Dear professor Tao:

To show the absolutely integrable density function in Exercise 23 can be approximated in L1 norm by finite linear combinations of intervals, are we supposed to use the definition in note 1 together with the dominated convergence theorem?

[*This will work – T.*]

Sam · 24 December 2024

Thank you professor and Merry Christmas.

Sam · 23 December 2024

Also, isn't the PDF necessarily absolutes integrable by definition? Thank you.

[*Yes, if it exists -T.*]

Sam · 15 January 2025

Dear professor Tao:

Can you give a little more justification on the statement in Exercise 26 that any gaussian random variable is sub-gaussian ? What inequality are we using here to derive the result?



Terence Tao · 16 January 2025

There are multiple ways to obtain this estimate, for instance by using upper bounds on the error function, see https://en.wikipedia.org/wiki/Error_function, or by dyadic decomposition.

Sam · 12 February 2025

Dear professor Tao:
Can you provide some hint on why (7) can be established for sub-gaussian variables in Exercise 26?

[Fubini's theorem will suffice here. -T]

Sam · 12 February 2025

Subgaussian variables can be shown to have finite moment of all orders, but how do we use the partial expansion to get the desired result? Or is it preferable to apply some convergence theorem instead?



Aditya Guha Roy · 13 February 2025

Hi, thanks for the question because I spent a lot of time on this same question when I was first learning the topic; I am glad someone asked it and I could write it here.

1. **Using MGFs**. The key thing that you use is that the MGF of a subgaussian random variable is entire and is of order $O(e^{ct^2})$ and you can prove that for an entire function which is of order $O(e^{ct^2})$ has all derivatives at 0 finite. In particular, the claim follows.

2. **The tail way of doing it**.
Thanks to the tail-representation of expected values we can write
$\mathbf{E}[|X|^p] = \int_0^\infty px^{p-1}\mathbf{P}|X| > xdx$
Now just replace the tail probability on the right hand side of the above line of display by the bound which we get due to the subgaussian property of X and that will give you $\mathbf{E}[|X|^p] < \infty$

Now I think, it's easier to come up with the 2nd approach because the tail probabilities in the definition of subgaussian random variables would ring a bell, but I really like the MGF approach too.

3. **DCT**
You also talk about using convergence theorems. Well I think one can modify the 2nd approach to find an excuse to use DCT, but I am not sure.

Sam · 14 February 2025

Thanks for replying. The 2nd approach is how I got the finite moment of all orders for subgaussian variables. It is hinted now that Fubini's theorem is enough here, can you elaborate a bit on how they directly lead to (7)?



Aditya Guha Roy · 14 February 2025

That's because Fubini allows us to interchange expectation and sum when things converge like here (because $\mathbf{E}|X|^p < \infty$ means the sum $\sum t^n/n!\mathbf{E}[X^n]$ converge for each t

You Taylor expand the Euler-exponential function e^{itX} and then take expectation. Then Fubini grants you to interchange the sum and the expectation, with impunity.

Sam · 14 February 2025

Thank you for the explanation. One issue only: Isn't the convergence of the said series depends not only on whether $\mathbf{E}X^n < \infty$, but the growth rate of such terms as $n \to \infty$ as well?

Typical displacement of a SSRW after n steps is like √n | Aditya Guha Roy's weblog · 13 February 2025

[...] Remark 4. Actually the claim for the case when serves as a motivation for the central limit theorem (see Proposition 1 here). [...]

Sam · 15 February 2025

Dear professor Tao:

From my understanding of your earlier hint on Exercise 26, one invokes the Fubini's theorem by treating the sum as an integral over $\mathbf{N}$ with respect to the counting measure mu', is this correct? If so, how does one show that the series expansion converges simply from the condition of finite moment of all orders?

[For the non-negative version of Fubini's theorem (Tonelli's theorem), convergence is not required. -T]

Sam · 16 February 2025

Dear professor Tao:

Thank you so much for the updated hint. Tonelli theorem justifies the interchange of sum and expectation for the unsigned $|X|$ (and $|t| \geq 0$), I'm still a bit at loss as to how it helps to justify the signed case. I guess I'm not using the condition $\mathbf{E}X^k < \infty$ correctly.

[The analysis for $|X|$ from Tonelli's theorem provides enough estimates to then justify the use of Fubini's theorem for X. This combination of the Fubini and Tonelli theorems is very common in these sorts of arguments. -T]

Sam · 17 February 2025

Dear professor Tao:

In Exercise 26 one applies Fubini's theorem to the positive and negative parts of X^k, and the finite moment condition ensures that one does not run into indeterminant form as $\infty - \infty$. Is this correct?

· 26 May 2026

In the Lindeberg method, we require the function to be continuous and compactly supported. Is this ultimately related to the requirement of continuity and compact support in vague convergence and convergence in distribution, or is there another reason? If so, why do we need the class of functions in Definition 2 to satisfy these specific properties?

Terence Tao · 28 May 2026

By the Riesz representation theorem, the space $C_c(\mathbf{R})$ is the predual of the space of finite Radon measures, so it is the natural space of test functions for which to define vague convergence. But, as mentioned in the text, the Lindeberg method actually works best for smoother test functions (I use C_c^∞ in the notes, but this is overkill – three degrees of regularity, for instance, would definitely suffice). But the specific class of test functions turns out to not be of major importance here, since one can use density arguments to pass from the stronger class to the weaker one.

· 7 June 2026

Dr Tao, In the first paragraph of the first section of Lindeberg's method, "depend the choice" should be "depend on the choice."

Also, can the Berry-Esséen theorem be proved using Lindeberg's method? Or is it that you chose the Fourier-analytic approach to prove it because the theorem is not suitable for proof via Lindeberg approach?

Terence Tao · 7 June 2026

Thanks for the correction.

A direct application of the Lindeberg method gives a weaker form of the Berry-Esseen theorem, in which the decay in the error term has a worse exponent, and more regularity is required on the test function, as compared against the Fourier analytic proofs. It is possible that one could work harder and make the Lindeberg approach more competitive with the Fourier analytic approach, but I doubt that it will match (or overtake) it completely.

· 22 June 2026

I've been struggling with Exercise 9 (ii), I don't know how Stirling's formula comes into play. Could you give us a bit hint on Exercise 9 (ii)?

Terence Tao · 25 June 2026

Use part (i) and express the binomial coefficient in terms of factorials.

· 31 August 2026

Dr Tao:
In the proof of (5) of Theorem 8, why is a telescoping series introduced?

Terence Tao · 1 September 2026

This is the heart of the exchange method: to approximate a statistic involving one set $X_1, \dots, X_n$ of random variables by the corresponding statistic involving a different set $Y_1, \dots, Y_n$ of random variables, replace the variables one at a time and track how much each individual swap changes the statistic. The telescoping identity merely records the fact that the total effect of performing all the swaps is simply the sum of the influence of each individual swap. (There is a technicality that the ordering in which one performs the swaps affects the precise shape of the intermediate terms; there are some variants of the exchange method in which one uses a random ordering which can be technically advantageous in some contexts.)