

Assignment 5 Presentation

Chirag Mehta

AI20BTECH11006

Question

Quadratic forms Q2.66

Find the point at which the line $(-1 \ 1)x = 1$ is a tangent to the curve $y^2 = 4x$

Theorem

Theorem

The solution to a under-determined system of equations $Ax = b$ is given by

$$x = A^+b + (I - A^+A)w \quad (1)$$

where A^+ is the pseudoinverse of the matrix A

Proof.

Let $Ax = b$ have at least one solution

$$Ax = b \quad (2)$$

$$AA^+(Ax) = AA^+(b) \quad (3)$$

Using property of pseudoinverse

$$Ax = A^+Ab = b \quad (4)$$

Therefore, A^+b is a specific solution.

The entire set of solution is given by $A^+b + k$, where k is a vector in kernel space or null space of A

$$A(A^+b + k) = b \quad (5)$$

$$b + Ak = b \quad (6)$$

$$Ak = 0 \quad (7)$$

Any vector in the null space of A can be written as

Lemma

For an idempotent matrix the pseudoinverse is the matrix itself

Proof.

For an symmetric idempotent matrix

$$A^n = A \quad (14)$$

$$AAA = A \quad (15)$$

Therefore, $A = A^+$ satisfies all the conditions of pseudoinverse which are listed as follows

- ① $AA^+A = A$
- ② $A^+AA^+ = A^+$
- ③ AA^+ is symmetric
- ④ A^+A is symmetric



Solution

The general form of a conic is given by

$$\mathbf{x}^T \mathbf{V} \mathbf{x} + 2\mathbf{u}^T \mathbf{x} + f = 0 \quad (16)$$

for $y^2 = 4x$

$$\mathbf{V} = \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix}, \mathbf{u} = \begin{pmatrix} -2 \\ 0 \end{pmatrix}, f = 0 \quad (17)$$

Solution Contd.

For any given conic such that V is non-invertible, the point of tangency is given by

$$Vq = kn - u \quad (18)$$

$$\text{where, } k = \frac{p_1^T u}{p_1^T n}, \quad Vp_1 = 0 \quad (19)$$

Clearly

$$p_1 = \begin{pmatrix} 1 \\ 0 \end{pmatrix} \quad (20)$$

$$\implies k = \frac{-2}{-1} = 2 \quad (21)$$

Solution Contd.

Using the obtained values

$$Vq = 2n - u \quad (22)$$

$$Vq = \begin{pmatrix} -2 \\ 2 \end{pmatrix} - \begin{pmatrix} -2 \\ 0 \end{pmatrix} = 2e_2 \quad (23)$$

Using (1) and (2)

$$q = 2Ve_2 + (I - V^2)w \quad (24)$$

$$q = 2e_2 + \lambda e_1 \quad (25)$$

Solution Contd.

Using (25) and (16), we get

$$2\mathbf{q}^T \mathbf{e}_2 - 4\mathbf{e}_1^T \mathbf{q} = 0 \quad (26)$$

Evaluating this, we get

$$4 - \lambda = 0 \quad (27)$$

$$\implies \mathbf{q} = \begin{pmatrix} 1 \\ 2 \end{pmatrix} \quad (28)$$

Solution Contd.




Figure: Plot of the line and parabola