

Entry and line mode

Christopher M. Baker Howard Bondell Nathaniel Bloomfield
Elena Tartaglia Andrew P. Robinson

February 7, 2022

Abstract

In the Australian border biosecurity system, data about shipping containers is recorded in one of two modes: *entry* mode or *line* mode. The key difference between the modes is how the *directions* are recorded, that is, data about whether entries were inspected or found to be non-compliant. In general, an entry contains multiple lines of data, where each line is a single type of item or commodity. Analysis is simple when the data is recorded in line mode: the directions are recorded individually for each line. The challenge comes when data is recorded in entry mode, because the same direction is recorded against each line in the entry. In other words, if at least one line in an entry has a non-compliant inspection result, then all lines in that entry are recorded as non-compliant. Therefore, entry mode data creates a challenge when we try to estimate the probability that certain items are non-compliant, because we do not know which records of non-compliance match up with which line. We develop a statistical model to use entry mode data to help inform biosecurity risk of items. We use asymptotic analysis to estimate the value of entry mode data compared to line mode data, do a simulation study to verify that we can accurately estimate parameters in a large dataset, and we apply our methods to a real dataset.

1 Introduction

Invasive species pose a multifaceted threat to society, leading to reductions in agricultural productivity, as well as damages to the environment, human health, and the economy [Kumar Rai and Singh, 2020]. A large amount of effort goes into managing invasive species [Jardine and Sanchirico, 2018], in either eradicating them [Baker and Bode, 2020, Holmes et al., 2019, Wenger et al., 2017, Helmstedt et al., 2016], or suppressing their numbers to reduce damages [Binny et al., 2021, Brook et al., 2012, Sharov et al., 2002]. The costs associated with managing invasive species provides governments with an incentive to manage biosecurity risks at their borders to prevent new species establishing.

Given the massive scale of global trade, biosecurity regulators need to be able to allocate their resources, but to do so effectively they must understand this risks associated with various entries. At the border, one of the most effective ways of achieving this is by using the outcome of previously conducted inspections as intelligence that

informs future operations. This allows regulators to identify high risk commodities and importers, and modify their inspection targets and policies accordingly.

However, gaining insight from border inspections is a massive logistical challenge, as data must be consistently recorded in a format that makes this analysis possible. Putting infrastructure into place to collect this data, and collecting it accurately can be expensive and challenging. Often, shortcuts may be taken that render the data less valuable in analysing patterns of risk.

In Australia systems have been put in place to capture biosecurity data since the early 90's. However, to reduce the administrative burden on border staff, two ways of recording cargo inspections were implemented: line mode and entry mode. In line mode, the details of which items within an entry are inspected and found to be compliant or non-compliant are fully recorded. However, in entry mode, the results of an inspection are applied to all items within an entry. Entry mode makes it much quicker for border staff to manage entries with a large number of items, but means that when the data are analysed, entries in entry mode are censored — in these cases, it is unknown which items were inspected and found to be compliant or non-compliant. This makes analysis of the data to identify trends in biosecurity risk challenging.

Australian biosecurity data in entry mode is closely related to pooled testing, which is often used for disease surveillance. Within the pooled testing literature there are two main branches: one aims to identify positives within a pool, while the other seeks to use pooled data to estimate quantities about the population. It is the latter — estimating quantities — that we are interested in. The fundamental problem is estimating a prevalence, p , within a population, when only pooled data is available [Thompson, 1962]. More recent work has focused on improving estimates by reducing bias, either through altering the sampling strategy [Schaarschmidt, 2007, Hepworth and Watson, 2009] or by incorporating bias correction into models [Hepworth and Biggerstaff, 2017, 2021]. There have also been extensions of the problem where p is not a constant, but it is estimated using linear regression using only the pooled data [Delaigle and Hall, 2015, Chatterjee and Bandyopadhyay, 2020, McMahan et al., 2017, Liu et al., 2020]. These papers have made significant progress in fitting increasingly complex models.

In practice, however, pooled testing is manifestly different from the biosecurity scenario. Pooled testing exists by design, as a way to gather information about a population while reducing testing. In biosecurity, the searching is done to every individual line, and the results are only pooled at the point of data capture. Hence, we are interested in how much information we are losing due to aggregating results as entry mode. Or, equivalently, how much better could we be at managing bio-security risk if we stopped using entry mode when an entry contains multiple lines?

In this paper we investigate how entry mode affects our ability to identify the biosecurity risk of items. We start with an asymptotic analysis, where we calculate the precision of estimates and determine the implications of mixing different item types in entry mode. We then , which allows us to understand how larger entries and more item types affect the precision of our estimates. Finally, we analyse border interception data provided

by the Australian Department of Agriculture, Water and the Environment (DAWE) to see the real-world differences between using the line-only data and including the entry mode data.

2 Model overview

2.1 Data

To make our language about the data more precise, we will explicitly define what we mean by entries, lines and directions. Entries are a collection of lines, and a line is a group of the same type of item or commodity being imported. When cargo enters the country, each line is given *directions*. These detail all of the activities undertaken by the biosecurity regulator to manage the biosecurity risk of each line, and also the outcome of those activities.

In line mode, the directions assigned to each line are recorded along with the outcome for that line. In entry mode, directions are only recorded per entry. This means that in entry mode, if any line in that entry has an inspection, and non-compliance is found, every line in that entry is recorded to be non-compliant. If all of the lines in the entry are compliant, then they are all marked compliant, so in this case line and entry mode are equivalent.

2.2 Modelling

Throughout this paper, we focus on estimating the probability that a line is non-compliant using information including the type of item, country of origin and whether it has complete documentation. The full model for the probability that a line is non-compliant, p_{ijkl} , is:

$$\text{logit}(p_{ijkl}) = \alpha_i + \beta \mathbb{I}_j + \delta_k + \gamma_\ell, \quad (1)$$

where the fixed effects are α_i , $\beta \mathbb{I}_j$ and δ_k : α_i represents the item type, the indicator variable \mathbb{I}_j denotes whether there is correct documentation, the coefficient β is the weight given when there is correct documentation, and δ_k represents the country of provenance. The random effect γ_ℓ represents the entry effect, which we include because there may be correlations between lines within an entry. The indices can take values

$$i = 1, \dots, a, \quad a \in \mathbb{N}, \quad a = \# \text{ items} \quad (2)$$

$$j = 1, 2, \quad \text{without and with documentation} \quad (3)$$

$$k = 1, \dots, d, \quad d \in \mathbb{N}, \quad d = \# \text{ countries} \quad (4)$$

$$\ell = 1, \dots, g, \quad g \in \mathbb{N}, \quad g = \# \text{ entries.} \quad (5)$$

The values of the indicator variable are

$$\mathbb{I}_1 = 0, \quad \text{without documentation} \quad (6)$$

$$\mathbb{I}_2 = 1, \quad \text{with documentation.} \quad (7)$$

The random effect γ_ℓ has distribution

$$\gamma_\ell | \sigma \sim \text{Normal}(0, \sigma), \quad \ell = 1, \dots, g, \quad g \in \mathbb{N}. \quad (8)$$

If all data were in line mode, the above model would be a fairly standard mixed effects logistic regression with categorical variables. However, because of entry mode, we don't observe outcomes for each line, as every line in the entry is marked as non-compliant if any line in the entry is found to be non-compliant. Therefore, the outcome is whether the entry is compliant and we need to calculate the probability that the entry is non-compliant, which is 1 minus the probability that every line in the entry is compliant:

$$\mathbb{P}(\text{Entry } l \text{ non-compliant}) = q_l = 1 - \prod_{n \in \text{lines}(l)} [1 - p_n], \quad (9)$$

where p_n is the probability that the n^{th} line is non-compliant, calculated from Eq. (1). Hence, for entries in entry mode, we treat the entry as a Bernoulli random variable with probability defined by Eq. (9), while for entries in line mode, we treat each line as a Bernoulli random variable with probability as defined in Eq. (1).

This paper includes three analyses: an asymptotic analysis, a simulation study, and a case study of Australian biosecurity data. For the asymptotic analysis we only consider different item types, so we rather than using Eq. (20), we just consider the probability that a line of item type i is non-compliant, p_i . The simulation study and the case study both use the full model, as defined above.

3 Asymptotic analysis

We use asymptotic analysis to investigate how the precision of estimates depends on entry size, the number of entries, the probability of interception and whether item types are mixed. This analysis is broken into two parts. The first part assumes that all items are a single type, which allows us to quantify how the amount of data, probability of interception and entry size affects precision. The second part assumes that there are two different item types, and it explores how changing the proportion of entries with both item types mixed affects precision.

Throughout this section we make two simplifications. Firstly, we do not separate line mode and entry mode because entry mode data with entry size 1 is mathematically equivalent to line mode data. Hence, throughout these analysis, entry size 1 means line mode and entry size greater than 1 implies entry mode. Secondly, we assume that each item has a fixed probability of interception. Therefore, we consider each line a Bernoulli trial, which only depends on the item type. When the entry size is greater than one, the relevant probability is whether at least one line was intercepted.

We estimate precision via calculation the Fisher information matrix, \mathcal{I} . The Fisher information matrix is the expected value of the negative of the Hessian matrix of the log-likelihood evaluated at the value of the parameter. We use the standard error estimate

as our measurement of precision, which are the square roots of the diagonal elements of the inverse of \mathcal{I} .

3.1 Single item type

For the single item type case, we set the probability of interception to be p , and define N as the total number of entries, I as the number of entries intercepted and S as the size (i.e. the number of lines) in each entry. The likelihood is a binomial distribution, where the outcome is the an entry being intercepted. The probability that an entry is not intercepted is

$$\mathbb{P}(\text{entry not intercepted}) = (1 - p)^S, \quad (10)$$

meaning that the binomial likelihood for a set entry size, S , is

$$\mathcal{L}_S = (1 - (1 - p)^S)^I (1 - p)^S)^{N-I}. \quad (11)$$

To generalise Eq. (11) to arbitrary entry sizes, we need the product over entry size:

$$\mathcal{L} = \prod_{S \in \mathbb{N}} (1 - (1 - p)^S)^{I_{E,S}} ((1 - p)^S)^{N_{E,S} - I_{E,S}}, \quad (12)$$

where S is the entry size, $I_{E,S}$ is the number of inteceptions of entry size S and $N_{E,S}$ is the number of entries of size S . The log-likelihood is

$$\log \mathcal{L} = \sum_{S \in \mathbb{N}} I_{E,S} \log (1 - (1 - p)^S) + (N_{E,S} - I_{E,S}) \log ((1 - p)^S). \quad (13)$$

As there is only one parameter, we calculate its second derivative rather than there being a Hessian matrix:

$$\left[\frac{\partial^2 \log \mathcal{L}}{\partial p^2} \right] = \sum_{S \in \mathbb{N}} \frac{S \left(N_{E,S} + \frac{I_{E,S}((1+S)(1-p)^S - 1)}{((1-p)^S - 1)^2} \right)}{(1 - p)^2}. \quad (14)$$

To calculate the Fisher information, we need the expected value of the number of interceptions, which depends on the size of the entry:

$$\mathbb{E}[I_{E,S}] = N_{E,S}(1 - (1 - p)^S). \quad (15)$$

Hence the Fisher information is

$$\mathcal{I} = -\mathbb{E} \left[\frac{\partial^2 \log \mathcal{L}}{\partial p^2} \right] = - \sum_{S \in \mathbb{N}} \frac{N_{E,S} S^2 (1 - p)^{S-2}}{(1 - p)^S - 1}, \quad (16)$$

and the standard error estimate is

$$SE = \sqrt{\frac{1}{-\sum_{S \in \mathbb{N}} \frac{N_{E,S} S^2 (1-p)^{S-2}}{(1-p)^S - 1}}}. \quad (17)$$

Using Eq. (17) we can understand how the probability of interception, the entry size and the number of lines affect the standard error, and we plot these relationships in

Figure 1. The left plot shows that the standard error depends on the probability of interception and that the relationship depends on the entry size. For all entry sizes, the standard error is small when the probability of interception is small (below ~ 0.3). However, for larger values of the probability of interception, the standard error increase significantly if the entry size is three or greater. The large p behaviour is driven by the $(1 - p)^{S-2}$ term in Eq. (17), which means SE goes to 0 if $S = 1$, while it diverges if $S \geq 3$. Figure 1 also shows how the standard error decreases as the number of lines of data increases. The lower the entry size is, the lower the standard error, and, as $SE \sim \sqrt{1/N_{E,S}}$, entry mode data with larger entry sizes require a large amount of data to reach the same standard error.




Figure 1: The standard error (Eq. (17)) as the probability of interception, p , is varied (left) and as the number of lines are varied (right). For the left plot the number of lines is held constant at 120. For the right plot, the probability of interception is held constant at 0.5

3.2 Two item types

In this section we consider a situation where there are two items with probabilities of interception of p_1 and p_2 , and we examine how these different items type probabilities interact. We focus on a scenario where every entry is of size two, meaning there are three types of entries: only type 1; only type 2; or mixed, with one line of type 1 and one of type 2. We denote the number of lines within a single entry of type 1 and 2 as S_1 and S_2 respectively, and $I(S_1, S_2)$ and $N(S_1, S_2)$ are the number of interceptions and total number of entries with S_1 type 1 lines and S_2 type 2 lines. For our case, we can have $S_1 = 2, S_2 = 0$, $S_1 = 1, S_2 = 1$ or $S_1 = 0, S_2 = 2$. Rewriting the log-likelihood from Eq. (13), we get

$$\begin{aligned} \log \mathcal{L} = & \sum_{S_1, S_2} I(S_1, S_2) \log \left(1 - (1 - p_1)^{S_1} (1 - p_2)^{S_2} \right) + \\ & (N(S_1, S_2) - I(S_1, S_2)) \log \left((1 - p_1)^{S_1} (1 - p_2)^{S_2} \right). \end{aligned} \quad (18)$$

We compute the Fisher information matrix and the standard error using Mathematica. The standard error for p_1 is

$$\frac{1}{2} \sqrt{-\frac{p_1 (p_1^2 - 3p_1 + 2) (N_{1,2}(p_1 - 1)(p_2 - 2)p_2 + 4N_{2,2}(p_2 - 1)(p_1(p_2 - 1) - p_2))}{N_{1,1}(p_1 - 1)(N_{1,2}(p_1 - 1)(p_2 - 2)p_2 + 4N_{2,2}(p_2 - 1)(p_1(p_2 - 1) - p_2)) + N_{1,2}N_{2,2}(p_1 - 2)p_1(p_2 - 1)^2}}, \quad (19)$$

where $N_{1,1}$ and $N_{2,2}$ are the number of entries with two type 1 lines and two type 2 lines respectively, while $N_{1,2}$ are the number of entries with both type 1 and type 2 lines. The standard error for p_2 is the same, with $N_{1,1}$ and $N_{2,2}$ switched and p_1 and p_2 switched.

By examining Eq. (19) we can see that the behavior of the standard errors is more complex than when we considered only one type of item. Notably, the number of entries of only type 2, $N_{2,2}$, is in the equation for the type 1 standard error, along with the probability of interception of type 2, p_2 .

Figure 2 shows how the standard error varies with the proportion of mixed entries, for different interception probabilities. In each case, we hold p_1 at 0.1, and we vary p_2 for 0.05 up to 0.9 across the four plots. Interestingly, how the standard error changes as a function of the proportion of mixed entries changes markedly, depending on the value of p_2 . In particular, when p_2 is 0.7 and 0.9, the standard error for the p_1 increases as the proportion of mixed entries increases, while the standard error for p_2 actually decreases, while the proportion of mixed entries is below $\sim 90\%$.




Figure 2: The standard error when for entry size 2, when the proportion of mixed entries is varied. In each plot there are 50 total entries and p_1 is held constant at 0.1, while p_2 goes from 0.05 up to 0.9.

3.3 Asymptotic analysis conclusions

From the asymptotic analysis we can conclude two key lessons:

1. Entry mode data is most useful when the probability of interception, p , is small. In the extreme case where $p \rightarrow 0$, entry mode data is equivalent to line mode data because whenever an entry is *not* intercepted, we know every line was not intercepted, regardless of whether it is entry mode or line mode.
2. Entries in entry mode with mixed entries can increase or decrease the precision of estimates, depending on the true probability of non-compliance.

4 Simulation study

As the first step in developing a method to analyse real-world data sets, we simulate a data set that contains key complexities including multiple item types, fixed effects and random effects. Once we have simulated data, we can fit a model to estimate parameters. The advantage of our approach is that we can (1) verify that our model behaves correctly and (2) explore how model precision varies in a more complex setting.

4.1 Data simulation

We simulate data from the full model (as described in Section 2.2). For our simulations, chose parameters which we expect to be similar to their values in real-world data. We set $\beta = -1$, choosing an negative effect of having correct documentation, because we expect lines with correct documentation have a higher chance of being compliant. We include $a = 5$ item types with α_i taking values of -6.91, -4.60, -3.89, -2.94, -1.39 (corresponding to interception probabilities of 0.001, 0.01, 0.02, 0.05 and .2, if all other effects were 0). We chose negative values for the item effect α_i , since the probability of detection is expected to be low for any item in real-world data. We used $d = 3$ countries and set their weights to be .5, -1 and .25. For within-entry correlation, we set $\sigma = 0.25$. For each entry we draw whether it is in line mode or not with probability 0.25, and for each line we set the probability of having correct documentation to be 0.2. We show an example of the format of the data in Table 1.

Table 1: Example simulated data. Lines 1-3 are all marked as non-compliant because entry 1 is in entry mode, even though only 1 or 2 of the lines were actually non-compliant.

Line	Entry	Type	Documentation	Mode	Non-compliant
1	1	3	1	Entry	1
2	1	2	0	Entry	1
3	1	1	0	Entry	1
4	2	5	0	Line	0
5	3	4	1	Line	1
\vdots	\vdots	\vdots	\vdots	\vdots	\vdots

4.2 STAN model

We model the system in a Bayesian framework, using the RSTAN package in R. The basic structure for fitting the simulated data is to define the probability of interception for each line (following Eq. (1))

$$\text{logit}(p_{ijk\ell}) = \alpha_i + \beta \mathbb{I}_j + \delta_k + \gamma_\ell, \quad (20)$$

where

$$\gamma_\ell | \sigma \sim \text{Normal}(0, \sigma), \quad \ell = 1, \dots, g, \quad g \in \mathbb{N}. \quad (21)$$

As we fit our model in a Bayesian framework, we set priors on each parameter:

$$\alpha_i \sim \text{Normal}(-4, 4), \quad i = 1, \dots, a, \quad a \in \mathbb{N}, \quad (22)$$

$$\beta \sim \text{Normal}(0, 0.5), \quad (23)$$

$$\delta_k \sim \text{Normal}(0, 0.5), \quad k = 1, \dots, d, \quad d \in \mathbb{N} \quad (24)$$

$$\sigma \sim \text{Normal}(0, 0.5) \quad (25)$$

The probability $p_{ijk\ell}$ is the probability of non-compliance of a line that has characteristics (i, j, k, ℓ) . We denote the probability that line n is non-compliant as $p_{(n)}$, where $p_{(n)} = p_{ijk\ell}$ if that line has characteristics (i, j, k, ℓ) .

4.3 Simulation results

We fit the STAN model to the simulated data for two reasons. Firstly we confirm that, with sufficient data, the STAN model can accurately estimate model parameters. Secondly we explore how changing the amount of data and the proportion of entry mode data affects precision, over a range of entry sizes.

Figure 3

5 Case study

We use border interception data from Australia to compare the diff

References

- C. M. Baker and M. Bode. Recent advances of quantitative modeling to support invasive species eradication on islands. *Conservation Science and Practice*, n/a(n/a):e246, 2020. ISSN 2578-4854. doi: 10.1111/csp2.246. URL <https://conbio.onlinelibrary.wiley.com/doi/abs/10.1111/csp2.246>. eprint: <https://conbio.onlinelibrary.wiley.com/doi/pdf/10.1111/csp2.246>.
- R. N. Binny, J. Innes, N. Fitzgerald, R. Pech, A. James, R. Price, C. Gillies, and A. E. Byrom. Long-term biodiversity trajectories for pest-managed ecological restorations: eradication vs. suppression. *Ecological Monographs*, n/a(n/a),




Figure 3: SUPPLEMENT FIGURE

2021. ISSN 1557-7015. doi: <https://doi.org/10.1002/ecm.1439>. URL <https://esajournals.onlinelibrary.wiley.com/doi/abs/10.1002/ecm.1439>. eprint: <https://esajournals.onlinelibrary.wiley.com/doi/pdf/10.1002/ecm.1439>.

L. A. Brook, C. N. Johnson, and E. G. Ritchie. Effects of predator control on behaviour of an apex predator and indirect consequences for mesopredator suppression. *Journal of Applied Ecology*, 49(6):1278–1286, Dec. 2012. ISSN 1365-2664. doi: 10.1111/j.1365-2664.2012.02207.x. URL <http://onlinelibrary.wiley.com.ezp.lib.unimelb.edu.au/doi/10.1111/j.1365-2664.2012.02207.x/abstract>.

A. Chatterjee and T. Bandyopadhyay. Regression models for group testing: Identifiability and asymptotics. *Journal of Statistical Planning and Inference*, 204:141–152, Jan. 2020. ISSN 0378-3758. doi: 10.1016/j.jspi.2019.05.003. URL <https://www.sciencedirect.com/science/article/pii/S0378375818301423>.




Figure 4: SD as amount of data increases, depending on p and the amount of entry data

A. Delaigle and P. Hall. Nonparametric methods for group testing data, taking dilution into account. *Biometrika*, 102(4):871–887, Dec. 2015. ISSN 0006-3444. doi: 10.1093/biomet/asv049. URL <https://doi.org/10.1093/biomet/asv049>.

K. J. Helmstedt, J. D. Shaw, M. Bode, A. Terauds, K. Springer, S. A. Robinson, and H. P. Possingham. Prioritizing eradication actions on islands: it's not all or nothing. *Journal of Applied Ecology*, 53:733–741, Jan. 2016. ISSN 1365-2664. doi: 10.1111/1365-2664.12599. URL <http://onlinelibrary.wiley.com.ezp.lib.unimelb.edu.au/doi/10.1111/1365-2664.12599/abstract>.

G. Hepworth and B. J. Biggerstaff. Bias Correction in Estimating Proportions by Pooled




Figure 5: SD as amount of data increases, depending on p and the amount of entry data

Testing. *Journal of Agricultural, Biological and Environmental Statistics*, 22(4):602–614, Dec. 2017. ISSN 1537-2693. doi: 10.1007/s13253-017-0297-2. URL <https://doi.org/10.1007/s13253-017-0297-2>.

G. Hepworth and B. J. Biggerstaff. Bias Correction in Estimating Proportions by Imperfect Pooled Testing. *Journal of Agricultural, Biological and Environmental Statistics*, 26(1):90–104, Mar. 2021. ISSN 1537-2693. doi: 10.1007/s13253-020-00411-5. URL <https://doi.org/10.1007/s13253-020-00411-5>.

G. Hepworth and R. Watson. Debaised estimation of proportions in group testing. *Journal of the Royal Statistical Society: Series C (Applied Statistics)*, 58(1):105–121, 2009. ISSN 1467-9876. doi: 10.1111/j.1467-9876.2008.00639.x. URL <https://rss.onlinelibrary.wiley.com/doi/abs/10.1111/j.1467-9876.2008.00639.x>. eprint: <https://rss.onlinelibrary.wiley.com/doi/pdf/10.1111/j.1467-9876.2008.00639.x>.

N. D. Holmes, D. R. Spatz, S. Oppel, B. Tershy, D. A. Croll, B. Keitt, P. Genovesi, I. J. Burfield, D. J. Will, A. L. Bond, A. Wegmann, A. Aguirre-Muñoz, A. F. Raine, C. R. Knapp, C.-H. Hung, D. Wingate, E. Hagen, F. Méndez-Sánchez, G. Rocamora, H.-W. Yuan, J. Fric, J. Millett, J. Russell, J. Liske-Clark, E. Vidal, H. Jourdan, K. Campbell, K. Springer, K. Swinnerton, L. Gibbons-Decherong, O. Langrand, M. d. L. Brooke, M. McMinn, N. Bunbury, N. Oliveira, P. Sposimo, P. Geraldese, P. McClelland, P. Hodum, P. G. Ryan, R. Borroto-Páez, R. Pierce, R. Griffiths, R. N. Fisher, R. Wanless, S. A. Pasachnik, S. Cranwell, T. Micol, and S. H. M. Butchart. Globally important islands where eradicating invasive mammals will ben-

Table 2: Parameter estimates that are only possible when using all data, as opposed to the line-only fits

	param_name	week	summary_mean_all	summary_sd_all	summary_rhat_all
1	p.intercept[72]	1.00	-1.70	1.68	1.00
2	p.intercept[59]	2.00	-1.34	1.71	1.01
3	p.intercept[60]	2.00	-2.07	1.63	1.04
4	p.intercept[61]	2.00	-1.18	1.92	1.00
5	country_effect[62]	2.00	-0.01	0.49	1.00
6	country_effect[63]	2.00	-0.01	0.47	1.00
7	country_effect[64]	2.00	-0.01	0.49	1.01
8	country_effect[65]	2.00	0.02	0.48	1.02
9	country_effect[66]	2.00	0.01	0.48	1.00
10	country_effect[67]	2.00	-0.01	0.48	1.00
11	p.intercept[72]	3.00	-1.43	1.78	1.00

efit highly threatened vertebrates. *PLOS ONE*, 14(3):e0212128, Mar. 2019. ISSN 1932-6203. doi: 10.1371/journal.pone.0212128. URL <https://journals.plos.org/plosone/article?id=10.1371/journal.pone.0212128>.

S. L. Jardine and J. N. Sanchirico. Estimating the cost of invasive species control. *Journal of Environmental Economics and Management*, 87:242–257, Jan. 2018. ISSN 0095-0696. doi: 10.1016/j.jeem.2017.07.004. URL <http://www.sciencedirect.com/science/article/pii/S0095069616301322>.

P. Kumar Rai and J. S. Singh. Invasive alien plant species: Their impact on environment, ecosystem services and human health. *Ecological Indicators*, 111:106020, Apr. 2020. ISSN 1470-160X. doi: 10.1016/j.ecolind.2019.106020. URL <https://www.sciencedirect.com/science/article/pii/S1470160X19310167>.

Y. Liu, C. S. McMahan, J. M. Tebbs, C. M. Gallagher, and C. R. Bilder. Generalized additive regression for group testing data. *Biostatistics*, (kxaa003), Feb. 2020. ISSN 1465-4644. doi: 10.1093/biostatistics/kxaa003. URL <https://doi.org/10.1093/biostatistics/kxaa003>.

C. S. McMahan, J. M. Tebbs, T. E. Hanson, and C. R. Bilder. Bayesian regression for group testing data. *Biometrics*, 73(4):1443–1452, 2017. ISSN 1541-0420. doi: 10.1111/biom.12704. URL <https://onlinelibrary.wiley.com/doi/abs/10.1111/biom.12704>. eprint: <https://onlinelibrary.wiley.com/doi/pdf/10.1111/biom.12704>.

F. Schaarschmidt. Experimental design for one-sided confidence intervals or hypothesis tests in binomial group testing. *Communications in Biometry and Crop Science*, 2007. ISSN 1896-0782. URL http://agrobiol.sggw.waw.pl/~cbcs/pobierz.php?plik=CBCS_2_1_5.pdf. Publisher: Faculty of Agriculture and Biology, Warsaw Agricultural University, Poland.

- A. Sharov, D. Leonard, A. Liebhold, E. Roberts, and W. Dickerson. “Slow The Spread”: A National Program to Contain the Gypsy Moth. *Journal of Forestry*, 100(5):30–36, July 2002.
- K. H. Thompson. Estimation of the Proportion of Vectors in a Natural Population of Insects. *Biometrics*, 18(4):568–578, 1962. ISSN 0006-341X. doi: 10.2307/2527902. URL <https://www.jstor.org/stable/2527902>. Publisher: [Wiley, International Biometric Society].
- A. S. Wenger, V. M. Adams, G. D. Iacona, C. Lohr, R. L. Pressey, K. Morris, and I. D. Craigie. Estimating realistic costs for strategic management planning of invasive species eradications on islands. *Biological Invasions*, pages 1–19, Nov. 2017. ISSN 1387-3547, 1573-1464. doi: 10.1007/s10530-017-1627-6. URL <https://link.springer.com/article/10.1007/s10530-017-1627-6>.