

NCCN Clinical Practice Guidelines in Oncology (NCCN Guidelines®)

Bladder Cancer

Version 2.2014

NCCN.org

Continue



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014 Panel Members

Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)

***Peter E. Clark, MD** ☞ Chair
Vanderbilt-Ingram Cancer Center

Philippe E. Spiess, MD, MS ☞ Vice chair
Moffitt Cancer Center

Neeraj Agarwal, MD ‡
Huntsman Cancer Institute
at the University of Utah

Matthew C. Biagioli, MD, MS §
Moffitt Cancer Center

Mario A. Eisenberger, MD † ☞
The Sidney Kimmel Comprehensive
Cancer Center at Johns Hopkins

Richard E. Greenberg, MD ☞
Fox Chase Cancer Center

Harry W. Herr, MD ☞
Memorial Sloan-Kettering Cancer Center

Brant A. Inman, MD, MSc ☞
Duke Cancer Institute

A. Karim Kader, MD, PhD ☞
UC San Diego Moores Cancer Center

Deborah A. Kuban, MD §
The University of Texas
MD Anderson Cancer Center

Timothy M. Kuzel, MD ‡
**Robert H. Lurie Comprehensive Cancer
Center of Northwestern University**

Subodh M. Lele, MD ≠
Fred & Pamela Buffett Cancer Center at
The Nebraska Medical Center

Jeff Michalski, MD, MBA §
Siteman Cancer Center at Barnes-
Jewish Hospital and Washington
University School of Medicine

Jeffrey S. Montgomery, MD, MHSA ☞
University of Michigan
Comprehensive Cancer Center

Lance C. Pagliaro, MD †
The University of Texas
MD Anderson Cancer Center

Sumanta K. Pal, MD †
City of Hope Comprehensive Cancer Center

Anthony Patterson, MD ☞
St. Jude Children's Research Hospital/
University of Tennessee Health Science Center

Elizabeth R. Plimack, MD, MS †
Fox Chase Cancer Center

Kamal S. Pohar, MD ☞
The Ohio State University Comprehensive
Cancer Center - James Cancer Hospital
and Solove Research Institute

Michael P. Porter, MD, MS ☞
Fred Hutchinson Cancer Research Center/
Seattle Cancer Care Alliance

Jerome P. Richie, MD ☞
Dana-Farber/Brigham and Women's
Cancer Center

Wade J. Sexton, MD ☞
Moffitt Cancer Center

William U. Shipley, MD § ☞
Massachusetts General Hospital Cancer Center

Eric J. Small, MD † ☞
UCSF Helen Diller Family
Comprehensive Cancer Center

Guru Sonpavde, MD †
University of Alabama at Birmingham
Comprehensive Cancer Center

Donald L. Trump, MD †
Roswell Park Cancer Institute

Jonathan Tward, MD, PhD §
Huntsman Cancer Institute
at the University of Utah

Geoffrey Wile, MD ☐
Vanderbilt-Ingram Cancer Center

Timothy G. Wilson, MD ☞
City of Hope Comprehensive Cancer Center

NCCN
Mary Dwyer, MS
Maria Ho, PhD

☞ Urology
† Medical oncology
‡ Hematology/Hematology oncology
§ Radiotherapy/Radiation oncology
☐ Diagnostic radiology
≠ Pathology
* Writing committee member

Continue

[NCCN Guidelines Panel Disclosures](#)



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014 Table of Contents

Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)

[NCCN Bladder Cancer Panel Members](#) [Summary of the Guidelines Updates](#)

Bladder Cancer:

- [Clinical Presentation and Initial Evaluation \(BL-1\)](#)
- [Noninvasive Disease or Tis, Workup, Primary Evaluation/Surgical Treatment \(BL-1\)](#)
 - ▶ [Secondary Surgical Treatment, Adjuvant Intravesical Treatment, Follow-up \(BL-2\)](#)
 - ▶ [Posttreatment cTa, cT1, Tis Recurrent or Persistent Disease \(BL-3\)](#)
- [Muscle Invasive or Metastatic, Workup, Primary Evaluation/Surgical Treatment \(BL-1\)](#)
 - ▶ [cT2 Primary and Adjuvant Treatment \(BL-4\)](#)
 - ▶ [cT3, cT4a Primary and Adjuvant Treatment \(BL-5\)](#)
 - ▶ [cT4b and Metastatic Disease, Additional Workup, Primary and Adjuvant Treatment \(BL-6\)](#)
 - ▶ [Follow-up, Recurrent or Persistent Disease \(BL-7\)](#)
- [Principles of Surgical Management \(BL-A\)](#)
- [Principles of Pathology Management \(BL-B\)](#)
- [Approximate Probability of Recurrence and Progression \(BL-C\)](#)
- [Non-Urothelial Cell Carcinoma of the Bladder \(BL-D\)](#)
- [Follow-up After Cystectomy and Bladder Preservation \(BL-E\)](#)
- [Principles of Intravesical Treatment \(BL-F\)](#)
- [Principles of Chemotherapy Management \(BL-G\)](#)
- [Principles of Radiation Management of Invasive Disease \(BL-H\)](#)

Upper GU Tract Tumors:

- [Renal Pelvis \(UTT-1\)](#)
- [Urothelial Carcinoma of the Ureter \(UTT-2\)](#)
- [Urothelial Carcinoma of the Prostate \(UCP-1\)](#)
- [Primary Carcinoma of the Urethra \(PCU-1\)](#)

Clinical Trials: NCCN believes that the best management for any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.

To find clinical trials online at NCCN member institutions, [click here: nccn.org/clinical_trials/physician.html](#).

NCCN Categories of Evidence and Consensus: All recommendations are Category 2A unless otherwise specified.

See [NCCN Categories of Evidence and Consensus](#).

[Staging \(ST-1\)](#)

The NCCN Guidelines® are a statement of evidence and consensus of the authors regarding their views of currently accepted approaches to treatment. Any clinician seeking to apply or consult the NCCN Guidelines is expected to use independent medical judgment in the context of individual clinical circumstances to determine any patient's care or treatment. The National Comprehensive Cancer Network® (NCCN®) makes no representations or warranties of any kind regarding their content, use or application and disclaims any responsibility for their application or use in any way. The NCCN Guidelines are copyrighted by National Comprehensive Cancer Network®. All rights reserved. The NCCN Guidelines and the illustrations herein may not be reproduced in any form without the express written permission of NCCN. ©2014.



NCCN Guidelines Version 2.2014 Updates

Bladder Cancer

Summary of the changes in the 2.2014 version of the Guidelines for Bladder Cancer from the 1.2014 version include:

[MS-1](#)

- The discussion was updated to reflect the algorithm changes.

Summary of the changes in the 1.2014 version of the Guidelines for Bladder Cancer from the 1.2013 version include:

[Bladder Cancer](#)

[BL-3](#)

- Footnote “I” was modified: “If not a cystectomy candidate, consider concurrent chemotherapy and radiation, *change in intravesical agent, or a clinical trial.*”

[BL-4](#)

- After primary treatment with bladder preservation, the evaluation was changed from “Evaluate after 40-50 Gy, at completion of RT, or at 3 mo with: Cystoscopy, prior tumor site rebiopsy or TURBT, cytology, and imaging of abdomen/pelvis” to “Reassess tumor status 3 weeks after 40-45 Gy or 2-3 months after full dose (60-65 Gy).” Also for BL-5.
 - ▶ After no tumor, “Consider adjuvant chemotherapy (category 2B)” was removed. Also for BL-5.
- After primary treatment for patients with extensive comorbid disease, the evaluation was changed from “Evaluate after 40-50 Gy, at completion of RT, or at 3 mo with: Cystoscopy, prior tumor site rebiopsy or TURBT, cytology, and imaging of abdomen/pelvis” to “Reassess tumor status 2-3 months after treatment.” Also for BL-5.
 - ▶ After no tumor, “Observation” was added.
- Adjuvant therapy
 - ▶ Pathologic risk was modified: “(pT3-4 or positive nodes.)” Also for BL-5.

[BL-5](#)

- Primary treatment
 - ▶ Radical cystectomy treatment option was revised: “...and **strongly consider recommend** neoadjuvant cisplatin-based combination chemotherapy (category 1).”

[BL-7](#)

- Chest imaging was moved from the first to the second bullet: “Imaging of *chest*, upper tracts, abdomen, and pelvis for recurrence every 3-6 mo for 2 y, then as clinically indicated.”

[BL-A](#)

- Segmental (partial) cystectomy, 2nd bullet was revised: “No carcinoma in situ *as determined by random biopsies.*”

[BL-E](#)

- Follow-up after a radical cystectomy
 - ▶ 1st bullet was revised by adding “liver function tests”
 - ▶ 2nd bullet was revised by changing: “Imaging of the chest, upper tracts, abdomen, and pelvis every 3-6 ~~42~~ mo...”

[BL-G 2 of 3](#)

Principles of Chemotherapy Management

- “Radiosensitizing chemotherapy regimens” heading was revised by adding “*for bladder-preserving chemoradiation following a maximal TURBT*”
 - ▶ For first-line chemotherapy, the 4th sub-bullet was added: “Cisplatin and paclitaxel (category 2B).”
- A new section was added: “Radiosensitizing chemotherapy given concurrently with conventionally fractionated radiation for palliation of metastases or for pelvic recurrence after cystectomy”
 - ▶ Cisplatin
 - ▶ Taxane (docetaxel or paclitaxel) (category 2B)
 - ▶ 5-FU (category 2B)
 - ▶ 5-FU and mitomycin C (category 2B)
 - ▶ Capecitabine (category 3)
 - ▶ Low-dose gemcitabine (category 3)”

[Continued on next page](#)



NCCN Guidelines Version 2.2014 Updates

Bladder Cancer

Summary of the changes in the 1.2014 version of the Guidelines for Bladder Cancer from the 1.2013 version include:

[BL-H 1 of 2](#)

Principles of Radiation Management, Carcinoma of the Bladder

- Bullet 5 was revised: ~~“External beam radiation is most successful for patients without hydronephrosis or with extensive invasive tumor-associated Tis. Concurrent chemotherapy and radiation therapy or radiation therapy alone is most successful for patients without hydronephrosis and without extensive carcinoma in situ associated with their muscle-invading tumor.”~~
- Bullet 6 was revised: “For patients with stage Ta, T1, or Tis, external beam radiation therapy (EBRT) alone is rarely appropriate. For patients with recurrent Ta-T1 disease ~~without extensive Tis who are not candidates for cystectomy~~, chemoradiation may be considered *usually following BCG therapy but without extensive Tis who are not candidates for cystectomy, concurrent chemotherapy and radiation therapy may be considered as a potentially curative alternative to radical cystectomy, which is the standard treatment by NCCN Guidelines.*”
- Bullet 9 was revised: ~~“External beam radiation can also be used as potentially curative therapy for medically inoperable patients or for local palliation in patients with metastatic disease. Concurrent chemotherapy with radiation therapy or radiation therapy alone should be considered as potentially curative therapy for medically inoperable patients or for local palliation in patients with metastatic disease.”~~
- Bullet 10 was added: “When giving palliative radiation for metastatic bladder cancer or for recurrent pelvic tumor, combining radiation with radiosensitizing chemotherapy should be considered. See BL-G 2 of 3 for agents. Chemotherapy should not be used concurrently with high-dose (>3 Gy per fraction) palliative radiation.”

[BL-H 2 of 2](#)

- Principles of Radiation Management, “Carcinoma of the Urethra” was added.

[Upper GU Tract Tumors](#)

[UTT-2](#)

- Urothelial carcinoma of ureter
 - ▶ The primary treatment option for mid, low grade was modified by adding: “Excision and ureteroureterostomy/ileal ureter *in highly selected patients.*”

[Primary Carcinoma of the Urethra](#)

[PCU-1](#)

- Treatment of “Primary Carcinoma of the Urethra” was added to the NCCN Guidelines.

[Staging](#)

[ST-5](#)

- Urethra cancer staging was added.




National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014

Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)



^aImaging may include one or more of the following: intravenous pyelogram (IVP), CT urography, renal ultrasound with retrograde pyelogram, ureteroscopy, or MRI urogram.

^b[See Principles of Surgical Management \(BL-A\).](#)

^cThe modifier “c” refers to clinical staging based on bimanual examination under anesthesia and endoscopic surgery (biopsy or transurethral resection) and imaging studies. The modifier “p” refers to pathologic staging based on cystectomy and lymph node dissection.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014

Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)



^aImaging may include one or more of the following: IVP, CT urography, renal ultrasound with retrograde pyelogram, ureteroscopy, or MRI urogram.

^b[See Principles of Surgical Management \(BL-A\)](#).

^cThe modifier “c” refers to clinical staging based on bimanual examination under anesthesia and endoscopic surgery (biopsy or transurethral resection) and imaging studies. The modifier “p” refers to pathologic staging based on cystectomy and lymph node dissection.

^dMontironi R, Lopez-Beltran A. The 2004 WHO classification of bladder tumors: A summary and commentary. *Int J Surg Pathol* 2005;13:143-153.
[See Principles of Pathology Management \(BL-B\)](#).

^e[See Probability of Recurrence and Progression \(BL-C\)](#) and [Non-Urothelial Cell Carcinoma of the Bladder \(BL-D\)](#).

^f[See Follow-Up After Cystectomy and Bladder Preservation \(BL-E\)](#).

^gIndications for adjuvant therapy: Based on probability of recurrence and progression to muscle-invasive disease, such as size, number, and grade.


^h[See Principles of Intravesical Treatment \(BL-F\)](#).

ⁱImmediate intravesical chemotherapy, not immunotherapy, may decrease recurrence.

^jAlthough there is no intravesical chemotherapy standard for cTa low grade, mitomycin is most commonly used.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




If not a cystectomy candidate, consider concurrent chemotherapy and radiation, change in intravesical agent, or a clinical trial.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer


^bSee Principles of Surgical Management (BL-A).

^cThe modifier "c" refers to clinical staging based on bimanual examination under anesthesia and endoscopic surgery (biopsy or transurethral resection) and imaging studies. The modifier "p" refers to pathologic staging based on cystectomy and lymph node dissection.

^fSee Follow-Up After Cystectomy and Bladder Preservation (BL-E).

^mSee Principles of Chemotherapy Management (BL-G).

ⁿSee Principles of Radiation Management of Invasive Disease (BL-H).

^oThere are data to support equivalent survival rates, but not uniform consensus about the role of these approaches. Not all institutions have experience with these multidisciplinary treatment approaches, which require a dedicated team.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer



^bSee Principles of Surgical Management (BL-A).

^cThe modifier "c" refers to clinical staging based on bimanual examination under anesthesia and endoscopic surgery (biopsy or transurethral resection) and imaging studies. The modifier "p" refers to pathologic staging based on cystectomy and lymph node dissection.

^fSee Follow-Up After Cystectomy and Bladder Preservation (BL-E).

^mSee Principles of Chemotherapy Management (BL-G).

ⁿSee Principles of Radiation Management of Invasive Disease (BL-H).

^oThere are data to support equivalent survival rates, but not uniform consensus about the role of these approaches. Not all institutions have experience with these multidisciplinary treatment approaches, which require a dedicated team.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer


^b[See Principles of Surgical Management \(BL-A\).](#)
^cThe modifier “c” refers to clinical staging based on bimanual examination under anesthesia and endoscopic surgery (biopsy or transurethral resection) and imaging studies. The modifier “p” refers to pathologic staging based on cystectomy and lymph node dissection.

^f[See Follow-Up After Cystectomy and Bladder Preservation \(BL-E\).](#)
^m[See Principles of Chemotherapy Management \(BL-G\).](#)
ⁿ[See Principles of Radiation Management of Invasive Disease \(BL-H\).](#)
^pIf technically possible.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer


^aImaging may include one or more of the following: IVP, CT urography, renal ultrasound with retrograde pyelogram, ureteroscopy, or MRI urogram.

^b[See Principles of Surgical Management \(BL-A\)](#).

^f[See Follow-Up After Cystectomy and Bladder Preservation \(BL-E\)](#).

^h[See Principles of Intravesical Treatment \(BL-F\)](#).

^m[See Principles of Chemotherapy Management \(BL-G\)](#).

ⁿ[See Principles of Radiation Management of Invasive Disease \(BL-H\)](#).

^qDepending on risk of recurrence.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF SURGICAL MANAGEMENT

Transurethral Resection for Papillary Appearing Tumor (likely non-muscle invasive)

- Adequate resection with muscle in specimen
- Early repeat TURBT (within six weeks) if
 - Incomplete initial resection
 - No muscle in original specimen for high-grade disease
 - Large or multi-focal lesions
 - Any T1 lesion

Transurethral Resection for Suspected or Known Carcinoma In Situ

- Multiple selective and/or random biopsies
- Additional biopsy adjacent to papillary tumor
- Consider prostate urethral biopsy

Transurethral Resection for Sessile or Invasive Appearing Tumor (likely muscle invasive)

- Perform EUA
- Repeat TURBT if
 - No muscle in specimen for high-grade disease
 - Any T1 lesion
 - First resection does not allow adequate staging/attribution of risk for treatment selection
 - Incomplete resection and considering tri-modality bladder preservation therapy

Segmental (Partial) Cystectomy

- Reserved for solitary lesion in location amenable to segmental resection with adequate margins
- No carcinoma in situ as determined by random biopsies
- Bilateral pelvic lymphadenectomy should be performed and include at a minimum common, internal iliac, external iliac, and obturator nodes.

Radical Cystectomy

- Bilateral pelvic lymphadenectomy should be performed and include at a minimum common, internal iliac, external iliac, and obturator nodes.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer

PRINCIPLES OF PATHOLOGY MANAGEMENT

- Tumors in many cases that would have been classified as grade 2 by the WHO 1973 grading system are now classified as high-grade using the WHO 2004 and the ISUP/WHO 1998 systems.
- The pathology report on biopsy/TURBT specimens should specify:
 - If muscularis propria (detrusor muscle) is present and, if present, whether this structure is invaded by tumor
 - Presence or absence of lymphovascular space invasion
 - Presence or absence of subjacent carcinoma in situ

Malignancy Grading of Bladder Carcinoma: Old and New Systems*

<u>Modified Bergkvist 1987</u>	<u>WHO 1973</u>	<u>WHO/ISUP 1998 Consensus WHO, 2004</u>
Papilloma grade 0	Papilloma	Papilloma
Papilloma with atypia grade 1	TCC grade 1	Papillary urothelial neoplasm of low malignant potential
Urothelial carcinoma grade 2A	TCC grade 1	Urothelial carcinoma, low-grade
Urothelial carcinoma grade 2B	TCC grade 2	Urothelial carcinoma, low-grade or high-grade
Urothelial carcinoma grade 3	TCC grade 3	Urothelial carcinoma, high-grade

*From Droller MJ: Bladder Cancer, Current Diagnosis and Treatment. Totowa, NJ, 2001. With kind permission of Springer Science + Business Media, LLC.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer

APPROXIMATE PROBABILITY OF RECURRENCE AND PROGRESSION

<u>Pathology</u>	<u>Approximate Probability of Recurrence in 5 years</u>	<u>Approximate Probability of Progression to Muscle Invasion</u>
Ta, low grade	50%	Minimal
Ta, high grade	60%	Moderate
T1, low grade (rare)	50%	Moderate
T1, high grade	50%-70%	Moderate-High
Tis	50%-90%	High

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NON-UROTHELIAL CELL CARCINOMA OF THE BLADDER

Same management as urothelial cell carcinoma with the following issues:

Mixed Histology:

- Urothelial carcinoma plus pure squamous, adenocarcinoma, micropapillary, nested, plasmacytoid, and sarcomatoid should be identified because of the potential to have a more aggressive natural history.
- These are usually treated in a similar fashion to pure urothelial carcinoma of the bladder except their generally worse prognosis must be taken into consideration.

Pure Squamous:

- Cystectomy, RT, or other agents commonly used with squamous cell carcinoma of other sites such as 5-FU, taxanes, and methotrexate.

Adenocarcinoma:

- Radical cystectomy or segmental (partial) cystectomy.
- Conventional chemotherapy (eg, MVAC) for urothelial carcinoma is not effective; however, the use of chemotherapy or RT should be individualized and may be of potential benefit in select patients.
- Consider alternative therapy or clinical trial.

Any Small-Cell Component (or neuroendocrine features):

- Neoadjuvant chemotherapy using small-cell regimens and local treatment (cystectomy or radiotherapy).
- Primary chemotherapy regimens similar to small cell lung cancer. [See NCCN Guidelines for Small Cell Lung Cancer.](#)

Urachal Carcinoma:

- Requires complete urachal resection.
- Conventional chemotherapy for urothelial carcinoma is not effective; however, the use of chemotherapy or RT should be individualized and may be of potential benefit in select patients.
- En-bloc resection of the urachal ligament with the umbilicus.

Primary Bladder Sarcoma:

- Treatment as per [NCCN Guidelines for Soft Tissue Sarcoma.](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



FOLLOW-UP AFTER CYSTECTOMY AND BLADDER PRESERVATION

After a radical cystectomy

- Urine cytology, liver function tests, creatinine, and electrolytes every 3 to 6 mo for 2 y and then as clinically indicated
- Imaging of the chest, upper tracts, abdomen, and pelvis every 3 to 6 mo for 2 y based on risk of recurrence and then as clinically indicated
- Urethral wash cytology every 6 to 12 mo, particularly if Tis was found within the bladder or prostatic urethra
- If a continent diversion was created, monitor for vitamin B12 deficiency annually

After a segmental (partial) cystectomy or bladder preservation

- Same follow-up as above, in addition to the following:
 - Cystoscopy and urine cytology ± selected mapping biopsy every 3 to 6 mo for 2 y, then increasing intervals as appropriate

For Recurrent or Persistent Disease ([See BL-7](#))

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF INTRAVESICAL TREATMENT

Indications: Based on probability of recurrence and progression to muscle-invasive disease, such as size, number, and grade.

Immediate Intravesical Chemotherapy

- Initiated within 24 hrs after resection
- Use after TUR lowers recurrence rate in Ta low-grade tumors
- Treatment should not be given if extensive TURBT or if suspected bladder perforation

Induction Intravesical Chemotherapy

- Initiated 3-4 wks after resection
- Maximum of 2 inductions without complete response
- Maintenance therapy is optional

Induction Intravesical Immunotherapy

- Initiated 3-4 wks after resection
- Withhold if traumatic catheterization, bacteriuria, persistent gross hematuria, persistent severe local symptoms, or systemic symptoms
- Maximum of 2 inductions without complete response
- Some data suggest benefit of maintenance therapy
- Dose reduction is encouraged if there are substantial local symptoms during maintenance therapy

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Bladder Cancer

PRINCIPLES OF CHEMOTHERAPY MANAGEMENT

Perioperative chemotherapy (neoadjuvant or adjuvant)

• Regimens

- ▶ DDMVAC (dose-dense methotrexate, vinblastine, doxorubicin, and cisplatin) with growth factor support for 3 or 4 cycles^{1,2}
- ▶ Gemcitabine and cisplatin for 4 cycles^{3,4}
- ▶ CMV (cisplatin, methotrexate, and vinblastine) for 3 cycles⁵

- Randomized trials and meta-analyses show a survival benefit for cisplatin-based neoadjuvant chemotherapy in patients with muscle-invasive bladder cancer.^{1,6,7}
- Meta-analysis suggests a survival benefit to adjuvant therapy for pathologic T3, T4 or N+ disease at cystectomy.⁷
- Neoadjuvant chemotherapy is preferred over adjuvant-based chemotherapy on a higher level of evidence data.
- DDMVAC is preferred over standard MVAC based on category 1 evidence showing DDMVAC to be better tolerated and more effective than conventional MVAC in advanced disease.^{2,8} Based on these data, the traditional dose and schedule for MVAC is no longer recommended.
- Perioperative gemcitabine and cisplatin is a reasonable alternative to DDMVAC based on category 1 evidence showing equivalence to conventional MVAC in the setting of advanced disease.^{4,9}
- For gemcitabine/cisplatin, both 21- and 28-day regimens are acceptable. Better dose compliance may be achieved with fewer delays in dosing using the 21-day schedule.¹⁰
- Neoadjuvant chemotherapy may be considered for select patients with upper tract urothelial carcinoma, particularly for higher stage and/or grade tumors, as renal function will decline after nephroureterectomy and may preclude adjuvant therapy.
- Carboplatin should not be substituted for cisplatin in the perioperative setting.
 - ▶ For patients with borderline renal function or minimal dysfunction, a split-dose administration of cisplatin may be considered (such as 35 mg/m² on days 1 and 2 or days 1 and 8) (category 2B). While safer, the relative efficacy of the cisplatin-containing combination administered with such modifications remains undefined.
 - ▶ For patients who are not candidates for cisplatin, there are no data to support a recommendation for perioperative chemotherapy.

[Continued on BL-G 2 of 3](#)

[References on BL-G 3 of 3](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF CHEMOTHERAPY MANAGEMENT

First-line chemotherapy for metastatic disease

- **Regimens**
 - Gemcitabine and cisplatin⁴ (category 1)
 - DDMVAC^{2,8} with growth factor support (category 1)
- **Alternative regimens**
 - Carboplatin- or taxane-based regimens, or single-agent chemotherapy (category 2B)
- The presence of both visceral metastases and ECOG performance score ≥ 2 strongly predict poor outcome with chemotherapy. Patients without these adverse prognostic factors have the greatest benefit from chemotherapy.
- For most patients, the risks of adding paclitaxel to gemcitabine and cisplatin outweigh the limited benefit seen in the randomized trial.¹¹
- A substantial proportion of patients cannot receive cisplatin-based chemotherapy due to renal impairment or other comorbidities.
 - Participation in clinical trials of new or more tolerable therapy is recommended.
 - Carboplatin- or taxane-based regimens, or single-agent therapy can be considered for these patients (category 2B).

Second-line chemotherapy for metastatic disease

- No standard therapy exists in this setting, thus participation in clinical trials of new agents is recommended.
- Depending on first-line treatment received, single-agent taxane or gemcitabine is preferred for palliation in this setting. Additional palliative options include single-agent cisplatin, carboplatin, doxorubicin, 5-FU, ifosfamide, pemetrexed, methotrexate, and vinblastine.

Radiosensitizing chemotherapy regimens for bladder-preserving chemoradiation following a maximal TURBT

- **First-line chemotherapy**
 - Cisplatin
 - Cisplatin and 5-FU¹²
 - 5-FU and mitomycin C¹³
 - Cisplatin and paclitaxel¹⁴ (category 2B)
 - Clinical trial

Radiosensitizing chemotherapy given concurrently with conventionally fractionated radiation for palliation of metastases or for pelvic recurrence after cystectomy

- Cisplatin
- Taxane (docetaxel or paclitaxel) (category 2B)
- 5-FU (category 2B)
- 5-FU and mitomycin C (category 2B)
- Capecitabine (category 3)
- Low-dose gemcitabine (category 3)

[References on BL-G 3 of 3](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF CHEMOTHERAPY MANAGEMENT

REFERENCES

- ¹Grossman HB, Natale RB, Tangen CM, et al. Neoadjuvant chemotherapy plus cystectomy compared with cystectomy alone for locally advanced bladder cancer. *N Engl J Med* 2003;349:859-866.
- ²Sternberg CN, de Mulder PH, Schornagel JH, et al. Randomized phase III trial of high-dose-intensity methotrexate, vinblastine, doxorubicin, and cisplatin (MVAC) chemotherapy and recombinant human granulocyte colony-stimulating factor versus classic MVAC in advanced urothelial tract tumors: European Organization for Research and Treatment of Cancer Protocol no. 30924. *J Clin Oncol* 2001;19:2638-2646.
- ³Dash A, Pettus JA, Herr HW, et al. A role for neoadjuvant gemcitabine plus cisplatin in muscle-invasive urothelial carcinoma of the bladder: a retrospective experience. *Cancer* 2008;113:2471-2477.
- ⁴Von der Maase H, Hansen SW, Roberts JT, et al. Gemcitabine and cisplatin versus methotrexate, vinblastine, doxorubicin, and cisplatin in advanced or metastatic bladder cancer: results of a large, randomized, multinational, multicenter, phase III study. *J Clin Oncol* 2000;18:3068-3077.
- ⁵Griffiths G, Hall R, Sylvester R, et al. International phase III trial assessing neoadjuvant cisplatin, methotrexate, and vinblastine chemotherapy for muscle-invasive bladder cancer: long-term results of the BA06 30894 trial. *J Clin Oncol* 2011;29:2171-2177.
- ⁶Advanced Bladder Cancer Meta-analysis Collaboration. Neoadjuvant chemotherapy in invasive bladder cancer: update of a systematic review and meta-analysis of individual patient data advanced bladder cancer (ABC) meta-analysis collaboration. *Eur Urol* 2005;48:202-205; discussion 205-206.
- ⁷Advanced Bladder Cancer Meta-analysis Collaboration. Adjuvant chemotherapy in invasive bladder cancer: a systematic review and meta-analysis of individual patient data Advanced Bladder Cancer (ABC) Meta-analysis Collaboration. *Eur Urol* 2005;48:189-199; discussion 199-201.
- ⁸Sternberg CN, de Mulder P, Schornagel JH, et al. Seven year update of an EORTC phase III trial of high-dose intensity M-VAC chemotherapy and G-CSF versus classic M-VAC in advanced urothelial tract tumours. *Eur J Cancer* 2006;42:50-54.
- ⁹von der Maase H, Sengelov L, Roberts JT, et al. Long-term survival results of a randomized trial comparing gemcitabine plus cisplatin, with methotrexate, vinblastine, doxorubicin, plus cisplatin in patients with bladder cancer. *J Clin Oncol* 2005;23:4602-4608.
- ¹⁰Soto Parra H, Cavina R, Latteri F, et al. Three-week versus four-week schedule of cisplatin and gemcitabine: results of a randomized phase II study. *Ann Oncol* 2002;13:1080-1086.
- ¹¹Bellmunt J, von der Maase H, Mead GM, et al. Randomized phase III study comparing paclitaxel/cisplatin/gemcitabine and gemcitabine/cisplatin in patients with locally advanced or metastatic urothelial cancer without prior systemic therapy: EORTC Intergroup Study 30987. *J Clin Oncol* 2012;30:1107-1113.
- ¹²James ND, Hussain SA, Hall E, et al; BC2001 Investigators. Radiotherapy with or without chemotherapy in muscle-invasive bladder cancer. *N Engl J Med* 2012;366:1477-1488.
- ¹³Mitin T, Hunt D, Shipley W, et al. Transurethral surgery and twice-daily radiation plus paclitaxel-cisplatin or fluorouracil-cisplatin with selective bladder preservation and adjuvant chemotherapy for patients with muscle invasive bladder cancer (RTOG 0233): a randomized multicentre phase 2 trial. *Lancet Oncol* 2013;14:863-872.
- ¹⁴Efstathiou JA, Spiegel DY, Shipley WU, et al. Long-term outcomes of selective bladder preservation by combined-modality therapy for invasive bladder cancer: The MGH experience. *Eur Urol* 2012; 61:705-711.

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF RADIATION MANAGEMENT OF INVASIVE DISEASE

Carcinoma of the Bladder

- Precede radiation therapy alone or concurrent chemotherapy and radiation by maximal TUR of the tumor when safely possible.
- Simulate and treat patients when they have an empty bladder.
- Use multiple fields from high-energy linear accelerator beams.
- For invasive tumors, consider low-dose preoperative radiation therapy prior to segmental cystectomy (category 2B).
- Concurrent chemotherapy and radiation therapy or radiation therapy alone is most successful for patients without hydronephrosis and without extensive carcinoma in situ associated with their muscle-invading tumor.
- For patients with stage Ta, T1, or Tis, external beam radiation therapy (EBRT) alone is rarely appropriate. For patients with recurrent Ta-T1 disease usually following BCG therapy but without extensive Tis who are not candidates for cystectomy, concurrent chemotherapy and radiation therapy may be considered as a potentially curative alternative to radical cystectomy, which is the standard treatment by NCCN Guidelines.
- Treat the whole bladder with or without pelvic lymph nodes with 40 to 45 Gy and then boost the bladder tumor to a total dose up to 66 Gy excluding, if possible, normal areas of the bladder from the high-dose volume.
- Concurrent chemotherapy with radiation therapy is encouraged for added tumor cytotoxicity, and can be given without increased toxicity over radiation therapy alone. Concurrent 5-FU and mitomycin C can be used instead of cisplatin in patients with low or moderate renal function. Such therapy is optimally given by dedicated multidisciplinary teams.
- Concurrent chemotherapy with radiation therapy or radiation therapy alone should be considered as potentially curative therapy for medically inoperable patients or for local palliation in patients with metastatic disease.
- When giving palliative radiation for metastatic bladder cancer or for recurrent pelvic tumor, combining radiation with radiosensitizing chemotherapy should be considered. See [BL-G 2 of 3](#) for agents. Chemotherapy should not be used concurrently with high-dose (>3 Gy per fraction) palliative radiation.

[Continued on
next page](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



PRINCIPLES OF RADIATION MANAGEMENT OF INVASIVE DISEASE

Carcinoma of the Urethra:

- **Data support the use of radiation therapy for urothelial carcinoma and squamous cell carcinoma of the urethra (case series and experience treating these carcinomas arising from other disease sites); radiation can also be considered for adenocarcinomas of the urethra.**
- **Definitive Radiation Therapy (organ preservation)**
 - ▶ **cT2 cN0**
 - ◊ **66 to 70 Gy EBRT delivered to gross disease with a margin to encompass areas of potential microscopic spread. Concurrent chemotherapy is encouraged for added tumor cytotoxicity.**
 - ◊ **Strongly consider prophylactic radiation treatment of regional-nodal basins (inguinal and low pelvic nodes for female and distal male tumors; pelvic lymph nodes for proximal male tumors).**
 - ▶ **cT3-T4, or lymph node positive**
 - ◊ **45 to 50.4 Gy EBRT delivered to gross disease with a margin to encompass areas of microscopic spread and to regional-nodal basins (inguinal and low pelvic nodes for female and distal male tumors; pelvic lymph nodes for proximal male tumors). Boost gross primary disease to 66 to 70 Gy and gross nodal disease to 54 to 66 Gy, if feasible. Dose delivered to gross nodal disease may be limited secondary to normal tissue dose constraints. Concurrent chemotherapy should be administered for added tumor cytotoxicity.**
 - ▶ **Postoperative Adjuvant Radiation Therapy**
 - ◊ **Treatment field should encompass areas at risk for harboring residual microscopic disease based on pathologic findings at resection and may include resection bed, inguinal lymph nodes, and pelvic lymph nodes. Areas at risk for harboring residual microscopic disease should receive 45 to 50.4 Gy EBRT. Involved resection margins and areas of extranodal extension should be boosted to 54 to 60 Gy if feasible based on normal tissue constraints. Areas of gross residual disease should be boosted to 66 to 70 Gy, if feasible based on normal tissue constraints. Concurrent chemotherapy should be considered for added tumor cytotoxicity.**

Note: All recommendations are category 2A unless otherwise indicated.


Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Upper GU Tract Tumors

WORKUP



^aImaging may include one or more of the following: IVP, CT urography, retrograde pyelogram, ureteroscopy, or MRI urogram.

^bMontironi R, Lopez-Beltran A. The 2004 WHO classification of bladder tumors: A summary and commentary. Int J Surg Pathol 2005;13:143-153.

[See Principles of Pathology Management \(BL-B\).](#)

^c[See Principles of Chemotherapy Management \(BL-G\).](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Upper GU Tract Tumors

WORKUP

Urothelial
carcinoma
of ureter

- Imaging of upper tract collecting system^a
- Cytology
- Cystoscopy
- Renal function tests
- Renal scan (optional)
- Chest x-ray
- CBC, chemistry profile
- Bone scan if abnormal enzymes or bone signs and symptoms

Upper

Mid

Distal

Metastatic

Low grade^b

High grade^b

PRIMARY TREATMENT

Nephroureterectomy with cuff of bladder and regional lymphadenectomy if high grade and consider neoadjuvant chemotherapy^c in selected patients

or
Endoscopic resection

Excision and ureteroureterostomy/ileal ureter in highly selected patients

or
Endoscopic resection

Nephroureterectomy with cuff of bladder and consider regional lymphadenectomy

Nephroureterectomy with cuff of bladder and regional lymphadenectomy and consider neoadjuvant chemotherapy^c in selected patients

Distal ureterectomy and regional lymphadenectomy if high grade and reimplantation of ureter (preferred if clinically feasible) and consider neoadjuvant chemotherapy^c in selected patients

or
Endoscopic resection (low grade)

or
Nephroureterectomy with cuff of bladder and regional lymphadenectomy if high grade and consider neoadjuvant chemotherapy^c in selected patients

Chemotherapy^c

[See
Adjuvant
Treatment
and Follow-
up \(UTT-3\)](#)

^aImaging may include one or more of the following: IVP, CT urography, retrograde pyelogram, ureteroscopy, or MRI urogram.

^bMontironi R, Lopez-Beltran A. The 2004 WHO classification of bladder tumors: A summary and commentary. Int J Surg Pathol 2005;13:143-153.

[See Principles of Pathology Management \(BL-B\).](#)

^c[See Principles of Chemotherapy Management \(BL-G\).](#)

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Upper GU Tract Tumors

PATHOLOGIC STAGING^d

ADJUVANT TREATMENT

FOLLOW-UP

Adjuvant treatment for
renal pelvis
and
urothelial carcinoma
of ureter

pT0, pT1

None

- Cystoscopy every 3 mo for 1 y, then at increasing intervals
- Imaging of upper tract collecting system^a at 3- to 12-mo intervals, if endoscopic resection
± CT scan or MRI
± Chest x-ray

pT2, pT3,
pT4, pN+

Consider adjuvant
chemotherapy^{c,e}

- Cystoscopy every 3 mo for 1 y, then at increasing intervals
- Imaging of upper tract collecting system^a at 3- to 12-mo intervals
± CT scan or MRI
± Chest x-ray

^aImaging may include one or more of the following: IVP, CT urography, retrograde pyelogram, ureteroscopy, or MRI urogram.

^c[See Principles of Chemotherapy Management \(BL-G\).](#)

^dThe modifier “p” refers to pathologic staging based on cystectomy and lymph node dissection.

^eFollow recommendations for adjuvant chemotherapy after ensuring that patient is fully staged to rule out metastatic disease.


Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Urothelial Carcinoma of the Prostate


^aImaging may include one or more of the following: IVP, CT urography, retrograde pyelogram, ureteroscopy, or MRI urogram.

^bSee [Principles of Chemotherapy Management \(BL-G\)](#).

Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.




NCCN Guidelines Version 2.2014

Primary Carcinoma of the Urethra

WORKUP

DIAGNOSIS




Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014

Primary Carcinoma of the Urethra



Note: All recommendations are category 2A unless otherwise indicated.

Clinical Trials: NCCN believes that the best management of any cancer patient is in a clinical trial. Participation in clinical trials is especially encouraged.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 1
**American Joint Committee on Cancer (AJCC)
TNM Staging System for Bladder Cancer (7th ed., 2010)**
Primary Tumor (T)

TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
Ta	Noninvasive papillary carcinoma
Tis	Carcinoma in situ: "flat tumor"
T1	Tumor invades subepithelial connective tissue
T2	Tumor invades muscularis propria
pT2a	Tumor invades superficial muscularis propria (inner half)
pT2b	Tumor invades deep muscularis propria (outer half)
T3	Tumor invades perivesical tissue
pT3a	Microscopically
pT3b	Macroscopically (extravesical mass)
T4	Tumor invades any of the following: prostatic stroma, seminal vesicles, uterus, vagina, pelvic wall, abdominal wall
T4a	Tumor invades prostatic stroma, uterus, vagina
T4b	Tumor invades pelvic wall, abdominal wall

ANATOMIC STAGE/PROGNOSTIC GROUPS

Stage 0a	Ta	N0	M0
Stage 0is	Tis	N0	M0
Stage I	T1	N0	M0
Stage II	T2a	N0	M0
	T2b	N0	M0
Stage III	T3a	N0	M0
	T3b	N0	M0
	T4a	N0	M0
Stage IV	T4b	N0	M0
	Any T	N1-3	M0
	Any T	Any N	M1

Regional Lymph Nodes (N)

Regional lymph nodes include both primary and secondary drainage regions. All other nodes above the aortic bifurcation are considered distant lymph nodes.

NX	Lymph nodes cannot be assessed
N0	No lymph node metastasis
N1	Single regional lymph node metastasis in the true pelvis (hypogastric, obturator, external iliac, or presacral lymph node)
N2	Multiple regional lymph node metastasis in the true pelvis (hypogastric, obturator, external iliac, or presacral lymph node metastasis)
N3	Lymph node metastasis to the common iliac lymph nodes

Distant Metastasis (M)

M0	No distant metastasis
M1	Distant metastasis

[Continued on next page](#)

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 1 (Continued)**American Joint Committee on Cancer (AJCC)
TNM Staging System for Bladder Cancer (7th ed., 2010)****Clinical Staging**

Primary tumor assessment includes bimanual examination under anesthesia before and after endoscopic surgery (biopsy or transurethral resection) and histologic verification of the presence or absence of tumor when indicated. Bimanual examination following endoscopic surgery is an indicator of clinical stage. The finding of bladder wall thickening, a mobile mass, or a fixed mass suggests the presence of T3 and/or T4 disease, respectively. Appropriate imaging techniques for extravesical extension of the primary tumor and lymph node evaluation should be incorporated into clinical staging. When indicated, evaluation for distant metastases includes imaging of the chest, biochemical studies, and isotopic studies to detect common metastatic sites.

Pathologic Staging

Microscopic examination and confirmation of extent are required. Total cystectomy and lymph node dissection generally are required for this staging; however, a pathologic staging classification should be given for partial cystectomy specimens. Laterality does not affect the N classification.

Histologic Grade (G)

For urothelial histologies, a low- and high-grade designation is used to match the current World Health Organization/International Society of Urological Pathology (WHO/ISUP) recommended grading system:

LG Low grade
HG High grade

If a grading system is not specified, generally the following system is used:

GX Grade cannot be assessed
G1 Well differentiated
G2 Moderately differentiated
G3 Poorly differentiated
G4 Undifferentiated

Histopathologic Type

The histologic types are as follows:

Urothelial (transitional cell) carcinoma

In situ
Papillary
Flat
With squamous differentiation
With glandular differentiation
With squamous and glandular differentiation

Squamous cell carcinoma**Adenocarcinoma****Undifferentiated carcinoma**

The predominant cancer is urothelial (transitional cell) carcinoma. Histologic variants include micropapillary and nested subtypes.

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 2

American Joint Committee on Cancer (AJCC)
TNM Staging System for Renal Pelvis and Ureter Cancer (7th ed., 2010)

Primary Tumor (T)

- TX** Primary tumor cannot be assessed
- T0** No evidence of primary tumor
- Ta** Papillary noninvasive carcinoma
- Tis** Carcinoma in situ
- T1** Tumor invades subepithelial connective tissue
- T2** Tumor invades the muscularis
- T3** (For renal pelvis only) Tumor invades beyond muscularis into peripelvic fat or the renal parenchyma T3. (For ureter only) Tumor invades beyond muscularis into periureteric fat
- T4** Tumor invades adjacent organs, or through the kidney into the perinephric fat.

Regional Lymph Nodes (N)*

- NX** Regional lymph nodes cannot be assessed
- N0** No regional lymph node metastasis
- N1** Metastasis in a single lymph node, 2 cm or less in greatest dimension
- N2** Metastasis in a single lymph node, more than 2 cm but not more than 5 cm in greatest dimension; or multiple lymph nodes, none more than 5 cm in greatest dimension
- N3** Metastasis in a lymph node, more than 5 cm in greatest dimension

* Note: Laterality does not affect the N classification.

Distant Metastasis (M)

- M0** No distant metastasis
- M1** Distant metastasis

ANATOMIC STAGE/PROGNOSTIC GROUPS

Stage 0a	Ta	N0	M0
Stage 0is	Tis	N0	M0
Stage I	T1	N0	M0
Stage II	T2	N0	M0
Stage III	T3	N0	M0
Stage IV	T4	N0	M0
	Any T	N1	M0
	Any T	N2	M0
	Any T	N3	M0
	Any T	Any N	M1

[Continued on next page](#)

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 2 (Continued)**American Joint Committee on Cancer (AJCC)****TNM Staging System for Renal Pelvis and Ureter Cancer (7th ed., 2010)****Histologic Grade (G)**

For urothelial histologies, a low- and high-grade designation is used to match the current World Health Organization/International Society of Urological Pathology (WHO/ISUP) recommended grading system:

LG Low grade
HG High grade

If a grading system is not specified, generally the following system is used:

GX Grade cannot be assessed
G1 Well differentiated
G2 Moderately differentiated
G3 Poorly differentiated
G4 Undifferentiated

Histopathologic Type

The histologic types are as follows:

Urothelial (transitional cell) carcinoma

In situ

Papillary

Flat

With squamous differentiation

With glandular differentiation

With squamous and glandular differentiation

Squamous cell carcinoma**Adenocarcinoma****Undifferentiated carcinoma**

The predominant cancer is urothelial (transitional cell) carcinoma.

Histologic variants include micropapillary and nested subtypes.

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 3
**American Joint Committee on Cancer (AJCC)
TNM Staging System for Urethral Carcinoma (7th ed., 2010)**
Primary Tumor (T) (Male and Female)

TX	Primary tumor cannot be assessed
T0	No evidence of primary tumor
Ta	Noninvasive papillary, polypoid, or verrucous carcinoma
Tis	Carcinoma in situ
T1	Tumor invades subepithelial connective tissue
T2	Tumor invades any of the following: corpus spongiosum, prostate, periurethral muscle
T3	Tumor invades any of the following: corpus cavernosum, beyond prostatic capsule, anterior vagina, bladder neck
T4	Tumor invades other adjacent organs

Regional Lymph Nodes (N)

NX	Regional lymph nodes cannot be assessed
N0	No regional lymph node metastasis
N1	Metastasis in a single lymph node 2 cm or less in greatest dimension
N2	Metastasis in a single node more than 2 cm in greatest dimension, or in multiple nodes

Distant Metastasis (M)

M0	No distant metastasis
M1	Distant metastasis

ANATOMIC STAGE/PROGNOSTIC GROUPS

Stage 0a	Ta	N0	M0
Stage 0is	Tis	N0	M0
	Tis pu	N0	M0
	Tis pd	N0	M0
Stage I	T1	N0	M0
Stage II	T2	N0	M0
Stage III	T1	N1	M0
	T2	N1	M0
	T3	N0	M0
	T3	N1	M0
Stage IV	T4	N0	M0
	T4	N1	M0
	Any T	N2	M0
	Any T	Any N	M1

[Continued on next page](#)
Urothelial (Transitional Cell) Carcinoma of the Prostate

Tis pu	Carcinoma in situ, involvement of the prostatic urethra
Tis pd	Carcinoma in situ, involvement of the prostatic ducts
T1	Tumor invades urethral subepithelial connective tissue
T2	Tumor invades any of the following: prostatic stroma, corpus spongiosum, periurethral muscle
T3	Tumor invades any of the following: corpus cavernosum, beyond prostatic capsule, bladder neck (extraprostatic extension)
T4	Tumor invades other adjacent organs (invasion of the bladder)

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.



NCCN Guidelines Version 2.2014 Staging Bladder Cancer

Table 3 (Continued)**American Joint Committee on Cancer (AJCC)
TNM Staging System for Urethral Carcinoma (7th ed., 2010)****Histologic Grade (G)**

For urothelial histologies, a low- and high-grade designation is used to match the current World Health Organization/International Society of Urological Pathology (WHO/ISUP) recommended grading system:

LG Low grade

HG High grade

If a grading system is not specified, generally the following system is used:

GX Grade cannot be assessed

G1 Well differentiated

G2 Moderately differentiated

G3 Poorly differentiated

G4 Undifferentiated

Histopathologic Type

The classification applies to urothelial (transitional cell), squamous, and glandular carcinomas of the urethra and to urothelial (transitional cell) carcinomas of the prostate and prostatic urethra. There should be histologic or cytologic confirmation of the disease.

Used with the permission of the American Joint Committee on Cancer (AJCC), Chicago, Illinois. The original and primary source for this information is the AJCC Cancer Staging Manual, Seventh Edition (2010) published by Springer Science+Business Media, LLC (SBM). (For complete information and data supporting the staging tables, visit www.springer.com.) Any citation or quotation of this material must be credited to the AJCC as its primary source. The inclusion of this information herein does not authorize any reuse or further distribution without the expressed, written permission of Springer SBM, on behalf of the AJCC.

Discussion

NCCN Categories of Evidence and Consensus

Category 1: Based upon high-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2A: Based upon lower-level evidence, there is uniform NCCN consensus that the intervention is appropriate.

Category 2B: Based upon lower-level evidence, there is NCCN consensus that the intervention is appropriate.

Category 3: Based upon any level of evidence, there is major NCCN disagreement that the intervention is appropriate.

All recommendations are category 2A unless otherwise noted.

Table of Contents

Overview	MS-2
Histology	MS-2
Clinical Presentation and Workup	MS-2
Pathology and Natural History	MS-3
Staging and Grading	MS-4
Non-Muscle-Invasive Disease	MS-4
Workup and Primary Surgical Treatment	MS-4
Intravesical Therapy	MS-5
cTa, Low-Grade Tumors	MS-5
cTa, High-Grade Tumors	MS-6
cT1 Tumors	MS-6
Tis	MS-7

Posttreatment Recurrent or Persistent cTa, cT1, & Tis Disease	MS-7
Muscle-Invasive Disease	MS-8
Workup and Primary Surgical Treatment	MS-8
Radical Cystectomy	MS-8
Partial Cystectomy	MS-9
Neoadjuvant Chemotherapy	MS-9
Adjuvant Chemotherapy	MS-9
Adjuvant Radiation	MS-10
Bladder-Preserving Options	MS-10
Chemotherapy for Advanced Disease	MS-13
T2, T3, and T4a Tumors	MS-15
T4b Disease or Positive Nodes	MS-16
Follow-up after Surgery	MS-16
Recurrence or Persistent Disease after Surgery	MS-17
Metastatic Disease	MS-17

Upper Genitourinary Tract Tumors	MS-17
Renal Pelvis Tumors	MS-18
Ureteral Tumors	MS-18
Urothelial (Transitional Cell) Carcinomas of the Prostate	MS-19
Primary Carcinoma of the Urethra	MS-20
Non-Urothelial Carcinomas of the Bladder	MS-21
Summary	MS-21
Table 1. Principles of Pathology Management: Malignancy Grading of Bladder Carcinoma: Old and New Systems^{a,b}	MS-23
Table 2. Combination Chemotherapy Regimens	MS-24
References	MS-25

Overview

An estimated 74,690 new cases of urinary bladder cancer will be diagnosed in the United States (56,390 men and 18,300 women) in 2014.¹ Bladder cancer, the fourth most common cancer, is three times more common in men than in women in the United States. During the same period, approximately 15,580 deaths (11,170 men and 4410 women) will result from bladder cancer. Bladder cancers are rarely diagnosed in individuals younger than 40 years. Because the median age of diagnosis is 65 years, medical comorbidities are a frequent consideration in patient management.

The clinical spectrum of bladder cancer can be divided into 3 categories that differ in prognosis, management, and therapeutic aims. The first category consists of non-muscle-invasive tumors, for which treatment is directed at reducing recurrences and preventing progression to a more advanced stage. The second group encompasses the muscle-invasive lesions, and the goal of therapy is to determine if the bladder should be removed or preserved without compromising survival, and to determine if the primary lesion can be managed independently or if patients are at high risk for distant spread requiring systemic approaches to improve the likelihood of cure. The critical concern of therapy for the third group, consisting of metastatic lesions, is how to prolong quantity and quality of life. Numerous agents with different mechanisms of action have antitumor effects in this disease. The issue has become how to use these agents to achieve the best possible outcome.

Histology

More than 90% of urothelial tumors originate in the urinary bladder, 8% originate in the renal pelvis, and the remaining 2% originate in the ureter and urethra. Urothelial (transitional cell) carcinomas, the most common histologic subtype in the United States, may develop anywhere

transitional epithelium is present, from the renal pelvis to the ureter, bladder, and proximal two thirds of the urethra. The distal third of the urethra is dominated by squamous epithelium. The diagnosis of squamous cell tumors, which constitute 3% of the urinary tumors diagnosed in the United States, requires the presence of keratinization in the pathologic specimen.²

Of the other histologic subtypes, 1.4% are adenocarcinomas and 1% are small-cell tumors (with or without an associated paraneoplastic syndrome). Adenocarcinomas often occur in the dome of the bladder in the embryonal remnant of the urachus, in the periurethral tissues, or with a signet ring-cell histology. Urothelial tumors often have a mixture of divergent histologic subtypes, such as urothelial (transitional cell) and squamous, adenocarcinoma, and more recently appreciated nested micropapillary and sarcomatoid subtypes.³ These should be treated as urothelial carcinomas.

The systemic chemotherapy regimens used to treat urothelial carcinomas (transitional cell tumors) are generally ineffective for tumors with pure non-urothelial (non-transitional cell) histology, such as adenocarcinoma or squamous carcinoma. In some cases with a mixed histology, only the non-urothelial component remains after systemic treatment.

Clinical Presentation and Workup

The most common presenting symptom in patients with bladder cancer is microscopic or gross hematuria, although urinary frequency from irritation or a reduced bladder capacity can also develop. Less commonly, a urinary tract infection is the presenting symptom, or upper tract obstruction or pain may occur in a more advanced lesion. Patients presenting with these symptoms should be evaluated with office cystoscopy to determine if a lesion is present. If one is documented, the

patient should be scheduled for a transurethral resection of the bladder tumor (TURBT) to confirm the diagnosis and determine the extent of disease within the bladder. Urine cytology may also be obtained around the time of cystoscopy.

If the cystoscopic appearance of the tumor is solid (sessile), high-grade, or suggests invasion into muscle, a CT scan or MRI of the abdomen and pelvis is recommended before the TURBT. Because the results of a CT scan rarely alter the management of tumors with a purely papillary appearance or in cases where only the mucosa appears to be abnormal, suggesting carcinoma in situ (CIS), a CT scan, or other upper tract imaging can be deferred until after surgery. Additional workup for all patients should include urine cytology if not already tested and evaluation of the upper tracts with an intravenous pyelogram (IVP), renal ultrasound with retrograde pyelogram, CT urography, ureteroscopy, or MRI urogram. CT urography is generally the preferred approach to upper tract imaging in patients who can safely receive intravenous contrast agents.

TURBT with a bimanual examination under anesthesia (EUA) is performed to resect visible tumor and to sample muscle within the area of the tumor to assess whether invasion has occurred. When a large papillary lesion is noted, more than one session may be needed to completely resect the tumor. With CIS, biopsy of sites adjacent to the tumor and multiple random biopsies may be performed to assess for a field change. A transurethral resection (TUR) biopsy of the prostate may also be considered. Finally, if an invasive tumor is noted, an adequate sample of muscle must be obtained. A small fragment of tumor with few muscle fibers is inadequate for assessing the depth of invasion and guiding treatment recommendations.

Additional diagnostic tests, such as a bone scan, should be performed if elevated levels of alkaline phosphatase are seen in the blood. Treatment decisions are then based on disease extent within the 3 general categories: non-muscle-invasive, muscle-invasive, or metastatic. Chest imaging is indicated if invasive disease is suspected.

Positive urinary cytology may indicate urothelial tumor anywhere in the urinary tract. In the presence of a positive cytology and a normal cystoscopy, the upper tracts and the prostate in men must be evaluated and ureteroscopy may be considered.

Management of bladder cancer is based on the pathologic findings of the biopsy specimen, with attention to histology, grade, and depth of invasion. These factors are used to estimate the probability of recurrence and progression to a more advanced stage. Consideration may be given to FDA-approved urinary biomarker testing by fluorescence in situ hybridization or nuclear matrix protein 22 in the monitoring for recurrence.^{4,5}

Pathology and Natural History

Approximately 70% of newly detected cases are non-muscle-invasive disease— exophytic papillary tumors confined largely to the mucosa (Ta) (70%) or, less often, to the submucosa (T1) (25%) or flat high-grade lesions (CIS, 5%).⁶ These tumors tend to be friable and have a high propensity for bleeding. Their natural history is characterized by a tendency to recur in the same portion or another part of the bladder, and these recurrences can be either at the same stage as the initial tumor or at a more advanced stage.

Papillary tumors confined to the mucosa or submucosa are generally managed endoscopically with complete resection. Progression to a

more advanced stage may result in local symptoms or, less commonly, symptoms related to metastatic disease.

An estimated 31% to 78% of patients with a tumor confined to the mucosa or submucosa will experience a recurrence or new occurrence of urothelial (transitional cell) carcinoma within 5 years.⁷ These probabilities of recurrence vary as a function of the initial stage and grade, size, and multiplicity. Refining these estimates for individual patients is an area of active research.

Staging and Grading

The most commonly used staging system is the TNM staging system⁸ by the AJCC, as shown in the algorithm.

Tumor grade has been recognized as an important prognostic indicator with regard to the potential for disease recurrence and progression. The most widely used classification for grading of non-muscle-invasive urothelial neoplasms has been the 1973 WHO classification. This system has designations for papilloma and grades 1, 2, and 3 carcinomas. In 2004, members of the WHO and International Society of Urological Pathology (ISUP) published and recommended a revised consensus classification for papillary neoplasms.⁹ A new category of papillary urothelial neoplasm of low malignant potential was created to describe lesions with an increased number of urothelial layers when compared with papilloma but without cytologic features of malignancy. Under the WHO 2004 system, some grade 2 lesions are classified as low-grade tumors, and others are classified as high-grade. This new system potentially allows for enhanced prognostic significance but is dependent on the pathologist for making these distinctions. The 2004 WHO classification is yet to be validated by clinical trials; therefore, tumors are graded using both the 1973 and the 2004 WHO classifications. The different classification systems are compared in

Table 1: *Principles of Pathology Management*. The 7th edition of the AJCC staging system has replaced the previous 4 grade system to match the current WHO/ISUP recommended grading system.⁸

After stage and grade have been determined, treatment decisions are based on the depth of invasion and extent of disease.

Non-Muscle-Invasive Disease

Workup and Primary Surgical Treatment

A physical examination usually does not reveal non-muscle-invasive disease. Non-muscle-invasive tumors are divided into noninvasive papillomas or carcinomas (Ta), tumors invading the lamina propria (T1), and CIS or Tis. These tumors have previously been referred to as *superficial*, which is an imprecise term that should be avoided. In some cases, a papillary or T1 lesion will be documented as having an associated in situ component (Tis).

Noninvasive disease may be diagnosed by initial cystoscopy and cytology. Once suspected, imaging of upper tract collecting systems is required. In addition, a pelvic CT scan must be performed before TURBT if sessile or high-grade disease is suspected.

Standard treatment for Ta, T1, and Tis is TURBT.¹⁰ It is used to diagnose, to stage, and to treat visible tumors. TURBT with a bimanual EUA is performed to resect visible tumor and to sample muscle within the area of the tumor to assess whether invasion has occurred. The involvement of the prostatic urethra and ducts in male patients with Ta, T1, and Tis bladder tumors has been reported. The risk is higher in the case of tumors in the bladder neck. Therefore, if the lesion is sessile or if Tis or high-grade disease is suspected, selected mapping biopsies and TUR biopsy of prostate must be considered.

Clinical investigation of the specimen obtained by TUR or biopsies is an important step in the diagnosis and subsequent management of bladder cancer. The modifier “c” before the stage refers to clinical staging based on bimanual EUA and endoscopic surgery (biopsy or TUR) and imaging studies. A modifier “p” would refer to pathologic staging based on cystectomy and lymph node dissection.

A second TUR is performed when a high-grade, T1 tumor and possibly a Ta has been detected at the initial TUR. This is especially critical in cases in which no muscularis propria was included in the resection.¹¹ However, depending on the depth of invasion and grade, intravesical therapy may be recommended. This suggestion is based on the estimated probability of recurrence (ie, new tumor formation within the bladder) and progression to a more advanced, usually muscle-invasive stage, which is an event that should be considered independently. Cystectomy is rarely considered for a Ta, low-grade lesion.

Intravesical Therapy

Intravesical therapy is used in two general settings: as prophylactic or adjuvant therapy after a complete endoscopic resection or, rarely, as therapy with the goal of eradicating residual disease that could not be completely resected. This distinction is important, because most published data reflect prophylactic or adjuvant use with the goal of preventing recurrence or delaying progression to a higher grade or stage. In many cases, intravesical therapy may be overused if given to patients who have a low probability of recurrence or progression. Bacillus Calmette-Guérin (BCG) has been shown to be as effective as prophylaxis to prevent bladder cancer recurrences following TURBT. Management of the different histologic subtypes of noninvasive bladder tumors of different grades is outlined in subsequent sections.

cTa, Low-Grade Tumors

TUR is the standard treatment for cTa, low-grade tumors. Although a complete TUR by itself can eradicate cTa low-grade tumors, these tumors have a relatively high risk for recurrence. Therefore, after TUR the panel recommends that, in addition to observation, the clinicians should consider administering a single dose of immediate intravesicular chemotherapy (not immunotherapy) within 24 hours of resection. A meta-analysis of 7 randomized trials confirmed that immediate intravesical therapy decreased risk of recurrence by 11% (from 48% down to 37%) in patients having either single or multiple tumors.¹² Later studies had mixed results, with two reporting a decrease in recurrence and one finding no advantage.¹³⁻¹⁵ The immediate intravesical chemotherapy may be followed by a 6-week induction of intravesical chemotherapy. Mitomycin C is the agent most commonly used. Immunotherapy is not recommended in these patients.

The need for adjuvant therapy depends on the patient prognosis. If the patient has a low risk for recurrence, a single immediate intravesical treatment may be sufficient. Factors to consider include the size, number, T category, and grade of the tumor(s), as well as concomitant CIS and prior recurrence.⁷ Meta-analyses have confirmed the efficacy of adjuvant intravesical chemotherapy in reducing the risk of recurrence.^{16,17} Immediate intravesical treatment should be avoided in the case that TURBT was extensive or if bladder perforation is suspected.

Close follow-up of all patients is needed, although the risk for progression to a more advanced stage is low. As a result, these patients are advised to undergo a cystoscopy at 3 months initially, and then at increasing intervals.

cTa, High-Grade Tumors

Tumors staged as cTa, high-grade lesions are papillary tumors with a relatively high risk for recurrence and progression towards more invasiveness. Restaging TUR detected residual disease in 27% of Ta patients when muscle was present in the original TUR.¹⁸ In the absence of muscularis propria in the initial TUR specimen, 49% of patients with superficial disease will be understaged versus 14% if muscle was present.¹¹ Repeat resection is recommended if there is incomplete resection, or is strongly considered if there is no muscle in the specimen.

After TUR, in addition to observation, patients with Ta, high-grade tumors may be treated with intravesical BCG or mitomycin C. In the literature, there are 4 meta-analyses confirming that BCG after TUR is superior to TUR alone or TUR and chemotherapy in preventing recurrences of high-grade Ta and T1 tumors.¹⁹⁻²² The NCCN Bladder Cancer Panel Members recommend BCG as the preferred option over mitomycin for adjuvant treatment of high-grade lesions. Observation is also an option.

Follow-up is recommended, with a urinary cytology and cystoscopy at 3- to 6-month intervals for the first 2 years, and at increasing intervals as appropriate thereafter. Imaging of the upper tract should be considered every 1 to 2 years for high-grade tumors. Urine molecular tests for urothelial tumor markers are now available.²³ Most of these tests have a better sensitivity for detecting bladder cancer than urinary cytology, but specificity is lower. However, it remains unclear whether these tests offer additional information that is useful for detection and management of non-muscle-invasive bladder tumors. Therefore, NCCN Panel Members consider this to be a category 2B recommendation.

cT1 Tumors

T1 tumors are those that invade subepithelial connective tissue (also referred to as lamina propria). Based on the histologic differentiation, most cT1 lesions are high grade and considered to be potentially dangerous with a higher risk for recurrence and progression. These tumors may occur as solitary lesions or as multifocal tumors with or without an associated in situ component.

These are also treated with a complete endoscopic resection. In patients with high-risk disease, especially if the complete resection is uncertain because of the tumor size and location, no muscle is shown in the specimen, lymphovascular invasion has occurred, or inadequate staging is speculated, repeat TURBT is strongly advised.²⁴ This is supported by a trial that prospectively randomized 142 patients with pT1 tumors to a second TURBT within 2 to 6 weeks of the initial TURBT or no repeat TURBT.²⁵ All patients received adjuvant intravesical therapy. Although overall survival was similar, the 3-year recurrence-free survival was significantly higher in the repeat TURBT arm versus the control arm (69% vs. 37%, respectively), especially among patients with high-grade tumors.

Within the category of T1 disease, a particularly high-risk stratum can be identified: multifocal lesions, tumors associated with vascular invasion, or lesions that recur after BCG treatment. There are data suggesting that early cystectomy may be preferred if residual disease is found, because of the high risk for progression to a more advanced stage.²⁶ Therefore, for high-risk tumors, cystectomy rather than repeat TURBT is recommended.

If residual disease is found after a second resection, immunotherapy with BCG (category 1 recommendation) or cystectomy is recommended. If no residual disease is found after the second

resection, intravesical therapy with BCG (preferred; category 1 recommendation) or mitomycin C is recommended. Follow-up is similar to that for high-grade Ta disease.

Tis

Primary CIS or Tis is a high-grade lesion that is believed to be a precursor of invasive bladder cancer. Standard therapy for this lesion is resection followed by intravesical therapy with BCG. This therapy is generally given once a week for 6 weeks, followed by a rest period of 4 to 6 weeks, with a full re-evaluation at week 12 (ie, 3 months) after the start of therapy. If the patient is unable to tolerate BCG, intravesical mitomycin C may be administered. Follow-up is similar to that for cT1 and cTa (high-grade) tumors.

Posttreatment Recurrent or Persistent cTa, cT1, and Tis Disease

Based on Cystoscopy Results

Patients who were under observation after initial TURBT, who show a documented recurrence by positive cystoscopy results should undergo another TURBT followed by adjuvant intravesical therapy based on the stage and grade of the recurrent lesion, and then followed at 3-month intervals.

Recurrence Following Intravesical Treatment

Patients with recurrent/persistent tumors that responded to induction intravesical therapy, after initial intravesical treatment and 12-week (3-month) evaluation, can be given a second induction course of BCG or mitomycin C induction therapy. No more than two consecutive induction courses should be given. If a second course of BCG is given and residual disease is seen at the second 12-week (3-month) follow-up, TURBT is performed. For patients who have Tis or cTa disease after TURBT, intravesical therapy with a different intravesical agent is an alternative to cystectomy. Valrubicin has been approved for

CIS that is refractory to BCG, although panelists disagree on its value.²⁷ For patients with recurrence of high-grade cT1 disease after TURBT and induction BCG, cystectomy is the main option.²⁸ However, non-surgical candidates might consider concurrent chemoradiation, change of the intravesical agent, or a clinical trial.

For patients showing no residual disease at the follow-up cystoscopy, whether 1 or 2 courses of induction therapy were administered, maintenance therapy with BCG is optional. This recommendation is based on findings that an induction course of intravesical therapy followed by a maintenance regimen produced better outcomes than intravesical chemotherapy.^{19,20,29-32} Malmstrom et al³³ performed a meta-analysis including 9 trials in 2820 patients with non-muscle-invasive bladder cancer. They report that mitomycin C is superior to BCG without maintenance in preventing recurrence, but inferior to BCG in trials with maintenance. While the optimal maintenance regimen has not been established, most patients undergo maintenance for 1 to 3 years. The duration is often limited by toxicity. A recent study of 1355 patients with a median follow-up of 7.1 years found no benefit in 3 years of maintenance BCG compared to 1 year for intermediate-risk patients.³⁴ In high-risk patients, 3-year maintenance BCG reduced recurrence compared to 1-year maintenance, but did not impact progression or survival. Although a few NCCN Member Institutions do not routinely administer maintenance BCG, panelists agree that it should be an option.

Based on Cytology Results

In patients without a documented recurrence but with cytology positive, cystoscopy and imaging negative, TUR must be performed with directed or selected mapping biopsies including TUR biopsies of the prostate. In addition, cytology of the upper tract must be evaluated and ureteroscopy may be considered for detecting tumors of the upper tract.



NCCN Guidelines Version 2.2014

Bladder Cancer

If the selected mapping biopsy of the bladder is positive, then the recommendation is to administer intravesical BCG treatment followed by maintenance BCG (optional) if a complete response is seen. For tumors that fail BCG or show an incomplete response, the subsequent management options include cystectomy, changing the intravesical agent, or participation in a clinical trial. Further investigation and validation of results is warranted for establishing the efficacy of alternative agents in second-line treatments.

If TUR biopsy of the prostate is positive, the treatment is described below under *Urothelial (Transitional Cell) Carcinomas of the Prostate*. If cytology of the upper tract and/or ureteroscopy results is positive, then the treatment is described below in *Upper Genitourinary Tract Tumors*.

If the TUR biopsies of the bladder and prostate are negative, then follow-up at 3-month intervals is recommended and maintenance therapy with BCG is optional. If the cytology of the upper tract and ureteroscopy is negative, follow-up at 3-month intervals is recommended.

Muscle-Invasive Disease

Workup and Primary Surgical Treatment

Before any treatment is advised, several workup procedures are recommended to accurately determine the clinical staging. Laboratory studies, such as complete blood cell count and chemistry profile, including alkaline phosphatase, must be performed, and the patient should be assessed for the presence of regional or distant metastases. This evaluation should include a cystoscopy, chest radiograph or CT scan, bone scan in patients with symptoms or elevated alkaline phosphatase, and imaging of the upper tracts with a CT or magnetic resonance scan of the abdomen and pelvis. Imaging studies help assess the extent of local tumor invasion and the spread to lymph

nodes or distant organs.³⁵ CT and MRI may be used to assess local invasion. Unfortunately, CT scans, ultrasound, and MRI cannot accurately predict the true depth of invasion.

TURBT is the initial treatment for all muscle-invasive disease. The goal of the TUR is to correctly identify the stage; therefore, bladder muscle must be included in the resection biopsies. The overwhelming majority of muscle-invasive tumors are high-grade urothelial carcinomas.

Further treatment following initial TURBT is required for muscle-invasive tumors. Different treatment modalities are discussed below. These include radical cystectomy, partial cystectomy, neoadjuvant or adjuvant therapy, bladder-preserving approaches, and chemotherapy for advanced disease.

Radical Cystectomy

The appropriate surgical procedure involves a cystoprostatectomy in men and a cystectomy and commonly a hysterectomy in women, followed by the formation of a urinary diversion. Forms of urinary diversion include an ileal conduit or directing urine to an internal urinary reservoir, with drainage to the abdominal wall or the urethra. Relative contraindications to urethral drainage include Tis in the prostatic ducts or positive urethral margin. Orthotopic diversion or a neobladder provides bladder function similar to that of a native bladder with some increased risk for nighttime incontinence or urinary retention requiring intermittent self-catheterization.

Unfortunately, the accuracy of the staging cystoscopy and TURBT is modest, with under-staging frequently encountered. A pelvic lymph node dissection (PLND) is considered an integral part of the surgical management of bladder cancer. A more extensive PLND, which may include the common iliac or even lower para-aortic or para-caval nodes,

yields more nodes to be examined, increases yield of positive nodes, and is associated with better survival and a lower pelvic recurrence rate.³⁶⁻⁴⁰ Patient factors that may preclude a PLND include severe scarring secondary to previous treatments or surgery, advanced age, or severe comorbidities.

Partial Cystectomy

In fewer than approximately 5% of cases, an initial invasive tumor develops in an area of the bladder where an adequate margin of soft tissue and a minimum of 2 cm of noninvolved urothelium can be removed along with the tumor without compromising continence or significantly reducing bladder capacity. Partial cystectomy is most frequently recommended for lesions that develop on the dome of the bladder and have no associated Tis in other areas of the urothelium. Relative contraindications to this procedure are lesions that occur in the trigone or bladder neck. The requirement for a ureteral reimplantation, however, is not an absolute contraindication.

Similar to radical cystectomy, partial cystectomy begins with a laparotomy (intraperitoneal) and resection of the pelvic lymph nodes. If the intraoperative findings preclude a partial cystectomy, a radical cystectomy is performed. The decision to recommend adjuvant radiation or chemotherapy is based on the pathologic stage (ie, positive nodes or perivesical tissue involvement), similar to that for patients who undergo a radical cystectomy.

Neoadjuvant Chemotherapy

Increasing data support the role of neoadjuvant chemotherapy before cystectomy for T2 and T3 lesions.⁴¹⁻⁴³ Two randomized trials showed a survival benefit with neoadjuvant chemotherapy, particularly in patients with clinical T3 disease (palpable mass during EUA or unequivocal mass on CT).^{41,42} Grossman et al⁴¹ randomized 307 patients with

muscle-invasive bladder cancer to radical cystectomy alone or 3 cycles of methotrexate, vinblastine, doxorubicin, and cisplatin (MVAC) followed by radical cystectomy. Neoadjuvant chemotherapy increased median survival (77 vs. 46 months, $P = .06$) and lowered the rate of residual disease (15% vs. 38%, $P < .001$) with no apparent increase in treatment-related morbidity or mortality. Another trial randomized 196 patients with invasive bladder cancer to 2 cycles of neoadjuvant MVAC before radical cystectomy or cystectomy only.⁴⁴ Neoadjuvant chemotherapy resulted in more patients achieving pT0 than cystectomy alone (34% vs. 9%; $P < .01$). Overall survival favored the neoadjuvant group, although it did not reach statistical significance. In a meta-analysis of 11 trials involving 3,005 patients, platinum-based neoadjuvant chemotherapy was associated with improved 5-year overall and disease-free survival (5% and 9% absolute improvement, respectively).⁴⁵

An international, multicenter, randomized trial (BA06 30894) investigated the effectiveness of neoadjuvant cisplatin, methotrexate, and vinblastine (CMV) in 976 patients.⁴⁶ At a median follow-up of 8 years, patients receiving CMV before surgery had a 16% reduction in mortality risk (HR, 0.84; 95% CI, 0.72–0.99; $P = .037$).

Adjuvant Chemotherapy

Data conflict regarding the role of adjuvant systemic chemotherapy in invasive bladder cancer, because no randomized comparisons of adequate sample size have definitively shown a survival benefit of such therapy.⁴⁷ Many trials showing a survival benefit were not randomized, raising the question of selection bias in the analysis of outcomes. A meta-analysis of 6 trials found a 25% mortality reduction with adjuvant chemotherapy, but the authors pointed out several limitations of the data and concluded that evidence is insufficient for treatment

decisions.⁴⁸ Studies showed a survival advantage from therapy with cyclophosphamide, doxorubicin, and cisplatin (CAP) and with MVAC or methotrexate, vinblastine, epirubicin, and cisplatin (MVEC).⁴⁹⁻⁵¹ However, methodologic issues have raised questions as to the applicability of these studies to all patients with urothelial tumors. In the MVEC trial, patients who experienced relapse in the control arm did not undergo chemotherapy, which is not typical of more contemporary series. A randomized phase III study in 194 patients reported no difference in overall or disease-free survival between patients receiving adjuvant gemcitabine and cisplatin (GC) and those receiving chemotherapy at relapse.⁵²

Although evidence for adjuvant therapy is not as strong as for neoadjuvant therapy, current data suggest that adjuvant chemotherapy may delay recurrences, which may justify the administration of chemotherapy in those at a high risk for relapse.⁵³ A minimum of three cycles of a cisplatin-based combination, such as MVAC, or more commonly now GC, may be used in patients undergoing adjuvant therapy. Regimen and dosing recommendations are mainly based on studies in advanced disease. Carboplatin should not be substituted for cisplatin in the perioperative setting. No data support the use of adjuvant chemotherapy for non-urothelial carcinomas, regardless of stage.

Patients with tumors that are pathologic stage T2 or less and have no nodal involvement or lymphovascular invasion are considered to have lower risk and do not necessarily require adjuvant chemotherapy. Some groups suggest stratifying patients based on the p53 status of the tumor, because tumors with more than 20% of positive cells seem to have a higher risk for systemic relapse. Determining the p53 status of the tumor is still considered an experimental procedure and is not part of routine management.

Adjuvant Radiation

Data on radiation or chemoradiation following cystectomy are scarce and further prospective studies are needed to evaluate their efficacy and potential toxicity. One older randomized study of 236 patients with pT3a to pT4a bladder cancer demonstrated improvement in 5-year disease-free survival and local control compared to surgery alone.⁵⁴ A retrospective series from Milan similarly demonstrated improved cancer-specific survival with adjuvant radiotherapy for patients with pT2-T4a disease.⁵⁵ Because local recurrence rates are high for some patients after cystectomy (32% for pT3-T4 patients and 68% for patients with positive surgical margins),³⁸ adjuvant radiation therapy is reasonable to consider in these patients. Radiotherapy to 40 to 45 Gy, with or without concurrent cisplatin, could be used. The safety of higher doses, especially in the setting of a neobladder, needs to be further studied. Since pT3a to pT4a patients are also at high risk of developing metastatic disease, they are also treated with first-line multidrug chemotherapy if their renal function is adequate for cisplatin. Radiation and multidrug chemotherapy should not be given concurrently.

Bladder-Preserving Options

Within the categories of T2 and T3a urothelial carcinomas, selected patients may be considered for bladder-preserving approaches.⁵⁶ Options include aggressive endoscopic TUR alone, TUR followed by chemotherapy alone, radiotherapy alone, or a combination of chemotherapy and radiotherapy. Partial cystectomy, also a form of bladder preservation, has been discussed above. No uniform consensus has been reached about the applicability of these approaches to the management of T2 tumors.

Bladder-preserving approaches are reasonable alternatives to cystectomy for patients who are medically unfit for surgery and those

seeking an alternative to radical cystectomy. It is also endorsed by the International Consultation on Urologic Diseases-European Association of Urology evidence-based guidelines.⁵⁷ There is an apparent underutilization of aggressive bladder-preserving therapies for non-cystectomy candidates, especially the elderly and racial minorities.⁵⁸ Between 23% and 50% of patients with muscle-invasive bladder cancer who are 65 years of age and older receive no treatment or non-aggressive therapy.

The decision to use a bladder-preserving approach is partially based on the location of the lesion, depth of invasion, size of the tumor, status of the “uninvolved” urothelium, and status of the patient (eg, bladder capacity, bladder function, comorbidities). Those patients who are medically fit for radical cystectomy but with hydronephrosis are poor candidates for bladder-sparing procedures.^{59,60} Patients for whom a bladder-sparing approach is considered should undergo as complete a TUR of the tumor as possible, EUA, and metastatic workup before therapy is initiated.

With any of the alternatives to cystectomy, a concern exists over the ability to determine with certainty which bladders that appear to be endoscopically free of tumor (T0), based on a clinical assessment that includes a repeat TURBT, are in fact pathologically free of tumor (pT0). Up to a third of bladders believed to be free of disease preoperatively after chemotherapy can have residual disease at cystectomy.⁶¹ On the other hand, one series reported that all patients who achieved a complete response after radiotherapy with concurrent cisplatin and 5-FU also had no pathologic residual disease on immediate cystectomy.⁶² The frequency of residual disease after cytotoxic agents (either radiation or chemotherapy) is lower for patients who present with T2 disease than with T3 disease, which must be considered when proposing a bladder-sparing approach. When possible, bladder-sparing

options should be chosen in the context of clinical trials. After maximal TUR, close cystoscopic observation alone, chemotherapy alone, radiotherapy alone, or chemotherapy combined with radiotherapy (all also followed by close cystoscopic observation and further treatment, if necessary) are all potential treatment options. However, only chemotherapy combined with radiotherapy has been formally evaluated in prospective randomized comparisons^{63,64}; the others are still considered to be investigational.

All bladder-sparing approaches are based on the principle that not all cases require an immediate cystectomy and that the decision to remove the bladder can be deferred until the response to therapy is assessed. When chemotherapy combined with radiotherapy is used, a cystoscopy with bladder biopsy is commonly performed midway through treatment (after the induction phase of treatment). If disease is seen, immediate cystectomy is recommended. For all of the other methods, repeat biopsy or TUR is performed 2 to 3 months after full-dose cytotoxic therapy (either chemotherapy alone or radiation alone). If persistent disease is observed, a prompt salvage cystectomy is recommended when possible.

TUR Alone

TUR alone may be curative in selected cases in which the lesion is solitary, is less than 2 cm in size, and has minimally invaded the muscle. These cases should also have no associated in situ component, palpable mass, or associated hydronephrosis.⁶⁵

If considered for TURBT alone, patients should undergo an aggressive re-resection of the site within 4 weeks of the primary procedure to ensure that no residual disease is present. If the repeat TURBT is negative for residual tumor, the patient can be managed conservatively with repeat endoscopic evaluations and cytologies every 3 months until



NCCN Guidelines Version 2.2014

Bladder Cancer

a relapse is documented. At that point, management would depend on the stage of the lesion documented at relapse.

Chemotherapy Following TUR

The use of chemotherapy alone is not considered to be adequate without additional treatment to the bladder and remains investigational. This view is based on studies showing that the proportions of complete pathologic response in the bladder using neoadjuvant chemotherapy alone were only up to 38%.⁴¹ A higher proportion of bladders can be rendered tumor-free and therefore preserved when chemotherapy is combined with concurrent radiotherapy.

Chemotherapy Followed by Partial Cystectomy

Less than 5% of invasive tumors present initially in a location and pattern that is amenable to curative resection with partial cystectomy.⁶⁶ Non-randomized studies reported 5- to 10-year overall survival of 69% to 73%; however, the rate of invasive recurrence was 23% to 33%.^{61,67} This approach is currently not widely used, but it has the advantages of surgically removing the diseased portion of the bladder and allowing for definitive lymph node staging.

Radiotherapy Following TUR

Radiotherapy alone is inferior to radiotherapy combined with chemotherapy for patients with an invasive bladder tumor, and is not considered standard for patients who can tolerate combined therapy.^{63,64} In a randomized trial of 360 patients, radiotherapy with concurrent mitomycin C and 5-FU improved 2-year locoregional disease-free survival from 54% (radiotherapy alone) to 67% ($P = .01$), and 5-year overall survival from 35% to 48% ($P = .16$), without increasing grade 3-4 acute or late toxicity.⁶⁴ Hence, radiotherapy alone is only indicated for those who cannot tolerate a cystectomy or chemotherapy because of medical comorbidities.

Radiotherapy with Concurrent Chemotherapy following TUR

Several groups have investigated the combination of concurrent or sequential chemotherapy and radiotherapy after TURBT. First, an endoscopic resection that is as complete as possible is performed. Incomplete resection is an unfavorable prognostic factor for the ability to preserve the bladder and for survival.^{68,69}

Radiation Therapy Oncology Group protocol 89-03 compared concurrent cisplatin and radiotherapy with versus without 2 cycles of induction MCV (methotrexate, cisplatin, and vinblastine) chemotherapy.⁶⁰ No difference in complete clinical response or 5-year overall survival was observed between the treatment arms. Other studies also reported no significant survival benefit for neoadjuvant chemotherapy before bladder-preserving chemotherapy with radiation therapy.^{69,70}

Radiotherapy with concurrent cisplatin-based chemotherapy as radiosensitizer is the most common and well-studied chemoradiation method used to treat muscle-invasive bladder cancer.^{56,59,60,62-64,68,69} After a complete TURBT, 40 Gy of external beam radiotherapy is administered, typically with a 4-field technique. Two doses of concurrent cisplatin are given on weeks 1 and 4. After this induction phase, an endoscopic re-evaluation is performed. If residual disease is noted, a cystectomy is advised. If no disease is visible and the cytology and biopsy are negative (T0), an additional 25 Gy of consolidation external-beam radiotherapy is administered along with one additional dose of cisplatin. The patient is then followed up with serial urine cytologies and cystoscopies as previously outlined.

Results from several prospective trials have demonstrated the effectiveness of this approach. In RTOG 89-03 in which 123 patients with clinical stage T2-T4a were treated with radiotherapy with

concurrent cisplatin, with or without induction MCV chemotherapy, 5-year overall survival was approximately 49% in both arms.⁶⁰ RTOG 95-06 treated 34 patients with twice-daily irradiation and concurrent cisplatin and 5-FU. Three-year overall survival was 83%.⁷¹ RTOG 97-06 treated 47 patients with twice-daily irradiation and concurrent cisplatin; patients also received adjuvant chemotherapy with methotrexate, vinblastine, and cisplatin.⁷² Three-year overall survival was 61%. RTOG 99-06 treated 80 patients using twice-daily irradiation plus cisplatin and paclitaxel, followed by adjuvant cisplatin and gemcitabine. Five-year overall survival was 56%.⁷³ In these trials, the complete response rate achieved ranged from 59% to 81%. An alternative approach involves twice-daily radiation with concurrent paclitaxel plus cisplatin or 5-FU plus cisplatin.⁷⁴

Currently, the following radiosensitizing regimens are reasonable for bladder-preserving chemoradiation following a maximal TURBT: cisplatin, cisplatin and 5-FU, 5-FU and mitomycin C, and cisplatin and paclitaxel (category 2B). Enrollment in a clinical trial is appropriate if available.

Up to about 80% of long-term survivors maintain an intact bladder, while other patients ultimately require radical cystectomy.^{59,60,68-73} A combined analysis of survivors from these 4 trials, with median follow-up of 5.4 years, showed that combined-modality therapy was associated with low rates of late grade 3 toxicity (5.7% genitourinary and 1.9% gastrointestinal).⁷⁵ No late grade 4 toxicities or treatment-related deaths were recorded.

Chemotherapy for Advanced Disease

The specific chemotherapy regimen recommended partially depends on the presence or absence of medical comorbidities, such as cardiac disease and renal dysfunction, along with the risk classification of the

patient based on disease extent. In general, long-term survival with combination chemotherapy alone has been reported only in good-risk patients, defined as those with good performance status, no visceral (ie, liver, lung) or bone disease, and normal alkaline phosphatase or lactic dehydrogenase levels. Poor-risk patients, defined as those with poor performance status or visceral disease, have consistently shown very poor tolerance to multiagent combination programs and few complete remissions, which are prerequisites for cure.

Currently, 3 drug types are active in the management of advanced bladder cancer: cisplatin, the taxanes, and gemcitabine. Combinations of 2 or 3 of these agents have shown clinical benefit (Table 2). Commonly used combinations include GC^{76,77} and dose-dense MVAC.^{78,79} A large, international, phase III study randomized 405 patients with locally advanced or metastatic disease to GC or standard MVAC.⁸⁰ At a median follow-up of 19 months, overall survival and time-to-progression were similar in the two arms. However, less toxic deaths were recorded among patients receiving GC compared to MVAC (1% vs. 3%), although this did not reach statistical significance. A 5-year update analysis confirmed that GC was not inferior to MVAC in terms of survival (overall survival, 13.0% vs. 15.3%; progression-free survival, 9.8% vs. 11.3%, respectively).⁷⁷ Another large, randomized, phase III trial compared dose-dense MVAC to standard MVAC.^{78,79} At a median follow-up of 7.3 years, 24.6 % of patients were alive in the dose-dense MVAC cohort compared with 13.2% in the standard MVAC cohort. There was one toxic death in each arm, but less overall toxicity was seen in the dose-dense group. From these data, standard MVAC is inferior to high-dose-intensity MVAC in terms of toxicity and efficacy, and is inferior to GC in terms of toxicity (and therefore no longer used). Both GC and dose-dense MVAC are category 1 recommendations for metastatic disease.

The performance status of the patient is a major determinant of which regimen is used, and regimens with lower toxicity profiles are recommended in patients with compromised liver or renal status or serious comorbid conditions. In patients with glomerular filtration rate (GFR) <60 mL/min, carboplatin may be substituted for cisplatin in the above mentioned regimen. A phase II/III study assessed 2 carboplatin-containing regimens in medically unfit patients (performance status 2).⁸¹ The overall response rate was 42% for gemcitabine plus carboplatin and 30% for methotrexate, carboplatin, and vinblastine. However, the response rates dropped to 26% and 20%, respectively, with increased toxicity among patients who were both unfit and had renal impairment (GFR <60 mL/min).

More recently, the taxanes have been shown to be active as both front-line and palliative therapies. Based on these results, several groups are exploring 2- and 3-drug combinations using these agents, with and without cisplatin, as initial therapy. The alternative regimens, including cisplatin/paclitaxel,⁸² gemcitabine/paclitaxel,⁸³ cisplatin/gemcitabine/paclitaxel,⁸⁴ carboplatin/gemcitabine/paclitaxel,⁸⁵ and cisplatin/gemcitabine/docetaxel,⁸⁶ have shown modest activity in bladder cancer in phase I-II trials. A randomized phase III trial was conducted to compare GC and GC plus paclitaxel in 626 patients with locally advanced or metastatic urothelial cancer.⁸⁷ The addition of paclitaxel to GC resulted in higher response rates and a borderline overall survival advantage, which was not statistically significant in the intent-to-treat analysis. Analysis of eligible patients only (92%) resulted in a small (3.2 months) statistically significant survival advantage in favor of the 3-drug regimen ($P = .03$). There was no difference in progression-free survival. The incidence of neutropenic fever was substantially higher with the 3-drug combination (13.2 vs. 4.3%; $P <$

.001). Panelists feel that the risk of adding paclitaxel outweighs the limited benefit reported from the trial.

Although current data are insufficient to recommend the above alternative regimens as routine first-line options, non-cisplatin-containing regimens may be considered in patients who cannot tolerate cisplatin because of renal impairment or other comorbidities (category 2B). The NCCN Panel recommends enrollment in clinical trials of potentially less toxic therapies.

The regimens effective for urothelial carcinoma histologies have limited efficacy for patients with non-urothelial carcinomas. These individuals are often treated based on the identified histology. For example, adenocarcinomas are managed surgically with radical or segmental cystectomy and individualizing the adjuvant chemotherapy and radiotherapy for maximum benefit. Pure squamous cell tumors are treated by cystectomy, radiation therapy, or agents commonly used for squamous cell carcinoma of other sites such as 5-FU or taxanes. However, overall experience with chemotherapy in non-urothelial carcinomas is limited.

Independent of the specific regimen used, patients with metastatic disease are re-evaluated after 2 to 3 cycles of chemotherapy, and treatment is continued for 2 more cycles in patients whose disease responds or remains stable. Surgery or radiotherapy may be considered in patients who show a major partial response in an unresectable primary tumor or who have a solitary site of residual disease that is resectable after chemotherapy. In selected series, this approach has been shown to afford a survival benefit. If disease is completely resected, 2 additional cycles of chemotherapy can be considered, depending on patient tolerance. Patients for whom surgery or radiotherapy are not considered options are generally treated with

chemotherapy for a maximum of 6 cycles, depending on their response. If no response is noted after 2 cycles or if significant morbidities are encountered, a change in therapy is advised, taking into account the patient's current performance status, extent of disease, and specific prior therapy administered. The same applies to patients who experience systemic relapse after adjuvant chemotherapy.

Second-line chemotherapy data are highly variable and unclear in this setting; therefore, no standard therapy exists. The NCCN Bladder Cancer Panel Members highly recommend enrollment in a clinical trial. The available options for palliative chemotherapy depend on what was offered as first line. Docetaxel, paclitaxel, or gemcitabine monotherapy is preferred.⁸⁸⁻⁹¹ Other options include: cisplatin, carboplatin, doxorubicin, 5-FU, ifosfamide, pemetrexed, methotrexate, and vinblastine—with modest benefit limited to small phase II trials.⁹²⁻⁹⁶

Chemoradiation for Advanced Disease

Chemotherapy is sometimes combined with palliative radiation to treat metastases or pelvic recurrence after cystectomy. However, concurrent chemotherapy is inappropriate if high-dose radiation (>3 Gy fractions) is used. The radiosensitizing chemotherapy regimen remains controversial in this setting. Possible options include cisplatin, docetaxel, or paclitaxel (category 2B); 5-FU (category 2B); 5-FU and mitomycin C (category 2B); capecitabine (category 3); and low-dose gemcitabine (category 3).

T2, T3, and T4a Tumors

The critical issues in the management and prognosis of these patients are whether a palpable mass is appreciated at EUA and if the tumor has extended through the bladder wall. Tumors that are organ-confined (T2) have a better prognosis than those that have extended through the bladder wall to the perivesical fat (T3) and beyond. T4a tumors involve

the prostatic stroma, uterus, or vagina and are typically surgically managed similar to T3 tumors.

Primary surgical treatment for clinical T2, T3, and T4a lesions with no nodal disease seen on abdominal/pelvic CT or MRI scan is a radical cystectomy and pelvic lymphadenectomy. Based on results from two randomized trials (category 1),^{41,42} NCCN panelists agree that there is stronger evidence to support neoadjuvant chemotherapy for T3 disease than for T2 disease. Therefore, neoadjuvant chemotherapy is recommended for patients with cT3 tumors (category 1) and should be strongly considered for those with cT2 tumors (category 1). If no neoadjuvant chemotherapy was given, postoperative adjuvant chemotherapy is considered based on pathologic risk, such as positive nodes or pathologic T3-T4 lesions (category 2B recommendation).

Partial cystectomy can be considered only in T2 patients with a single tumor in a suitable location and no presence of Tis along with consideration of neoadjuvant chemotherapy. Partial cystectomy is not an option for T3 or T4a patients. Adjuvant radiotherapy or chemotherapy based on pathologic risk, such as positive nodes, positive margin, high-grade lesions, and pathologic T3-T4 lesions, may be considered (category 2B recommendation). As in the case for radical cystectomy, patients who have received neoadjuvant chemotherapy should not receive adjuvant chemotherapy following partial cystectomy.

Bladder preservation strategy with concurrent chemotherapy and radiation (category 2B recommendation) is an option in highly selected patients. Candidates for bladder-sparing approaches are those patients whose tumors present without hydronephrosis and those tumors that allow a visibly complete or a maximally debulking TURBT. The overall tumor status should be reassessed 3 weeks after radiation if 40 to 45

Gy was initially administered or 2 to 3 months after if the full dose of 60 to 65 Gy was delivered. If no residual tumor is detected, appropriate options include observation or completion of radiation up to 66 Gy. If tumor is present, cystectomy is the preferred option.

In patients with extensive comorbid disease or poor performance status, treatment options include TURBT alone, concurrent chemoradiation, or chemotherapy alone. Based on high-level evidence, only cisplatin alone or 5-FU and mitomycin C together are the drugs proven to be radiosensitizing with radiation and are superior to radiation alone.^{63,64} The overall tumor status should be reassessed 2 to 3 months after treatment. If no tumor is evident, the patient should be observed. If tumor remains, cystectomy is the preferred choice if feasible. Patients who are not surgical candidates should consider completion of radiation with alternative radiosensitizing chemotherapy and/or alternative chemotherapy.

T4b Disease or Positive Nodes

For patients who show no nodal disease on abdominal/pelvic CT or MRI scans or biopsy, the primary treatment recommendation includes 2 to 3 courses of chemotherapy with or without radiotherapy followed by evaluation with TURBT, cystoscopy, and CT scan of the abdomen and pelvis. In highly selected T4a node-negative patients, cystectomy with or without chemotherapy is another primary treatment option. If no evidence of tumor is present after primary chemotherapy, one may consider the consolidation chemotherapy regimen with or without radiation. Alternatively, cystectomy may also be considered as the subsequent management option for these patients. However, upon evaluation after primary therapy, if residual disease is noted, chemotherapy with or without radiation can be used. A change in

chemotherapy regimen is reasonable. Cystectomy, if feasible, is again an option for patients who respond and those who do not respond to primary therapy.

For patients with positive nodes, documented on imaging, a biopsy is considered if possible to confirm nodal spread. Patients with positive nodes should receive chemotherapy with or without radiation and should be evaluated with cystoscopy, TURBT, and abdomen/pelvis imaging. If no residual tumor is detected, patients may receive a radiation boost or a cystectomy. If cancer is still present following primary therapy, patients should follow the pathway for metastatic disease.

Follow-up after Surgery

Results from a meta-analysis of 13,185 patients who have undergone cystectomy reported 0.75% to 6.4% prevalence of upper tract recurrence in these patients.⁹⁷ Surveillance by urine cytology detected 7% and upper urinary tract imaging detected 30% of these recurrences.

Follow-up after a cystectomy should include urine cytology, liver function tests, creatinine, and electrolytes every 6 to 12 months for 2 years and then as clinically indicated. Imaging of the chest, upper tracts, abdomen, and pelvis should be conducted every 3 to 6 months for 2 years based on the risk of recurrence and then as clinically indicated. Patients should be monitored annually for vitamin B₁₂ deficiency if a continent diversion was created. Urethral wash cytology every 6 to 12 months is advised, particularly if Tis was found within the bladder or prostatic urethra.

Follow-up after a partial cystectomy is similar to that for a radical cystectomy, with the addition of monitoring for relapse in the bladder by serial cytologic examinations and cystoscopies (may include selected



NCCN Guidelines Version 2.2014

Bladder Cancer

mapping biopsy) at 3- to 6-month intervals for the first 2 years, then at increasing intervals according to clinician discretion.

For patients who have undergone bladder preservation, attention to the bladder as a site of recurrence is only one part of the overall management, because these individuals remain at risk for recurrence elsewhere in the urothelial tract and distantly. Imaging studies and laboratory testing should be performed as outlined under post-cystectomy follow-up. Additionally, continued monitoring of the urothelium with cystoscopy and urinary cytologies with or without mapping biopsy is a routine part of the management of all cases in which the bladder is preserved. Follow-up intervals are typically every 3 to 6 months for the first 2 years, and then at increasing intervals as appropriate.

Recurrence or Persistent Disease after Surgery

Metastatic disease or local recurrence following cystectomy may be managed with palliative chemotherapy, radiation, or a combination of the two.

A positive cytology with no evidence of disease in the bladder should prompt selective washings of the upper tracts and a biopsy of the prostatic urethra. If the results are positive, patients are managed as described in the sections below.

For patients who have their bladders preserved, a local recurrence or persistent disease should be evaluated as a new cancer. Recurrences are treated based on the extent of disease at relapse, with consideration of prior treatment. Tis, Ta, or T1 tumors are generally managed with intravesical BCG therapy or cystectomy. If no response is noted following BCG treatment, a cystectomy is advised. Invasive disease is generally managed with radical cystectomy, and a second

attempt at bladder preservation is not advisable. Cystectomy may not be possible in a patient who has undergone a full course of external-beam radiotherapy and has bulky residual disease. For these patients, palliative chemotherapy is advised, generally with a regimen that is not cross-resistant to the one previously received. If the patient has not undergone radiotherapy, a course of radiotherapy is an alternative. Palliative TURBT is also an option.

Metastatic Disease

About half of all patients relapse after cystectomy depending on the pathologic stage of the tumor and nodal status. Local recurrences account for about 10% to 30% of relapses, whereas distant metastases are more common.

If metastasis is suspected, additional workup to evaluate the extent of the disease is necessary. This includes a chest CT and a bone scan if enzyme levels are abnormal or the patient shows signs or symptoms of skeletal involvement.

If the evidence of spread is limited to nodes, nodal biopsy should be considered and patients should be managed as in the case for T4 disease. Patients who present with disseminated metastatic disease are generally treated with systemic chemotherapy. Management of persistent disseminated disease may involve chemotherapy, radiation, or a combination of the two. Details on the choice of regimens have been discussed above.

Upper Genitourinary Tract Tumors

Upper tract tumors, including those that originate in the renal pelvis or in the ureter, are relatively uncommon.



Renal Pelvis Tumors

Tumors that develop in the renal pelvis may be identified during evaluation of hematuria or a renal mass. In the latter case, renal pelvic tumors must be distinguished from the more typical adenocarcinomas that originate in the renal parenchyma. These tumors may also be detected during an assessment to pinpoint the source of a positive cytology in the setting of a negative cystoscopy with a retrograde pyelogram.

Workup

The evaluation of a patient with a suspected renal pelvic tumor should include cystoscopy and imaging of the upper tract collecting system with IVP, CT urography, retrograde pyelogram, ureteroscopy, MRI urogram, or a combination of techniques. A chest radiograph can help evaluate for possible metastatic disease and assess any comorbid diseases that may be present. Urine cytology obtained from a urine sample or during a cystoscopy may help identify carcinoma cells. Hematologic, renal, and hepatic function should also be evaluated. Additional imaging studies, such as renal scan or bone scan, may be needed if indicated by the results of these tests or the presence of specific symptoms.

Primary Treatment

In general, the primary form of treatment for renal pelvic tumors is surgery.

Well-differentiated tumors of low grade may be managed with a nephroureterectomy with a cuff of bladder, a nephron-sparing procedure through a transureteroscopic approach, or a percutaneous approach with or without postsurgical intrapelvic chemotherapy or BCG. High-grade tumors or those that are large and invade the renal parenchyma are managed through nephroureterectomy with a cuff of bladder and regional lymphadenectomy. Decline in renal function

following surgery may preclude adjuvant therapy. Hence, in selected patients, neoadjuvant chemotherapy may be considered based on extrapolation of data from bladder cancer series.⁴¹⁻⁴³ If metastatic disease is documented or associated comorbid conditions are present, treatment should include systemic chemotherapy with regimens similar to those used for urothelial (transitional cell) bladder tumors.

In the settings of positive upper tract cytology but negative imaging and biopsy studies, treatment remains controversial and appropriate management is currently poorly defined. Frequent monitoring for disease is necessary for these patients.

Follow-up

Subsequent management is dictated by the extent of disease at surgery. Tumors that are pathologic stage pT0 or pT1 should be followed up with serial cystoscopies at 3-month intervals for the first year and, if negative, every 6 months thereafter. Such tumors should also be followed up with ureteroscopy and upper tract imaging (such as IVP; retrograde pyelogram; CT, if available; or MRI urogram) at 3- to 12-month intervals if endoscopic resection is considered.

Patients with pathologic stage pT2, pT3, pT4, or nodal disease should be considered for adjuvant chemotherapy. Serial evaluations of the urothelial tract, along with imaging studies to exclude metastatic disease, should also be performed.

Ureteral Tumors

Ureteral tumors may develop de novo or in patients who have undergone successful treatment for superficial tumors that originate in the bladder. The presentation varies as a function of disease extent. Ureteral tumors may be identified in patients who have a positive cytology with a negative cystoscopy in whom selective catheterization of

the ureters is performed. More extensive lesions may result in pain or obstruction.

Workup

The evaluation is similar to that outlined for tumors that originate in the renal pelvis.

Treatment

For resectable ureteral tumors, the primary management is surgery. The specific procedure required varies depending on the location of the tumor (upper, mid, or distal location) and disease extent. Neoadjuvant chemotherapy may be considered in selected patients, such as when the degree of invasiveness is established before definitive surgery.⁹⁸

Tumors that originate in the upper ureter occasionally can be managed endoscopically but more commonly are treated with nephroureterectomy with a cuff of bladder plus regional lymphadenectomy for high-grade tumors. A portion of the bladder is removed to ensure complete removal of the entire intramural ureter. Tumors that originate in the mid portion can be divided by grade and size. Small, low-grade tumors can be managed with excision and ureteroureterostomy or complete ureterectomy and ileal ureter in highly selected patients, endoscopic resection, or nephroureterectomy with a cuff of bladder and consideration of regional lymphadenectomy. Larger, high-grade lesions are managed with nephroureterectomy with a cuff of bladder and regional lymphadenectomy. Distal ureteral tumors may be managed with a distal ureterectomy and reimplantation of the ureter (preferred if clinically feasible), endoscopic resection, or, in some cases, a nephroureterectomy with a cuff of bladder, with the addition of regional lymphadenectomy recommended for high-grade tumors.

Follow-up

The final pathologic stage is used to guide subsequent management, as is the case for tumors that originate in other sites. No adjuvant therapy is advised for lesions that are pT1 or less, but serial follow-up of the urothelial tracts or remaining unit (as previously described under *Renal Pelvis Tumors*) is recommended.

Patients with more extensive disease are advised to consider systemic adjuvant treatment with chemotherapy, depending on the patient's anticipated tolerance to the regimen based on comorbidities. The reasons for considering adjuvant therapy are similar to those for tumors that originate in the bladder.

Urothelial (Transitional Cell) Carcinomas of the Prostate

Urothelial (transitional cell) carcinomas of the prostate represent a distinct entity with a unique staging system. In this respect, they must be distinguished from urothelial (transitional cell) carcinomas of bladder origin that invade into the prostate through the bladder wall. Urothelial (transitional cell) carcinomas of the prostate may occur de novo or, more typically, concurrently or after treatment of a bladder cancer. As in the case with tumors originating in other sites of the urothelium, management of prostate urothelial (transitional cell) carcinomas is based on extent of disease with particular reference to the urethra, duct, acini, and stroma.

Workup

The evaluation of a suspected urothelial carcinoma of the prostate includes a digital rectal examination (DRE), cystoscopy with bladder biopsy, and a TUR biopsy of the prostate that includes the prostatic stroma. Multiple stromal biopsies are also advised and, if the DRE is abnormal, determination of the prostate-specific antigen level and

additional needle biopsies may be required in selected patients to exclude primary adenocarcinoma of the prostate. Upper tract collecting system imaging is also recommended.

Primary Treatment

Pending histologic confirmation, tumors that are limited to the prostatic urethra with no acinar or stromal invasion can be managed with BCG and transurethral resection of the prostate (TURP), with follow-up similar to that for superficial disease of the bladder. Patients with tumors that invade the ducts, acini, or stroma should undergo an additional workup with chest radiograph, or CT if necessary, to exclude metastatic disease, and then a cystoprostatectomy with or without urethrectomy should be performed. Neoadjuvant chemotherapy may be considered in patients with stromal invasion, based on extrapolation of data from bladder cancer therapy.⁴¹⁻⁴³ Alternatively, TURP and BCG may be offered to patients with only ductal and acini invasion. Adjuvant chemotherapy may be advised for stromal invasion after primary treatment. Local recurrences in patients undergoing TURP and BCG therapy are treated with cystoprostatectomy with or without urethrectomy.

Primary Carcinoma of the Urethra

Primary carcinoma that arises in the urethra is rare. Unlike for bladder cancer, squamous cell carcinoma is the most common histologic subtype for urethral cancer.⁹⁹ The 5-year overall survival is 42%.^{100,101} Stage and disease location are the most important prognostic factors for male patients, while tumor size and histology are also prognostically significant for female patients.^{99,101} Unfortunately, there is a lack of robust, prospective data to support treatment decisions due to disease rarity.

Workup

A cystourethroscopy should be performed if carcinoma of the urethra is suspected. This includes EUA and TUR or transvaginal biopsy. Chest x-ray and MRI of the pelvis are also recommended to evaluate the extent of the disease.

If palpable inguinal lymph nodes are present, a chest/abdominal/pelvic CT and lymph node biopsy should be performed.

Treatment

Patients with Tis, Ta, or T1 disease should have a repeat TUR. In select cases, TUR is followed by intraurethral therapy. A total urethrectomy may be considered if the patient has undergone a radical cystectomy or cutaneous diversion.

Treatment for T2 disease is based on patient gender and tumor location. For male patients with pendulous urethra, distal urethrectomy or partial penectomy are viable options. Patients may consider neoadjuvant chemotherapy (category 2B) before a urethrectomy. Those who have positive margins may undergo additional surgery, radiation, or chemoradiation. At recurrence, options include chemotherapy, total penectomy, radiation, or a combination.

Male patients with T2 tumors in the bulbar urethra should undergo urethrectomy with or without cystoprostatectomy. Adjuvant chemotherapy or chemoradiation may be considered for pathologic T3, T4, or nodal disease. Recurrent cases may be treated with chemotherapy and/or radiation.

Initial treatment options for female patients with T2 tumors include chemoradiation or urethrectomy with or without cystectomy. Partial urethrectomy was associated with a high urethral recurrence rate.¹⁰² At



NCCN Guidelines Version 2.2014

Bladder Cancer

recurrence, the patient may receive chemotherapy or pelvic exenteration (category 2B).

A multimodal treatment approach (ie, surgery, chemotherapy, radiation) is common for advanced disease. If chemotherapy is used, the choice of regimen should be based on histology. A cohort study reported a 72% response rate with the following treatment scheme before surgery: cisplatin, gemcitabine, and ifosfamide for squamous cell carcinoma; 5-FU, gemcitabine, and cisplatin-based regimens for adenocarcinoma; and MVAC for urothelial tumors.¹⁰³ Combined chemoradiation with 5-FU and mitomycin C has shown efficacy in a series of male patients with squamous cell carcinoma of the urethra.¹⁰⁴ Patients receiving salvage surgery after chemoradiation had a higher 5-year disease-free survival rate (72%) than those receiving chemoradiation alone (54%).

Patients with T3 or T4 disease but no clinical nodes should either receive neoadjuvant chemotherapy followed by consolidative surgery or radiation, or radiation with or without chemotherapy. If positive nodes are present, radiation with or without chemotherapy is the preferred treatment for squamous cell carcinoma. Chemotherapy followed by consideration of consolidative surgery is an alternative. At recurrence, the patient may undergo pelvic exenteration (category 2B) with or without en bloc ilioinguinal lymphadenectomy. Chemotherapy alone or in addition to surgery is a category 2B option.

Patients with distant metastases should receive chemotherapy based on histology.

Non-Urothelial Carcinomas of the Bladder

Approximately 10% of bladder tumors are non-urothelial (non-transitional cell) carcinoma. These pathologic entities include mixed histology, pure squamous, adenocarcinoma, small cell tumors, urachal

carcinoma, or primary bladder sarcoma. Depending on the pathologic findings, adjuvant chemotherapy may or may not be recommended. Patients with non-urothelial invasive disease are generally treated with cystectomy, although those with certain urachal tumors require complete urachal resection (en bloc resection of the urachal ligament with the umbilicus) or may be appropriately treated with partial cystectomy. In patients with non-urothelial carcinomas of any stage, no data support the use of adjuvant chemotherapy, although the risk for relapse may be high. Some of the general principles of management applicable to urothelial (transitional cell) carcinomas are appropriate with minor variations. These variations are documented in the algorithm.

Patients with small cell carcinoma of the bladder are best treated with initial chemotherapy (see [NCCN Guidelines for Small Cell Lung Cancer](#)) followed by either radiation therapy or cystectomy as consolidation, if there is no metastatic disease. Primary bladder sarcomas are treated as per the [NCCN Guidelines for Soft Tissue Sarcoma](#).

Summary

Urothelial tumors represent a spectrum of diseases with a range of prognoses. After a tumor is diagnosed anywhere within the urothelial tract, the patient remains at risk for developing a new lesion at a different or same location and with a similar or more advanced stage. Continued monitoring for recurrence is an essential part of management, because most recurrences are superficial and can be treated endoscopically. Within each category of disease, more refined methods to determine prognosis and guide management, based on molecular staging, are under development with the goal of optimizing each patient's likelihood of cure and chance for organ preservation.

For patients with more extensive disease, newer treatments typically involve combined modality approaches using recently developed



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014 Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)

surgical procedures, or 3-dimensional treatment planning for more precise delivery of radiation therapy. Although these are not appropriate in all cases, they offer the promise of an improved quality of life and prolonged survival.

Finally, within the category of metastatic disease, several new agents have been identified that seem superior to those currently considered standard therapies. Experts believe, therefore, that the treatment of urothelial tumors will evolve rapidly over the next few years, with improved outcomes for patients at all stages of disease.

Table 1. Principles of Pathology Management: Malignancy Grading of Bladder Carcinoma: Old and New Systems^{a,b}

Modified Bergkvist 1987	WHO 1973	WHO/ISUP 1998 Consensus WHO, 2004
Papilloma grade 0	Papilloma	Papilloma
Papilloma with atypia grade 1	TCC grade 1	Papillary urothelial neoplasm of low malignant potential
Urothelial carcinoma grade 2A	TCC grade 1	Urothelial carcinoma, low-grade
Urothelial carcinoma grade 2B	TCC grade 2	Urothelial carcinoma, low-grade or high-grade
Urothelial carcinoma grade 3	TCC grade 3	Urothelial carcinoma, high-grade

^aFrom Droller MJ. Bladder Cancer, Current Diagnosis and Treatment. Totowa (NJ): Humana Press, 2001.

^bSeveral classifications have been proposed for grading of tumors of the bladder epithelium. Because they are in general usage, the current NCCN Guidelines for Bladder and Upper Tract Cancers continue to use the WHO histologic classification of tumors of the urinary tract from 1973. However, a revised classification has been adopted by numerous organizations, including the WHO in its most recent publication in 2004. This classification has also been adopted by the College of American Pathologists, the American Society of Clinical Pathology, and the International Society of Urological Pathology.

Please note several major changes in this classification. First, the term *transitional cell* is changed to *urothelial*. Also, dysplastic changes of the urothelium without invasion are now classified either as carcinoma in situ or as dysplasia without specification of mild, moderate, or severe. Any dysplastic, flat, noninvasive lesion that does not meet the criteria of CIS is referred to as *dysplasia*.

The criteria used for the new classification system are more specific than those for the 1973 WHO classification system. The entire classification system, including the range of types of tumors, is presented on pages 90–91 of the new WHO classification of tumors.

References

- Busch C, Hawes D, Johansson S, Cote R. Pathologic assessment of bladder cancer and pitfalls in staging. In: Droller MJ, ed. Bladder Cancer, Current Diagnosis and Treatment. Totowa (NJ): Humana Press, 2001:149–182.
- Busch C, Algaba F. The WHO/ISUP 1998 and WHO 1999 systems for malignancy grading of bladder cancer. Scientific foundation and translation to one another and previous systems. Virchows Arch 2002;441:105–108.
- Check W. Bladder biopsies in step with clinical side. CAP Today (College of American Pathologists) 2004;18:43–54.
- Eble JN, Sauter G, Epstein JI, et al. Pathology and Genetics of Tumours of the Urinary System and Male Genital Organs. Lyon, France: IARC (International Agency for Research on Cancer) Press, 2004.
- Epstein JI, Amin MB, Reuter VR, Mostofi FK. The World Health Organization/International Society of Urological Pathology consensus classification of urothelial (transitional cell) neoplasms of the urinary bladder. Bladder Consensus Conference Committee. Am J Surg Pathol 1998;22:1435–1448.
- Murphy WM, Grignon DJ, Perlman EJ. Tumors of the Kidney, Bladder, and Related Urinary Structures. AFIP Atlas of Tumor Pathology. 4th series. Washington (DC): American Registry of Pathology, 2004.



NCCN Guidelines Version 2.2014

Bladder Cancer

Table 2. Combination Chemotherapy Regimens

Regimen	Dosage	
Gemcitabine/ Cisplatin ^{77,87,105,106}	Gemcitabine*	1000 mg/m ² on days 1, 8, 15 of a 28-day cycle or 1000 mg/m ² on days 1, 8 of a 21-day cycle
	Cisplatin	70 mg/m ² on day 2
Dose-Dense MVAC ^{78,79}	Methotrexate	30 mg/m ² on day 1 or day 2 of a 14-day cycle
	Vinblastine	3 mg/m ² on day 1 or day 2
	Doxorubicin	30 mg/m ² on day 1 or day 2
	Cisplatin	70 mg/m ² on day 1
CMV ⁴⁶	Methotrexate	30 mg/m ² on days 1, 8 of a 21-day cycle
	Vinblastine	4 mg/m ² on days 1, 8
	Cisplatin	100 mg/m ² on day 2 before hydration
	Folinic acid	15 mg every 6 hours on days 2, 9 after hydration

*This dose should not be combined with radiation.



NCCN Guidelines Version 2.2014

Bladder Cancer

References

1. Siegel R, Naishadham D, Jemal A. Cancer statistics, 2012. *CA Cancer J Clin* 2012;62:10-29. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22237781>.
2. Chalasani V, Chin JL, Izawa JI. Histologic variants of urothelial bladder cancer and nonurothelial histology in bladder cancer. *Can Urol Assoc J* 2009;3:S193-198. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/20019984>.
3. Wasco MJ, Daignault S, Zhang Y, et al. Urothelial carcinoma with divergent histologic differentiation (mixed histologic features) predicts the presence of locally advanced bladder cancer when detected at transurethral resection. *Urology* 2007;70:69-74. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17656211>.
4. Kamat AM, Dickstein RJ, Messetti F, et al. Use of fluorescence in situ hybridization to predict response to bacillus Calmette-Guerin therapy for bladder cancer: results of a prospective trial. *J Urol* 2012;187:862-867. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22245325>.
5. Grossman HB, Soloway M, Messing E, et al. Surveillance for recurrent bladder cancer using a point-of-care proteomic assay. *JAMA* 2006;295:299-305. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16418465>.
6. American Urological Association. Guideline for the Management of Nonmuscle Invasive Bladder Cancer: (Stages Ta,T1, and Tis). 2007. Available at: <http://www.auanet.org/content/clinical-practice-guidelines/clinical-guidelines.cfm>. Accessed November 27,2012.
7. Sylvester RJ, van der Meijden AP, Oosterlinck W, et al. Predicting recurrence and progression in individual patients with stage Ta T1 bladder cancer using EORTC risk tables: a combined analysis of 2596 patients from seven EORTC trials. *Eur Urol* 2006;49:466-465; discussion 475-467. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16442208>.
8. Edge S, Byrd D, Compton C, eds. *AJCC Cancer Staging Manual*, 7th ed. New York: Springer; 2010.
9. Montironi R, Lopez-Beltran A. The 2004 WHO classification of bladder tumors: a summary and commentary. *Int J Surg Pathol* 2005;13:143-153. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15864376>.
10. Babjuk M, Oosterlinck W, Sylvester R, et al. EAU guidelines on non-muscle-invasive urothelial carcinoma of the bladder, the 2011 update. *Eur Urol* 2011;59:997-1008. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21458150>.
11. Herr HW. The value of a second transurethral resection in evaluating patients with bladder tumors. *J Urol* 1999;162:74-76. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10379743>.
12. Sylvester RJ, Oosterlinck W, van der Meijden AP. A single immediate postoperative instillation of chemotherapy decreases the risk of recurrence in patients with stage Ta T1 bladder cancer: a meta-analysis of published results of randomized clinical trials. *J Urol* 2004;171:2186-2190, quiz 2435. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15126782>.
13. Berrum-Svennung I, Granfors T, Jahnson S, et al. A single instillation of epirubicin after transurethral resection of bladder tumors prevents only small recurrences. *J Urol* 2008;179:101-105; discussion 105-106. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17997459>.
14. Bohle A, Leyh H, Frei C, et al. Single postoperative instillation of gemcitabine in patients with non-muscle-invasive transitional cell carcinoma of the bladder: a randomised, double-blind, placebo-controlled phase III multicentre study. *Eur Urol* 2009;56:495-503. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19560257>.
15. Gudjonsson S, Adell L, Merdasa F, et al. Should all patients with non-muscle-invasive bladder cancer receive early intravesical chemotherapy after transurethral resection? The results of a



prospective randomised multicentre study. Eur Urol 2009;55:773-780. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19153001>.

16. Huncharek M, McGarry R, Kupelnick B. Impact of intravesical chemotherapy on recurrence rate of recurrent superficial transitional cell carcinoma of the bladder: results of a meta-analysis. Anticancer Res 2001;21:765-769. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11299841>.

17. Huncharek M, Geschwind JF, Witherspoon B, et al. Intravesical chemotherapy prophylaxis in primary superficial bladder cancer: a meta-analysis of 3703 patients from 11 randomized trials. J Clin Epidemiol 2000;53:676-680. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10941943>.

18. Grimm MO, Steinhoff C, Simon X, et al. Effect of routine repeat transurethral resection for superficial bladder cancer: a long-term observational study. J Urol 2003;170:433-437. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12853793>.

19. Bohle A, Jocham D, Bock PR. Intravesical bacillus Calmette-Guerin versus mitomycin C for superficial bladder cancer: a formal meta-analysis of comparative studies on recurrence and toxicity. J Urol 2003;169:90-95. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12478111>.

20. Han RF, Pan JG. Can intravesical bacillus Calmette-Guerin reduce recurrence in patients with superficial bladder cancer? A meta-analysis of randomized trials. Urology 2006;67:1216-1223. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16765182>.

21. Shelley MD, Kynaston H, Court J, et al. A systematic review of intravesical bacillus Calmette-Guerin plus transurethral resection vs transurethral resection alone in Ta and T1 bladder cancer. BJU Int 2001;88:209-216. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11488731>.

22. Shelley MD, Wilt TJ, Court J, et al. Intravesical bacillus Calmette-Guerin is superior to mitomycin C in reducing tumour recurrence in high-risk superficial bladder cancer: a meta-analysis of randomized trials. BJU Int 2004;93:485-490. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15008714>.

23. Lokeshwar VB, Habuchi T, Grossman HB, et al. Bladder tumor markers beyond cytology: International Consensus Panel on bladder tumor markers. Urology 2005;66:35-63. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16399415>.

24. Ramirez-Backhaus M, Dominguez-Escrig J, Collado A, et al. Restaging transurethral resection of bladder tumor for high-risk stage Ta and T1 bladder cancer. Curr Urol Rep 2012;13:109-114. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22367558>.

25. Divrik RT, Yildirim U, Zorlu F, Ozen H. The effect of repeat transurethral resection on recurrence and progression rates in patients with T1 tumors of the bladder who received intravesical mitomycin: a prospective, randomized clinical trial. J Urol 2006;175:1641-1644. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16600720>.

26. Herr HW, Sogani PC. Does early cystectomy improve the survival of patients with high risk superficial bladder tumors? J Urol 2001;166:1296-1299. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11547061>.

27. Steinberg G, Bahnson R, Brosman S, et al. Efficacy and safety of valrubicin for the treatment of Bacillus Calmette-Guerin refractory carcinoma in situ of the bladder. The Valrubicin Study Group. J Urol 2000;163:761-767. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10687972>.

28. Raj GV, Herr H, Serio AM, et al. Treatment paradigm shift may improve survival of patients with high risk superficial bladder cancer. J Urol 2007;177:1283-1286; discussion 1286. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/17382713>.



29. Bohle A, Bock PR. Intravesical bacille Calmette-Guerin versus mitomycin C in superficial bladder cancer: formal meta-analysis of comparative studies on tumor progression. *Urology* 2004;63:682-686; discussion 686-687. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15072879>.

30. Sylvester RJ, van der MA, Lamm DL. Intravesical bacillus Calmette-Guerin reduces the risk of progression in patients with superficial bladder cancer: a meta-analysis of the published results of randomized clinical trials. *J Urol* 2002;168:1964-1970. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12394686>.

31. Sylvester RJ, van der Meijden AP, Witjes JA, Kurth K. Bacillus calmette-guerin versus chemotherapy for the intravesical treatment of patients with carcinoma in situ of the bladder: a meta-analysis of the published results of randomized clinical trials. *J Urol* 2005;174:86-91; discussion 91-92. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15947584>.

32. Lamm DL, Blumenstein BA, Crissman JD, et al. Maintenance bacillus Calmette-Guerin immunotherapy for recurrent TA, T1 and carcinoma in situ transitional cell carcinoma of the bladder: a randomized Southwest Oncology Group Study. *J Urol* 2000;163:1124-1129. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10737480>.

33. Malmstrom PU, Sylvester RJ, Crawford DE, et al. An individual patient data meta-analysis of the long-term outcome of randomised studies comparing intravesical mitomycin C versus bacillus Calmette-Guerin for non-muscle-invasive bladder cancer. *Eur Urol* 2009;56:247-256. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19409692>.

34. Oddens J, Brausi M, Sylvester R, et al. Final results of an EORTC-GU cancers group randomized study of maintenance bacillus Calmette-Guerin in intermediate- and high-risk Ta, T1 papillary carcinoma of the urinary bladder: one-third dose versus full dose and 1 year versus 3 years of maintenance. *Eur Urol* 2013;63:462-472. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23141049>.

35. Verma S, Rajesh A, Prasad SR, et al. Urinary bladder cancer: role of MR imaging. *Radiographics* 2012;32:371-387. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22411938>.

36. Leissner J, Hohenfellner R, Thuroff JW, Wolf HK. Lymphadenectomy in patients with transitional cell carcinoma of the urinary bladder; significance for staging and prognosis. *BJU Int* 2000;85:817-823. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10792159>.

37. Herr HW, Bochner BH, Dalbagni G, et al. Impact of the number of lymph nodes retrieved on outcome in patients with muscle invasive bladder cancer. *J Urol* 2002;167:1295-1298. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11832716>.

38. Herr HW, Faulkner JR, Grossman HB, et al. Surgical factors influence bladder cancer outcomes: a cooperative group report. *J Clin Oncol* 2004;22:2781-2789. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15199091>.

39. Konety BR, Joslyn SA, O'Donnell MA. Extent of pelvic lymphadenectomy and its impact on outcome in patients diagnosed with bladder cancer: analysis of data from the Surveillance, Epidemiology and End Results Program data base. *J Urol* 2003;169:946-950. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12576819>.

40. Wright JL, Lin DW, Porter MP. The association between extent of lymphadenectomy and survival among patients with lymph node metastases undergoing radical cystectomy. *Cancer* 2008;112:2401-2408. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/18383515>.

41. Grossman HB, Natale RB, Tangen CM, et al. Neoadjuvant chemotherapy plus cystectomy compared with cystectomy alone for locally advanced bladder cancer. *N Engl J Med* 2003;349:859-866. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/12944571>.

42. Sherif A, Holmberg L, Rintala E, et al. Neoadjuvant cisplatin based combination chemotherapy in patients with invasive bladder



NCCN Guidelines Version 2.2014

Bladder Cancer

cancer: a combined analysis of two Nordic studies. *Eur Urol* 2004;45:297-303. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15036674>.

43. Winkquist E, Kirchner TS, Segal R, et al. Neoadjuvant chemotherapy for transitional cell carcinoma of the bladder: a systematic review and meta-analysis. *J Urol* 2004;171:561-569. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14713760>.

44. Kitamura H, Tsukamoto T, Shibata T, et al. Randomised phase III study of neoadjuvant chemotherapy with methotrexate, doxorubicin, vinblastine and cisplatin followed by radical cystectomy compared with radical cystectomy alone for muscle-invasive bladder cancer: Japan Clinical Oncology Group study JCOG0209. *Ann Oncol* 2014. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/24669010>.

45. Neoadjuvant chemotherapy in invasive bladder cancer: update of a systematic review and meta-analysis of individual patient data advanced bladder cancer (ABC) meta-analysis collaboration. *Eur Urol* 2005;48:202-205; discussion 205-206. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15939524>.

46. Griffiths G, Hall R, Sylvester R, et al. International phase III trial assessing neoadjuvant cisplatin, methotrexate, and vinblastine chemotherapy for muscle-invasive bladder cancer: long-term results of the BA06 30894 trial. *J Clin Oncol* 2011;29:2171-2177. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21502557>.

47. Hussain MH, Wood DP, Bajorin DF, et al. Bladder cancer: narrowing the gap between evidence and practice. *J Clin Oncol* 2009;27:5680-5684. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19858384>.

48. Adjuvant chemotherapy in invasive bladder cancer: a systematic review and meta-analysis of individual patient data Advanced Bladder Cancer (ABC) Meta-analysis Collaboration. *Eur Urol* 2005;48:189-199; discussion 199-201. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/15939530>.

49. Lehmann J, Franzaring L, Thuroff J, et al. Complete long-term survival data from a trial of adjuvant chemotherapy vs control after radical cystectomy for locally advanced bladder cancer. *BJU Int* 2006;97:42-47. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16336326>.

50. Stockle M, Wellek S, Meyenburg W, et al. Radical cystectomy with or without adjuvant polychemotherapy for non-organ-confined transitional cell carcinoma of the urinary bladder: prognostic impact of lymph node involvement. *Urology* 1996;48:868-875. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/8973669>.

51. Skinner DG, Daniels JR, Russell CA, et al. The role of adjuvant chemotherapy following cystectomy for invasive bladder cancer: a prospective comparative trial. *J Urol* 1991;145:459-464; discussion 464-457. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1997689>.

52. Cignetti F, Ruggeri EM, Felici A, et al. Adjuvant chemotherapy with cisplatin and gemcitabine versus chemotherapy at relapse in patients with muscle-invasive bladder cancer submitted to radical cystectomy: an Italian, multicenter, randomized phase III trial. *Ann Oncol* 2011. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/21859900>.

53. Millikan R, Dinney C, Swanson D, et al. Integrated therapy for locally advanced bladder cancer: final report of a randomized trial of cystectomy plus adjuvant M-VAC versus cystectomy with both preoperative and postoperative M-VAC. *J Clin Oncol* 2001;19:4005-4013. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11600601>.

54. Zaghloul MS, Awwad HK, Akoush HH, et al. Postoperative radiotherapy of carcinoma in bilharzial bladder: improved disease free survival through improving local control. *Int J Radiat Oncol Biol Phys* 1992;23:511-517. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/1612951>.

55. Cozzarini C, Pellegrini D, Fallini M, et al. Reappraisal of the role of adjuvant radiotherapy in muscle-invasive transitional cell carcinoma of the bladder. *International Journal of Radiation Oncology, Biology,*



NCCN Guidelines Version 2.2014

Bladder Cancer

Physics 1999;45(Suppl):221-222. Available at:

<http://linkinghub.elsevier.com/retrieve/pii/S0360301699901621>.

56. Mak RH, Zietman AL, Heney NM, et al. Bladder preservation: optimizing radiotherapy and integrated treatment strategies. *BJU Int* 2008;102:1345-1353. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/19035903>.

57. Gakis G, Efstathiou J, Lerner SP, et al. ICUD-EAU International Consultation on Bladder Cancer 2012: Radical Cystectomy and Bladder Preservation for Muscle-Invasive Urothelial Carcinoma of the Bladder. *Eur Urol* 2012. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22917985>.

58. Fedeli U, Fedewa SA, Ward EM. Treatment of muscle invasive bladder cancer: evidence from the National Cancer Database, 2003 to 2007. *J Urol* 2011;185:72-78. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/21074192>.

59. Shipley WU, Kaufman DS, Zehr E, et al. Selective bladder preservation by combined modality protocol treatment: long-term outcomes of 190 patients with invasive bladder cancer. *Urology* 2002;60:62-67; discussion 67-68. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12100923>.

60. Shipley WU, Winter KA, Kaufman DS, et al. Phase III trial of neoadjuvant chemotherapy in patients with invasive bladder cancer treated with selective bladder preservation by combined radiation therapy and chemotherapy: initial results of Radiation Therapy Oncology Group 89-03. *J Clin Oncol* 1998;16:3576-3583. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9817278>.

61. Herr HW, Bajorin DF, Scher HI. Neoadjuvant chemotherapy and bladder-sparing surgery for invasive bladder cancer: ten-year outcome. *J Clin Oncol* 1998;16:1298-1301. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9552029>.

62. Housset M, Maulard C, Chretien Y, et al. Combined radiation and chemotherapy for invasive transitional-cell carcinoma of the bladder: a prospective study. *J Clin Oncol* 1993;11:2150-2157. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8229129>.

63. Coppin CM, Gospodarowicz MK, James K, et al. Improved local control of invasive bladder cancer by concurrent cisplatin and preoperative or definitive radiation. The National Cancer Institute of Canada Clinical Trials Group. *J Clin Oncol* 1996;14:2901-2907. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/8918486>.

64. James ND, Hussain SA, Hall E, et al. Radiotherapy with or without chemotherapy in muscle-invasive bladder cancer. *N Engl J Med* 2012;366:1477-1488. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22512481>.

65. Herr HW. Conservative management of muscle-infiltrating bladder cancer: prospective experience. *J Urol* 1987;138:1162-1163. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/3669160>.

66. Sweeney P, Kursh ED, Resnick MI. Partial cystectomy. *Urol Clin North Am* 1992;19:701-711. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/1441027>.

67. Sternberg CN, Pansadoro V, Calabro F, et al. Can patient selection for bladder preservation be based on response to chemotherapy? *Cancer* 2003;97:1644-1652. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12655521>.

68. Rodel C, Grabenbauer GG, Kuhn R, et al. Combined-modality treatment and selective organ preservation in invasive bladder cancer: long-term results. *J Clin Oncol* 2002;20:3061-3071. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/12118019>.

69. Efstathiou JA, Spiegel DY, Shipley WU, et al. Long-term outcomes of selective bladder preservation by combined-modality therapy for invasive bladder cancer: the MGH experience. *Eur Urol* 2012;61:705-711. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22101114>.



NCCN Guidelines Version 2.2014

Bladder Cancer

70. Zapatero A, Martin De Vidales C, Arellano R, et al. Long-term Results of Two Prospective Bladder-sparing Trimodality Approaches for Invasive Bladder Cancer: Neoadjuvant Chemotherapy and Concurrent Radio-chemotherapy. *Urology* 2012. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/22999456>.

71. Kaufman DS, Winter KA, Shipley WU, et al. The initial results in muscle-invasive bladder cancer of RTOG 95-06: phase I/II trial of transurethral surgery plus radiation therapy with concurrent cisplatin and 5-fluorouracil followed by selective bladder preservation or cystectomy depending on the initial response. *Oncologist* 2000;5:471-476. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11110598>.

72. Hagan MP, Winter KA, Kaufman DS, et al. RTOG 97-06: initial report of a phase I-II trial of selective bladder conservation using TURBT, twice-daily accelerated irradiation sensitized with cisplatin, and adjuvant MCV combination chemotherapy. *Int J Radiat Oncol Biol Phys* 2003;57:665-672. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/14529770>.

73. Kaufman DS, Winter KA, Shipley WU, et al. Phase I-II RTOG study (99-06) of patients with muscle-invasive bladder cancer undergoing transurethral surgery, paclitaxel, cisplatin, and twice-daily radiotherapy followed by selective bladder preservation or radical cystectomy and adjuvant chemotherapy. *Urology* 2009;73:833-837. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19100600>.

74. Mitin T, Hunt D, Shipley WU, et al. Transurethral surgery and twice-daily radiation plus paclitaxel-cisplatin or fluorouracil-cisplatin with selective bladder preservation and adjuvant chemotherapy for patients with muscle invasive bladder cancer (RTOG 0233): a randomised multicentre phase 2 trial. *Lancet Oncol* 2013;14:863-872. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/23823157>.

75. Efstathiou JA, Bae K, Shipley WU, et al. Late pelvic toxicity after bladder-sparing therapy in patients with invasive bladder cancer: RTOG 89-03, 95-06, 97-06, 99-06. *J Clin Oncol* 2009;27:4055-4061. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19636019>.

76. Kaufman D, Raghavan D, Carducci M, et al. Phase II trial of gemcitabine plus cisplatin in patients with metastatic urothelial cancer. *J Clin Oncol* 2000;18:1921-1927. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/10784633>.

77. von der Maase H, Sengelov L, Roberts JT, et al. Long-term survival results of a randomized trial comparing gemcitabine plus cisplatin, with methotrexate, vinblastine, doxorubicin, plus cisplatin in patients with bladder cancer. *J Clin Oncol* 2005;23:4602-4608. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16034041>.

78. Sternberg CN, de Mulder PH, Schornagel JH, et al. Randomized phase III trial of high-dose-intensity methotrexate, vinblastine, doxorubicin, and cisplatin (MVAC) chemotherapy and recombinant human granulocyte colony-stimulating factor versus classic MVAC in advanced urothelial tract tumors: European Organization for Research and Treatment of Cancer Protocol no. 30924. *J Clin Oncol* 2001;19:2638-2646. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11352955>.

79. Sternberg CN, de Mulder P, Schornagel JH, et al. Seven year update of an EORTC phase III trial of high-dose intensity M-VAC chemotherapy and G-CSF versus classic M-VAC in advanced urothelial tract tumours. *Eur J Cancer* 2006;42:50-54. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/16330205>.

80. von der Maase H, Hansen SW, Roberts JT, et al. Gemcitabine and cisplatin versus methotrexate, vinblastine, doxorubicin, and cisplatin in advanced or metastatic bladder cancer: results of a large, randomized, multinational, multicenter, phase III study. *J Clin Oncol* 2000;18:3068-3077. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/11001674>.

81. De Santis M, Bellmunt J, Mead G, et al. Randomized phase II/III trial assessing gemcitabine/ carboplatin and methotrexate/carboplatin/vinblastine in patients with advanced urothelial cancer "unfit" for cisplatin-based chemotherapy: phase II--results of EORTC study 30986. *J Clin Oncol* 2009;27:5634-5639. Available at: <http://www.ncbi.nlm.nih.gov/pubmed/19786668>.



82. Burch PA, Richardson RL, Cha SS, et al. Phase II study of paclitaxel and cisplatin for advanced urothelial cancer. J Urol 2000;164:1538-1542. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11025699>.

83. Meluch AA, Greco FA, Burris HA, 3rd, et al. Paclitaxel and gemcitabine chemotherapy for advanced transitional-cell carcinoma of the urothelial tract: a phase II trial of the Minnie pearl cancer research network. J Clin Oncol 2001;19:3018-3024. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11408496>.

84. Bellmunt J, Guillem V, Paz-Ares L, et al. Phase I-II study of paclitaxel, cisplatin, and gemcitabine in advanced transitional-cell carcinoma of the urothelium. Spanish Oncology Genitourinary Group. J Clin Oncol 2000;18:3247-3255. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10986057>.

85. Hussain M, Vaishampayan U, Du W, et al. Combination paclitaxel, carboplatin, and gemcitabine is an active treatment for advanced urothelial cancer. J Clin Oncol 2001;19:2527-2533. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11331332>.

86. Pectasides D, Glotsos J, Bountouroglou N, et al. Weekly chemotherapy with docetaxel, gemcitabine and cisplatin in advanced transitional cell urothelial cancer: a phase II trial. Ann Oncol 2002;13:243-250. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11886001>.

87. Bellmunt J, von der Maase H, Mead GM, et al. Randomized phase III study comparing paclitaxel/cisplatin/gemcitabine and gemcitabine/cisplatin in patients with locally advanced or metastatic urothelial cancer without prior systemic therapy: EORTC Intergroup Study 30987. J Clin Oncol 2012;30:1107-1113. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22370319>.

88. Lorusso V, Pollera CF, Antimi M, et al. A phase II study of gemcitabine in patients with transitional cell carcinoma of the urinary tract previously treated with platinum. Italian Co-operative Group on

Bladder Cancer. Eur J Cancer 1998;34:1208-1212. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9849481>.

89. McCaffrey JA, Hilton S, Mazumdar M, et al. Phase II trial of docetaxel in patients with advanced or metastatic transitional-cell carcinoma. J Clin Oncol 1997;15:1853-1857. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9164195>.

90. Papamichael D, Gallagher CJ, Oliver RT, et al. Phase II study of paclitaxel in pretreated patients with locally advanced/metastatic cancer of the bladder and ureter. Br J Cancer 1997;75:606-607. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9052419>.

91. Vaughn DJ, Broome CM, Hussain M, et al. Phase II trial of weekly paclitaxel in patients with previously treated advanced urothelial cancer. J Clin Oncol 2002;20:937-940. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/11844814>.

92. Sweeney CJ, Roth BJ, Kabbinnavar FF, et al. Phase II study of pemetrexed for second-line treatment of transitional cell cancer of the urothelium. J Clin Oncol 2006;24:3451-3457. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16849761>.

93. Galsky MD, Iasonos A, Mironov S, et al. Phase II trial of dose-dense doxorubicin plus gemcitabine followed by paclitaxel plus carboplatin in patients with advanced urothelial carcinoma and impaired renal function. Cancer 2007;109:549-555. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17200962>.

94. Galsky MD, Mironov S, Iasonos A, et al. Phase II trial of pemetrexed as second-line therapy in patients with metastatic urothelial carcinoma. Invest New Drugs 2007;25:265-270. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/17146733>.

95. Witte RS, Elson P, Bono B, et al. Eastern Cooperative Oncology Group phase II trial of ifosfamide in the treatment of previously treated advanced urothelial carcinoma. J Clin Oncol 1997;15:589-593.

Available at: <http://www.ncbi.nlm.nih.gov/pubmed/9053481>.



National
Comprehensive
Cancer
Network®

NCCN Guidelines Version 2.2014 Bladder Cancer

[NCCN Guidelines Index](#)
[Bladder Cancer TOC](#)
[Discussion](#)

96. Han KS, Joung JY, Kim TS, et al. Methotrexate, vinblastine, doxorubicin and cisplatin combination regimen as salvage chemotherapy for patients with advanced or metastatic transitional cell carcinoma after failure of gemcitabine and cisplatin chemotherapy. *Br J Cancer* 2008;98:86-90. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18087289>.

97. Picozzi S, Ricci C, Gaeta M, et al. Upper urinary tract recurrence following radical cystectomy for bladder cancer: a meta-analysis on 13,185 patients. *J Urol* 2012;188:2046-2054. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/23083867>.

98. Audenet F, Yates DR, Cussenot O, Roupret M. The role of chemotherapy in the treatment of urothelial cell carcinoma of the upper urinary tract (UUT-UCC). *Urol Oncol* 2010. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/20884249>.

99. Dayyani F, Hoffman K, Eifel P, et al. Management of Advanced Primary Urethral Carcinomas. *BJU Int* 2014. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/24447439>.

100. Dalbagni G, Zhang ZF, Lacombe L, Herr HW. Male urethral carcinoma: analysis of treatment outcome. *Urology* 1999;53:1126-1132. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10367840>.

101. Grigsby PW. Carcinoma of the urethra in women. *Int J Radiat Oncol Biol Phys* 1998;41:535-541. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/9635699>.

102. Dimarco DS, Dimarco CS, Zincke H, et al. Surgical treatment for local control of female urethral carcinoma. *Urol Oncol* 2004;22:404-409. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/15464921>.

103. Dayyani F, Pettaway CA, Kamat AM, et al. Retrospective analysis of survival outcomes and the role of cisplatin-based chemotherapy in patients with urethral carcinomas referred to medical oncologists. *Urol Oncol* 2013;31:1171-1177. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/22534087>.

104. Cohen MS, Triaca V, Billmeyer B, et al. Coordinated chemoradiation therapy with genital preservation for the treatment of primary invasive carcinoma of the male urethra. *J Urol* 2008;179:536-541; discussion 541. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/18076921>.

105. Lorusso V, Manzione L, De Vita F, et al. Gemcitabine plus cisplatin for advanced transitional cell carcinoma of the urinary tract: a phase II multicenter trial. *J Urol* 2000;164:53-56. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/10840423>.

106. Roberts JT, von der Maase H, Sengelov L, et al. Long-term survival results of a randomized trial comparing gemcitabine/cisplatin and methotrexate/vinblastine/doxorubicin/cisplatin in patients with locally advanced and metastatic bladder cancer. *Ann Oncol* 2006;17 Suppl 5:v118-122. Available at:

<http://www.ncbi.nlm.nih.gov/pubmed/16807438>.