

Sexual size dimorphism in parasitoid wasps

BETHIA HURLBUTT

*Department of Biological Sciences, Purdue University,
West Lafayette, IN 47907, U.S.A.*

Received 19 May 1986, accepted for publication 23 September 1986

Sexual dimorphism in body length and proportion of overlap between the ranges of body length for males and females were estimated for 361 species of parasitoid wasps from 21 families. In most species, females are generally larger than males, though the range of male and female sizes overlap. Species in the family Ichneumonidae differ significantly from species in other families in three ways: (1) ichneumonids on average are larger, (2) in most species, females are generally smaller than males, and (3) on average, proportion overlap between the ranges of body length for males and females is greater. At present, there is a paucity of life history data on parasitoid wasp species for which size dimorphism is known. Thus it is not clear why ichneumonids differ from species in other families. Possible evolutionary explanations for variation in dimorphism among parasitoid wasp species are discussed.

KEY WORDS:—sexual dimorphism – sexual selection – parasitoid wasps – Ichneumonidae

CONTENTS

Introduction	63
Methods	64
Results	65
Dimorphism	65
Overlap	66
Discussion	67
Competition avoidance	67
Differential effect of size on reproductive success	67
Future study	69
Acknowledgements	70
References	70
Appendix	71

INTRODUCTION

Sexual size dimorphism has been reviewed for a range of vertebrate taxa. Interspecific differences in degree of dimorphism have been related to differences in mating systems (e.g. Alexander, Hoogland, Howard, Noonan & Sherman, 1979; Stamps, 1983; Carothers, 1984) body size (Clutton-Brock, Harvey & Rudder, 1977), and habitat type (Clutton-Brock *et al.*, 1977; Berry & Shine, 1980). However, little attention has been given to sexual size dimorphism in invertebrate groups except zooplankton (Gilbert, 1983) and spiders (Jocque, 1983).

In this paper, I quantitatively assess the degree of sexual size dimorphism in more than 360 species of parasitoid wasps from 21 families. I examine taxonomic patterns of dimorphism and discuss possible evolutionary causes of dimorphism. Parasitoid wasps are a large group within the order Hymenoptera. They consist of more than 100 000 identified species and probably numerous other as yet undiscovered ones (Hopper, 1984). A parasitoid wasp lays one or more eggs in or on a host. When the wasp eggs hatch, the emerging larvae feed on their hosts, killing the hosts by the time the wasp larvae pupate and emerge as adults. For parasitoid wasps as a group, females are reported to be larger than males (e.g. van den Assem, 1976). However, the extent to which this generalization holds across the various taxa of parasitoid wasps has not been examined.

Parasitoid wasps exhibit a variety of mating systems, and I intended to examine relationships between degree of size dimorphism and mating attributes. However, I was unable to do this as there are few species for which data are available both on mating system and on male and female size. I hope that this paper will serve as a source and stimulus for further study.

METHODS

Data on length, weight, and headwidth of adult male and female wasps were compiled from published records (specifically from *U.S. National Museum Proceedings*, 1921–1968; *Bulletin of Entomological Research*, 1960 to date; *Canadian Entomologist*, 1965 to date; *Annals of the Entomological Society of America*, 1972 to date; and *Entomologica Scandinavica*, 1973 to date; as well as from other publications). I also took measurements of length (five males, 12 females) and headwidth (289 males, 95 females) on *Hyposoter exiguae* (Ichneumonidae) from a laboratory colony. Species whose hosts are not known were included only if they belong to families or subfamilies composed primarily of parasitoid species. Species were classified taxonomically following Krombein, Hurd, Smith & Burks (1979).

Size measures are reported for males and females separately as ranges or as averages. The degree of dimorphism was estimated by dividing the midpoint size value for the female by that for the male or by dividing the average female size by the average male size. My analyses focus on ranges of lengths because this is how size is most often reported. I assume that all lengths were measured from the front of the head to the tip of the abdomen, excluding the ovipositor (as by P. DeBach, pers. comm.; C. M. Yoshimoto, pers. comm.; N. F. Johnson, pers. comm.; Samson, 1984).

The proportion of overlap between the range of length for males and females was also estimated for each species. It was calculated as the difference between the maximum size of the smaller sex and the minimum size of the larger sex, divided by the difference between the maximum size of the larger sex and the minimum size of the smaller sex. Overlap values greater than zero indicate that there is overlap between the male and the female size ranges. Values less than zero indicate no overlap, and values of zero indicate that the ranges abut.

Nonparametric statistics were used when variables were not normally distributed as determined from tests of skewness and kurtosis (Sokal & Rohlf, 1981).

RESULTS

Dimorphism

Sizes for males and for females, dimorphism, and proportion of overlap are presented in the Appendix. Length data are given in terms of the range for 361 species. The dimorphism values of these species vary from 0.64 to 1.77 (Fig. 1). In most species, females are larger than males. The mean dimorphism (female: male) is 1.12 ± 0.19 (standard deviations are used throughout).

One obvious exception to the general trend of female-biased size dimorphism is the subfamily Ichneumoninae in the family Ichneumonidae. In most species of Ichneumoninae, females are smaller than males. The average dimorphism is 0.91 ± 0.08 ($N = 81$). It should be noted that all the size data on Ichneumoninae are from Florida and neighboring states (Heinrich, 1977). Size data on species from other subfamilies of Ichneumonidae are few but are from Canada, the U.S.A. and the Virgin Islands. Although the average dimorphism of these species (1.06 ± 0.05 , $N = 5$) is greater than the average dimorphism for the subfamily Ichneumoninae (Mann-Whitney $U = 19.0$, $P = 0.001$), it is still less than the average dimorphism for species outside the family Ichneumonidae (1.19 ± 0.17 , $N = 275$; Mann-Whitney $U = 293.0$, $P = 0.028$).

Two other families besides Ichneumonidae which have average dimorphism values which are male-biased are Mymaridae and Scelionidae (Table 1). However, these values are based on only two and one species, respectively.

The lower (more male-biased) dimorphism of ichneumonids may somehow be related to their larger size. Within the family Ichneumonidae, the average midlength of males and females for species of Ichneumoninae (13.06 ± 4.48 mm) is larger than that for species of non-Ichneumoninae (7.07 ± 2.95 mm; Mann-Whitney $U = 36.0$, $P = 0.002$). The non-Ichneumoninae are in turn larger than species belonging to other families (2.54 ± 2.17 mm; Mann-Whitney $U = 92.0$, $P = 0.001$).

In addition to variation in dimorphism among families, subfamilies, and




Figure 1. Histogram of sexual dimorphism in length: □ = species in the subfamily Ichneumoninae, ■ = species not in the subfamily Ichneumoninae (includes species not in the family Ichneumonidae).

Table 1. The average (s.d. in parentheses) sexual dimorphism in length for 21 families

Family	Number of species	Dimorphism (female/male)
Ichneumonidae	86	0.92 (0.09)
Mymaridae	2	0.98 (0.07)
Scelionidae	1	0.99 (0.00)
Aphidiidae	2	1.05 (0.03)
Chrysididae	1	1.11 (0.00)
Sclerogibbidae	1	1.11 (0.00)
Bethylidae	7	1.12 (0.21)
Chalcididae	21	1.12 (0.13)
Eucoilidae	11	1.13 (0.08)
Braconidae	33	1.15 (0.17)
Trichogrammatidae	2	1.16 (0.02)
Pteromalidae	53	1.18 (0.15)
Platygastridae	2	1.19 (0.25)
Encyrtidae	70	1.20 (0.17)
Diapriidae	3	1.23 (0.12)
Eulophidae	49	1.25 (0.17)
Tiphiidae	8	1.25 (0.27)
Dryinidae	3	1.28 (0.20)
Torymidae	3	1.32 (0.29)
Eurytomidae	2	1.34 (0.34)
Eupelmidae	1	1.45 (0.00)

species, there is also intraspecific variation in dimorphism (see Appendix). This intraspecific variation is associated with differences in (1) character measured (e.g. *Muscidifurax* (Pteromalidae), *Pachycrepoideus vindemiae* (Pteromalidae), *Trichogramma minutum* (Trichogrammatidae)), (2) host species (e.g. *Invreia deceptor* (Chalcididae), *Dolchomitus* sp., *Pimpla turionellae* (Ichneumonidae), *Poropoea morimotoi* (Trichogrammatidae)), (3) host size (e.g. *H. exiguae* (Ichneumonidae), *Roptrocerus xylophagorum* (Torymidae)), (4) rearing temperature (Nealis, Jones & Wellington 1984), and (5) geographic location of the wasp population (e.g. *Muscidifurax* (Pteromalidae)).

Overlap

Proportion of overlap shows much more interspecific variation than dimorphism (coefficient of variation = 117% *vs.* 17%). It ranges from -0.64 to 1.00 (Appendix), with a mean of 0.30 ± 0.35 . The sexes overlap in length in 313 of the 361 species examined. Overlap is significantly greater for ichneumonids than for non-ichneumonids (0.38 ± 0.33 *vs.* 0.27 ± 0.35 ; $t = 2.58$, $P = 0.01$). This difference in overlap is related to the difference in dimorphism. Overlap will approach its maximum as dimorphism approaches 1.00. The average dimorphism of non-ichneumonid species (1.19) is about twice as far from a dimorphism of 1.00 as is that of ichneumonids (0.92). Overlap and average midlength of males and females are not correlated (Spearman rank coefficient = 0.05, $P = 0.16$).

DISCUSSION

Several hypotheses have been proposed to explain the evolution of sexual size dimorphisms in vertebrates (Alexander *et al.*, 1979; Carothers, 1984). The competition avoidance hypothesis states that dimorphism is selected for as a means of reducing competition between males and females of the same species by allowing differential use of resources, e.g. food (Selander, 1966). Alternatively, sexual size dimorphism may result from differences in maternal *vs.* paternal reproductive effort or from sexual selection either through mate competition or mate choice. These latter two hypotheses can be included under a more general hypothesis that sexual size dimorphism results from differences between males and females in the effect of size on reproductive success.

Competition avoidance

In parasitoid wasps, selection for differential food usage between the sexes probably does not contribute to sexual size dimorphism. Adults feed on nectar or honeydew. There do not seem to be sex-specific differences in nectar usage. At least for *Scambus buolianae*, males and females exhibit the same relative preferences for four different kinds of flowers (Leius, 1967). In some species, females, but not males, also feed on host body fluids. This trophic difference between the sexes is probably a result of nutritional requirements of females for egg production (e.g. *Bracon hebetor*, *B. serinopae* (Clark, 1963), *Muscidifurax zaraptor* (Whiting, 1967), *Spalangia cameroni* (Gerling & Legner, 1968)), rather than a means of reducing competition for food resources. Furthermore, this could not be a universal explanation for sexual size dimorphism because not all dimorphic species exhibit host feeding, e.g. *H. exiguae* (personal observation). Also, host feeding does not explain both male-biased and female-biased dimorphisms: no species are known in which males, but not females, host feed. In some species, males do not feed at all, e.g. *Melittobia chalybii* (Buckell, 1928) and thus do not compete with females for food.

Differential effect of size on reproductive success

Differences between males and females in the effect of size on reproductive success is a more likely explanation for sexual size dimorphisms in parasitoid wasps. For parasitoid wasps, these differences may result, not only from differences in maternal and paternal reproductive effort and from sexual selection, as has been suggested for vertebrates, but also from sib-mating and from the haplodiploid genetics of parasitoid wasps. The combined effects of all four of these factors will determine a species' degree of sexual size dimorphism.

Female-biased size dimorphisms are expected in species for which size has a more positive effect on the reproductive success of females than of males. There is usually a positive relationship between female body size and egg production (e.g. Iwata, 1966; Lawrence, 1981; Narasimham, 1984). However, there are no data for parasitoid wasps on how male size affects sperm production.

Large size may also be beneficial to females in obtaining hosts. (1) In species of *Trichogramma*, large females travel faster and further than small females (Boldt, 1974). Thus, large females may be better able to find hosts. (2) In species in which females must penetrate a hard covering to parasitize their hosts,

large females may be more capable of piercing the covering than smaller females (van den Assem, 1976). (3) In species which parasitize larval hosts or host eggs defended by the parents, large females may be less susceptible than small females to mortality related to host defence (e.g. biting). (4) In species in which the females fight among themselves for hosts, large females would be expected to have an advantage in fights, as in *Biosteres longicaudatus* (Lawrence, 1981). Thus, species exhibiting female-female combat would be expected to have more female-biased dimorphism values than species without female-female combat. At present, information on both size dimorphism and the occurrence of female-female combat is known for only a few species.

Sib-mating is an additional factor which may contribute to female-biased sexual size dimorphisms in parasitoid wasps. For those species for which information on mating is available, the usual pattern seems to be that mating among siblings is common but some outcrossing may still occur (Matthews, 1975; Gordh & Hawkins, 1981). When all mating is among siblings, females will be expected to produce only enough sons to fertilize their daughters (Hamilton, 1967). If a small male can produce enough sperm to fertilize all his sisters and if the cost of producing a son increases with its size, then females of sib-mating species should produce small males. In solitary species (species which produce one offspring per host, as opposed to gregarious species which produce multiple offspring per host), the cost of producing an offspring may increase with its size if a wasp's size is positively correlated with its host size (e.g. Jowyk & Smilowitz, 1978; Nealis *et al.*, 1984; Samson, 1984), and large hosts are scarce and/or parasitizing large hosts is difficult or dangerous. In gregarious species the cost of producing an offspring also may increase with the offspring's size because of larval competition. The larger the son, the more host resources he consumes, and the less resources are available for siblings developing in the same host.

Sib-mating is probably more common in gregarious species than in solitary species. For example, in some gregarious species, mating occurs within the host, and males may never even leave the host (e.g. Suzuki & Hiehata, 1985). Thus one would predict that, all else being equal, gregarious species would have more female-biased dimorphism values than solitary species.

The haplodiploid genetic system of parasitoid wasps provides another explanation for female-biased dimorphisms in gregarious species (Pickering, 1980). This genetic system results in asymmetries in relatedness among family members: a male is more closely related to his sister than she is to him, and sisters are more closely related than are brothers. Based on these asymmetries, inclusive fitness theory predicts that selection will result in females which are more selfish to their brothers than males are to their sisters. Thus, female larvae may be more aggressive than their brothers in competing for host resources which are needed to reach maximum size.

Another factor which may affect sexual size dimorphism is sexual selection through mate competition. Competition for mates seems to be limited to males in parasitoid wasps. Males will mate repeatedly throughout their lives, whereas females generally mate only once (Matthews, 1975; Gordh & DeBach, 1978) or repeatedly only for a brief period shortly after emergence (e.g. Sary, 1970).

Sexual selection will result in a female-biased size dimorphism when small size is indirectly beneficial to male reproductive success through selection for early emergence. Both sexes tend to mate at or near their emergence sites (Matthews,

1975). Males frequently emerge before females. By emerging early, a male may increase the number of virgin females with which he can potentially mate. This will be especially true for species in which males defend emergence sites and in which there is a prior ownership advantage (as in *Asolcus basilis*; Wilson, 1961). If male and female larvae grow at the same rate, then to emerge first, males must emerge smaller. This explanation for female-biased size dimorphisms differs from the other explanations I have discussed in that it suggests a benefit to small male size. Some evidence that small size and/or short development time has a positive effect on male reproductive success comes from the solitary species *H. exiguae*. In *H. exiguae*, small male size is not simply a result of mothers ovipositing sons in small hosts, and sons thus not having sufficient food to grow as large as females. Males are smaller and/or emerge earlier than females at a given size host (Jowyk & Smilowitz, 1978).

Alternatively, sexual selection through mate competition may cause a more male-biased size dimorphism. In some species, of parasitoid wasps, males fight for ownership of female emergence sites. Assuming a positive correlation between size and fighting ability, species with male-male aggression would be expected to exhibit a lower degree of size dimorphism than species without male-male aggression. Male-male combat could be the cause of the male-biased dimorphism found in most species of Ichneumoninae; however, the prevalence of male-male combat in this subfamily is not known.

Another aspect of sexual size dimorphism, besides interspecific variation, which deserves attention is intraspecific variation. Variation in the extent of dimorphism in different characters may result from how selection pressures influence different body parts. For example, selection for increased egg production may result in greater dimorphism in abdominal size than in headwidth. Geographic variation in dimorphism may result from different environmental pressures. For example, geographic variation in host distribution may result in interpopulation variation in the occurrence of female-female combat. When hosts are abundant, it may not be worthwhile for females to defend them. Populations with female-female combat are expected to have greater female bias in size dimorphism than populations without female-female combat. For the two species in which dimorphism is known to change with host size (see Results), dimorphism is greater in larger hosts. Perhaps food limitation constrains dimorphism in small hosts by preventing the larger sex from reaching its maximum size. This explanation for intraspecific variation in sexual size dimorphism may also help to explain some of the interspecific differences in sexual size dimorphism. Do species which parasitize smaller hosts exhibit less dimorphism than those parasitizing larger hosts?

FUTURE STUDY

I have discussed some factors which potentially may contribute to sexual size dimorphism in parasitoid wasps. As more data becomes available, a better understanding of the forces acting on the size of males relative to females may be obtained by applying an inter- and an intraspecific approach. The interspecific approach involves comparing the behaviour, physiology and ecology of closely related species which exhibit differences in degree and/or direction of size dimorphism, e.g. *Pteromalinae* species (van den Assem, 1976). As discussed

previously, do species with male-male aggression exhibit more male-biased size dimorphisms than species without male-male aggression? Do species with female-female aggression exhibit more female-biased size dimorphism than species without female-female aggression? How do species of Ichneumoninae, which tend to have male-biased dimorphisms, differ from other species of parasitoid wasps? The intraspecific approach involves examining, for individual species, the effect of size on reproductive success of males relative to females. To date this has been done for only one species (*Heterospilis prosopidis*; Charnov, Losden Hartogh, Jones & Assem, 1981; Jones, 1982). Size had a greater effect on the reproductive success of females than of males. Thus one would predict a female-biased size dimorphism. Indeed females generally are larger than males (W. T. Jones, pers. comm.).

ACKNOWLEDGEMENTS

I thank R. D. Howard for encouraging me to write this paper. I thank R. D. Howard, R. B. King, and anonymous reviewers for their comments and suggestions. Support during preparation of this manuscript was provided by a National Science Foundation Graduate Fellowship.

REFERENCES

- ALEXANDER, R. D., HOOGLAND, J. L., HOWARD, R. D., NOONAN, K. M. & SHERMAN, P. W., 1979. Sexual dimorphisms and breeding systems in pinnipeds, ungulates, primates, and humans. In A. Chagnon & W. Irons (Eds), *Evolutionary Biology and Human Social behavior*: 402-435. North Scituate, Massachusetts: Duxbury Press.
- ASSEM, J. VAN DEN, 1976. Male courtship behaviour, female receptivity signal, and size differences between the sexes in Pteromalinae (Hym., Chalcidoidea, Pteromalidae), and comparative notes on other chalcidoids. *Netherlands Journal of Zoology*, 26: 535-548.
- BERRY, J. F. & SHINE, R., 1980. Sexual size dimorphism and sexual selection in turtles (Order Testudines). *Oecologia*, 44: 185-191.
- BOLDT, P. E., 1974. Temperature, humidity, and host: effect on rate of search of *Trichogramma evanescens* and *T. minutum* Auctt. (not Riley, 1971). *Annals of the Entomological Society of America*, 67: 706-708.
- BUCKELL, E. R., 1928. Notes on the life history and habits of *Melittobia chalybii* Ashmead (Chalcidoidea: Elachertidae). *Pan-Pacific Entomologist*, 5: 14-22.
- CAROTHERS, J. H., 1984. Sexual selection and sexual dimorphism in some herbivorous lizards. *American Naturalist*, 124: 244-254.
- CHARNOV, E. L., LOS-DEN HARTOGH, R. L., JONES, W. T. & ASSEM, J. VAN DEN, 1981. Sex ratio evolution in a variable environment. *Nature*, 289: 27-33.
- CLARK, A. M., 1963. The influence of diet upon the adult life span of two species of *Bracon*. *Annals of the Entomological Society of America*, 56: 616-619.
- CLUTTON-BROCK, T. H., HARVEY, P. H. & RUDDER, B., 1977. Sexual dimorphism, sociometric sex ratio and body weight in primates. *Nature*, 269: 797-800.
- GERLING, D. & LEGNER, E. F., 1968. Developmental history and reproduction of *Spalangia cameroni*, parasite of synanthropic flies. *Annals of the Entomological Society of America*, 61: 1436-1443.
- GILBERT, J. J., 1983. Sexual dimorphism in zooplankton (copepoda, cladocera, and rotifera). *Annual Review of Ecology and Systematics*, 14: 1-33.
- GORDH, G. & DEBACH, P., 1978. Courtship behavior in the *Aphytis lingnanensis* group, its potential usefulness in taxonomy, and a review of sexual behavior in the parasitic Hymenoptera (Chalcidoidea: Aphelinidae). *Hilgardia*, 46: 37-75.
- GORDH, G. & HAWKINS, B., 1981. *Goniozus emigratus* (Rohwer), a primary external parasite of *Paramyelois transitella* (Walker), and comments on bethylids attacking Lepidoptera (Hymenoptera: Bethyidae; Lepidoptera: Pyralidae). *Journal of Kansas Entomological Society*, 54: 787-803.
- HAMILTON, W. D., 1967. Extraordinary sex ratios. *Science*, 156: 477-488.
- HEINRICH, G. H., 1977. Ichneumoninae of Florida and neighboring states. In *Arthropods of Florida and Neighboring Land Areas*, 9. Gainesville, Florida: Florida Department of Agriculture and Consumer Services.
- HOPPER, K. R., 1984. The effects of host-finding and colonization rates on abundances of parasitoids of a gall midge. *Ecology*, 65: 20-27.

- IWATA, K., 1966. The comparative anatomy of the ovary in hymenoptera. Supplement on Ichneumonidae, *Coccygomimus luctuosa* Smith, *C. parnaae* Viereck and *C. pluto* Ashmead. *Acta Hymenopterologica*, 2: 133–135.
- JOCQUE, R., 1983. A mechanism explaining sexual size dimorphism in spiders. *Biologisch Jaarboek Dodonaea*, 51: 104–115.
- JONES, W. T., 1982. Sex ratio and host size in a parasitoid wasp. *Behavioral Ecology and Sociobiology*, 10: 207–210.
- JOWYK, E. A. & SMILOWITZ, Z., 1978. A comparison of growth and developmental rates of the parasite *Hyposoter exiguae* reared from two instars of its host, *Trichoplusia ni*. *Annals of the Entomological Society of America*, 71: 467–472.
- KROMBEIN, K. V., HURD, P. D., SMITH, D. R. & BURKS, B. D., 1979. *Catalog of Hymenoptera in America North of Mexico*. Washington D. C.: Smithsonian Institution Press.
- LAWRENCE, P. O., 1981. Interference competition and optimal host selection in the parasitic wasp, *Biosteres longicaudatus*. *Annals of the Entomological Society of America*, 74: 540–544.
- LEIUS, K., 1967. Food sources and preferences of adults of a parasite, *Scambus buolianae* (Hym.: Ichn.), and their consequences. *Canadian Entomologist*, 99: 865–871.
- MATTHEWS, R. W., 1975. Courtship in parasitic wasps. In P. W. Price (Ed.), *Evolutionary Strategies of Parasitic Insects and Mites*: 66–86. New York: Plenum Press.
- NARASIMHAM, A. U., 1984. Comparative studies on *Tetrastichus hagenowii* (Ratzeburg) and *T. asthenognmus* (Waterston), two primary parasites of cockroach oothecae, and on their hyperparasite *Tetrastichus* sp. (*T. miser* (Nees) group) (Hymenoptera: Eulophidae). *Bulletin of Entomological Research*, 74: 175–189.
- NEALIS, V. G., JONES, R. E. & WELLINGTON, W. G., 1984. Temperature and development in host–parasite relationships. *Oecologia*, 61: 224–229.
- PICKERING, J., 1980. Larval competition and brood sex ratios in the gregarious parasitoid *Pachysomoides stupidus*. *Nature*, 283: 291–292.
- SAMSON, P. R., 1984. The biology of *Raptrocercus xylophagorum* (Hym.: Torymidae), with a note on its taxonomic status. *Entomophaga*, 29: 287–298.
- SELANDER, R. K., 1966. Sexual dimorphism and differential niche utilization in birds. *Condor*, 68: 113–151.
- SOKAL, R. R. & ROHLF, F. J., 1981. *Biometry*, 2. San Francisco: Freeman.
- STAMPS, J. A., 1983. Sexual selection, sexual dimorphism and territoriality in lizards. In R. B. Huey, E. R. Pianka & T. W. Schoener (Eds), *Lizard Ecology: Studies on a Model Organism*: 169–204. Cambridge, Massachusetts: Harvard University Press.
- STARY, P., 1970. *Biology of Aphid Parasites with respect to Integrated Control*. The Hague: Junk Press.
- SUZUKI, Y. & HIEHATA, K., 1985. Mating systems and sex ratios in the egg parasitoids, *Trichogramma dendrolimi* and *T. papilionis* (Hymenoptera: Trichogrammatidae). *Animal Behaviour*, 33: 1223–1227.
- WHITING, A. R., 1967. The biology of the parasitic wasp *Mormoniella vitripennis* (= *Nasonia brevicornis*) (Walker). *Quarterly Review of Biology*, 42: 333–406.
- WILSON, F., 1961. Adult reproductive behaviour in *Asolcus basalis* (Hymenoptera: Scelionidae). *Australian Journal of Zoology*, 9: 739–751.

APPENDIX

Sexual size dimorphism and proportion of overlap of size ranges (see text for definitions)

Wasp	Measure*	Dimorphism (female/male)	Size range or average		
			male	female	Overlap
APHIDIIDAE					
<i>Binodoxys nearctaphidis</i> (Mackauer, 1965)	Length	1.03	1.8–2.0	1.8–2.1	0.67
<i>Praon callaphis</i> (Mackauer <i>et al.</i> 1982)	Length	1.06	1.3–1.8	1.3–2.0	0.71
BETHYLIDAE					
<i>Anisepyrus aeneus</i> (= <i>Rhabdepyris bridwelli</i>) (Evans, 1959a)	Length	1.14	4–5	5.0–5.3	0.00
<i>A. columbianus</i>	Length	1.25	3–5	4–6	0.33
<i>A. occidentalis</i>	Length	1.20	3.5–5.0	4.0–6.2	0.37
<i>A. subviolaceus</i>	Length	1.19	3.3–4.8	4.0–5.6	0.35
<i>A. williamsi</i> (Evans, 1959b)	Length	1.19	3–5	4–5.5	0.40

*Length and headwidth in millimetres. Weight in milligrams.

B. HURLBUTT
Appendix (*cont.*)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>Dissonophalus xanthopus</i> (Evans, 1954)	Length	0.64	1.7–3.3	1.2–2.0	0.14
<i>Goniozus aethiops</i> (Gordh & Evans, 1976)	Length	1.23	2.0–2.4	2.4–3.0	0.00
BRACONIDAE					
<i>Acrophasmus ferrugineus</i>	Length	1.46	2.4–5.5	4.5–7	0.22
<i>A. immigrans</i> (= <i>A. lycti</i>) (Marsh, 1968a)	Length	1.30	2–3	2.5–4	0.25
<i>Agathirsia gibbosa</i> (= <i>Agathis gibbosa</i>) (Odebiyi & Oatman, 1972)	Length: range average	0.96 0.97	3.4–4.5 3.9	3.1–4.5 3.8	0.79
<i>Alysia cariosa</i> (Marsh, 1968b)	Length	0.79	5.5–9	4–7.5	0.40
<i>Apanteles absonus</i>	Length	1.06	2.3–2.8	2.5–2.9	0.50
<i>A. fumiferanae</i>	Length	0.96	2.0–2.9	2.0–2.7	0.78
<i>A. milleri</i>	Length	1.07	2.7–2.8	2.8–3.1	0.00
<i>A. morrissi</i>	Length	1.08	2.5–2.8	2.8–2.9	0.00
<i>A. petrovae</i> (Mason, 1974)	Length	1.29	2.2–2.3	2.8–3.0	–0.63
<i>A. dignus</i> (Cardona & Oatman, 1971)	Length average	1.04	2.14	2.23	
<i>A. glomeratus</i>		>			
<i>A. rubecula</i> (Nealis <i>et al.</i> , 1984)		>			
<i>Asobara citri</i>		=			
<i>A. tabida</i> (Bouletreau & David, 1981)		=			
<i>Biosteres longicaudatus</i> (Lawrence <i>et al.</i> , 1976)	Length	1.42	2.5–2.8	3.5–4.0	–0.47
<i>Bracon caulicola</i> (= <i>Microbracon caulicola</i>) (Gahan, 1922)		>			
<i>B. compressitarsis</i> (Wharton, 1983)	Length	1.42	2.0–2.8	3.0–3.8	–0.11
<i>B. hebetor</i> (Benson, 1973)	Headwidth	>			
<i>B. rhyacioniae</i> (Goyer & Schenk, 1970)	Length: average	1.53	3.0	4.6	
<i>Calaphidius elegans</i> (Gardenfors, 1983)	Length	1.08	1.6–2.2	1.7–2.4	0.63
<i>Cantharoctonus canadensis</i> (Mason, 1968b)	Length	1.00	2.2–3.0	2.4–2.8	0.50
<i>Coeloides crocator</i>	Length	1.29	3.5–6.0	3.8–8.5	0.44
<i>C. rossicus betulae</i>	Length	1.17	4.5–7.0	4.5–9	0.56
<i>C. scolytivorus</i>	Length	1.00	2.5–3.5	2.5–3.5	1.00
<i>C. symptiys</i>	Length	1.14	3.0–7.5	3.3–8.7	0.74
<i>C. tsugatorus</i> (Mason, 1978)	Length	0.99	3.5–8.5	3.4–8.5	0.98
<i>Colastes polypori</i> (Mason, 1968b)	Length	1.06	1.7–3.5	2.0–3.5	0.83
<i>Diospilus fomitis</i> (Mason, 1968b)	Length	1.11	2.8–3.5	3.0–4.0	0.42
<i>Glyptocolastes caryae</i> (Marsh, 1968a)	Length	1.31	2.5–4.0	2.5–6	0.43

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>Heterospilus annulatus</i> (Marsh, 1982)	Length	1.20	2-3	2.5-3.5	0.33
<i>Hormius tsugae</i> (Mason, 1968b)	Length	1.03	1.7-2.2	1.5-2.5	0.50
<i>Illidops terrestris</i> (Wharton, 1983)	Length	1.12	1.8-2.4	2.2-2.5	0.29
<i>Leluthia floridensis</i>	Length	1.29	3-4	4-5	0.00
<i>L. mexicana</i> (Marsh, 1967)	Length	1.29	2.5-4.5	3-6	0.43
<i>Macrocentrus ancyliivorus</i> (Finney <i>et al.</i> , 1947)		>			
<i>Meteorus betulini</i> (Mason, 1968b)	Length	1.00	3.5-4.5	3.5-4.5	1.00
<i>Myosoma durango</i>	Length	1.55	4.5-5.5	7.0-8.5	-0.38
<i>M. eumystax</i> (Mason, 1978)	Length	1.24	4.5-7.5	6.0-8.9	0.34
<i>Opius bellus</i> (Gahan, 1930)	Length	1.07	3.5-4	3.5-4.5	0.50
<i>Perilitus areolaris</i> (Gerdin & Hedqvist, 1984)	Length	1.05	2.8-3.5	2.9-3.7	0.67
<i>Rhacontus</i> (Marsh, 1976)		>			
<i>Rogas hyphantriae</i> (Gahan, 1922)		=			
<i>Stenocorse bruchiivora</i> (Marsh, 1968a)	Length	1.07	2.5-4.5	3-4.5	0.75
<i>Zelee daurica</i> (Watanabe, 1969)	Length	1.13	7-8	8-9	0.00
CHALCIDIDAE					
<i>Brachymeria intermedia</i> (Minot & Leonard, 1976)	Weight	>			
<i>Ceratostigma debilis</i>	Length	1.00	4-4.5	4-4.5	1.00
<i>C. meteori</i> (Burks, 1940)	Length	1.27	3.5-4	4.5-5	-0.33
<i>Chalcis barbara</i> (Burks, 1940)	Length	1.04	5-7	5-7.5	0.80
<i>C. barbara</i>	Length	1.08	5.0-7.0	5.0-8.0	0.67
<i>C. divisa</i>	Length	1.03	6.5-8.0	7.0-8.0	0.67
<i>C. neptis</i> (Burks, 1977)	Length	0.96	5.5-6.0	5.0-6.0	0.50
<i>C. canadensis</i>	Length	1.11	4.5-5	5-5.5	0.00
<i>C. flebilis</i>	Length	1.11	4-5.5	4.5-6	0.50
<i>C. microgaster</i> (Burks, 1940)	Length	1.13	3.5-4.5	4-5	0.33
<i>Inureia deceptor</i> (Grissel & Schauf, 1981)					
from host:					
<i>Elasmopalpus lignosellus</i>	Length	1.10	2.3-3.8	2.9-3.8	0.60
<i>Stegasta bosqueella</i>	Length	0.86	2.3-2.7	2.0-2.3	0.00
combined	Length	0.95	2.3-3.8	2.0-3.8	0.83
<i>Metadontia amoena</i> (Burks, 1940)	Length	1.29	4.5-6	6-7.5	0.00
<i>Spilochalcis arcana</i>	Length	0.95	4.5-5	4-5	0.50
<i>S. dorsata</i>	Length	1.16	4.5-5	5-6	0.00
<i>S. hirtifemora</i>	Length	1.27	2.5-3	3-4	0.00
<i>S. igneoides</i>	Length	1.24	4.5-6	4.5-8.5	0.38

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; >denotes female size > male size; <denotes female size < male size.

Appendix (*cont.*)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>S. juxta</i>	Length	1.12	4-4.5	4.5-5	0.00
<i>S. mariae</i>	Length	1.32	4-7	4.5-10	0.42
<i>S. nigricornis</i>	Length	1.07	5.5-8.5	6-9	0.71
<i>S. phais</i>	Length	1.28	4-5	5-6.5	0.00
<i>S. phoenica</i>	Length	1.29	5-5.5	6.5-7	-0.50
<i>S. subobsoleta</i>	Length	0.85	4.5-5.5	4-4.5	0.00
<i>S. transitiva</i> (Burks, 1940)	Length	1.10	4-6.5	5-6.5	0.60
CHRYSIDIDAE					
<i>Muesebeckidium occidentale</i> (Krombein, 1969)	Length	1.11	4-5	4.5-5.5	0.33
DIAPRIIDAE					
<i>Trichopria fumipennis</i>	Length	1.18	2.4-2.6	2.8-3.1	-0.29
<i>T. stratiomyiae</i> (Huggert, 1982)	Length	1.14	1.6-2.0	2.0-2.1	0.00
<i>T. popei</i> (= <i>Phaenopria popei</i>) (Knutson & Berg, 1963)	Length	1.37	0.84-1.20	1.20-1.59	0.00
<i>Trichopria</i> sp. (Bouletreau & David, 1981)	Length	=			
DRYINIDAE					
<i>Neogonatopus agropyrus</i>	Length	1.11	2-2.25	2-3	0.50
<i>N. ombrodes</i> (Freytag, 1977)	Length	1.22	2-2.5	2.5-3	0.00
<i>Pachygonatopus minimus</i> (Barrett <i>et al.</i> , 1965)	Length	1.50	1.8-2.0	2.7-3.0	-0.58
ENCYRTIDAE					
<i>Adektitopus gordhi</i> (Noyes & Hayat, 1984)	Length	1.47	0.87-1.03	1.24-1.56	-0.30
<i>Aleurodipphilus americanus</i> (DeBach & Rose, 1981)	Length	1.16	0.65-0.81	0.74-0.96	0.23
<i>Aphytis acrenulatus</i>	Length	1.56	0.62-0.74	0.86-1.25	-0.19
<i>A. africanus</i>	Length	1.16	0.49-0.80	0.61-0.89	0.48
<i>A. capensis</i>	Length	1.27	0.80-1.19	1.22-1.30	-0.06
<i>A. capillatus</i>	Length	1.01	0.64-0.97	0.60-1.03	0.77
<i>A. cercinus</i>	Length	1.09	0.79-1.19	0.93-1.22	0.60
<i>A. chilensis</i>	Length	1.07	0.54-0.93	0.50-1.08	0.67
<i>A. chrysomphali</i>	Length	1.08	0.56-0.86	0.55-0.99	0.68
<i>A. cochereaui</i>	Length	1.34	0.59-1.08	0.89-1.35	0.25
<i>A. coheni</i>	Length	1.26	0.66-0.90	0.86-1.11	0.09
<i>A. columbi</i>	Length	1.21	0.86-0.95	0.96-1.23	-0.03
<i>A. confusus</i>	Length	1.26	0.75-0.85	0.94-1.07	-0.28
<i>A. costalimai</i>	Length	1.29	0.68-0.86	0.81-1.18	0.10
<i>A. cylindratus</i>	Length	1.40	0.66-0.85	0.76-1.36	0.13
<i>A. debachi</i>		>			
<i>A. diaspidis</i>	Length	1.09	0.61-1.11	0.61-1.26	0.77
<i>A. equatorialis</i>	Length	1.12	0.60-0.74	0.70-0.80	0.20
<i>A. fabresi</i>	Length	1.27	0.59-0.92	0.85-1.07	0.15
<i>A. fisheri</i>	Length	1.31	0.61-0.86	0.68-1.24	0.29
<i>A. funicularis</i>	Length	1.17	0.61-0.93	0.71-1.09	0.46
<i>A. gordoni</i>	Length	1.12	0.55-0.77	0.65-0.83	0.43
<i>A. griseus</i>	Length	1.05	0.72-1.03	0.67-1.17	0.62
<i>A. hispanicus</i>	Length	1.05	0.62-1.02	0.73-1.00	0.67
<i>A. holoxanthus</i>	Length	1.29	0.71-0.92	0.92-1.18	0.00
<i>A. ignotus</i>	Length	1.46	0.62-1.22	1.23-1.45	-0.01

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (*cont.*)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>A. immaculatus</i>	Length	1.06	0.73–1.04	0.84–1.03	0.61
<i>A. japonicus</i>	Length	1.10	0.64–0.83	0.73–0.89	0.40
<i>A. lepidosaphes</i>	Length	1.26	0.70–0.93	0.90–1.15	0.07
<i>A. lingnanensis</i>	Length	1.08	0.69–0.96	0.73–1.05	0.64
<i>A. longicaudus</i>	Length	1.77	0.77–0.96	1.45–1.61	–0.58
<i>A. luteus</i>	Length	1.10	0.91–1.25	0.87–1.50	0.54
<i>A. maculatipennis</i>	Length	1.43	0.57–0.72	0.88–0.96	–0.41
<i>A. margaretae</i>	Length	1.14	0.69–0.97	0.78–1.11	0.45
<i>A. mazalae</i>	Length	1.12	0.65–0.79	0.79–0.82	0.00
<i>A. melanostictus</i>	Length	1.10	0.55–1.10	0.67–1.14	0.73
<i>A. melinus</i>	Length	1.17	0.73–1.01	0.83–1.21	0.38
<i>A. merceti</i>	Length	1.41	0.76–1.16	1.02–1.68	0.15
<i>A. mimosae</i>	Length	1.06	0.74–0.83	0.81–0.85	0.18
<i>A. mytilaspidis</i>	Length	0.83	1.14–1.27	0.75–1.26	0.23
<i>A. notialis</i>	Length	1.29	0.66–0.91	0.81–1.22	0.18
<i>A. obscurus</i>	Length	1.06	0.73–1.03	0.76–1.11	0.71
<i>A. opuntiae</i>	Length	1.04	0.84–0.98	0.70–1.20	0.28
<i>A. philippinensis</i>	Length	1.28	0.76–0.90	0.94–1.18	–0.10
<i>A. pilosus</i>	Length	1.06	0.73–0.80	0.65–0.97	0.22
<i>A. pinnaspidis</i>	Length	1.22	0.54–0.80	0.63–1.00	0.37
<i>A. proclia</i>	Length	1.07	0.82–1.06	0.82–1.20	0.63
<i>A. roseni</i>	Length	1.51	0.68–1.16	1.18–1.59	–0.02
<i>A. salvadorensis</i>	Length	1.26	0.75–0.92	0.98–1.12	–0.16
<i>A. sensorius</i>	Length	1.15	0.43–0.67	0.52–0.74	0.48
<i>A. setosus</i>	Length	1.18	0.58–0.73	0.62–0.92	0.32
<i>A. taylora</i>	Length	1.30	0.62–0.84	0.75–1.15	0.17
<i>A. theae</i>	Length	0.98	0.55–0.78	0.53–0.77	0.88
<i>A. tucumani</i>	Length	1.11	0.85–1.03	0.95–1.14	0.28
<i>A. ulianovi</i>	Length	1.19	0.41–0.87	0.50–1.02	0.61
<i>A. yasumatsui</i> (Rosen & De Bach, 1979)	Length	1.21	0.62–1.02	0.81–1.18	0.38
<i>Chrysoplatycerus ferrisi</i> (Timberlake, 1922)	Length	1.67	0.84–1.18	1.51–1.87	–0.32
<i>Copidosoma innocuellae</i> (Barron, 1970)	Length	1.16	1.0–1.5	1.3–1.6	0.33
<i>Cremesina aquilonaris</i> (Noyes & Hayat, 1984)	Length	1.52	0.67–1.05	1.02–1.59	0.03
<i>Encarsia acaudaleyrodia</i> (Hayat, 1976)	Length	1.13	0.45–0.52	0.47–0.63	0.28
<i>E. pergandiella</i> (Girault, 1908)	Length: range	1.21	0.500–0.580	0.620–0.690	–0.21
	average	1.20	0.546	0.6545	
(Gerling, 1966a)	range	1.08	0.370–0.700	0.485–0.667	0.55
	average	1.03	0.529	0.544	
<i>Eretmocerus californicus</i> (Gerling, 1966b)	Length: range	1.09	0.565–0.825	0.720–0.795	0.29
	average	1.09	0.704	0.767	
<i>E. rajasthanicus</i> (Hayat, 1976)	Length	1.00	0.5–0.62	0.51–0.61	0.83
<i>Hengata spinosa</i> (Noyes & Hayat, 1984)	Length	1.01	0.71–0.94	0.71–0.95	0.96
<i>Hexacladia hilaris</i> (Burks, 1972)	Length	1.10	2.25–2.75	2.5–3.0	0.33
<i>Kataka mudigerensis</i> (Noyes & Hayat, 1984)	Length	1.22	0.89–0.92	0.98–1.22	–0.18
<i>Parapyrus manihoti</i> (Noyes, 1984)	Length	1.13	0.52–0.97	0.57–1.11	0.68

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>Protaphelinus nikolskajae</i> (Mackauer, 1972)	Length	1.09	1.03–1.14	0.87–1.50	0.17
<i>Stemmatosteres kuchari</i> (Yoshimoto, 1972)	Length	1.07	0.65–0.70	0.65–0.8	0.33
<i>Tongyus nesus</i> (Noyes & Hayat, 1984)	Length	1.27	0.96–1.30	1.11–1.75	0.24
<i>Zeteticontus utilis</i> (Noyes, 1982)	Length	1.15	1.39–2.08	1.58–2.40	0.50
EUCOILIDAE					
<i>Ganaspidium pusillae</i> (Weld, 1955)	Length: range average	0.98 1.02	0.95–1.05 0.98	0.8–1.15 1.0	0.29
<i>Leptopilina boulardi</i>	Length	1.15	1.0–1.6	1.2–1.8	0.50
<i>L. clavigipes</i>	Length	1.07	1.2–1.5	1.2–1.7	0.60
<i>L. fimbriata</i>	Length	1.04	1.2–1.4	1.2–1.5	0.67
<i>L. heterotoma</i>	Length	1.16	1.3–1.8	1.5–2.1	0.37
<i>L. longipes</i>	Length	1.22	1.1–1.6	1.5–1.8	0.14
<i>L. victoriae</i> (Nordlander, 1980)	Length	1.13	1.4–1.6	1.5–1.9	0.20
<i>Rhoptromeris heptoma</i>	Length: range average	1.24 1.15	1.0–1.5 1.3	1.2–1.9 1.5	0.33
<i>R. nigriventris</i>	range average	1.16	1.1–1.4 1.2	1.3–1.6 1.4	0.20
<i>R. villosa</i> (Nordlander, 1978)	Length	1.12	1.0–1.4	1.3–1.4	0.25
<i>Zamischus elongatus</i> (Yoshimoto, 1971)	Length	1.10	5.0–5.5	5.5–6.0	0.00
EULOPHIDAE					
<i>Astichus notus</i> (Yoshimoto, 1970a)	Length	1.27	1.1–1.5	1.5–1.8	0.00
<i>Chrysocharis caribea</i> (Boucek, 1977b)	Length	1.20	0.9–1.4	1.3–1.45	0.18
<i>Colpixys gigas</i> (Boucek, 1972)	Length	1.73	3.5–3.9	6–6.8	–0.62
<i>Dicladocerus nearcticus</i>	Length	1.45	1.8–2.0	2.5–3.0	–0.42
<i>D. pacificus</i>	Length	1.42	1.2–1.4	1.5–2.2	–0.10
<i>D. westwoodi</i> (Yoshimoto, 1976b)	Length	1.26	2.0–2.2	2.5–2.8	–0.37
<i>Dicladocerus</i> (= <i>Diglyphus</i>) (Gordh & Hendrickson, 1979)		>			
<i>Edovum putleri</i> (Grissell, 1981)	Length	1.07	1.2–1.5	1.3–1.6	0.50
<i>Elachertus lasiodermæ</i> (Hedqvist, 1977a)	Length	1.29	1.3–1.8	1.8–2.2	0.00
<i>Encarsia versicolor</i> (Girault, 1908)	Length: range average	1.21 1.20	0.500–0.580 0.546	0.620–0.690 0.6545	–0.21
<i>Horismenus bruchophagus</i>	Length	1.20	1.2–1.8	1.6–2.0	0.25
<i>H. carolinensis</i>	Length	1.14	1.5–2.0	1.8–2.2	0.29
<i>H. latrodecti</i> (Burks, 1971)	Length	1.65	0.7–1.3	1.5–1.8	–0.18
<i>Mesocharis nearctica</i> (Yoshimoto, 1976a)	Length	1.20	1.5–2.0	2.0–2.2	0.00

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (*cont.*)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>Myrmokata diparoides</i> (Boucek, 1972)	Length	1.06	1.5–1.8	1.6–1.9	0.50
<i>Pediobius balyanae</i>	Length	1.35	0.8–0.9	1.1–1.2	–0.50
<i>P. derroni</i>	Length	1.15	1.3–1.4	1.5–1.6	–0.33
<i>P. madas</i>	Length	1.28	1.1–1.8	1.6–2.1	0.20
<i>P. multisetis</i> (Boucek, 1976)	Length	1.53	1.05–1.3	1.5–2.1	–0.19
<i>P. lonchaeae</i>	Length	1.12	1.0–1.6	1.1–1.8	0.63
<i>P. testaceipes</i> (Burks, 1966)	Length	1.30	0.9–1.1	1.2–1.4	–0.20
<i>Psigalio ternatus</i> (Askew, 1984)	Length	1.40	2.0–2.5	2.8–3.5	–0.20
<i>Syntomosphyrum blattae</i>	Length	1.05	1.0–1.2	0.8–1.5	0.29
<i>S. esurus</i> (Burks, 1952)	Length	1.27	1.0–1.2	1.2–1.6	0.00
<i>Tachinobia repanda</i> (Boucek, 1977a)	Length	0.86	1.15–1.30	0.7–1.4	0.21
<i>Tetrastichus ainsliei</i>	Length	1.17	1.1–1.2	1.2–1.5	0.00
<i>T. baldufi</i>	Length	1.09	1.0–1.2	1.0–1.4	0.50
<i>T. cornutus</i>	Length	1.32	1.5–1.9	2.0–2.5	–0.10
<i>T. crino</i> (= <i>T. oecanthivorus</i>)	Length	1.08	1.2–1.3	1.3–1.4	0.00
<i>T. gigas</i>	Length	1.40	2.0–3.0	3.0–4.0	0.00
<i>T. hagenowii</i>	Length	1.03	1.2–1.9	1.2–2.0	0.87
<i>T. hibus</i>	Length	1.18	1.4–2.0	1.8–2.2	0.25
<i>T. malacosomae</i>	Length	1.37	0.8–1.1	1.1–1.5	0.00
<i>T. marylandensis</i>	Length	1.24	0.8–1.5	1.0–1.85	0.48
<i>T. melanis</i>	Length	1.19	1.3–1.4	1.3–1.9	0.17
<i>T. minutus</i>	Length	1.29	0.8–1.3	0.9–1.8	0.40
<i>T. pandora</i>	Length	1.03	1.1–1.8	1.4–1.6	0.29
<i>T. pattersonae</i>	Length	1.08	1.8–2.0	1.9–2.2	0.25
<i>T. racemariae</i>	Length	1.14	1.5–2.0	1.8–2.2	0.29
<i>T. rapo</i>	Length	1.23	1.1–1.5	1.4–1.8	0.14
<i>T. sobrius</i>	Length	1.56	1.0–1.25	1.5–2.0	–0.25
<i>T. spilopteris</i>	Length	1.23	1.1–1.5	1.0–2.2	0.33
<i>T. strobilus</i>	Length	1.09	1.5–1.9	1.5–2.2	0.57
<i>T. tesserus</i>	Length	1.12	1.6–1.7	1.6–2.1	0.20
<i>T. tineivorus</i> (= <i>T. carpatus</i>)	Length	1.29	0.8–1.3	1.1–1.6	0.25
<i>T. triozae</i>	Length	1.25	0.7–0.9	0.95–1.05	–0.14
<i>T. turionum</i>	Length	1.29	1.0–1.4	1.1–2.0	0.30
<i>T. verrucarii</i> (Burks, 1943)	Length	1.21	1.0–1.4	1.3–1.6	0.17
<i>T. inferens</i> (Yoshimoto, 1970b)	Length	1.52	1.4–1.7	1.7–3.0	0.00
<i>Wichmannia pictipennis</i> (Boucek, 1972)	Length	1.21	0.8–1.1	1.0–1.3	0.20
EUPELMIDAE					
<i>Anastatus umae</i> (Boucek, 1979)	Length	1.45	2.7–3.1	3.4–5.0	–0.13
EURYTOMIDAE					
<i>Buresium naso</i>	Length	1.58	1.45–3.1	3.0–4.2	0.04
<i>B. tenue</i> (Boucek, 1983)	Length	1.10	1.7–2.5	2.0–2.6	0.56

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism†	Size range or average		Overlap
		(female/male)	male	female	
ICHNEUMONIDAE					
Anomaloniae (subfamily):					
<i>Agrypon caribbaeum</i> (Bland, 1984)	Length	1.02	11.0–12.9	11.0–13.4	0.79
Campopleginae (subfamily):					
<i>Hyposoter exiguae</i> (Hurlbutt, unpubl.)	Length: range	1.08	4.3–7.5	5.6–7.1	0.47
	average s.d.	1.04	6.14 1.17	6.38 0.52	
	Headwidth: range	1.00	0.85–1.55	0.95–1.45	0.71
	average s.d.	1.06	1.18 0.12	1.24 0.11	
(Jowyk & Smilowitz, 1978)	Weight:				
on 2nd instar host	average s.d.	1.07	4.81 0.23	5.17 0.34	
on 4th instar host	average s.d.	1.11	6.25 0.46	6.91 0.49	
Cryptinae (subfamily):					
<i>Anurotropus minutus</i> (Cushman, 1924)		>			
<i>Cryptus leechi</i> (Mason, 1968a)	Length	1.08	4.5–7.5	5–8	0.71
<i>Diapetimorpha alabama</i> (Cushman, 1929)		>			
<i>Mastrus laplantei</i> (Mason, 1968a)	Length	1.00	3.5–5	3.5–5	1.00
<i>Messatoporus discoidalis</i> (Cushman, 1929)		>			
Ichneumoninae (subfamily):					
<i>Apaeleticus americanus</i>	Length	1.00	7.5–8.5	7.5–8.5	1.00
<i>Barichneumon carolinensis</i>	Length	0.79	6–8	5–6	0.00
<i>B. flaviscuta</i>	Length	0.88	7–9	6–8	0.33
<i>B. libens</i>	Length	0.92	6–7	5–7	0.50
<i>B. neosorex</i>	Length	0.94	7–9	7–8	0.50
<i>B. peramoenus calliandros</i>	Length	0.88	7–10	7–8	0.33
<i>B. peramoenus peramoenus</i>	Length	0.79	9–10	7–8	–0.33
<i>Carinodes havanensis</i>	Length	1.08	12–14	13–15	0.33
<i>Chasmias scelestus</i>	Length	0.86	16–19	13–17	0.17
<i>Chratichneumon</i>					
<i>C. anisotae</i>	Length	0.82	15–18	12–15	0.00
<i>C. annulatipes annulatipes</i>	Length	0.64	9–13	5–9	0.00
<i>C. facetus facetus</i>	Length	0.89	8–10	7–9	0.33
<i>C. flavipectus flavipectus</i>	Length	0.80	6–9	4–8	0.40
<i>C. floridensis</i>	Length	0.89	6–12	5–11	0.71
<i>C. georgius</i>	Length	0.94	15–17	14–16	0.33
<i>C. horani</i>	Length	0.90	9–11	8–10	0.33
<i>C. insulae</i>	Length	0.82	10–12	8–10	0.00
<i>C. louisianae</i>	Length	0.87	11–12	9–11	0.00
<i>C. paraparatus</i>	Length	0.87	9–14	8–12	0.50
<i>C. paratus paratus</i>	Length	0.68	9–13	5–10	0.13
<i>C. proximus</i>	Length	0.90	14–16	13–14	0.00
<i>C. pseudanisotae</i>	Length	0.97	12–17	13–15	0.40
<i>C. ritus</i>	Length	0.96	11–14	10–14	0.75
<i>C. scitulus</i>	Length	0.94	6–10	7–8	0.25
<i>C. subfilatus</i>	Length	1.00	11–15	11–15	1.00
<i>C. sublatus</i>	Length	0.77	14–17	9–15	0.13
<i>C. unifasciatorius</i>					
<i>C. unifasciatorius</i>	Length	0.91	14–18	12–17	0.50
<i>C. variegatus variegatus</i>	Length	0.83	16–19	14–15	–0.20
<i>C. vesus</i>	Length	0.73	9–13	7–9	0.00

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>C. vinnulus</i>	Length	1.04	9-14	9-15	0.83
<i>C. w-album w-album</i>	Length	0.89	12-15	11-13	0.25
<i>Diphyus comes</i>	Length	0.80	16-19	13-15	-0.17
<i>Eutanyacra pycnopus</i>	Length	1.00	12-13	12-13	1.00
<i>E. succincta</i>	Length	0.90	14-16	13-14	0.00
<i>Gnamptopelta obsidianator</i>					
<i>G. o. austrina</i>	Length	0.98	21-26	21-25	0.80
<i>G. o. obsidianator</i>	Length	0.90	25-27	21-26	0.17
<i>Hemihoplitis teres</i>	Length	0.95	9-12	9-11	0.67
<i>Homotherus townesi</i>	Length	0.88	8-9	7-8	0.00
<i>Ichneumon</i> (= <i>Coelichneumon</i>)					
<i>I. azotus</i>	Length	0.97	14-18	15-16	0.25
<i>I. eximius</i>	Length	0.94	15-18	14-17	0.50
<i>I. jejunos</i>	Length	1.04	11-16	13-15	0.40
<i>I. maurus</i>	Length	1.03	18-22	19-22	0.75
<i>I. navus albidior</i>	Length	0.95	8-13	8-12	0.80
<i>I. navus navus</i>	Length	0.95	8-13	8-12	0.80
<i>I. pulcher</i>	Length	1.03	16-18	17-18	0.50
<i>I. punctifer</i>	Length	0.97	14-18	14-17	0.75
<i>I. sassacus</i>	Length	0.94	15-17	13-17	0.50
<i>I. viola</i>	Length	0.92	16-21	16-18	0.40
<i>Limonethe maurator</i>	Length	0.93	12-18	12-16	0.67
<i>Linyx</i>					
<i>L. exhortator thoracicus</i>	Length	1.00	7-8	7-8	1.00
<i>Melanichneumon</i>					
<i>M. disparilis disparilis</i>	Length	0.90	9-11	8-10	0.33
<i>M. heiligbrodtii</i>	Length	0.96	10-14	10-13	0.75
<i>M. leviculus</i>	Length	0.86	10-11	8-10	0.00
<i>Orgichneumon</i>					
<i>O. calcatorius albidior</i>	Length	0.91	16-19	14-18	0.40
<i>O. calcatorius calcatorius</i>	Length	0.97	13-16	11-17	0.50
<i>Phaogenes trianguliferens</i>	Length	1.00	7-8	7-8	1.00
<i>Platylabus clarus</i>	Length	0.93	14-16	13-15	0.33
<i>P. hyperetis</i>	Length	0.93	10.5-11	9.5-10.5	0.00
<i>Probolus detritus</i>	Length	0.90	14-16	12-15	0.25
<i>Protichneumon ambiguus</i>	Length	1.00	20-25	20-25	1.00
<i>P. grandis grandis</i>	Length	0.94	26-28	24-27	0.25
<i>P. radlkeorum</i>	Length	0.94	25-29	25-26	0.25
<i>Protopelmus atrocaeruleus</i>	Length	1.09	20-24	22-26	0.33
<i>Pterocormus</i> (= <i>Ichneumon</i>)					
<i>P. ambulatrix</i>	Length	0.74	14-17	10-13	-0.14
<i>P. devinctus</i>	Length	0.95	20-21	19-20	0.00
<i>P. fuscifrons fuscifrons</i>	Length	1.00	12-15	13-14	0.33
<i>P. lewisii</i>	Length	0.94	17-18	14-19	0.20
<i>P. mendax</i>	Length	0.96	11-13	10-13	0.67
<i>P. versabilis</i>	Length	0.85	12-14	10-12	0.00
<i>P. weemsi</i>	Length	0.89	13-15	11-14	0.25
<i>P. weemsi</i> (= <i>I. tritus</i>)	Length	0.81	13-14	10-12	-0.25
<i>Rictichneumon belfragei</i>	Length	0.86	13-15	10-14	0.20
<i>Rubicundiella annulicornis</i>	Length	0.88	7-10	6-9	0.50
<i>R. mucronata</i>	Length	0.81	10-11	7-10	0.00
<i>R. perturbatrix</i>	Length	0.82	7-10	6-8	0.25
<i>Setanta compta compta</i>	Length	0.93	13-15	12-14	0.33
<i>Spilichneumon</i>					
<i>S. provancheri provancheri</i>	Length	0.93	14-15	13-14	0.00
<i>Tricholabus adventicus</i>	Length	0.96	12-14	11-14	0.67

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

B. HURLBUTT
Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>T. mitchelli</i>	Length	1.00	12-13	12-13	1.00
<i>Tricyphus apicalis</i>	Length	0.90	20-21	17-20	0.00
<i>T. floridanus</i>	Length	0.95	20-21	19-20	0.00
<i>Trogomorpha trogiformis</i>	Length	0.93	13-16	12-15	0.50
<i>Virgichneumon</i>					
<i>V. albomarginatus</i>	Length	0.89	13-15	12-13	0.00
<i>V. subcyaneus subcyaneus</i>	Length	0.86	12-16	10-14	0.33
<i>Virgichneumon brevicinctor</i>	Length	0.96	9-14	10-12	0.40
<i>V. terminalis terminalis</i> (Heinrich, 1977)	Length	0.83	7-11	7-8	0.25
Mesosteninae (subfamily):					
<i>Lagarosoma assilum</i> (Gupta & Gupta, 1984)		>			
Pimplinae (subfamily):					
<i>Dolichomitus</i> sp. (Kishi, 1970) on host species (lightest to heaviest):					
<i>Shirahoshizo</i> sp.	Pupal weight	1.33	10-50	30-50	0.50
<i>Niphaes variegatus</i>	Pupal weight	1.67	10-50	20-80	0.43
<i>Gregopimpla himalayensis</i> (Shiga & Nakanishi, 1968)	Weight	2.00	1-4	2-8	0.29
<i>Itopectis cristatae</i> (Nozato, 1969)	Pupal length	1.27	3.79-7.33	5.47-8.67	0.38
<i>Pimpla turionellae</i> (Arthur & Wylie, 1959) on host species (lightest to heaviest)					
<i>Rhyacionia buoliana</i>	Average weight	1.33	12	16	
<i>Pyrausta nubilalis</i>	Average weight	1.61	18	29	
<i>Galleria mellonella</i>	Average weight	1.36	22	30	
<i>Pieris rapae</i>	Average weight	1.29	21	27	
<i>Trichoplusia ni</i>	Average weight	1.79	24	43	
<i>Algaia milberti</i>	Average weight	1.29	31	40	
<i>Malacosoma americanum</i>	Average weight	1.52	29	44	
<i>Danaus plexippus</i>	Average weight	2.11	36	76	
Combined	Average weight	1.58	24	38	
<i>Rhyssa persuasoria</i> (Spradbery & Ratowsky, 1974)		>			
Tryphoninae (subfamily):					
<i>Phytodietus rutilus</i> (Krebs, 1969)	Length	1.11	5.9-6.7	6.7-7.3	0.00
MYMARIDAE					
<i>Anneckia oophaga</i> (Subba Rao, 1968)	Length	1.04	1.3-1.5	1.2-1.7	0.40
<i>Polynema ema</i> (Schauff & Grissell, 1982)	Length: range average	0.93 0.97	0.60-0.75 0.66	0.60-0.66 0.64	0.40
PLATYGASTRIDAE					
<i>Platygaster gracilipes</i> (Huggert, 1975)	Length	1.36	1.0-1.2	1.1-1.9	0.11
<i>P. tubulosa</i> (Huggert, 1980)	Length	1.02	1.1-1.3	0.74-1.70	0.21

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
PTEROMALIDAE					
<i>Arthrolytus muesebecki</i> (Burks, 1969b)	Length	1.31	3.0–3.5	3.5–5.0	0.00
<i>A. oezbeki</i> (Doganlar, 1978)	Length	1.47	1.13–2.81	2.29–3.52	0.22
<i>Echthrodape africana</i> (Burks, 1969a)	Length	0.94	3.5–4.5	3.5–4.0	0.50
<i>Euperilampus krombeini</i> (Burks, 1969a)	Length	1.16	4.0–5.5	5.0–6.0	0.25
<i>Habrocytus actinopterae</i> (Hedqvist, 1977b)	Length	1.06	1.5–2.0	1.5–2.2	0.71
<i>Halticoptera rosae</i> (Burks, 1955a)	Length	1.15	3.0–3.5	3.5–4.0	0.00
<i>Herbertia indica</i>	Length	1.08	1.1–1.4	1.2–1.5	0.50
<i>H. wallacei</i> (Burks, 1959)	Length	1.33	1.0–1.25	1.25–1.75	0.00
<i>Lariophagus distinguendus</i> (van den Assem, 1971)	Length	1.21	0.67–2.3	1.1–2.5	0.66
<i>Macromesus javensis</i> (Hedqvist, 1968)	Length	1.10	1.9–2.2	2.0–2.5	0.33
<i>Mesopolobus finlaysoni</i> (Doganlar, 1979)	Length	1.31	1.61–1.80	2.06–2.42	–0.32
<i>Monacon platypodis</i> (Boucek, 1980)	Length	0.97	2.8–3.2	2.8–3.0	0.50
<i>Muesebeckisia mandibularis</i> (Hedqvist, 1969)	Length	1.21	3.2–4.0	4.2–4.5	–0.15
<i>Muscidifurax raptor</i> (Legner, 1976)	Average weight	1.44	0.09	0.13	
(Kogan & Legner, 1970)	Headwidth	1.11	0.61	0.68	
	Length:				
(Rueda & Axtell, 1985)	range	1.13	1.70–2.20	1.90–2.50	0.38
	average	1.15	1.83	2.10	
(Kogan & Legner, 1970)	average	1.35	1.73	2.33	
(Legner, 1969)					
from Chile	Weight	1.76	0.62	1.09	
from South Africa	Weight	1.27	0.94	1.19	
from Israel	Weight	1.32	0.99	1.31	
from Orlando, Florida	Weight	1.57	0.86	1.35	
from s.w. U.S.					
(small form)	Weight	1.48	0.92	1.36	
from Costa Rica	Weight	1.51	0.90	1.36	
from Puerto Rico	Weight	2.45	0.60	1.47	
from Uruguay	Weight	1.81	0.85	1.54	
from Mexico	Weight	1.73	1.20	2.07	
from s.w. U.S.					
(giant form)	Weight	1.32	1.80	2.38	
<i>M. raptorellus</i> (Kogan & Legner, 1970)					
from Chile	Length	1.16	1.82	2.11	
	Headwidth	1.03	0.60	0.62	
from Uruguay	Length	1.24	1.85	2.29	
	Headwidth	1.08	0.62	0.67	
<i>M. raptoroides</i> (Kogan & Legner, 1970)					
from Costa Rica	Length	1.30	1.78	2.31	
	Headwidth	1.06	0.63	0.67	

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

B. HURLBUTT
Appendix (*cont.*)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
from Mexico	Length	1.21	1.98	2.39	
	Headwidth	1.06	0.66	0.70	
<i>M. zaraptor</i>					
(Legner, 1976)	Average weight	1.43	0.14	0.20	
(Kogan & Legner, 1970)	Headwidth	1.06	0.71	0.75	
	Length	1.30	2.18	2.84	
(Rueda & Axtell, 1985)	Length:				
	range	1.26	1.80–2.20	1.95–3.10	0.19
	average	1.41	2.03	2.87	
<i>Nasonia vitripennis</i>	Length:				
(Rueda & Axtell, 1985)	range	1.20	1.00–1.50	1.10–1.90	0.44
	average	1.50	1.15	1.73	
<i>Obtusiclava oryzae</i>					
(Subba Rao, 1973)	Length	1.35	1.75–2.5	2.5–3.25	0.00
<i>Pachycrepoideus vindemiae</i>					
(van Alphen & Thunissen, 1983)	Weight	1.35	0.025–0.075	0.028–1.075	0.58
(Rueda & Axtell, 1985)	Headwidth	1.02	0.002–0.246	0.002–0.240	0.98
	Length:				
	range	1.19	1.10–2.00	1.50–2.20	0.45
	average	1.08	1.81	1.96	
<i>Perilampus anomocerus</i>	Length	1.06	1.75–2.25	1.75–2.5	0.67
<i>P. canadensis</i>	Length	1.25	2.5–3.5	3–4.5	0.25
<i>P. carinifrons</i>	Length	1.18	2–2.25	1.75–3.25	0.17
<i>P. carolinensis</i>	Length	1.33	3.5–4	4.5–5.5	–0.25
<i>P. chrysopae</i>	Length	1.13	1.5–2.5	1.5–3	0.67
<i>P. crawfordi</i>	Length	1.00	2.5–3	2.5–3	1.00
<i>P. fulvicornis</i>	Length	1.11	1.5–3	1.5–3.5	0.75
<i>P. granulatus</i>	Length	1.06	1.75–2.5	1.75–2.75	0.75
<i>P. hyalinus</i>	Length	1.25	2–4	2–5.5	0.57
<i>P. muesebecki</i>	Length	1.21	1.5–2	1.5–2.75	0.40
<i>P. ocellatus</i>	Length	1.04	2.75–3.25	3–3.25	0.50
<i>P. platygaster</i>	Length	1.14	2.25–3	2.5–3.5	0.40
<i>P. robertsoni</i>	Length	1.10	2.25–2.75	2.5–3	0.33
<i>P. rohweri</i>	Length	0.91	2.5–3	2–3	0.50
<i>P. similis</i>	Length	1.22	1.75–2.75	1.75–3.75	0.50
<i>P. stygicus</i>	Length	1.07	1.75–2	1.75–2.25	0.50
<i>P. subcarinatus</i>	Length	1.04	2.5–3.5	2.5–3.75	0.80
<i>P. tristis</i>					
(= <i>P. capitatus</i>)					
(Smulyan, 1936)	Length	1.11	1.75–2.75	2–3	0.60
<i>P. philembia</i>					
(Burks, 1969a)	Length	1.13	1.75–2.0	1.75–2.50	0.33
<i>Pteromalus osmiae</i>	Length	1.42	2.0–3.0	3.0–4.1	0.00
<i>P. sylveni</i>					
(Hedqvist, 1979)	Length	1.28	2.0–2.3	2.0–3.5	0.20
<i>P. puparum</i>					
(Bouletreau & David, 1981)		>			
(Nealis <i>et al.</i> , 1984)		>			
<i>Spalangia cameroni</i>	Length:				
(Boucek, 1963)	range	1.07	2.4–3	2.5–3.3	0.56
(Rueda & Axtell, 1985)	range	1.13	1.80–2.60	2.10–2.85	0.48
	average	1.13	2.29	2.59	
<i>S. drosophilae</i>	Length:				
(Boucek, 1963)	range	1.07	1.3–1.6	1.4–1.7	0.50
(Rueda & Axtell, 1985)	range	1.22	0.80–1.25	0.90–1.60	0.44
	average	1.24	1.05	1.30	

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; > denotes female size > male size; <denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>S. endius</i> (Legner, 1976)	Average weight	1.58	0.12	0.19	
	Length:				
(Boucek, 1963)	range	1.11	1.9–2.6	2–3	0.55
(Rueda & Axtell, 1985)	range	1.21	1.60–2.30	1.80–2.90	0.38
	average	1.25	2.06	2.57	
<i>S. erythromera</i>	Length	1.07	1.9–2.7	1.7–3.2	0.53
<i>S. fuscipes</i>	Length	1.13	1.3–1.8	1.5–2	0.43
<i>S. nigra</i>	Length:				
(Boucek, 1963)	range	1.21	2.5–3.7	3–4.5	0.35
(Rueda & Axtell, 1985)	range	1.09	2.00–2.70	2.20–2.90	0.55
	average	1.10	2.36	2.59	
<i>S. nigripes</i>	Length	1.21	2–3.1	2.5–3.7	0.35
<i>S. nigroaenea</i>	Length:				
(Boucek, 1963)	range	1.12	2.5–3.5	2.9–3.8	0.46
(Rueda & Axtell, 1985)	range	1.08	2.10–2.80	2.20–3.10	0.60
	average	1.07	2.48	2.66	
<i>S. subpunctata</i> (Boucek, 1963)	Length	1.02	2.1–2.7	1.8–3.1	0.46
<i>Theocolaxia bifasciata</i> (Hedqvist, 1969)	Length	1.05	3.0–3.2	3.1–3.4	0.25
<i>Urolepis rufipes</i>	Length:				
(Rueda & Axtell, 1985)	range	1.56	1.25–2.08	2.18–3.02	0.00
	average	1.56	1.75	2.73	
<i>Zatropis capitis</i>	Length	1.07	1.2–1.8	1.2–2.0	0.75
<i>Z. chalcis</i>	Length	1.23	1.2–1.8	1.5–2.2	0.30
<i>Z. rosaecolis</i> (Burks, 1955b)	Length	1.71	1.5–2.0	2.0–4.0	0.00
SCELIONIDAE					
<i>Telenomus fariai</i> (Bosque & Rabinovich, 1979)	Length	>			
<i>T. phymatae</i> (Masner & Johnson, 1979)	Length	0.99	1.21–1.31	1.09–1.40	0.32
<i>T. solitus</i> (Johnson, 1983)	Length: average s.d.	1.04	1.26 0.07	1.31 0.06	
SCLEROGIBBIDAE					
<i>Sclerogibba citipes</i> (Krombein, 1979)	Length	1.11	3.0–3.5	3.2–4.0	0.30
TIPHIIDAE					
Miziniinae (subfamily):					
<i>Hylomesa longiceps</i> (Krombein, 1968)	Length	1.31	12–17	15–23	0.18
Thynninae (subfamily):					
<i>Eirone salteri</i> (Brown, 1984)	Length	0.77	14–16	11–12	–0.40
Tiphiiinae (subfamily):					
<i>Krombeinia evansi</i> (Allen, 1969)	Length	1.10	6.6–11.2	8.4–11.2	0.61
<i>Paratiphia</i> <i>P. aequalis aequalis</i>	Length	1.18	6.5–10	9–10.5	0.25
<i>P. nevadensis</i> (= <i>P. intermedia</i>) (Allen, 1962)	Length	1.13	7–9	7–11	0.50
<i>Tiphia grenada</i> (Allen, 1970)	Length	1.53	6.0–7.4	9.5–11.0	–0.42

*Length and headwidth in millimetres. Weight in milligrams.

† = denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

Appendix (cont.)

Wasp	Measure*	Dimorphism† (female/male)	Size range or average		
			male	female	Overlap
<i>T. hispaniolae</i> (Allen & Krombein, 1961)	Length	1.65	7–10	12–16	–0.22
<i>T. irifa</i> (Allen, 1961)	Length	1.32	9–10	12–13	–0.50
Methochinae (subfamily): (Borror <i>et al.</i> , 1976)		<			
Myrmosinae (subfamily): (Borror <i>et al.</i> , 1976)		<			
TORYMIDAE					
<i>Liodontomerus nitens</i> (Boucck, 1982)	Length	1.61	1.5–2.1	2.0–3.8	0.04
<i>Microdontomerus crassipes</i> (Boucck, 1982)	Length	1.33	2.0–2.6	2.5–3.6	0.06
<i>Monodontomerus mexicanus</i> (Gahan, 1941)	Length	>			
<i>Roptrocercus xylophagorum</i> (Samson, 1984)					
on 2nd instar host	Length	1.23	2.19 ± 0.05	2.70 ± 0.09	
on 3rd instar host	Length	1.27	2.63 ± 0.04	3.35 ± 0.04	
<i>Torymus canariensis</i> (Hedqvist, 1977a)	Length	1.03	1.2–1.7	1.2–1.8	0.83
TRICHOGRAMMATIDAE					
<i>Paropoea morimotoi</i> on host (smallest to largest):	Length	1.14	0.80–1.19	0.91–1.36	0.50
<i>Paracentrocorynus</i> <i>nigricollis</i>	Length	1.15	0.80–0.86	0.91–1.00	
<i>Phialodes rufipennis</i>	Length	1.21	1.00–1.19	1.29–1.36	
combined	Length	1.14	0.80–1.19	0.91–1.36	
<i>P. reticulata</i> (Hirose, 1963)	Length	1.17	0.93–1.00	1.07–1.19	–0.27
<i>Trichogramma minutum</i> (Southard <i>et al.</i> , 1982)	average s.d.				
wild population:					
on <i>Choristoneura</i>	Length	0.77	2.16 0.47	1.67 0.27	
<i>fumiferana</i> (host)	Headwidth	0.85	4.63 0.90	3.95 0.47	
lab populations:					
12 generations on	Length	0.90	2.09 0.37	1.88 0.25	
<i>C. fumiferana</i>	Headwidth	0.96	4.51 0.75	4.35 0.65	
52 generations on					
<i>Sitotroga cerealella</i>	Length	0.92	2.64 0.25	2.42 0.25	
(host)	Headwidth	1.01	5.33 0.23	5.40 0.48	
on <i>C. fumiferana</i> ,					
after 60 generations	Length	0.88	2.41 0.38	2.14 0.23	
on <i>S. cerealella</i>	Headwidth	0.93	4.89 0.53	4.54 0.46	
on <i>S. cerealella</i> ,					
after 12 generations	Length	0.97	2.43 0.33	2.35 0.23	
on <i>C. fumiferana</i>	Headwidth	1.07	5.15 0.47	5.52 0.34	

*Length and headwidth in millimetres. Weight in milligrams.

†=denotes female size = male size; > denotes female size > male size; < denotes female size < male size.

References

- ALLEN, H. W., 1961. *Tiphia* (Hymenoptera) of America north of Mexico: key to species, descriptions of six new species, and notes on synonymy. *Transactions of the American Entomological Society*, 87: 1–20.
- ALLEN, H. W., 1962. Species of *Paratiphia* (Hymenoptera: Tiphidae) with dark mandibles in the male. *Transactions of the American Entomological Society*, 88: 225–234.

- ALLEN, H. W., 1969. Three species of Mexican *Krombeinia* having an elongated tegula with an extended apical angle (Hymenoptera: Tiphidae). *Proceedings of the Entomological Society of Washington*, 71: 99–102.
- ALLEN, H. W., 1970. *Tiphia* of the West Indies (Hymenoptera: Tiphidae). *Proceedings of the Entomological Society of Washington*, 72: 252–269.
- ALLEN, H. W. & KROMBEIN, K. V., 1961. *Tiphia* of Cuba and Hispaniola (Hymenoptera: Tiphidae). *Transactions of the American Entomological Society*, 87: 57–66.
- ALPHEN, J. J. VAN & THUNISSEN, I., 1983. Host selection and sex allocation by *Pachycrepoideus vindemiae* Rondani (Pteromalidae) as a facultative hyperparasitoid of *Asobara tabida* Nees (Braconidae; Alysiinae) and *Leptopilina heterotoma* Thomson (Eulophidae). *Netherlands Journal of Zoology*, 33: 497–514.
- ARTHUR, A. P. & WYLIE, H. G., 1959. Effects of host size on sex ratio, development time and size of *Pimpla turionellae* (L.) (Hymenoptera: Ichneumonidae). *Entomophaga*, 4: 297–301.
- ASKEW, R. R., 1984. Species of *Phygadeuon* and *Chrysocharis* (Hymenoptera: Eulophidae) parasitic on Tischeriidae (Lepidoptera) with the description of a new species. *Entomologist's Gazette*, 35: 103–109.
- ASSEM, J. VAN DEN, 1971. Some experiments on sex ratio and sex regulation in the pteromalid *Lariophagus distinguendus*. *Netherlands Journal of Zoology*, 21: 373–402.
- BARRETT, C. F., WESTDAL, P. H. & RICHARDSON, H. P., 1965. Biology of *Pachygonatopus minimus* Fenton (Hymenoptera: Dryinidae) a parasite of the six-spotted leafhopper, *Macrostes fasciatus* (Stål), in Manitoba. *Canadian Entomologist*, 97: 216–221.
- BARRON, J. R., 1970. A new species of *Copidosoma* (Hymenoptera: Encyrtidae) parasitizing larvae of *Anacampsis innocella* (Lepidoptera: Gelechiidae). *Canadian Entomologist*, 102: 1337–1339.
- BENSON, J. F., 1973. Intraspecific competition in the population dynamics of *Bracon hebetor* Say (Hymenoptera: Braconidae). *Journal of Animal Ecology*, 42: 105–124.
- BLAND, R. G., 1984. *Agrypon caribbaeum*, a new species of ichneumon wasp (Hymenoptera: Ichneumonidae) from the U.S. Virgin Islands. *Annals of the Entomological Society of America*, 77: 29–31.
- BORROR, D. J., DELONG, D. M. & TRIPLEHORN, C. A., 1976. *An Introduction to the Study of Insects*, 4. New York: Holt, Rinehart and Winston.
- BOSQUE, C. & RABINOVICH, J. E., 1979. Population dynamics of *Telenomus fariai* (Hymenoptera: Scelionidae), a parasite of Chagas' disease vectors. VII. Oviposition behaviour and host discrimination. *Canadian Entomologist*, 111: 171–180.
- BOUCEK, Z., 1963. A taxonomic study in *Spalangia* Latr. (Hymenoptera, Chalcidoidea). *Acta Entomologica Musei Nationalis Pragae*, 35: 429–481.
- BOUCEK, Z., 1972. Descriptions of new eulophid parasites (Hym., Chalcidoidea) from Africa and the Canary Islands. *Bulletin of Entomological Research*, 62: 199–205.
- BOUCEK, Z., 1976. Taxonomic studies on some Eulophidae (Hym.) of economic interest, mainly from Africa. *Entomophaga*, 21: 401–414.
- BOUCEK, Z., 1977a. Descriptions of *Tachinobia* gen. n. and three new species of Tetrastichinae (Hymenoptera: Eulophidae), with a tentative key to genera. *Bulletin of Entomological Research*, 67: 17–30.
- BOUCEK, Z., 1977b. Descriptions of two new species of Neotropical Eulophidae (Hymenoptera) of economic interest, with taxonomic notes on related species and genera. *Bulletin of Entomological Research*, 67: 1–15.
- BOUCEK, Z., 1979. Description of a new eupelmid parasite (Hymenoptera: Chalcidoidea) of cockroaches in India. *Bulletin of Entomological Research*, 69: 93–96.
- BOUCEK, Z., 1980. A revision of the genus *Monacon* Waterston (Hymenoptera: Chalcidoidea: Perilampinae), parasites of ambrosia beetles (Coleoptera: Platypodidae). *Bulletin of Entomological Research*, 70: 73–96.
- BOUCEK, Z., 1982. Four new Mediterranean Torymidae (Hymenoptera). *Entomologist's Gazette*, 33: 183–191.
- BOUCEK, Z., 1983. On *Buresium*, *Masneroma* (n. gen.) and some other Eurytomidae (Hymenoptera). *Entomologica Scandinavica*, 14: 186–194.
- BOULETREAU, M. & DAVID, J. R., 1981. Sexually dimorphic response to host habitat toxicity in *Drosophila* parasitic wasps. *Evolution*, 35: 395–399.
- BROWN, G. R., 1984. New species of *Eirone* (Hymenoptera: Tiphidae) from New Caledonia and Australia, with a review of the previously known New Caledonian species. *International Journal of Entomology*, 26: 249–259.
- BURKS, B. D., 1940. Revision of the Chalcid-flies of the tribe Chalcidini in America north of Mexico. *Proceedings of the United States National Museum*, 88: 237–354.
- BURKS, B. D., 1943. The North American parasitic wasps of the genus *Tetrastichus*—a contribution to biological control of insect pests. *Proceedings of the United States National Museum*, 93: 505–608.
- BURKS, B. D., 1952. The North American species of *Syntomosphyrum* (Hymenoptera, Chalcidoidea). *Proceedings of the Entomological Society of Washington*, 54: 258–264.
- BURKS, B. D., 1955a. A new species of the genus *Halticoptera* (Hymenoptera, Pteromalidae). *Proceedings of the Entomological Society of Washington*, 57: 81–82.
- BURKS, B. D., 1955b. A redefinition of the genus *Zatropis*, with descriptions of three new species (Hymenoptera, Pteromalidae). *Proceedings of the Entomological Society of Washington*, 57: 21–37.
- BURKS, B. D., 1959. The species of the genus *Herbertia* Howard (Hymenoptera, Pteromalidae). *Proceedings of the Entomological Society of Washington*, 61: 249–255.
- BURKS, B. D., 1966. The North American species of *Pediobius* Walker (Hymenoptera: Eulophidae). *Proceedings of the Entomological Society of Washington*, 68: 33–43.

- BURKS, B. D., 1969a. New Perilampidae (Hymenoptera: Chalcidoidea). *Proceedings of the Entomological Society of Washington*, 71: 73-81.
- BURKS, B. D., 1969b. The North American species of *Arthrolytus* Thomson (Hymenoptera: Pteromalidae). *Proceedings of the Entomological Society of Washington*, 71: 289-303.
- BURKS, B. D., 1971. The Nearctic species of *Horismenus* Walker (Hymenoptera: Eulophidae). *Proceedings of the Entomological Society of Washington*, 73: 68-83.
- BURKS, B. D., 1972. The genus *Hexacladia* Ashmead (Hymenoptera: Encyrtidae). *Proceedings of the Entomological Society of Washington*, 74: 363-371.
- BURKS, B. D., 1977. The Mexican species of *Chalcis* Fabricius (Hymenoptera: Chalcididae). *Proceedings of the Entomological Society of Washington*, 79: 383-399.
- CARDONA, C. & OATMAN, E. R., 1971. Biology of *Apanteles dignus* (Hymenoptera: Braconidae), a primary parasite of the tomato pinworm. *Annals of the Entomological Society of America*, 64: 996-1007.
- CUSHMAN, R. A., 1924. New genera and species of ichneumon-flies. *Proceedings of the United States National Museum*, 64: 1-16.
- CUSHMAN, R. A., 1929. A revision of the North American ichneumon-flies of the genus *Mesostenus* and related genera. *Proceedings of the United States National Museum*, 74: 1-58.
- DEBACH, P. & ROSE, M., 1981. A new genus and species of Aphelinidae with some synonymies, a rediagnosis of *Aspidiotiphagus* and a key to pentamerous heteromerous Prospaltellinae (Hymenoptera: Chalcidoidea: Aphelinidae). *Proceedings of the Entomological Society of Washington*, 83: 658-679.
- DOGANLAR, M., 1978. A new species of Pteromalinae (Hymenoptera: Pteromalidae) from western North America. *Canadian Entomologist*, 110: 1111-1115.
- DOGANLAR, M., 1979. Two new species of *Mesopolobus* Westwood (Hymenoptera: Pteromalidae) from western Canada. *Canadian Entomologist*, 111: 649-659.
- EVANS, H. E., 1954. The North American species of *Dissomphalus* (Hymenoptera, Bethyridae). *Proceedings of the Entomological Society of Washington*, 56: 288-309.
- EVANS, H. E., 1959a. The genus *Anisephyris* in America north of Mexico (Hymenoptera, Bethyridae). *Proceedings of the Entomological Society of Washington*, 61: 97-120.
- EVANS, H. E., 1959b. The Nearctic species of *Lophepyris*, a new subgenus of *Rhabdepyris* (Hymenoptera, Bethyridae). *Proceedings of the Entomological Society of Washington*, 61: 201-204.
- FINNEY, G. L., FLANDERS, S. E. & SMITH, H. S., 1947. Mass culture of *Macrocentrus ancylivorus* and its host, the potato tuber moth. *Hilgardia*, 17: 437-483.
- FREYTAG, P. H., 1977. A review of the genus *Neogonatopus* for North America (Hymenoptera: Dryinidae). *Annals of the Entomological Society of America*, 70: 569-576.
- GAHAN, A. B., 1922. Descriptions of miscellaneous new reared parasitic hymenoptera. *Proceedings of the United States National Museum*, 61: 1-24.
- GAHAN, A. B., 1930. Synonymical and descriptive notes on parasitic hymenoptera. *Proceedings of the United States National Museum*, 77: 1-12.
- GAHAN, A. B., 1941. A revision of the chalcid flies of the genus *Monodontomerus* in the United States National Museum. *Proceedings of the United States National Museum*, 90: 461-482.
- GARDENFORS, U., 1983. Re-description, distribution and host range of *Calaphidius* Mackauer (Hymenoptera: Braconidae, Aphidinae). *Entomologica Scandinavica*, 14: 77-80.
- GERDIN, S. & HEDQVIST, K.-J., 1984. *Perilitus areolaris* sp. n. (Hymenoptera: Braconidae), an imago-parasitoid of the large pine weevil, *Hylobius abietis* (Linnaeus), and its reproductive behaviour. *Entomologica Scandinavica*, 15: 363-369.
- GERLING, D., 1966a. Studies with whitefly parasites of southern California. I. *Encarsia pergandiella* Howard (Hymenoptera: Aphelinidae). *Canadian Entomologist*, 98: 707-724.
- GERLING, D., 1966b. Studies with whitefly parasites of southern California II. *Eretmocerus californicus* Howard (Hymenoptera: Aphelinidae). *Canadian Entomologist*, 98: 1316-1329.
- GIRAULT, A. A., 1908. *Encarsia versicolor* species novum, an eulophid parasite of the greenhouse whitefly *Aleyrodes vaporariorum* Westwood. *Psyche*, 15: 53-57.
- GORDH, G. & EVANS, H. E., 1976. A new species of *Goniozus* imported into California from Ethiopia for the biological control of pink bollworm and some notes on the taxonomic status of *Parasierola* and *Goniozus* (Hymenoptera: Bethyridae). *Proceedings of the Entomological Society of Washington*, 78: 479-489.
- GORDH, G. & HENDRICKSON, R. Jr., 1979. New species of *Diglyphus*, a world list of the species, taxonomic notes and a key to new world species of *Diglyphus* and *Diaulinopsis* (Hymenoptera: Eulophidae). *Proceedings of the Entomological Society of Washington*, 81: 666-684.
- GOYER, R. A. & SCHENK, J. A., 1970. Notes on the biology and habits of insect parasites associated with *Eucosma rescissoriana* (Lepidoptera: Olethreutidae) in northern Idaho. *Canadian Entomologist*, 102: 1546-1553.
- GRISSELL, E. E., 1981. *Edovum putleri*, n.g., n. sp. (Hymenoptera: Eulophidae), an egg parasite of the Colorado potato beetle (Chrysomelidae). *Proceedings of the Entomological Society of Washington*, 83: 790-796.
- GRISSELL, E. E. & SCHAUFF, M. E., 1981. New Nearctic *Inureia* (Hymenoptera: Chalcididae) from lepidopterous pests of peanut. *Proceedings of the Entomological Society of Washington*, 83: 1-12.
- GUPTA, S. & GUPTA, V., 1984. *Lagarosoma*, a new Brazilian genus of the tribe Gabuniini (Hymenoptera: Ichneumonidae). *International Journal of Entomology*, 26: 235-240.
- HAYAT, M., 1976. Two new species of Aphelinidae (Hym.: Chalcidoidea) parasitic on *Acaudaleyrodes rhachipora* (Hom.: Aleyrodidae) from India. *Entomophaga*, 21: 157-162.

- HEDQVIST, K. J., 1968. Studies in Pteromalidae in the U.S. National Museum (Hymenoptera). *Proceedings of the United States National Museum*, 124: 1-11.
- HEDQVIST, K. J., 1969. Notes on Cerocephalini with descriptions of new genera and species (Hymenoptera: Chalcidoidea: Pteromalidae). *Proceedings of the Entomological Society of Washington*, 71: 449-467.
- HEDQVIST, K. J., 1977a. Two new reared species of Torymidae and Eulophidae from Tenerife, Canary Islands (Hymenoptera: Chalcidoidea). *Entomologica Scandinavica*, 8: 235-237.
- HEDQVIST, K. J., 1977b. Notes on Chalcidoidea XI (Hymenoptera). A new species of *Habrocytus* Thomson from Sweden and a lectotype selection for *Pteromalus groenlandicus* Holmgren. *Entomologica Scandinavica*, 8: 237-238.
- HEDQVIST, K. J., 1979. Two new species of the genus *Pteromalus* Swed. from Sweden (Hymenoptera: Chalcidoidea, Pteromalidae). *Entomologica Scandinavica*, 10: 155-157.
- HEINRICH, G. H., 1977. Ichneumoninae of Florida and neighboring states. *Arthropods of Florida and Neighboring Land Areas*. Volume 9. Florida Department of Agriculture and Consumer Services, Gainesville, Florida.
- HIROSE, Y., 1963. Discovery of the genus *Poropoea* parasitic on the eggs of attelabid beetles in Japan (Hymenoptera, Trichogrammatidae). *Acta Hymenopterologica*, 2: 29-35.
- HUGGERT, L., 1975. Descriptions of Platygastriinae from Sweden (Hymenoptera: Platygastriidae). *Entomologica Scandinavica*, 6: 61-66.
- HUGGERT, L., 1980. Taxonomical studies on some genera and species of Platygastriinae (Hymenoptera: Proctotrupoidea). *Entomologica Scandinavica*, 11: 97-112.
- HUGGERT, L., 1982. Descriptions and redescrptions of *Trichopria* species from Africa and the Oriental and Australian regions (Hymenoptera, Proctotrupoidea: Diapriidae). *Entomologica Scandinavica*, 13: 109-122.
- JOHNSON, N. F., 1983. Description of *Telenomus solitus*, n. sp. (Hymenoptera: Scelionidae), a noctuid egg parasitoid. *Proceedings of the Entomological Society of Washington*, 85: 226-228.
- JOWYK, E. A. & SMILOWITZ, Z., 1978. A comparison of growth and developmental rates of the parasite *Hyposoter exiguae* reared from two instars of its host, *Trichoplusia ni*. *Annals of the Entomological Society of America*, 71: 467-472.
- KISHI, Y., 1970. Differences in the sex ratio of the pine bark weevil parasite, *Dolichomitus* sp. (Hymenoptera: Ichneumonidae), emerging from different host species. *Applied Entomology and Zoology*, 5: 126-132.
- KNUTSON, L. V. & BERG, C. O., 1963. *Phaenopria popei* (Hymenoptera: Diapriidae) reared from puparia of sciomyzid flies. *Canadian Entomologist*, 95: 724-726.
- KOGAN, M. & LEGNER, E. F., 1970. A biosystematic revision of the genus *Muscidifurax* (Hymenoptera: Pteromalidae) with descriptions of four new species. *Canadian Entomologist*, 102: 1268-1290.
- KREBS, C. F., 1969. A new Nearctic species of *Phytodietus* (Hymenoptera: Ichneumonidae). *Canadian Entomologist*, 101: 548-550.
- KROMBEIN, K. V., 1968. Studies in the Tiphidae, X *Hylomesa*, a new genus of myzinine wasp parasitic on larvae of longicorn beetles (Hymenoptera). *Proceedings of the United States National Museum*, 124: 1-22.
- KROMBEIN, K. V., 1969. The generic placement of two nearctic *Holopyga* with biological notes (Hymenoptera: Chrysididae). *Proceedings of the Entomological Society of Washington*, 71: 351-361.
- KROMBEIN, K. V., 1979. Biosystematic studies of Ceylonese wasps, VI. Notes on the Sclerogibbidae with descriptions of two new species (Hymenoptera: Chryridoidea). *Proceedings of the Entomological Society of Washington*, 81: 465-474.
- LAWRENCE, P. O., BARANOWSKI, R. M. & GREANY, P. D., 1976. Effect of host age on development of *Biosteres* (= *Opius*) *Longicaudatus*, a parasitoid of the Caribbean fruit fly, *Anastrepha suspensa*. *Florida Entomologist*, 59: 33-39.
- LEGNER, E. F., 1969. Reproductive isolation and size variation in the *Muscidifurax raptor* complex. *Annals of the Entomological Society of America*, 62: 382-385.
- LEGNER, E. F., 1976. Low storage temperature effects on the reproductive potential of three parasites of *Musca domestica*. *Annals of the Entomological Society of America*, 69: 435-441.
- MACKAUER, M., 1965. Notes on and additions to the *Trioxys* and *Binodoxys* species (Hymenoptera: Aphidiidae) of North America. *Canadian Entomologist*, 97: 225-231.
- MACKAUER, M., 1972. The aphid-attacking genera of Aphelinidae (Hymenoptera), including the description of a new genus. *Canadian Entomologist*, 104: 1771-1779.
- MACKAUER, M., SULLIVAN, D. J., & SULLIVAN, S. J., 1982. A new species of *Praon* (Hymenoptera: Aphidiidae), parasitic on the dusky-veined walnut aphid, *Callaphis juglandis* (Homoptera: Aphididae). *Canadian Entomologist*, 114: 1159-1161.
- MARSH, P. M., 1967. The Nearctic Doryctinae, V. The genus *Leluthia* and comments on the status of the tribe Hecabolini (Hymenoptera, Braconidae). *Proceedings of the Entomological Society of Washington*, 69: 359-364.
- MARSH, P. M., 1968a. The Nearctic Doryctinae, VI. The genera *Acrophasmus*, *Glyptocolastes*, *Doryctinus*, and a new genus *Stenocorse* (Hymenoptera: Braconidae). *Proceedings of the Entomological Society of Washington*, 70: 101-113.
- MARSH, P. M., 1968b. A new species of *Alysia* parasitic on carrion-feeding flies (Hymenoptera: Braconidae). *Proceedings of the Entomological Society of Washington*, 70: 208-210.
- MARSH, P. M., 1976. The Nearctic Doryctinae, X. The genus *Rhaconotus* Ruthe (Hymenoptera: Braconidae). *Proceedings of the Entomological Society of Washington*, 78: 389-403.

- MARSH, P. M., 1982. Two new species of *Heterospilus* (Hymenoptera: Braconidae) from Mexico being introduced against the cotton boll weevil, *Anthonomus grandis* (Coleoptera: Curculionidae). *Proceedings of the Entomological Society of Washington*, 84: 849–854.
- MASNER, L. & JOHNSON, N. F., 1979. A new species of *Telenomus* (Hymenoptera: Scelionidae), an egg parasite of the ambush bug, *Phymata* sp. (Heteroptera: Phymatidae). *Canadian Entomologist*, 111: 1115–1119.
- MASON, W. R., 1968a. New Canadian Cryptinae (Ichneumonidae: Hymenoptera). *Canadian Entomologist*, 100: 17–23.
- MASON, W. R., 1968b. New Canadian Braconidae (Hymenoptera). *Canadian Entomologist*, 100: 715–728.
- MASON, W. R., 1974. The *Apanteles* species (Hymenoptera: Braconidae) attacking Lepidoptera in the micro-habitat of the spruce budworm (Lepidoptera: Tortricidae). *Canadian Entomologist*, 106: 1087–1102.
- MASON, W. R., 1978. A synopsis of the Nearctic Braconini, with revisions of Nearctic species of *Coeloides* and *Myosoma* (Hymenoptera: Braconidae). *Canadian Entomologist*, 110: 721–768.
- MINOT, M. C. & LEONARD, D. E., 1976. Host preference and development of the parasitoid *Brachymeria intermedia* in *Lymantria dispar*, *Galleria mellonella*, and *Choristoneura fumiferana*. *Environmental Entomology*, 5: 527–532.
- NEALIS, V. G., JONES, R. E. & WELLINGTON, W. G., 1984. Temperature and development in host-parasite relationships. *Oecologia*, 61: 224–229.
- NORDLANDER, G., 1978. Revision of the genus *Rhoptromeris* Forster, 1869 with reference to north-western European species. Studies on Eucoilidae (Hym.: Cynipoidea) II. *Entomologica Scandinavica*, 9: 47–62.
- NORDLANDER, G., 1980. Revision of the genus *Leptopilina* Forster, 1869, with notes on the status of some other genera (Hymenoptera, Cynipoidea: Eucoilidae). *Entomologica Scandinavica*, 11: 428–453.
- NOYES, J. S., 1982. A new species of *Zeteticontus* Silvestri (Hymenoptera: Encyrtidae) from Israel and Kenya, a parasite of *Carpophilus hemipterus* (L.) (Coleoptera: Nitidulidae). *Bulletin of Entomological Research*, 72: 457–460.
- NOYES, J. S., 1984. A new genus and species of encyrtid (Hymenoptera: Chalcidoidea) parasitic on the casava mealybug, *Phenacoccus manihoti* Matile-Ferrero (Hemiptera: Pseudococcidae). *Bulletin of Entomological Research*, 74: 529–533.
- NOYES, J. S. & HAYAT, M., 1984. A review of the genera of Indo Pacific Encyrtidae (Hymenoptera: Chalcidoidea). *Bulletin of the British Museum (Natural History) Entomology*, 48.
- NOZATO K., 1969. Biology of *Itopectis cristatae* (Hym.: Ichneumonidae), a pupal parasite of pine shoot moths in Japan. *Kontyu*, 37: 75–82.
- ODEBIYI, J. A. & OATMAN, E. R., 1972. Biology of *Agathis gibbosa* (Hymenoptera: Braconidae), a primary parasite of the potato tuberworm. *Annals of the Entomological Society of America*, 65: 1104–1114.
- ROSEN, D. & DEBACH, P., 1979. *Species of Aphytis of the World* (Hymenoptera: Aphelinidae). The Hague: Junk Press.
- RUEDA, L. M. & AXTELL, R. C., 1985. Guide to common species of pupal parasites (Hymenoptera: Pteromalidae) of the house fly and other muscoid flies associated with poultry and livestock manure. Technical Bulletin 278. North Carolina Agricultural Research Service, North Carolina State University.
- SAMSON, P. R., 1984. The biology of *Roptrocercus xylophagorum* (Hym.: Torymidae), with a note on its taxonomic status. *Entomophaga*, 29: 287–298.
- SCHAUFF, M. E. & GRISELLE, E. E., 1982. Nomenclatural notes on *Polynema* (Hymenoptera: Mymaridae), with descriptions of a new species. *Proceedings of the Entomological Society of Washington*, 84: 529–534.
- SHIGA, M. & NAKANISHI, A., 1968. Intraspecific competition in a field population of *Gregopimpla himalayensis* (Hymenoptera: Ichneumonidae) parasitic on *Malacosoma neustria testacea* (Lepidoptera: Lasiocampidae). *Research Population Ecology (Kyoto)*, 10: 69–86.
- SMULYAN, M. T., 1936. A revision of the chalcid flies of the genus *Perilampus* Latreille occurring in America north of Mexico. *Proceedings of the United States National Museum*, 83: 369–412.
- SOUTHARD, S. G., HOUSEWEART, M. W., JENNINGS, D. T. & HALTEMAN, W. A., 1982. Size differences of laboratory reared and wild populations of *Trichogramma minutum* (Hymenoptera: Trichogrammatidae). *Canadian Entomologist*, 114: 693–698.
- SPRADBERY, J. P. & RATOWSKY, D. A., 1974. An analysis of geographical variation in the parasitoid *Rhyssa persuasoria* (L.) (Hymenoptera, Ichneumonidae). *Bulletin of Entomological Research*, 64: 653–668.
- SUBBA RAO, B. R., 1968. Descriptions of new genera and species of Mymaridae (Hymenoptera) from the Far East and the Ethiopian region. *Bulletin of Entomological Research*, 59: 659–670.
- SUBBA RAO, B. R., 1973. Description of a new species and genus of Pteromalidae (Hymenoptera) parasitic on *Pachydiplosis oryzae* (Wood-Mason) (Diptera, Cecidomyiidae). *Bulletin of Entomological Research*, 62: 627–629.
- TIMBERLAKE, P. H., 1922. A revision of the chalcid-flies of the encyrtid genus *Chrysoplatycerus*. *Proceedings of the United States National Museum*, 61: 1–10.
- WATANABE, C., 1969. Notes on the genera *Zelet* Curtis and *Xiphozela* Cameron with special reference to the species in Japan (Hymenoptera: Braconidae). *Proceedings of the Entomological Society of Washington*, 71: 318–328.
- WELD, L. H., 1955. A new genus and species of North American Cynipoidea (Hymenoptera). *Proceedings of the Entomological Society of Washington*, 57: 274.

- WHARTON, R. A., 1983. New species of *Illidops* and *Bracon* (Hymenoptera: Braconidae) of potential use in biological control. *Canadian Entomologist*, 115: 667–672.
- YOSHIMOTO, C. M., 1970a. A new species of *Astichus* (Hymenoptera: Eulophidae) associated with the birch bracket fungus *Polyporus betulinus* and woody fungus *Ganoderma appalanatum* in eastern Canada. *Canadian Entomologist*, 102: 656–659.
- YOSHIMOTO, C. M., 1970b. A new species of *Tetrastichus* (Hymenoptera: Eulophidae) parasitizing pupae of *Sesamia inferens* (Lepidoptera: Noctuidae). *Canadian Entomologist*, 102: 1607–1609.
- YOSHIMOTO, C. M., 1971. A new species of the genus *Zamischus* and description of the male *Z. brazillensis* (Hymenoptera, Cynipidae: Eucoilinae). *Canadian Entomologist*, 103: 979–982.
- YOSHIMOTO, C. M., 1972. A new species of the genus *Stemmatosteres* (Encyrtidae: Chalcidoidea). *Canadian Entomologist*, 104: 1837–1839.
- YOSHIMOTO, C. M., 1976a. Synopsis of the genus *Mestocharis* Forster in America north of Mexico (Chalcidoidea: Eulophidae). *Canadian Entomologist*, 108: 755–758.
- YOSHIMOTO, C. M., 1976b. Revision of the genus *Dicladocerus* (Eulophidae: Chalcidoidea) of America north of Mexico, with particular reference to species attacking larch casebearer (Lepidoptera: Coleophoridae). *Canadian Entomologist*, 108: 1173–1206.