

# Vectors and Linear Transformations

Ling 282/482: Deep Learning for Computational Linguistics

C.M. Downey

Fall 2025

# Linear Algebra Objects

# Linear Algebra Objects

- **Scalars**
  - Single numbers
  - What you're used to elsewhere in math
  - examples: 0, 1, 3.14,  $\pi$ , 7/22

# Linear Algebra Objects

- **Scalars**

- Single numbers
- What you're used to elsewhere in math
- examples: 0, 1, 3.14,  $\pi$ , 7/22

- **Vectors**

- *Lists* of scalars

# Linear Algebra Objects

- **Scalars**

- Single numbers
- What you're used to elsewhere in math
- examples: 0, 1, 3.14,  $\pi$ , 7/22

- **Vectors**

- *Lists* of scalars

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

# Linear Algebra Objects

- **Scalars**

- Single numbers
- What you're used to elsewhere in math
- examples: 0, 1, 3.14,  $\pi$ , 7/22

- **Vectors**

- *Lists* of scalars

- **Matrices**

- Lists of vectors

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

# Linear Algebra Objects

- **Scalars**

- Single numbers
- What you're used to elsewhere in math
- examples: 0, 1, 3.14,  $\pi$ , 7/22

- **Vectors**

- *Lists* of scalars

- **Matrices**

- Lists of vectors

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad A = \begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix}$$

# Vectors



# Vectors

- By default, vectors are considered to be *columns*

# Vectors

- By default, vectors are considered to be *columns*

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

# Vectors

- By default, vectors are considered to be ***columns***
  - *Transposed* vectors are rows

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix}$$

# Vectors

- By default, vectors are considered to be ***columns***
  - *Transposed* vectors are rows

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad x^T = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$$

# Vectors


- By default, vectors are considered to be ***columns***
  - *Transposed* vectors are rows
  - Often visualized as **arrows** or **points** in space

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad x^T = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$$

# Vectors

- By default, vectors are considered to be ***columns***
  - *Transposed* vectors are rows
  - Often visualized as **arrows** or **points** in space

$$x = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad x^T = \begin{bmatrix} 1 & 2 & 3 \end{bmatrix}$$



# Vector Properties

# Vector Properties

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{bmatrix}$$



# Vector Properties

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 - y_1 \\ x_2 - y_2 \\ x_3 - y_3 \end{bmatrix}$$

# Vector Properties

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} + \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 + y_1 \\ x_2 + y_2 \\ x_3 + y_3 \end{bmatrix}$$

$$\begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} - \begin{bmatrix} y_1 \\ y_2 \\ y_3 \end{bmatrix} = \begin{bmatrix} x_1 - y_1 \\ x_2 - y_2 \\ x_3 - y_3 \end{bmatrix}$$

$$c \begin{bmatrix} x_1 \\ x_2 \\ x_3 \end{bmatrix} = \begin{bmatrix} cx_1 \\ cx_2 \\ cx_3 \end{bmatrix}$$

(c is a scalar)

# Dot Product

# Dot Product

- Vectors have an operation called the **dot product** that isn't as intuitive

# Dot Product

- Vectors have an operation called the **dot product** that isn't as intuitive
- Denoted:  $a \cdot b$  or  $a^T b$  where  $a$  and  $b$  are **vectors** of the **same length**

# Dot Product

- Vectors have an operation called the **dot product** that isn't as intuitive
- Denoted:  $a \cdot b$  or  $a^T b$  where  $a$  and  $b$  are **vectors** of the **same length**
- Formula:
  - $a_1 b_1 + a_2 b_2 + \dots a_n b_n$
  - (corresponding elements are multiplied then summed)

# Dot Product

- Vectors have an operation called the **dot product** that isn't as intuitive
- Denoted:  $a \cdot b$  or  $a^T b$  where  $a$  and  $b$  are **vectors** of the **same length**
- Formula:
  - $a_1 b_1 + a_2 b_2 + \dots a_n b_n$
  - (corresponding elements are multiplied then summed)
- The dot product returns a **single value!!** (not a vector)

# Dot Product

- Vectors have an operation called the **dot product** that isn't as intuitive
- Denoted:  $a \cdot b$  or  $a^T b$  where  $a$  and  $b$  are **vectors** of the **same length**
- Formula:
  - $a_1 b_1 + a_2 b_2 + \dots a_n b_n$
  - (corresponding elements are multiplied then summed)
- The dot product returns a **single value!!** (not a vector)
- Intuition: the **strength** with which the two vectors **go in the same direction**
  - (Not super important to remember this right now)
  - Important fact: **perpendicular vectors** have a dot product of **zero!**



# Vector Spans and Spaces

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

$$c_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} + c_2 \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

$$c_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} + c_2 \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- ...except for  $c_1 = c_2 = 0$  (which always gives the zero vector)

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

$$c_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} + c_2 \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- ...except for  $c_1 = c_2 = 0$  (which always gives the zero vector)
- Otherwise the vectors are **independent**

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

$$c_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} + c_2 \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- ...except for  $c_1 = c_2 = 0$  (which always gives the zero vector)
- Otherwise the vectors are **independent**
- Definition applies to **any number** of vectors and constants

# Vector Independence

- Two vectors are linearly **dependent** iff there are scalars  $c_1, c_2$ :

$$c_1 \begin{bmatrix} a_1 \\ a_2 \\ a_3 \end{bmatrix} + c_2 \begin{bmatrix} b_1 \\ b_2 \\ b_3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

- ...except for  $c_1 = c_2 = 0$  (which always gives the zero vector)
- Otherwise the vectors are **independent**
- Definition applies to **any number** of vectors and constants
- Note:  $a = \mathbf{0}$  is used to indicate a vector of zeros

# Vector (In)dependence Examples

- What constants solve this equation?

$$c_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$



# Vector (In)dependence Examples

- What constants solve this equation?

$$c_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} 2 \\ 4 \\ 6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

# Vector (In)dependence Examples

- What constants solve this equation?

$$c_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + c_3 \begin{bmatrix} 5 \\ 7 \\ 9 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$

# Vector (In)dependence Examples

- What constants solve this equation?

$$c_1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + c_2 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} = \begin{bmatrix} 0 \\ 0 \\ 0 \end{bmatrix}$$


# Vector (In)dependence Examples

# Vector (In)dependence Examples

- Vectors are dependent if they are **colinear**


# Vector (In)dependence Examples

- Vectors are dependent if they are **colinear**





# Vector (In)dependence Examples

- Vectors are dependent if they are **colinear**
- Non-colinear vectors can also be dependent



# Vector (In)dependence Examples



- Vectors are dependent if they are **colinear**
- Non-colinear vectors can also be dependent





# Vector (In)dependence Examples

- Vectors are dependent if they are **colinear**
- Non-colinear vectors can also be dependent



(this is what  
adding  
vectors looks  
like)

# Vector Spans

# Vector Spans


- Adding together vectors with different constants is called a **linear combination** of those vectors

# Vector Spans

- Adding together vectors with different constants is called a **linear combination** of those vectors
- The set of **all linear combinations** of some vectors is called their **span**

# Vector Spans

- Adding together vectors with different constants is called a **linear combination** of those vectors
- The set of **all linear combinations** of some vectors is called their **span**



# Spaces and Spans

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$



# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on
- Two vectors of size 2 **span**  $R^2$  iff they are **independent**

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on
- Two vectors of size 2 **span**  $R^2$  iff they are **independent**
- Three vectors of size 3 **span**  $R^3$  iff they are **independent**

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on
- Two vectors of size 2 **span**  $R^2$  iff they are **independent**
- Three vectors of size 3 **span**  $R^3$  iff they are **independent**
- If the num of independent vectors is **less than the vector dimension**, they span a **(hyper)plane** within the larger space

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on
- Two vectors of size 2 **span**  $R^2$  iff they are **independent**
- Three vectors of size 3 **span**  $R^3$  iff they are **independent**
- If the num of independent vectors is **less than the vector dimension**, they span a **(hyper)plane** within the larger space

$$a = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad b = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$$

# Spaces and Spans

- The entirety of 1-dimensional space is called  $R^1$ 
  - 2-dimensional space is called  $R^2$
  - and so on
- Two vectors of size 2 **span**  $R^2$  iff they are **independent**
- Three vectors of size 3 **span**  $R^3$  iff they are **independent**
- If the num of independent vectors is **less than the vector dimension**, they span a **(hyper)plane** within the larger space
  - Ex: **a** and **b** above span a **2-D plane** in  $R^3$

$$a = \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} \quad b = \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix}$$

# Basis of a Space

# Basis of a Space

- A set of independent vectors that span a space are called a **basis** for that space



# Basis of a Space

- A set of independent vectors that span a space are called a **basis** for that space
- The simplest bases for  $R^2$  and  $R^3$  are known as the **Standard Basis**

# Basis of a Space

- A set of independent vectors that span a space are called a **basis** for that space
- The simplest bases for  $R^2$  and  $R^3$  are known as the **Standard Basis**

$$R^2$$
$$\dot{i} = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad \dot{j} = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

# Basis of a Space

- A set of independent vectors that span a space are called a **basis** for that space
- The simplest bases for  $R^2$  and  $R^3$  are known as the **Standard Basis**

$$R^2$$
$$i = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad j = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$R^3$$
$$i = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad j = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad k = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$

# Basis of a Space


- A set of independent vectors that span a space are called a **basis** for that space
- The simplest bases for  $R^2$  and  $R^3$  are known as the **Standard Basis**
- These are not the only bases for these spaces

$$R^2$$
$$i = \begin{bmatrix} 1 \\ 0 \end{bmatrix} \quad j = \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$

$$R^3$$
$$i = \begin{bmatrix} 1 \\ 0 \\ 0 \end{bmatrix} \quad j = \begin{bmatrix} 0 \\ 1 \\ 0 \end{bmatrix} \quad k = \begin{bmatrix} 0 \\ 0 \\ 1 \end{bmatrix}$$




# Span Video





# Span Video





# Matrix Multiplication

# Quick reminder: Dot Product

$$a \cdot b = a^T b = a_1 b_1 + a_2 b_2 \dots + a_n b_n$$



# Quick reminder: Dot Product

$$a \cdot b = a^T b = a_1 b_1 + a_2 b_2 \dots + a_n b_n$$

(vectors need to be the same length)

# Matrix-Vector Multiplication

$$A = \begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \quad x = \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

$$Ax = ?$$

# Matrix Multiplication Rules

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns



# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

$$\begin{array}{c} 4 \text{ rows} \\ \begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix} \\ 2 \text{ columns} \\ \text{“4x2 matrix”} \end{array} \quad \begin{array}{c} \begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \\ 3 \times 2 \end{array} \quad \begin{array}{c} \begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix} \\ 2 \times 3 \end{array}$$

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

3x2

$$\begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$$

2x3

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

3x2

$$\begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$$

2x3

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

3x2

$$\begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$$

3x2

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

3x2

$$\begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$$

2x3

✗

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$$

3x2

$$\begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$$

3x2

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$$

3x2      2x3

✗

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$$

3x2      3x2

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \end{bmatrix}$$

3x2      2x1

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$

3x2      2x3

✗  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$

3x2      3x2

✓  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 \\ 8 \end{bmatrix}$

3x2      2x1

# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$$

3x2      2x3

✗

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$$

3x2      3x2

✓

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \end{bmatrix}$$

3x2      2x1

$$\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix} \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix}$$

3x2      3x1



# Matrix Multiplication Rules

- Matrix multiplication is **not** commutative:  $Ax \neq xA$
- The “inner” sizes of the matrices must match

4 rows

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

2 columns

“4x2 matrix”

✓  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 & 9 & 11 \\ 8 & 10 & 12 \end{bmatrix}$  ✗  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 & 10 \\ 8 & 11 \\ 9 & 12 \end{bmatrix}$

3x2      2x3      3x2      3x2

✓  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 \\ 8 \end{bmatrix}$  ✗  $\begin{bmatrix} 1 & 4 \\ 2 & 5 \\ 3 & 6 \end{bmatrix}$   $\begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix}$

3x2      2x1      3x2      3x1

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} - \\ - \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} - \\ - \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ - \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ - \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ - \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ - \end{bmatrix}$$



# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ - \end{bmatrix}$$

# Matrix-Vector Multiplication

The Traditional Way

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Matrix-Vector Multiplication

# Matrix-Vector Multiplication

- Alternative way to think about this multiplication

# Matrix-Vector Multiplication

- Alternative way to think about this multiplication

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

# Matrix-Vector Multiplication

- Alternative way to think about this multiplication
  - The matrix consists of **column vectors**

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

# Matrix-Vector Multiplication

- Alternative way to think about this multiplication
  - The matrix consists of **column vectors**
  - The vector provides the **constants** for a **linear combination of the columns**

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix}$$

# Matrix-Vector Multiplication

- Alternative way to think about this multiplication
  - The matrix consists of **column vectors**
  - The vector provides the **constants** for a **linear combination of the columns**

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{1} \\ \textcolor{green}{1} \end{bmatrix} \longrightarrow \textcolor{red}{1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \textcolor{blue}{1} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + \textcolor{green}{1} \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$



# Matrix-Vector Multiplication

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \longrightarrow 1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 1 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + 1 \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Matrix-Vector Multiplication

- What is the significance of this alternate view?

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{1} \\ \textcolor{green}{1} \end{bmatrix} \longrightarrow \textcolor{red}{1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \textcolor{blue}{1} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + \textcolor{green}{1} \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Matrix-Vector Multiplication

- What is the significance of this alternate view?
  - For all  $Ax = b$ ,  $b$  is expressed as a **linear combination** of  $A$ 's columns, and so...

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \longrightarrow 1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 1 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + 1 \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Matrix-Vector Multiplication

- What is the significance of this alternate view?
  - For all  $Ax = b$ ,  $b$  is expressed as a **linear combination** of  $A$ 's columns, and so...
  - ... $b$  is always in the **span of  $A$ 's columns**

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} \textcolor{red}{1} \\ \textcolor{blue}{1} \\ \textcolor{green}{1} \end{bmatrix} \longrightarrow \textcolor{red}{1} \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + \textcolor{blue}{1} \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + \textcolor{green}{1} \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Matrix-Vector Multiplication

- What is the significance of this alternate view?
  - For all  $Ax = b$ ,  $b$  is expressed as a **linear combination** of  $A$ 's columns, and so...
  - ... $b$  is always in the **span of  $A$ 's columns**
  - This is called the **Column Space of  $A$** ,  $C(A)$

$$\begin{bmatrix} 1 & 4 & 7 \\ 2 & 5 & 8 \\ 3 & 6 & 9 \end{bmatrix} \begin{bmatrix} 1 \\ 1 \\ 1 \end{bmatrix} \longrightarrow 1 \begin{bmatrix} 1 \\ 2 \\ 3 \end{bmatrix} + 1 \begin{bmatrix} 4 \\ 5 \\ 6 \end{bmatrix} + 1 \begin{bmatrix} 7 \\ 8 \\ 9 \end{bmatrix} = \begin{bmatrix} 12 \\ 15 \\ 18 \end{bmatrix}$$

# Column Space

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?
  - 3 independent columns

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$



# Column Space

- What can you tell about the Column Space of this matrix?
  - 3 independent columns
  - $C(A)$  spans  $R^3$

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?
  - 3 independent columns
  - $C(A)$  spans  $R^3$
  - $Ax$  spans  $R^3$

$$\begin{bmatrix} 1 & 3 & 0 \\ 2 & 4 & 0 \\ 0 & 0 & 5 \end{bmatrix}$$

# Column Space

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?
  - **2** independent columns

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix?
- 2 independent columns

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

third column not independent of first two

# Column Space

- What can you tell about the Column Space of this matrix?
  - 2 independent columns
  - $C(A)$  spans a **2D plane** in  $R^3$

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

third column not independent of first two

# Column Space

- What can you tell about the Column Space of this matrix?
  - 2 independent columns
  - $C(A)$  spans a **2D plane** in  $R^3$
  - $Ax$  spans a **2D plane** in  $R^3$

$$\begin{bmatrix} 1 & 4 & 5 \\ 2 & 5 & 7 \\ 3 & 6 & 9 \end{bmatrix}$$

third column not independent of first two



# Column Space

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix? What is the size of “input” vector  $x$ ?

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix? What is the size of “input” vector  $x$ ?
- **2** independent columns

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix? What is the size of “input” vector  $x$ ?
- **2** independent columns
- $x$  is length **2**

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix? What is the size of “input” vector  $x$ ?
- **2** independent columns
- $x$  is length **2**
- $C(A)$  spans a **2D plane** in  $R^4$

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Column Space

- What can you tell about the Column Space of this matrix? What is the size of “input” vector  $x$ ?
- **2** independent columns
- $x$  is length **2**
- $C(A)$  spans a **2D plane** in  $R^4$
- $Ax$  spans a **2D plane** in  $R^4$

$$\begin{bmatrix} 1 & 5 \\ 2 & 6 \\ 3 & 7 \\ 4 & 8 \end{bmatrix}$$

# Matrix Rank

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**



# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**
  - Rank 2: **plane**

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**
  - Rank 2: **plane**
  - Rank 3: **3D hyperplane**

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**
  - Rank 2: **plane**
  - Rank 3: **3D hyperplane**
  - etc.

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**
  - Rank 2: **plane**
  - Rank 3: **3D hyperplane**
  - etc.
- MxN matrix can be considered a **function** from  $R^N$  to  $R^M$

# Matrix Rank

- The number of independent columns in a matrix is called the **rank**
- The rank determines the **dimension of the column space**
  - Rank 1: **line**
  - Rank 2: **plane**
  - Rank 3: **3D hyperplane**
  - etc.
- MxN matrix can be considered a **function** from  $R^N$  to  $R^M$ 
  - **However**, the function's range **may not span**  $R^M$ , unless it is **rank M**

# Linear Transformations



# Identity Matrix

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$
  - Where have we seen these columns before?

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$
  - Where have we seen these columns before?

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$i \qquad j$

$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

# Identity Matrix

- The Identity Matrix  $I$  always returns the **same vector/matrix** it's multiplied with
  - e.g.  $Ix = x$  and  $IA = A$
  - Where have we seen these columns before?

$$\begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix}$$

$i$                    $j$


$$\begin{bmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix}$$

$i$                    $j$                    $k$



# Identity Matrix as a Basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 0 \end{bmatrix} + 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \begin{bmatrix} 0 \\ 1 \end{bmatrix}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis

$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$






# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis

$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$






# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$



# Identity Matrix as a Basis

- Vectors can be viewed as being **composed of the Standard Basis vectors**
- A vector is a linear combination of this basis


$$\begin{bmatrix} 3 \\ 5 \end{bmatrix} = \begin{bmatrix} 1 & 0 \\ 0 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \underset{i}{\begin{bmatrix} 1 \\ 0 \end{bmatrix}} + 5 \underset{j}{\begin{bmatrix} 0 \\ 1 \end{bmatrix}}$$





# Linear Transformation


$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$



# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$






# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$






# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$






# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**
  - This is called a **linear transformation**

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$






# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**
  - This is called a **linear transformation**
  - This matrix **rotates** the space by  $45^\circ$  and stretches it

$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$




# Linear Transformation

- Multiplying by a matrix converts a vector to a **new basis**
  - The basis consists of the **matrix columns**
  - This is called a **linear transformation**
  - This matrix **rotates** the space by  $45^\circ$  and stretches it



$$\begin{bmatrix} 1 & -1 \\ 1 & 1 \end{bmatrix} \begin{bmatrix} 3 \\ 5 \end{bmatrix} = 3 \begin{bmatrix} 1 \\ 1 \end{bmatrix} + 5 \begin{bmatrix} -1 \\ 1 \end{bmatrix} =$$

new basis  $i'$   $j'$







# Linear Transformation



# Visualizing Linear Transformations



# Visualizing Linear Transformations





# Why do we need this?

# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**

# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**
- A Feedforward NN layer or Perceptron:

# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**
- A Feedforward NN layer or Perceptron:
  - Takes in an **input vector**

# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**
- A Feedforward NN layer or Perceptron:
  - Takes in an **input vector**
  - Applies a **linear transformation**



# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**
- A Feedforward NN layer or Perceptron:
  - Takes in an **input vector**
  - Applies a **linear transformation**
  - Adds a **non-linear activation function** (we'll cover this later)

# Why do we need this?

- Linear transformations (aka matrix multiplications) are one of the **basic operations of neural networks**
- A Feedforward NN layer or Perceptron:
  - Takes in an **input vector**
  - Applies a **linear transformation**
  - Adds a **non-linear activation function** (we'll cover this later)
- TLDR: Neural Nets transform vectors and vector spaces

# Quiz next session!

- You should be able to:
  - **Draw vectors** on a grid
  - **Add and subtract** vectors
  - Identify when two vectors are **independent**
  - Take the **dot product** of two vectors
  - **Multiply** a vector by a matrix
  - **Multiply** two matrices
  - Give the form and discuss properties of the **Identity Matrix**
- Bring a **pen or pencil!** I will provide **calculators**