Aristotle University of Thessaloniki

Faculty of Sciences — School of Physics

Computational Physics Master's Programme

M.Sc. Thesis
Computational Methods on Black Hole Imaging

Author:
Christos Mermigkas

Supervisor:
George Pappas
Assistant Professor, AUTH

November 11, 2026

Thessaloniki, Greece

Abstract

In this thesis, we present a robust and efficient computational framework for the numerical simulation and visualization of relativistic ray tracing in the strong gravitational regime of rotating Kerr black holes. Developed entirely in the Julia programming language [1], our methodology employs a backward ray-tracing paradigm to map the observer's celestial sphere to the spacetime geometry surrounding the event horizon. To address the extreme numerical sensitivity inherent in near-horizon photon dynamics, we implement two advanced computational techniques.

First, we utilize Forward-Mode Automatic Differentiation to compute exact metric derivatives and Christoffel symbols to machine precision, entirely eliminating the truncation and round-off errors associated with traditional finite difference methods.

Second, we couple a high-order adaptive Runge-Kutta integrator with a geometric Manifold Projection algorithm that strictly enforces the physical invariants of the phase space - the null condition, energy, and axial angular momentum - suppressing invariant drift even for highly unstable trajectories.

We systematically validate the numerical stability of our integrator by resolving fundamental photon orbits, including scattering, direct capture, and bound spirals near the photon sphere, demonstrating that conserved quantities are maintained at the machine-precision limit. On a standard 16-core laptop, the framework processes computationally demanding trajectories at an average rate of one million rays per hour.

Finally, we apply the framework to astrophysical imaging, successfully reproducing the characteristic asymmetric "D-shaped" shadow of a near-extremal Kerr black hole induced by frame-dragging. By incorporating a geometrically thin accretion disk model with relativistic radiative transfer, we capture key phenomena such as Doppler effect, gravitational lensing, and the hierarchy of higher-order photon sub-rings. The framework's metric-agnostic nature and rigorous geometric integration make it a powerful, general-purpose tool for interpreting horizon-scale observations from current and future interferometric campaigns, such as the Event Horizon Telescope (EHT).

Περίληψη

Στην παρούσα εργασία, παρουσιάζουμε ένα εύρωστο και αποδοτικό υπολογιστικό πλαίσιο για την αριθμητική προσομοίωση και οπτικοποίηση της σχετικιστικής ιχνηλάτησης ακτίνων (relativistic ray tracing) στο ισχυρό βαρυτικό πεδίο περιστρεφόμενων μελανών οπών Kerr. Αναπτυγμένο εξ ολοκλήρου στη γλώσσα προγραμματισμού Julia, το πλαίσιο υιοθετεί την αντίστροφη ιχνηλάτηση ακτίνων (backward ray-tracing) για την αντιστοίχιση της ουράνιας σφαίρας του παρατηρητή με τη γεωμετρία του χωροχρόνου γύρω από τον ορίζοντα γεγονότων. Για την αντιμετώπιση της ακραίας αριθμητικής ευαισθησίας που χαρακτηρίζει τη δυναμική των φωτονίων κοντά στον ορίζοντα, εφαρμόζουμε δύο προηγμένες υπολογιστικές τεχνικές.
Πρώτον, αξιοποιούμε την Αυτόματη Παραγώγιση προς τα εμπρός (Forward-Mode Automatic Differentiation) για τον ακριβή υπολογισμό των παραγώγων του μετρικού τανυστή και των συμβόλων Christoffel με ακρίβεια μηχανής, εξαλείφοντας πλήρως τα σφάλματα αποκοπής και στρογγυλοποίησης που συνοδεύουν τις παραδοσιακές μεθόδους πεπερασμένων διαφορών.
Δεύτερον, συνδυάζουμε έναν ολοκληρωτή Runge-Kutta υψηλής τάξης και προσαρμοστικού βήματος με έναν γεωμετρικό αλγόριθμο Προβολής Πολλαπλότητας (Manifold Projection). Η μέθοδος αυτή επιβάλλει αυστηρά τα φυσικά αναλλοίωτα του χώρου φάσεων -τη φωτονική συνθήκη (null condition), την ενέργεια και την αξονική στροφορμή- καταστέλλοντας την αριθμητική απόκλιση (drift) ακόμη και για εξαιρετικά ασταθείς τροχιές.
Επικυρώνουμε συστηματικά την αριθμητική ευστάθεια του ολοκληρωτή επιλύοντας θεμελιώδεις τροχιές φωτονίων, συμπεριλαμβανομένων της σκέδασης, της άμεσης σύλληψης και των δεσμευμένων ελικοειδών τροχιών κοντά στη σφαίρα των φωτονίων, αποδεικνύοντας ότι τα διατηρούμενα μεγέθη διατηρούνται στα όρια της ακρίβειας της μηχανής. Σε έναν τυπικό φορητό υπολογιστή με 16 πυρήνες, το πλαίσιο επεξεργάζεται υπολογιστικά απαιτητικές τροχιές με μέσο ρυθμό ένα εκατομμύριο ακτίνες ανά ώρα.
Τέλος, εφαρμόζουμε το πλαίσιο στην αστροφυσική απεικόνιση, αναπαράγοντας επιτυχώς τη χαρακτηριστική ασύμμετρη σκιά σχήματος «D» μιας σχεδόν ακραίας (near-extremal) μαύρης τρύπας Kerr, η οποία οφείλεται στη συμπαράσυρση πλαισίου (frame-dragging). Ενσωματώνοντας ένα μοντέλο γεωμετρικά λεπτού δίσκου προσαύξησης με σχετικιστική διάδοση ακτινοβολίας, αναδεικνύουμε κρίσιμα φαινόμενα όπως το φαινόμενο Doppler (Doppler effect), η βαρυτική εστίαση (gravitational lensing) και η ιεραρχία των φωτονικών υπο-δακτυλίων ανώτερης τάξης. Η ανεξαρτησία από τη μορφή της μετρικής (metric-agnostic nature) και η αυστηρή γεωμετρική ολοκλήρωση το καθιστούν ένα ισχυρό εργαλείο γενικής χρήσης για την ερμηνεία παρατηρήσεων σε κλίμακα ορίζοντα από τρέχουσες και μελλοντικές συμβολομετρικές εκστρατείες, όπως το Τηλεσκόπιο Ορίζοντα Γεγονότων (Event Horizon Telescope - EHT).

Acknowledgements

I would like to express my deepest gratitude towards my supervisor, Assistant Prof. George Pappas, for his invaluable guidance, patience, and support throughout this journey. His insights and mentorship have been fundamental to the completion of this work.

My sincere thanks go to my friend Giorgos, for the countless hours we spent together in the office writing code and discussing results. Having a study buddy who constantly pushed me made the most demanding parts of this process not only manageable but genuinely enjoyable.

My heartfelt appreciation goes to Ntina, who has been by my side from the very beginning. Her unwavering support, understanding, and belief in me have been my greatest source of strength during the most challenging moments of this endeavor.

Finally, I owe everything to my parents, for their endless love, sacrifices, and for providing me with every opportunity to pursue my dreams.

This achievement is as much theirs as it is mine. Thank you.

Do not go gentle into that good night

Dylan Thomas

"Do not go gentle into that good night,
Old age should burn and rave at close of day;
Rage, rage against the dying of the light.

Wild men who caught and sang the sun in flight,
And learn, too late, they grieved it on its way,
Do not go gentle into that good night.

Grave men, near death, who see with blinding sight
Blind eyes could blaze like meteors and be gay,
Rage, rage against the dying of the light."

Contents

1 Theoretical Foundations of Spacetime and Geodesic Motion **9**

1.1 Differential Geometry and Tensor Calculus . . . 9

1.2 Curvature Tensors and the Einstein Field Equations . . . 10

1.3 The Geodesic Equation and the Variational Principle . . . 11

1.3.1 Killing's Equation and Exact Conservation Laws . . . 11

1.4 Stationary, Axisymmetric Spacetimes and the Hamiltonian Formulation . . . 12

1.5 The Kerr Geometry . . . 13

1.5.1 Horizon Structure and Ergosphere . . . 13

1.5.2 Null Geodesics and the Carter Constant . . . 13

1.6 Light Rings and the Mathematical Shadow . . . 14

1.7 Strong-Field Gravitational Lensing and Multiple Imaging . . . 14

1.8 Radiative Transfer Formulation . . . 15

2 Numerical Methods and Computational Algorithms **17**

2.1 Theoretical Background: Well-Posedness and Error in ODE Integration . . . 17

2.1.1 The Initial Value Problem . . . 17

2.1.2 Local Error, Global Error, and Convergence . . . 17

2.1.3 Truncation Error, Round-off, and the Accuracy Floor . . . 17

2.2 Automatic Differentiation vs. Finite Differences . . . 18

2.2.1 Dual Numbers and the Chain Rule . . . 18

2.3 Geometric Integration via Manifold Projection . . . 19

2.3.1 The Problem of Invariant Drift . . . 19

2.3.2 Projection via Lagrange Multipliers . . . 19

2.3.3 Multidimensional Newton-Raphson Iterative Solver . . . 20

2.3.4 Convergence of the Newton–Raphson Iteration . . . 21

2.3.5 Regularity of the Constraints . . . 21

2.3.6 Projection Versus Alternative Stabilization Strategies . . . 21

2.4 High-Order Adaptive Integration . . . 22

2.4.1 Butcher Structure and Order Conditions . . . 22
2.4.2 Embedded Pairs and Local Error Estimation . . . 22

3 Setup **25**
3.1 Backward Ray Tracing and Initial Conditions . . . 25
3.1.1 Choice of Coordinate System . . . 25
3.1.2 Grid Construction and Initial Conditions . . . 25
3.1.3 UV Mapping and Calibration . . . 26

4 Results **27**
4.1 Validation of Numerical Stability . . . 27
4.1.1 Fundamental Photon Orbits and Dynamics . . . 28
4.1.2 Scattering Orbits . . . 28
4.1.3 Bound Spiral Orbits (Transient Capture) . . . 29
4.1.4 Direct Capture Orbits . . . 30
4.2 Black Hole Shadow and Astrophysical Imaging . . . 31
4.2.1 Gravitational Lensing of a Point Source . . . 31
4.2.2 Shadow Morphology and Frame Dragging . . . 33
4.2.3 Relativistic Accretion Disk and Radiative Transfer . . . 34

5 Conclusions and Future Prospects **39**
5.1 Summary of Contributions . . . 39
5.2 Future Research Directions . . . 40
5.2.1 GPU Acceleration and Real-Time Rendering . . . 40
5.2.2 Extension to Alternative Metrics and Beyond-GR Theories . . . 40
5.2.3 Template Generation and Parameter Estimation . . . 40

A Simulation Configurations **41**

Chapter 1

Theoretical Foundations of Spacetime and Geodesic Motion

1.1 Differential Geometry and Tensor Calculus

The mathematical formulation of the General Theory of Relativity posits that gravitational interactions are not mediated by conventional Newtonian forces, but are manifestations of the intrinsic curvature of a four-dimensional pseudo-Riemannian spacetime manifold $\mathcal{M}$ [12, 16]. A differentiable manifold $\mathcal{M}$ of dimension $n = 4$ is a topological space locally homeomorphic to Euclidean space $\mathbb{R}^4$, equipped with an atlas of overlapping charts (U_α, ϕ_α).

Let $x^\mu = (x^0, x^1, x^2, x^3) \equiv (t, r, \theta, \phi)$ denote a local coordinate system. When transitioning to another overlapping chart $\tilde{x}^\mu$, the coordinate transformation functions are smooth (C^∞) and possess a non-singular Jacobian matrix. The fundamental geometric object dictating distances and causality on $\mathcal{M}$ is the symmetric metric tensor $g_{\mu\nu} = g_{\nu\mu}$. The invariant infinitesimal line element ds^2 is given by:

$$ds^2 = g_{\mu\nu} dx^\mu dx^\nu, \tag{1.1}$$

where we adopt the standard relativistic signature $(-, +, +, +)$. The inverse metric tensor $g^{\mu\nu}$ satisfies the canonical orthogonality relation $g^{\mu\sigma} g_{\sigma\nu} = \delta^\mu_{\ \nu}$, and serves as the index-raising and lowering operator.

To describe the parallel transport of vectors on a curved manifold, we introduce the covariant derivative ∇_α. The standard partial derivative $\partial_\alpha A^\mu$ of a vector field fails to transform as a tensor. To correct this, we introduce an affine connection $\Gamma^\mu_{\ \alpha\beta}$:

$$\nabla_\alpha A^\mu = \partial_\alpha A^\mu + \Gamma^\mu_{\ \alpha\beta} A^\beta. \tag{1.2}$$

Assuming the conditions of zero torsion ($\Gamma^\mu_{\ \alpha\beta} = \Gamma^\mu_{\ \beta\alpha}$) and metric compatibility ($\nabla_\alpha g_{\mu\nu} = 0$), the affine connection is uniquely determined as the Christoffel symbols of the second kind:

$$\Gamma^\mu_{\ \alpha\beta} = \frac{1}{2} g^{\mu\sigma} \left(\partial_\alpha g_{\beta\sigma} + \partial_\beta g_{\alpha\sigma} - \partial_\sigma g_{\alpha\beta} \right). \tag{1.3}$$

We note that the accurate evaluation of Eq. (1.3) is crucial for numerical relativity, as it requires the computation of metric derivatives $\partial_\sigma g_{\mu\nu}$. This is a challenge we will systematically address in the following chapter.

1.2 Curvature Tensors and the Einstein Field Equations

The covariant derivative does not commute. Acting on an arbitrary vector field, the commutator defines the *Riemann curvature tensor*:

$$\nabla_{[\alpha}\nabla_{\beta]}V^{\mu} \equiv R^{\mu}{}_{\nu\alpha\beta}\,V^{\nu}. \tag{1.4}$$

If $\mathcal{M}$ is flat, a vector can be compared with itself after transport around an infinitesimal loop and Eq. (1.4) vanishes identically; on a curved manifold it does not. Curvature is therefore precisely the local obstruction to parallel transport being path-independent.

The Riemann tensor obeys the algebraic symmetries

$$R_{\mu\nu\alpha\beta} = -R_{\nu\mu\alpha\beta} = -R_{\mu\nu\beta\alpha} = R_{\alpha\beta\mu\nu}, \qquad R_{\mu[\nu\alpha\beta]} = 0, \tag{1.5}$$

where the last relation is the (first) Bianchi identity. In four dimensions these reduce the number of independent components from $4^4 = 256$ to 20. Two successive contractions produce the *Ricci tensor* and *Ricci scalar*,

$$R_{\mu\nu} \equiv R^{\alpha}{}_{\mu\alpha\nu}, \qquad R \equiv g^{\mu\nu}R_{\mu\nu}, \tag{1.6}$$

which enter the Einstein field equations (in geometrized units $G = c = 1$):

$$R_{\mu\nu} - \tfrac{1}{2}R\,g_{\mu\nu} = 8\pi\,T_{\mu\nu}. \tag{1.7}$$

The contracted Bianchi identity $\nabla^{\mu}G_{\mu\nu} = 0$ guarantees consistency of Eq. (1.7) with local energy–momentum conservation, $\nabla^{\mu}T_{\mu\nu} = 0$. In vacuum, $T_{\mu\nu} = 0$, the field equations reduce to $R_{\mu\nu} = 0$: ten independent conditions. The remaining ten components of the Riemann tensor reside in the *Weyl tensor*

$$C_{\mu\nu\alpha\beta} = R_{\mu\nu\alpha\beta} - \tfrac{1}{2}\left(g_{\mu\alpha}R_{\nu\beta} - g_{\mu\beta}R_{\nu\alpha} - g_{\nu\alpha}R_{\mu\beta} + g_{\nu\beta}R_{\mu\alpha}\right) + \tfrac{R}{6}\left(g_{\mu\alpha}g_{\nu\beta} - g_{\mu\beta}g_{\nu\alpha}\right), \tag{1.8}$$

which shares all symmetries of $R_{\mu\nu\alpha\beta}$ and is additionally traceless. The Weyl tensor represents the *free* gravitational field: it encodes tidal forces and gravitational waves and propagates through vacuum. The Kerr spacetime is Ricci-flat but strongly curved: all of its gravity lives in $C_{\mu\nu\alpha\beta}$.

Curvature has a direct operational meaning through *geodesic deviation*. Let ξ^{μ} be the separation vector between two neighbouring, infinitesimally close geodesics with tangent u^{μ}. Then, to leading order,

$$\frac{D^2\xi^{\mu}}{D\lambda^2} = -R^{\mu}{}_{\nu\alpha\beta}\,u^{\nu}\xi^{\alpha}u^{\beta}. \tag{1.9}$$

In the Newtonian limit this reproduces $\ddot{\xi}^{i} = -\left(\partial_i\partial_j\Phi\right)\xi^{j}$: the Riemann tensor is the relativistic generalization of the tidal matrix. A uniform gravitational field (equivalence principle) can always be transformed away at a point; Eq. (1.9) is what remains and what cannot.

For Kerr, the quadratic curvature invariant (Kretschmann scalar) evaluates to

$$K = \frac{48\,M^2}{\Sigma^6}\left(r^6 - 15\,a^2\,r^4\cos^2\theta + 15\,a^4\,r^2\cos^4\theta - a^6\cos^6\theta\right), \qquad \Sigma = r^2 + a^2\cos^2\theta. \tag{1.10}$$

which is perfectly finite at the horizons $r = r_{\pm}$. The only genuine (curvature) singularity sits at $\Sigma = 0$, i.e. at the ring $r = 0$, $\theta = \pi/2$. The pathologies of the metric components at $\Delta = 0$ are therefore purely *coordinate* effects — a fact we exploit when designing termination criteria for numerical ray tracing.

1.3 The Geodesic Equation and the Variational Principle

The trajectories of freely falling particles are extrema of the world-line action

$$S[x] = \frac{1}{2}\int g_{\mu\nu}(x)\,\dot{x}^{\mu}\dot{x}^{\nu}\,\mathrm{d}\lambda, \qquad \dot{x}^{\mu} \equiv \frac{\mathrm{d}x^{\mu}}{\mathrm{d}\lambda}. \tag{1.11}$$

Varying S with fixed endpoints and integrating by parts yields the Euler–Lagrange equations

$$\frac{\mathrm{d}}{\mathrm{d}\lambda}\left(g_{\mu\nu}\dot{x}^{\nu}\right) - \frac{1}{2}\,\partial_{\mu}g_{\alpha\beta}\,\dot{x}^{\alpha}\dot{x}^{\beta} = 0. \tag{1.12}$$

Expanding the total derivative, symmetrizing the connection coefficients via Eq. (1.3), and raising the index with $g^{\mu\nu}$ produces the *geodesic equation*

$$\ddot{x}^{\mu} + \Gamma^{\mu}_{\alpha\beta}\,\dot{x}^{\alpha}\dot{x}^{\beta} = 0. \tag{1.13}$$

The quadratic nonlinearity in the velocities is the mathematical signature of gravity self-interacting: a geodesic is *not* a straight line in any fixed background, but the straightest possible path on a curved manifold.

The parameter λ is called *affine* if Eq. (1.13) holds with vanishing right-hand side. Under a reparametrization $\lambda \to \lambda' = f(\lambda)$ the equation acquires a friction-like term,

$$\ddot{x}^{\mu} + \Gamma^{\mu}_{\alpha\beta}\,\dot{x}^{\alpha}\dot{x}^{\beta} = \kappa(\lambda)\,\dot{x}^{\mu}, \qquad \kappa = -\frac{f''(\lambda)}{\left[f'(\lambda)\right]^{2}}, \tag{1.14}$$

which can be removed iff f is affine, $f'' = 0$. The remaining freedom is a global rescaling $\lambda \to c\,\lambda$, which rescales $\dot{x}^{\mu} \to \dot{x}^{\mu}/c$ but leaves the *orbit* in $\mathcal{M}$ invariant. The normalization $g_{\mu\nu}\dot{x}^{\mu}\dot{x}^{\nu} = \kappa_0$ with $\kappa_0 \in \{-1, 0\}$ for timelike/null motion fixes this residual freedom: $\kappa_0 = -1$ selects proper time for massive particles, while for photons the condition is homogeneous,

$$g_{\mu\nu}\,\dot{x}^{\mu}\dot{x}^{\nu} = 0, \tag{1.15}$$

and every rescaling $\lambda \to c\lambda$ remains admissible. This degeneracy has an important consequence: for null geodesics the conserved quantities

$$b \equiv \frac{L_z}{E}, \qquad \bar{q} \equiv \frac{Q}{E^{2}}, \tag{1.16}$$

are invariant under the rescaling and thus label the orbit itself, independent of parametrization. They are the natural initial data for ray tracing: a photon is fully determined by its emission point plus the pair $(b, \bar{q})$, and the entire question of whether it is captured by the black hole or escapes — i.e. the shadow — is a question about the partition of the $(b, \bar{q})$ plane.

1.3.1 Killing's Equation and Exact Conservation Laws

A vector field ξ^{μ} is a *Killing vector* of the geometry if the metric is invariant along its flow,

$$\mathcal{L}_{\xi}\,g_{\mu\nu} = \nabla_{\mu}\xi_{\nu} + \nabla_{\nu}\xi_{\mu} = 0. \tag{1.17}$$

Killing vectors generate isometries; in maximally symmetric spaces (Minkowski, de Sitter) there are $n(n+1)/2 = 10$ of them in four dimensions, while a generic stationary, axisymmetric spacetime possesses exactly two: the timelike $\xi^{\alpha} = \partial_t$ and the axial $\eta^{\alpha} = \partial_{\phi}$. The dynamical relevance of Killing vectors is the following exact conservation theorem.

If ξ^{μ} is a Killing vector and u^{μ} the tangent of an affinely parametrized geodesic, then $\xi_{\mu}u^{\mu}$ is constant along the geodesic.

This follows by using metric compatibility and the geodesic equation,

$$\frac{\mathrm{d}}{\mathrm{d}\lambda}\left(\xi_\mu u^\mu\right) = u^\nu \nabla_\nu \left(\xi_\mu u^\mu\right) = u^\nu u^\mu \nabla_\nu \xi_\mu + \xi_\mu \underbrace{u^\nu \nabla_\nu u^\mu}_{=0} = u^\mu u^\nu \nabla_{(\mu}\xi_{\nu)} = 0, \tag{1.18}$$

because $u^\mu u^\nu$ is symmetric in (μ, ν) while only the antisymmetric part of $\nabla_\mu \xi_\nu$ survives the contraction.

Applied to ξ^α and η^α, this reproduces precisely the constants E and L_z defined previously. In spherically symmetric spacetimes the three rotational Killing vectors provide L_z and two further constants (the total angular momentum squared); in Kerr, spherical symmetry is broken, only two Killing vectors survive, and yet the geodesic system remains integrable, by virtue of a hidden symmetry discussed below.

1.4 Stationary, Axisymmetric Spacetimes and the Hamiltonian Formulation

In this thesis, we focus on stationary and axisymmetric spacetimes. Such spacetimes admit two Killing vector fields. The first is the axial η^α, which is spacelike and associated with rotations about the symmetry axis. The second is the timelike ξ^α, associated with time translations. This practically means that a stationary and axisymmetric spacetime does not change with time and looks the same after rotations about the axis.

The metric of such a spacetime has generally the form:

$$ds^2 = g_{tt}dt^2 + g_{rr}dr^2 + g_{\theta\theta}d\theta^2 + 2g_{t\phi}dtd\phi + g_{\phi\phi}d\phi^2. \tag{1.19}$$

We define the quantity $\mathcal{D} \equiv g_{t\phi}^2 - g_{tt}g_{\phi\phi}$, which will prove to be crucial.

The two primary Killing vector fields correspond to two invariant quantities, the energy E and the axial angular momentum L_z:

$$E \equiv -\xi^\alpha u_\alpha = -\left(g_{tt}\frac{dt}{d\lambda} + g_{t\phi}\frac{d\phi}{d\lambda}\right), \tag{1.20}$$

$$L_z \equiv \eta^\alpha u_\alpha = g_{t\phi}\frac{dt}{d\lambda} + g_{\phi\phi}\frac{d\phi}{d\lambda}, \tag{1.21}$$

where λ is the affine parameter. Solving these equations for the coordinate velocities $\dot{t}$ and $\dot{\phi}$, we obtain:

$$\dot{t} = \frac{Eg_{\phi\phi} + L_z g_{t\phi}}{\mathcal{D}}, \tag{1.22}$$

$$\dot{\phi} = -\frac{L_z g_{tt} + Eg_{t\phi}}{\mathcal{D}}. \tag{1.23}$$

Using the Lagrangian $\mathcal{L} = \frac{1}{2}g_{\alpha\beta}\dot{x}^\alpha\dot{x}^\beta$, the associated generalized momenta are $p_\alpha = \partial\mathcal{L}/\partial\dot{x}^\alpha$. Performing a Legendre transformation, the Hamiltonian is:

$$\mathcal{H} = \frac{1}{2}\left(\frac{p_r^2}{g_{rr}} + \frac{p_\theta^2}{g_{\theta\theta}} + V_{\text{eff}}\right), \tag{1.24}$$

where the effective potential V_{eff} governs the allowed regions of motion:

$$V_{\text{eff}} = -\frac{L_z^2 g_{tt} + 2EL_z g_{t\phi} + E^2 g_{\phi\phi}}{\mathcal{D}}. \tag{1.25}$$

The contour $V_{\text{eff}} = 0$ marks the region of allowed motion (the separatrix). For massless photons, the Hamiltonian vanishes identically along the trajectory ($\mathcal{H} = 0 \iff g_{\mu\nu}\dot{x}^\mu\dot{x}^\nu = 0$).

For a deeper understanding of the allowed regions and the structure of light rings, it is useful to factor the effective potential in terms of the impact parameter $b \equiv L_z/E$:

$$V_{\text{eff}} = -\frac{E^2 g_{tt}}{\mathcal{D}}(b - h_+)(b - h_-), \tag{1.26}$$

where the critical potentials $h_\pm$ are defined as:

$$h_\pm(r, \theta) = \frac{-g_{t\phi} \pm \sqrt{\mathcal{D}}}{g_{tt}}. \tag{1.27}$$

The functions $h_\pm$ play a central role in determining the existence and stability of circular photon orbits, as they represent the boundaries where the effective potential vanishes.

1.5 The Kerr Geometry

The exterior geometry of a rotating, uncharged black hole is uniquely determined by its mass M and spin angular momentum J (No-Hair Theorem) [3]. In standard Boyer-Lindquist coordinates (t, r, θ, ϕ) [10, 2], the Kerr metric takes the explicit form:

$$ds^2 = -\left(1 - \frac{2Mr}{\Sigma}\right)dt^2 - \frac{4Mar\sin^2\theta}{\Sigma}dtd\phi + \frac{\Sigma}{\Delta}dr^2 + \Sigma d\theta^2 + \left(r^2 + a^2 + \frac{2Ma^2 r\sin^2\theta}{\Sigma}\right)\sin^2\theta d\phi^2, \tag{1.28}$$

where the fundamental metric functions are defined by:

$$\Sigma(r, \theta) \equiv r^2 + a^2\cos^2\theta, \tag{1.29}$$

$$\Delta(r) \equiv r^2 - 2Mr + a^2 = (r - r_-)(r - r_+). \tag{1.30}$$

Here, $a \equiv J/M$ is the Kerr spin parameter, with the physical sub-extremal bound $0 \leq |a| \leq M$. When $a = 0$, the metric reduces to the spherically symmetric Schwarzschild solution.

1.5.1 Horizon Structure and Ergosphere

The event horizons are located at the coordinate singularities where $\Delta = 0$, yielding:

$$r_\pm = M \pm \sqrt{M^2 - a^2}. \tag{1.31}$$

The outer boundary r_+ represents the absolute event horizon, while r_- is the inner Cauchy horizon. The surface where $g_{tt} = 0$ defines the ergosurface:

$$r_E(\theta) = M + \sqrt{M^2 - a^2\cos^2\theta}. \tag{1.32}$$

In the bounded region between the event horizon and the ergosurface (the ergosphere), $g_{tt} > 0$. In this region, no physical particle can remain static with respect to an observer at infinity. This gives rise to the phenomenon of *frame-dragging*, where the spacetime itself is dragged along the rotation direction.

1.5.2 Null Geodesics and the Carter Constant

Due to the presence of the Carter constant $\mathcal{Q}$, the Kerr spacetime represents a completely integrable system. Using the Hamilton-Jacobi formulation, we assume an additively separable ansatz for the principal

action function, which allows the decoupling of the radial and polar dependencies. The system of first-order geodesic equations for a photon is given by:

$$\Sigma \frac{dr}{d\lambda} = \pm\sqrt{\mathcal{R}(r)}, \tag{1.33}$$

$$\Sigma \frac{d\theta}{d\lambda} = \pm\sqrt{\Theta(\theta)}, \tag{1.34}$$

$$\Sigma \frac{d\phi}{d\lambda} = -\left(aE - \frac{L_z}{\sin^2\theta}\right) + \frac{a}{\Delta}P, \tag{1.35}$$

$$\Sigma \frac{dt}{d\lambda} = -a\left(aE\sin^2\theta - L_z\right) + \frac{r^2+a^2}{\Delta}P, \tag{1.36}$$

where $P = E(r^2 + a^2) - L_z a$, and the radial and polar potentials are defined as:

$$\mathcal{R}(r) = \left[E(r^2+a^2) - aL_z\right]^2 - \Delta\left[\mathcal{Q} + (L_z - aE)^2\right], \tag{1.37}$$

$$\Theta(\theta) = \mathcal{Q} - \cos^2\theta\left[-a^2E^2 + \frac{L_z^2}{\sin^2\theta}\right]. \tag{1.38}$$

For numerical ray tracing, it is often more convenient to reformulate the problem using the normalized impact parameters:

$$b \equiv -\frac{p_\phi}{p_t} = \frac{L_z}{E}, \quad \text{and} \quad \alpha = \frac{p_\theta}{p_t}. \tag{1.39}$$

These parameters uniquely define the initial conditions for a photon at infinity.

1.6 Light Rings and the Mathematical Shadow

Bound null geodesics that remain confined to a constant radial coordinate $r = r_{\rm ph}$ without escaping to infinity or plunging into the event horizon satisfy the simultaneous orbital conditions:

$$\mathcal{R}(r) = 0 \quad \text{and} \quad \frac{d\mathcal{R}}{dr}(r) = 0. \tag{1.40}$$

This can be recast in terms of the effective potentials $h_\pm$ introduced in Section 1.4. The light rings (unstable circular photon orbits) correspond to the extrema or saddle points of the $h_\pm$ functions. For the Kerr metric, the radius of the prograde and retrograde photon rings, as well as the corresponding critical impact parameters b_c, can be derived analytically. The locus of these critical impact parameters on the observer's plane defines the mathematical boundary of the black hole shadow [15, 6, 7].

In the absence of a Carter constant (i.e., in non-Kerr metrics), this analytical formulation breaks down. Therefore, to compute the shadow and its self-similar structures, we must resort to extensive numerical integration of the geodesic equations, which we will describe in detail in the following chapter.

1.7 Strong-Field Gravitational Lensing and Multiple Imaging

While weak-field gravitational lensing results in minor image distortions and small deflection angles, the strong gravitational field in the immediate vicinity of a black hole fundamentally alters the optical geometry of the spacetime. The key feature governing this regime is the existence of the photon sphere, which acts as a critical boundary in the phase space of the geometry.

As a photon's trajectory approaches the critical impact parameter associated with this sphere, the deflection angle does not merely increase linearly; it diverges logarithmically. Physically, this means that photons

passing very close to this critical threshold do not just bend slightly-they can execute partial or even complete orbits around the black hole before eventually escaping to the observer. This logarithmic divergence is the fundamental mechanism responsible for the creation of multiple images from a single source.

Because the deflection angle can exceed 2π, a single source can produce an infinite sequence of images. A photon can orbit the black hole n times before escaping. We define the winding number n as the number of half-orbits the photon completes around the black hole. The primary image ($n = 0$) and the secondary image ($n = 1$) are formed by photons that pass the black hole on either side without completing a full loop. Photons with $n \geq 2$ correspond to higher-order images that have looped around the black hole multiple times.

As the winding number increases, the required impact parameter converges exponentially to the critical value. Consequently, the higher-order images are located exponentially closer to the mathematical shadow boundary, forming a self-similar, infinite hierarchy of photon sub-rings. Furthermore, near the critical curve, the magnification factor diverges. In practice, for a source of finite size, this extreme magnification stretches the source tangentially along the critical curve, transforming a point-like or spherical source into highly magnified, thin crescent arcs.

Capturing this infinite hierarchy of sub-rings and the extreme tangential stretching requires exceptional numerical precision. Standard integrators accumulate phase-space errors that smear out the higher-order images, rendering them invisible. This theoretical expectation motivates the implementation of the geometric Manifold Projection detailed in Chapter 2, ensuring that the numerical resolution matches the theoretical predictions of strong-field lensing.

1.8 Radiative Transfer Formulation

The propagation of radiation through the curved spacetime around a black hole is governed by the radiative transfer equation. In the geometric optics approximation, the Lorentz-invariant specific intensity $\mathcal{I} = I_\nu/\nu^3$ evolves along each photon geodesic according to

$$\frac{d\mathcal{I}}{d\lambda} = \gamma^{-1}\frac{j_0}{\nu_0^3}, \tag{1.41}$$

where λ is the affine parameter along the null geodesic, j_0 is the emissivity measured in the local fluid rest frame, ν_0 is the photon frequency in the rest frame, and γ is the redshift factor defined as

$$\gamma \equiv \frac{\nu}{\nu_0}, \tag{1.42}$$

with ν being the frequency measured by a distant static observer. In this formulation, we assume an optically thin medium with zero absorption ($\alpha_\nu = 0$).

To obtain the observable specific intensity I_ν at the detector, we transform the invariant quantity back to the observer's frame using $I_\nu = \mathcal{I}\,\nu_{\rm obs}^3$. The contribution to the intensity from a differential segment $d\lambda$ of the geodesic is then

$$\begin{aligned} dI = d\mathcal{I}\ \nu_{\rm obs}^3 &= \gamma^{-1}\frac{j_0}{\nu_0^3}\nu_{\rm obs}^3\, d\lambda \\ &= \gamma^2 j_0\, d\lambda, \end{aligned} \tag{1.43}$$

where we have used $\nu_{\rm obs}/\nu_0 = \gamma$. This transformation must be applied at each point along the geodesic before accumulating the intensity, since the observed intensity is affected by the relativistic effects encountered when transitioning from the emission frame to the observer's frame.

The total specific intensity at frequency ν is obtained by integrating along the entire photon trajectory:

$$I_\nu = \int_{\rm geodesic} \gamma^2 j_0 \, d\lambda. \tag{1.44}$$

To construct direct intensity images, we compute the frequency-integrated intensity by summing contributions across all frequency bins:

$$I = \int I_\nu \, d\nu \approx \sum_i I_{\nu_i} \Delta\nu_i, \tag{1.45}$$

where I_{ν_i} is the specific intensity at frequency ν_i and $\Delta\nu_i$ is the corresponding frequency bandwidth. This approach allows us to produce broadband images that account for the spectral distribution of the emitted radiation and the frequency-dependent relativistic effects along each line of sight.

Chapter 2

Numerical Methods and Computational Algorithms

2.1 Theoretical Background: Well-Posedness and Error in ODE Integration

2.1.1 The Initial Value Problem

The geodesic ray-tracing problem is an initial value problem (IVP)

$$\frac{\mathrm{d}\boldsymbol{y}}{\mathrm{d}\lambda} = \boldsymbol{f}(\boldsymbol{y}), \qquad \boldsymbol{y}(\lambda_0) = \boldsymbol{y}_0, \tag{2.1}$$

with $\boldsymbol{f}$ analytic wherever $\Delta \neq 0$. The Picard–Lindelöf theorem then guarantees existence and uniqueness of the solution, and continuous dependence on the initial data: perturbations grow at most like $e^{L|\Delta\lambda|}$, with L a Lipschitz constant of $\boldsymbol{f}$. Away from the photon region $L = O(M/r^3)$ there is no problem, but near the unstable photon orbits the effective sensitivity is exponentially large: two initially adjacent rays separate by $e^{\gamma\Delta\lambda}$ with γ the Lyapunov exponent. Machine-precision initialization and integration are thus necessities for resolving the photon-ring structure.

2.1.2 Local Error, Global Error, and Convergence

A one-step integrator has *order* p if its local error (one step started on the exact solution) satisfies $\|\boldsymbol{d}_n\| = O(h^{p+1})$. Under standard stability assumptions the global error after $N = T/h$ steps accumulates coherently to

$$\|\boldsymbol{e}_N\| = O(h^p), \tag{2.2}$$

a gain of one power of h over the local error, because $N \propto 1/h$ errors partially cancel for stable problems. Consistency (small local error) plus stability (bounded error propagation) is equivalent to convergence, the discrete analogue of the Lax equivalence principle.

2.1.3 Truncation Error, Round-off, and the Accuracy Floor

On a finite-precision machine the total error behaves as

$$E(h) \;\approx\; C_1\, h^p \;+\; C_2\, \frac{\epsilon_{\mathrm{mach}}}{h}, \tag{2.3}$$

where the first term is truncation error and the second is round-off accumulated over $\sim 1/h$ steps. Minimizing (2.3) gives an optimal step $h_* \propto (\epsilon_{\text{mach}})^{1/(p+1)}$ and an irreducible accuracy floor

$$E_{\min} = O\left(\epsilon_{\text{mach}}^{p/(p+1)}\right) \approx 10^{-14} \quad \text{for } p = 7 \text{ in double precision.} \tag{2.4}$$

This has a design consequence that is often misunderstood: requesting time-integration tolerances far below $\sim 10^{-14}$ buys no further trajectory accuracy, because round-off dominates. Exact conservation of the invariants ($g_{\mu\nu}u^\mu u^\nu = 0$, E, L_z) is therefore not to be won by tightening the integrator but by exploiting properties of the system under consideration.

2.2 Automatic Differentiation vs. Finite Differences

A critical requirement for evaluating the acceleration vector $\boldsymbol{a} = -\Gamma^\mu_{\alpha\beta}\, u^\alpha u^\beta$ is the precise computation of the metric tensor derivatives $\partial_\sigma g_{\mu\nu}$. Traditional implementations often rely on numerical finite differences:

$$\partial_\sigma g_{\mu\nu}(\boldsymbol{x}) \approx \frac{g_{\mu\nu}(\boldsymbol{x} + h\boldsymbol{e}_\sigma) - g_{\mu\nu}(\boldsymbol{x} - h\boldsymbol{e}_\sigma)}{2h}. \tag{2.5}$$

This approach introduces two fundamental sources of error: a truncation error $O(h^2)$ and a round-off error $O(\epsilon_{\text{mach}}/h)$. This results in the characteristic "U"-shaped error curve, where the optimal step size must be delicately balanced. Crucially, this approach is also metric-agnostic, allowing us to deal with any metric (whether it admits a Carter constant or not) the same way without any extra cost.

2.2.1 Dual Numbers and the Chain Rule

To completely eliminate these numerical vulnerabilities, we implement the derivatives using Forward-Mode Automatic Differentiation [17, 8, 13]. This method relies on the algebraic properties of dual numbers. We define the ring of dual numbers $\mathbb{D}$ such that an element is represented as $a + b\epsilon$, where ϵ satisfies the fundamental property:

$$\epsilon^2 = 0, \quad \text{with} \quad \epsilon \neq 0. \tag{2.6}$$

By extending a function $f(x)$ to the dual numbers, we can evaluate its Taylor expansion at $x + \epsilon$:

$$f(x + \epsilon) = f(x) + f'(x)\epsilon + \frac{1}{2}f''(x)\epsilon^2 + \dots \tag{2.7}$$

Because of the defining property $\epsilon^2 = 0$, all higher-order terms vanish identically. Therefore, by evaluating the function with the dual seed $x + \epsilon\boldsymbol{v}$, we obtain, in a single forward pass, both the function value and its exact directional derivative:

$$f(x + \epsilon\boldsymbol{v}) = f(x) + \epsilon\left(\nabla f(x) \cdot \boldsymbol{v}\right). \tag{2.8}$$

Crucially, AD applies the chain rule to every elementary operation performed during the evaluation. Therefore, there is no step size to tune, no truncation error, and no round-off cancellation; the result is the exact derivative to machine precision ($\sim 10^{-16}$). Notably, because this process is entirely algebraic and metric-agnostic, we can automatically compute the Christoffel symbols for the Boyer-Lindquist metric without ever manually deriving their analytic form.




Figure 2.1: Comparison of the error in computing derivatives using finite differences (red, U-shaped) and automatic differentiation (blue, flat at machine precision).

2.3 Geometric Integration via Manifold Projection

2.3.1 The Problem of Invariant Drift

Standard explicit Runge-Kutta integrators are non-geometric in nature: they do not preserve the algebraic invariants of the phase space. Along a physical null geodesic, the state vector $\boldsymbol{y}$ must remain strictly confined to a sub-manifold $\mathcal{N}$ embedded in the 8-dimensional phase space. This sub-manifold is defined by the simultaneous vanishing of three invariant constraint functions:

1. **The Lagrangian Null Condition:** $g_1(\boldsymbol{y}) \equiv g_{\mu\nu}(\boldsymbol{x})u^\mu u^\nu = 0$.
2. **Conservation of Energy:** $g_2(\boldsymbol{y}) \equiv -(g_{tt}(\boldsymbol{x})u^t + g_{t\phi}(\boldsymbol{x})u^\phi) - E = 0$.
3. **Conservation of Axial Angular Momentum:** $g_3(\boldsymbol{y}) \equiv g_{\phi\phi}(\boldsymbol{x})u^\phi + g_{t\phi}(\boldsymbol{x})u^t - L_z = 0$.

During long-term propagation, accumulated floating-point errors drive the unconstrained solution $\tilde{\boldsymbol{y}}$ off this sub-manifold. In strong gravitational environments, even microscopic deviations trigger exponential trajectory divergence.

2.3.2 Projection via Lagrange Multipliers

To suppress this drift, we implement the Manifold Projection method as described in [9]. Let $\tilde{\boldsymbol{y}}$ denote the provisional state vector computed by the solver at step $n+1$. We seek a corrected state $\boldsymbol{y}^*$ that satisfies the zero-level constraint equations. Using the method of Lagrange multipliers, we construct the auxiliary objective functional:

$$\mathcal{S}(\boldsymbol{y}^*, \boldsymbol{\lambda}) = \frac{1}{2}\|\boldsymbol{y}^* - \tilde{\boldsymbol{y}}\|^2 + \boldsymbol{\lambda}^T \boldsymbol{g}(\boldsymbol{y}^*), \tag{2.9}$$

Taking the functional gradient with respect to the state vector $\boldsymbol{y}^*$, we obtain:

$$\boldsymbol{y}^* = \tilde{\boldsymbol{y}} + \boldsymbol{J}(\boldsymbol{y}^*)^T \boldsymbol{\lambda}, \tag{2.10}$$

where $\boldsymbol{J}(\boldsymbol{y}) \equiv \frac{\partial \boldsymbol{g}}{\partial \boldsymbol{y}}$ is the constraint Jacobian matrix. This equation possesses profound geometric meaning: the correction vector $\delta\boldsymbol{y} = \boldsymbol{y}^* - \tilde{\boldsymbol{y}} = -\boldsymbol{J}^T\boldsymbol{\lambda}$ lies entirely within the subspace normal to the constraint sub-manifold. Thus, the projection alters the numerical state strictly along directions orthogonal to the physical invariant surface, ensuring the trajectory is not dissipated or artificially damped.

2.3.3 Multidimensional Newton-Raphson Iterative Solver

Substituting the projection equation into the constraint condition yields a non-linear algebraic system for the unknown multipliers $\boldsymbol{\lambda}$. To solve this efficiently, we employ a multidimensional Newton-Raphson scheme. Expanding $\boldsymbol{g}$ in a first-order Taylor series, we define the symmetric Gramian matrix $\boldsymbol{G} \equiv \boldsymbol{J}\boldsymbol{J}^T \in \mathbb{R}^{3\times 3}$. The iterative Newton increment $\Delta\boldsymbol{\lambda}^{(k)}$ is computed via:

$$\boldsymbol{G}\,\Delta\boldsymbol{\lambda}^{(k)} = \boldsymbol{g}(\boldsymbol{y}^{(k)}) \quad \Longrightarrow \quad \Delta\boldsymbol{\lambda}^{(k)} = \boldsymbol{G}^{-1}\boldsymbol{g}(\boldsymbol{y}^{(k)}). \tag{2.11}$$

The step-by-step execution flow of the Manifold Projection proceeds as follows:

1. **Initialization:** Set $\boldsymbol{\lambda}^{(0)} = 0$ and $\boldsymbol{y}^{(0)} = \tilde{\boldsymbol{y}}$. Evaluate $\boldsymbol{J}$ using the exact forward-mode AD introduced in Section 2.3.

2. **Iterative Correction Loop:** For $k = 1, 2, \ldots$:

$$\boldsymbol{u}^{(k)} = \tilde{\boldsymbol{y}} + \boldsymbol{J}^T\boldsymbol{\lambda}^{(k)}, \tag{2.12}$$
$$\boldsymbol{R}^{(k)} = \boldsymbol{g}(\boldsymbol{u}^{(k)}), \tag{2.13}$$
$$\Delta\boldsymbol{\lambda}^{(k)} = -\boldsymbol{G}^{-1}\boldsymbol{R}^{(k)}, \tag{2.14}$$
$$\boldsymbol{\lambda}^{(k+1)} = \boldsymbol{\lambda}^{(k)} + \Delta\boldsymbol{\lambda}^{(k)}. \tag{2.15}$$

3. **Convergence Criterion:** Terminate the loop when the increment norm satisfies:

$$\|\Delta\boldsymbol{\lambda}^{(k)}\| < \text{tol}, \tag{2.16}$$

   where we set the tolerance to a constant value equal to the machine precision limit, tol = 10^{-16}.

   > *We note that while a step-dependent tolerance, tol = $c \cdot \Delta t^{p+1}$, could be adopted to reduce the number of Newton-Raphson iterations during large integration steps, we have deliberately chosen a fixed, extremely strict tolerance of 10^{-16}. This choice does not alter the underlying physics or the mathematical formulation of the projection, but ensures that the state vector is re-projected onto the constraint sub-manifold with the absolute maximum accuracy achievable by the machine, regardless of the local step size. Consequently, the only cost incurred is an increase in computational power, which is negligible compared to the total ODE integration time, while completely guaranteeing the conservation of all physical quantities.*

   To safeguard against potential infinite loops in the event of numerical singularities (e.g., extremely close to the Boyer-Lindquist coordinate singularity), we additionally set a maximum limit of 500 iterations, possibly sacrificing some accuracy.

4. **State Replacement:** Overwrite the integrator state with $\boldsymbol{y}^*$ and evaluate the termination triggers.

2.3.4 Convergence of the Newton–Raphson Iteration

The projection reduces to solving $\boldsymbol{F}(\boldsymbol{\lambda}) \equiv \boldsymbol{g}\big(\tilde{\boldsymbol{y}} + \boldsymbol{J}^T\boldsymbol{\lambda}\big) = 0$, whose exact Jacobian is $\boldsymbol{F}' = \boldsymbol{J}\boldsymbol{J}^T \equiv \boldsymbol{G}$. The multidimensional Newton–Raphson method enjoys the following classical guarantee.

Theorem (quadratic convergence). *Let $\boldsymbol{F} \in C^2$, $\boldsymbol{F}(\boldsymbol{\lambda}_*) = 0$, and $\boldsymbol{G}(\boldsymbol{\lambda}_*)$ nonsingular. Then for $\boldsymbol{\lambda}_0$ sufficiently close to $\boldsymbol{\lambda}_*$ the iteration converges and*

$$\|\boldsymbol{\lambda}_{k+1} - \boldsymbol{\lambda}_*\| \;\le\; \frac{M_2}{2\,\sigma_{\min}}\,\|\boldsymbol{\lambda}_k - \boldsymbol{\lambda}_*\|^2, \tag{2.17}$$

where M_2 bounds $\|\boldsymbol{F}''\|$ and $\sigma_{\min}$ is the smallest singular value of $\boldsymbol{G}(\boldsymbol{\lambda}_*)$.

The proof is a second-order Taylor expansion of $\boldsymbol{F}(\boldsymbol{\lambda}_k)$ about $\boldsymbol{\lambda}_*$: the linear terms cancel exactly by construction of the Newton step, leaving a residual quadratic in the error — the number of correct digits roughly doubles per iteration, which is why the observed iteration counts are tiny whenever $\tilde{\boldsymbol{y}}$ is already close to the constraint manifold (i.e. always, when the integrator is accurate). Two practical refinements have a precise theory. First, *freezing* the Jacobian ($\tilde{\boldsymbol{G}} = \boldsymbol{G}$ held fixed) yields the chord method, which converges only linearly, with rate $\rho = \|\boldsymbol{I} - \tilde{\boldsymbol{G}}^{-1}\boldsymbol{G}\|$; periodically recomputing $\boldsymbol{G}$ restores the full quadratic rate. This is the exact trade-off exploited when the Jacobian refresh is amortized over several iterations: the projection is dominated by the cost of the constraint Jacobian, and the refresh cadence converts a per-iteration cost into a per-K-iteration cost at negligible loss of convergence speed. Second, since $\boldsymbol{G}$ is symmetric positive semidefinite, it is positive definite — and the linear system uniquely solvable — iff $\boldsymbol{J}$ has full row rank $m = 3$.

2.3.5 Regularity of the Constraints

The Lagrange construction presumes that the three constraint gradients $\partial_{\boldsymbol{y}} g_1, \partial_{\boldsymbol{y}} g_2, \partial_{\boldsymbol{y}} g_3$ are linearly independent on a neighbourhood of the constraint manifold $\mathcal{N}$. This is generically the case: $\mathrm{d}E$ and $\mathrm{d}L_z$ are the contractions with the mutually independent Killing covectors ξ_μ and η_μ, and the null constraint gradient $g_{\mu\nu}u^\nu$ is not a linear combination of them at any physical point of phase space. By the implicit function theorem the projection is then locally unique and depends smoothly on $\tilde{\boldsymbol{y}}$. Near $r = r_+$ the metric derivatives entering $\boldsymbol{J}$ diverge as $1/\Delta$; the constraints remain regular (their defining functions stay finite) but the Newton system becomes ill-conditioned, which is why a maximum-iteration safeguard accompanies the solver. This degradation is confined to a region from which no observable photon escapes, so it cannot contaminate the image.

2.3.6 Projection Versus Alternative Stabilization Strategies

Projection belongs to the broader family of *geometric integrators*, which are designed to respect structure of the flow rather than only its pointwise accuracy. The alternatives are instructive to contrast:

1. **Constraint damping (Baumgarte stabilization):** augment the vector field with terms that exponentially pull the solution back to $\mathcal{N}$. This modifies the equations actually integrated and introduces a free damping time scale; the invariants are restored only approximately and at the price of systematically perturbing the dynamics.

2. **Phase-space reduction:** eliminate degrees of freedom by solving the constraints analytically (e.g. determining u^t from the null condition). Exact by construction, but the reduced system loses the symmetries that make adaptive control and error estimation uniform across the phase space, and it is metric-specific — precisely what a general-purpose tracer must avoid.

3. **Symplectic integrators:** for Hamiltonian systems they conserve a modified Hamiltonian over exponentially long times (backward error analysis), giving superb long-term energy behaviour. However, geodesic Hamiltonians are non-separable in general coordinates (forcing implicit methods), adaptive step sizes destroy the symplectic property, and symplecticity preserves a *nearby* energy rather than the exact invariants E, L_z demanded here.

Projection is distinguished by three properties that make it the method of choice for ray tracing: it is *exact* (the invariants hold to machine precision at every accepted step, with no secular drift), *unbiased* (the correction lies entirely in the normal space to $\mathcal{N}$, so the tangential, physical dynamics is that of the underlying integrator to its full order, no order reduction occurs), and *parameter-free* (no damping constants, no reduced coordinates). For reversible problems one may additionally symmetrize the projection to inherit long-term reversibility; for the present purpose the standard projection suffices, since the observables are fixed by the invariant values and the tangential accuracy, both of which are controlled as described above.

2.4 High-Order Adaptive Integration

2.4.1 Butcher Structure and Order Conditions

An explicit s-stage Runge–Kutta method is specified by its Butcher tableau:

$$\boldsymbol{k}_i = f\left(\boldsymbol{y}_n + h\sum_{j=1}^{i-1} a_{ij}\,\boldsymbol{k}_j\right), \qquad \boldsymbol{y}_{n+1} = \boldsymbol{y}_n + h\sum_{i=1}^{s} b_i\,\boldsymbol{k}_i. \tag{2.18}$$

Matching the Taylor expansion of $\boldsymbol{y}_{n+1}$ with that of the exact solution through order p imposes a set of algebraic *order conditions*, one per rooted tree; their number grows as $1, 2, 4, 8, 17, 37, 78, 167$ for $p = 1, \ldots, 8$. The combinatorial explosion explains both why very high orders require many stages and why the construction of efficient coefficients (as in Verner's embedded pairs) is a non-trivial numerical-analysis achievement. For geodesic integration the right-hand side is analytic (away from $\Delta = 0$), so high order pays off without the order-reduction pathologies familiar from stiff or non-smooth problems.

2.4.2 Embedded Pairs and Local Error Estimation

Practical step-size control requires an error estimate per step. An *embedded pair* shares the internal stages $\boldsymbol{k}_i$ between two solutions of orders p and $p-1$,

$$\boldsymbol{y}_{n+1} = \boldsymbol{y}_n + h\sum_i b_i\boldsymbol{k}_i, \qquad \hat{\boldsymbol{y}}_{n+1} = \boldsymbol{y}_n + h\sum_i \hat{b}_i\boldsymbol{k}_i, \qquad \boldsymbol{e}_{n+1} = \boldsymbol{y}_{n+1} - \hat{\boldsymbol{y}}_{n+1} \approx O(h^p), \tag{2.19}$$

so the difference $\boldsymbol{e}_{n+1}$ estimates the error of the *lower*-order result at negligible extra cost, and the higher-order result is propagated (local extrapolation). The scalar error measure is taken in a component-wise weighted norm,

$$\mathrm{err} = \max_i \frac{\left|(\boldsymbol{e}_{n+1})_i\right|}{\mathrm{atol} + \mathrm{rtol}\cdot\max(|y_i|,\,|\hat{y}_i|)}, \tag{2.20}$$

which makes the tolerance scale-invariant across state components of disparate magnitude (here: coordinates versus four-velocity components).

A Note on Computational Units

Throughout the computational implementation and the visualization of the results, we adopt a system of dimensionless geometrized units. Specifically, we set the gravitational constant, the speed of light, and the black hole mass to unity ($G = c = M = 1$). Consequently, all spatial distances, including the affine parameter and impact parameters, are expressed in units of M, while time is measured in units of GM/c^3. This convention simplifies the numerical integration and scales the phase-space variables naturally.

Chapter 3

Setup

Having established the theoretical framework of stationary axisymmetric spacetimes and the geodesic equations governing null trajectories, we now proceed to describe the algorithms deployed to perform the black hole imaging.

3.1 Backward Ray Tracing and Initial Conditions

To generate images of the shadow and the surrounding strongly lensed region, we adopt a backward ray-tracing paradigm. Unlike the standard forward approach, where photons are emitted from a source and tracked towards an observer, backward ray-tracing treats each pixel of the observer's screen as an independent initial condition for a null geodesic. This is possible due to the fact that, in the absence of scattering, light rays are reversible.

3.1.1 Choice of Coordinate System

We begin by defining the observer's position. In our framework, we work in any coordinates spherical (t, r, θ, ϕ) or Cartesian t, x, y, z, since our methodology allows for any metric expressed in any coordinate system. Our AD machinery alongside the geometric high-order integration scheme we use allow us to do so. Thus, our testbed became the Kerr black hole since it allows both coordinate expression (Boyer-Lindquist and Kerr-Schild) as well as the static Schwarzschild counterpart by setting $a = 0$.

In every case, there exists an event horizon which we need to deal with numerically. Therefore, our integrations are terminated at a radial distance of approximately $r \approx 1.15M$ to $2M$ (depending on the case), ensuring that the physical quantities remain well-defined and machine-precision accuracy is strictly maintained before the coordinate breakdown occurs.

3.1.2 Grid Construction and Initial Conditions

The observer is assumed to be located at a fixed radial distance $r_0 \gg M$ and at an inclination angle θ_0 relative to the black hole's spin axis. The initial position vector is given by:

$$\boldsymbol{x}_{\text{obs}} = (t_0, r_0, \theta_0, \phi_0), \tag{3.1}$$

where we typically set $t_0 = 0$ and $\phi_0 = \pi/2$ for simplicity.

We then construct a two-dimensional grid of initial conditions $N \times M$, where N and M correspond to the number of different values for our two impact parameters β and α, respectively. These parameters define

the apparent position of the ray on the observer's celestial sphere and are linearly mapped to the grid's horizontal and vertical axes.

For each point (β_i, α_j) on this grid, we construct the initial four-velocity $\boldsymbol{u}^\mu$. To ensure that the photons are physical (massless), we enforce the null condition via the normalization of the initial four-momentum:

$$g_{\mu\nu} u^\mu u^\nu = 0. \tag{3.2}$$

In practice, this is achieved by taking an unnormalized spatial velocity vector, solving for the time component of the velocity that satisfies the null condition, and scaling it appropriately. The conserved energy E and axial angular momentum L_z are calculated directly from the metric tensor at the observer's location.

3.1.3 UV Mapping and Calibration

To generate an image, we place a virtual camera at a fixed radial distance r_0 and inclination angle θ_0. We map the impact parameters (b, α) onto the two-dimensional observer's screen, forming a grid $N \times M$ of initial conditions. For every pixel, the photon is traced backwards as described before.

For photons that escape to infinity, we map their final angular coordinates (θ, ϕ) onto an equirectangular background image. The mapping is given by:

$$u = 1 - \frac{\mod\ (\phi - \pi, 2\pi)}{2\pi}, \tag{3.3}$$

$$v = \frac{\mod\ (\theta, \pi)}{\pi}, \tag{3.4}$$

where (u, v) correspond to the normalized horizontal and vertical pixel coordinates of the background image. To ensure the correctness of this mapping and the proper handedness of the celestial sphere, we perform a rigorous calibration step using a dedicated four-colour test image (with distinct quadrants of red, green, blue, and yellow). By verifying the correct orientation of the colours in the rendered shadow, we guarantee that our subsequent simulations using realistic Milky Way backgrounds produce physically accurate gravitational lensing. However different choices of the UV mapping result in different outputs and thus conventions.




Figure 3.1: Four-colour calibration image used to verify the UV mapping and handedness of the celestial sphere projection.

Chapter 4

Results

4.1 Validation of Numerical Stability

Before presenting the physical results and the astrophysical imaging, we systematically validate the robustness of our proposed numerical integrator. The primary objective of this section is to demonstrate that the coupling of the high-order adaptive Runge-Kutta scheme with the Manifold Projection technique successfully constrains the drift of the conserved quantities to machine precision, even for highly unstable or long-lived photon trajectories.

Furthermore, we demonstrate the effectiveness of the projection algorithm through a controlled pedagogical example of circular motion. We integrate a simple harmonic oscillator constrained to a unit circle in the Euclidean plane, using a standard 4th-order Runge-Kutta scheme with a fixed step size. Without projection, the unconstrained solution doesn't preserve energy, which leads to errors compounding over time. With the implementation of the Manifold Projection, the orbit remains exactly on the circle indefinitely, illustrating the general applicability of the method. The success of this simple example confirms the mathematical robustness of the projection algorithm we have implemented for the relativistic case.




Figure 4.1: Demonstration of the Manifold Projection algorithm on a simple harmonic oscillator ($\dot{x} = y, \dot{y} = -x$) constrained to a unit circle. **Left:** The phase-space trajectory integrated using a standard 4th-order Runge-Kutta scheme (red) exhibits inward spiraling (amplitude decay) due to unconstrained numerical drift, whereas the manifold-projected RK4 (blue) remains exactly on the constraint surface. **Right:** The evolution of the radial distance over time ($t \in [0, 10000]$ with step $dt = 0.3$), explicitly showing the amplitude decay in the unconstrained method versus the exact conservation achieved by projection.

Our findings are presented in Fig. 4.1. In the absence of the geometric projection (red curve), the unconstrained integration leads to a systematic accumulation of truncation and round-off errors. While the trajectory may initially appear accurate, the errors in the conserved quantity grow significantly over time. On the contrary, the geometric integration technique (blue curve) preserves the expected behavior exactly. Furthermore, this happens even in an almost trivial right-hand side expression. This numerical drift is particularly severe in our main problem, i.e. when the photons approach the unstable photon sphere, where the highly non-linear behaviour of the metric amplifies these deviations, eventually leading to unphysical trajectories.

4.1.1 Fundamental Photon Orbits and Dynamics

Now moving on to the relativistic trajectories in the near extremal ($a = 0.998$) Kerr manifold, we monitor the evolution of the relative errors in the conserved energy $\Delta e/e_0$, the axial angular momentum $\Delta l/l_0$, and the error in the null constraint $g_{\mu\nu}u^\mu u^\nu$, along the affine parameter λ.

Figures 4.2, 4.3 and 4.4 demonstrate the effectiveness of our method, and the layout is the following

Top Left: On the top left we see the behavior of the radius as a percentage of the initial one, in order to show that there is no sudden jump (either ejection or plunge to the singularity).

Top Middle: The radius.

Top Right: The full 3D trajectory of the evolution.

Bottom Left: The relative error of the energy for each step.

Bottom Middle: The relative error of the angular momentum.

Bottom Right: The null condition error.

4.1.2 Scattering Orbits

For values of the impact parameter satisfying $|b| > b_c$, the effective potential V_{eff} prevents the photon from crossing the centrifugal barrier. The photon approaches the black hole, reaches a minimum radial distance $r_{\min} > r_{\text{ph}}$, and subsequently escapes to spatial infinity. This behaviour gives rise to the characteristic "V" shape in the radial coordinate as a function of the affine parameter, $r(\lambda)$, which we display in Fig. 4.2.

This class of orbits is particularly relevant for studies of gravitational lensing. The exact radius of closest approach, $r_{\min}$, depends sensitively on the impact parameter and the black hole's spin. By resolving these trajectories to machine precision, our framework accurately captures the strong deflection experienced by photons passing in the vicinity of the photon sphere.




Figure 4.2: Numerical integration of a scattering orbit by a near-extremal Kerr black hole ($a = 0.998, M = 1$). The initial conditions correspond to an impact parameter slightly larger than the critical value ($|b| > b_c$). **Top Row (Left to Right):** Relative radial deviation showing the absence of discontinuous jumps, the radial distance $r(\lambda)$ exhibiting the characteristic "V" shape turnaround, and the full 3D spatial trajectory. **Bottom Row (Left to Right):** Relative errors in conserved energy ($\Delta E/E_0$), axial angular momentum ($\Delta L_z/L_{z0}$), and the absolute error of the null constraint ($g_{\mu\nu}u^\mu u^\nu$). The integration was performed using a 7th-order Verner method with Manifold Projection (tolerances set to 10^{-15}), strictly bounding all errors near machine precision.

4.1.3 Bound Spiral Orbits (Transient Capture)

In the limit where the impact parameter approaches the critical value ($|b| \to b_c$), the photon can orbit the black hole multiple times in the vicinity of the unstable photon sphere before either plunging into the event horizon or escaping to infinity. These transient capture orbits are inherently unstable, characterized by an exponential sensitivity to initial conditions, governed by the Lyapunov exponent of the spacetime.

Because of this extreme sensitivity, the accurate numerical integration of these orbits represents one of the most stringent tests of our Manifold Projection algorithm. Even a minute amount of numerical drift or round-off error would artificially eject the photon from its bound trajectory. Figure 4.3 demonstrates that our algorithm maintains the invariants below the 10^{-12} threshold even after falling in the black hole, after staying in a highly unstable region, allowing us to precisely resolve the number of windings before the photon eventually escapes or is captured.




Figure 4.3: Numerical integration of a bound spiral orbit (transient capture) near the critical impact parameter ($|b| \approx b_c$) for a Kerr black hole ($a = 0.998, M = 1$). **Top Row:** Relative radial deviation, monotonic radial plunge $r(\lambda)$, and the 3D trajectory showing multiple revolutions around the unstable photon sphere before horizon penetration. **Bottom Row:** Evolution of errors in energy, angular momentum, and the null constraint. Despite the exponential sensitivity (Lyapunov divergence) in this region, the Manifold Projection algorithm maintains all invariant errors strictly below the 10^{-12} threshold until the trajectory reaches the event horizon.

4.1.4 Direct Capture Orbits

Finally, for impact parameters satisfying $|b| < b_c$, the photon lacks sufficient angular momentum to overcome the effective potential barrier. It plunges inwards, with the radial coordinate $r(\lambda)$ monotonically decreasing until it crosses the event horizon at $r = r_+$.

In our ray-tracing framework, these photons will correspond to the dark pixels of the black hole shadow. Their trajectories terminate when the radial coordinate reaches (near) the horizon, at which point the integration is halted. The robustness of the projection ensures that the energy and angular momentum remain conserved even during this rapid infall, preventing unphysical numerical oscillations near the coordinate singularity.




Figure 4.4: Numerical integration of a direct capture orbit ($|b| < b_c$) in a Kerr spacetime ($a = 0.998, M = 1$). The photon lacks sufficient angular momentum to overcome the effective potential barrier. **Top Row:** Relative radial change, the rapidly decreasing radial distance $r(\lambda)$ towards the event horizon, and the 3D plunge trajectory. **Bottom Row:** Constraint errors along the affine parameter. The physical invariants remain conserved to machine precision, avoiding unphysical numerical oscillations near the coordinate singularity.

Finally, these three fundamental trajectory classes provide the complete mapping between the observer's impact parameters and the resulting image structure. They form the underlying physical basis for the astrophysical imaging we present in the following section. Moreover, we have demonstrated that our method is capable of handling all of them indiscriminately.

4.2 Black Hole Shadow and Astrophysical Imaging

Having systematically classified and resolved the fundamental photon orbits, we now proceed to construct the images of the black hole as seen by a distant observer. This bridges the gap between the theoretical geodesic structure and observable astrophysical signatures, a key objective of modern strong-field gravity research.

To ensure full reproducibility and to provide the exact numerical context for our renderings, a comprehensive summary of the specific initial conditions, observer coordinates, grid resolutions, and integration tolerances utilized for each individual figure in this chapter is provided in Appendix A.

4.2.1 Gravitational Lensing of a Point Source

Before proceeding to complex astrophysical backgrounds and continuous accretion disks, we validate the imaging capabilities of our framework by simulating the gravitational lensing of a simple, monochromatic point source. This scenario serves as a stringent, isolated test of the code's ability to resolve the extreme light bending and the exponential sensitivity of photon trajectories in the strong-field regime, free from the complexities of radiative transfer models.

We place a point source (represented as a red dot) directly behind the black hole relative to the distant observer. The resulting lensed image, presented in Fig. 4.5, reveals the rich and highly non-linear topology of strong-field gravitational lensing.



Figure 4.5: Gravitational lensing of a point source (red dot) by a near-extremal Kerr black hole ($a = 0.998$). The two large crescent arcs represent the primary and secondary images, while the tiny red dots near the center correspond to higher-order photon sub-rings (photons that have orbited the black hole multiple times).

The image clearly exhibits the distinct families of images generated by the strong-field deflection mechanism:

- **Primary and Secondary Images (The Arcs):** The two large, highly magnified crescent arcs located on opposite sides of the shadow correspond to the primary ($n = 0$) and secondary ($n = 1$) images. These are formed by photons that pass the black hole on either side. The extreme stretching of the originally point-like source into long, thin arcs is a direct consequence of the divergent magnification factor. As the impact parameter approaches the critical curve, the source is stretched tangentially along the boundary of the shadow.
- **Higher-Order Sub-rings (The Central Dots):** The tiny red dots clustered near the center of the image, hugging the boundary of the black hole shadow, represent the higher-order images ($n \geq 2$). These are photons that have completed one or more full orbits around the black hole before escaping.

The formation of these higher-order sub-rings is governed by an exponential demagnification process. For a photon to complete an additional loop around the black hole, its initial trajectory must be exponentially closer to the exact critical impact parameter. Consequently, each successive image in the hierarchy is exponentially smaller, fainter, and located exponentially closer to the mathematical shadow boundary than the previous one.

From a computational perspective, resolving these higher-order images represents the ultimate stress test for any relativistic ray-tracing algorithm. Photons that form the central dots spend a significantly long affine parameter time navigating the highly unstable region near the photon sphere. In this region, the Lyapunov exponent of the spacetime dictates that any microscopic numerical error will grow exponentially, artificially ejecting the photon to infinity or causing it to plunge prematurely into the event horizon.

Standard numerical integrators, which suffer from secular drift in conserved quantities, inevitably smear out or completely erase these higher-order images. The sharp, distinct resolution of the sub-rings in Fig. 4.5 provides direct, visual confirmation of the efficacy of our Manifold Projection algorithm. By strictly enforcing the null condition and the conservation of energy and angular momentum at every integration step, our framework suppresses this numerical drift, allowing us to faithfully capture the theoretically predicted infinite hierarchy of photon rings that would otherwise be lost to numerical noise.

4.2.2 Shadow Morphology and Frame Dragging

The mathematical shadow of the black hole is defined by the boundary between the photons that are captured and those that escape.



Figure 4.6: Simulated shadow of a near-extremal Kerr black hole ($a = 0.998, M = 1$) illuminated by the four-colour calibration celestial sphere. The pronounced "D" shape of the mathematical shadow is a direct manifestation of frame-dragging, which breaks the equatorial symmetry. The distinct colour quadrants verify the correct orientation and handedness of the backward ray-tracing mapping.

Our numerical simulations accurately reproduce the well-known dependence of the shadow morphology on the spin parameter a. For a non-rotating Schwarzschild black hole ($a = 0$), the shadow is perfectly circular. As the spin parameter increases towards the extreme Kerr limit ($a \to 0.998$), the shadow exhibits a pronounced asymmetry, taking on a characteristic "D" shape. This deformation is a direct consequence of frame-dragging: photons co-rotating with the black hole are pulled closer to the horizon before escaping, thus shifting the critical impact parameter and breaking the equatorial circularity. Our imaging framework naturally captures this transition without any ad hoc assumptions, purely from the numerical solution of the geodesic equations. This allows us to now use any background image as a celestial sphere and have (more) realistic images.




Figure 4.7: Astrophysical visualization of the near-extremal Kerr black hole ($a = 0.998$) shadow, utilizing a high-resolution Milky Way equirectangular panorama as the background celestial sphere. The image captures the extreme gravitational light bending, the primary and secondary image arcs, and the characteristic asymmetric boundary of the photon capture region.

4.2.3 Relativistic Accretion Disk and Radiative Transfer

To create astrophysically meaningful images, we extend our framework to include a geometrically thin accretion disk [14, 11] located in the equatorial plane. As described in the theoretical introduction to radiative transfer, the observed intensity I_{obs} accumulated along a ray is given by the integral:

$$I_{\text{obs}} = \int_{\lambda_{\text{source}}}^{\lambda_{\text{obs}}} \gamma^2 \, j_0(r) \, d\lambda, \tag{4.1}$$

where γ is the relativistic redshift factor and $j_0(r)$ is the local rest-frame emissivity, modelled here by an exponential density profile.

The resulting images, presented in Fig. 4.10, highlight several key relativistic phenomena:

1. **Doppler Effect:** The approaching side of the disk (where the plasma rotates towards the observer) is strongly boosted in intensity, appearing much brighter than the receding side. This asymmetric brightness distribution is a direct consequence of the g^4 scaling.

2. **Gravitational Lensing:** The light from the far side of the disk is bent over and under the black hole, forming a narrow, bright photon ring that closely traces the mathematical shadow.

3. **Higher-Order Sub-rings:** The photons that wind multiple times around the black hole create an hierarchy of sub-rings. Due to our rigorous geometric projection, these sub-rings are resolved with exceptional sharpness, faithfully reproducing the predicted exponential demagnification.

To explicitly visualize the impact of the spin parameter a on the spacetime geometry within a realistic astrophysical context, we contrast the emission from an accretion disk around a non-rotating Schwarzschild black hole ($a = 0$) with that of a near-extremal Kerr black hole ($a = 0.998$).




Figure 4.8: Simulated emission from a geometrically thin accretion disk around a non-rotating Schwarzschild black hole ($a = 0, M = 1$). The observed intensity exhibits strong left-right asymmetry due to relativistic Doppler boosting of the approaching plasma on the left side. In the absence of spacetime rotation, the central mathematical shadow bounded by the primary photon ring remains perfectly circular.

As shown in Fig. 4.8, in the absence of frame-dragging, the mathematical shadow bounded by the photon ring remains perfectly circular. While the kinematic Doppler effect still causes a strong left-right brightness asymmetry in the accretion disk, the underlying geometric boundary of the photon capture region is strictly isotropic. In stark contrast, Fig. 4.9 demonstrates how the extreme rotation of the Kerr spacetime drags the surrounding geometry, compressing the photon ring on the co-rotating side and elongating it on the counter-rotating side, thereby manifesting the characteristic ”D” shape [4, 5].




Figure 4.9: Simulated emission from a geometrically thin accretion disk around a near-extremal Kerr black hole ($a = 0.998$). In contrast to the Schwarzschild geometry, the extreme frame-dragging drastically deforms the spacetime. This results in the characteristic ”D”-shaped compression of both the photon ring and the inner edge of the accretion disk on the co-rotating side.

The combination of an near-extremal Kerr geometry, relativistic Doppler effect, and a realistic background demonstrates the full capability of our framework. These images not only serve as powerful visual tools for understanding strong-field gravity, but also provide a robust, numerically accurate platform for interpreting the horizon-scale observations from current and future interferometric campaigns such as the Event Horizon Telescope (EHT).




Figure 4.10: Comprehensive astrophysical rendering of a near-extremal Kerr black hole ($a = 0.998$) featuring a geometrically thin accretion disk and a realistic Milky Way background. The visualization accurately resolves the primary disk emission subject to Doppler boosting, the strongly lensed image of the disk's far side wrapping over the poles, and the sharp hierarchy of higher-order photon sub-rings preserved by the geometric integration scheme.

Chapter 5

Conclusions and Future Prospects

5.1 Summary of Contributions

In this thesis, we have developed, implemented, and validated a robust computational framework for the numerical simulation and visualization of relativistic ray tracing in the strong-field regime of rotating Kerr black holes. Motivated by the recent observational breakthroughs of the Event Horizon Telescope (EHT), our work addresses the fundamental challenge of generating high-fidelity synthetic images, which requires tracing millions of photon trajectories through an environment characterized by extreme numerical sensitivity and unstable dynamics.

The core methodological achievement of this work lies in the synthesis of two advanced computational techniques designed to overcome the limitations of traditional numerical relativity codes. First, to address the bottleneck of computing exact metric derivatives and Christoffel symbols, we implemented Forward-Mode Automatic Differentiation (AD). By extending the arithmetic of the Julia programming language, we completely eliminated the truncation and round-off errors inherent to finite difference methods. This approach not only guarantees derivatives to machine precision ($\sim 10^{-16}$) but also renders the framework entirely metric-agnostic, allowing it to be applied to any spacetime geometry without manual algebraic derivation.

Second, to mitigate the severe invariant drift exhibited by standard explicit Runge-Kutta integrators near the unstable photon sphere, we coupled a high-order adaptive integrator with a geometric Manifold Projection algorithm. By formulating the physical constraints-the null condition, energy conservation, and axial angular momentum conservation-as a constraint manifold $\mathcal{N}$, we employed Lagrange multipliers and a multidimensional Newton-Raphson solver to project the numerical state back onto $\mathcal{N}$ at every integration step. Crucially, this correction is strictly orthogonal to the physical flow, ensuring that the trajectory is stabilized without introducing artificial numerical damping.

We systematically validated the numerical stability of this framework across the three fundamental classes of photon orbits: scattering, direct capture, and bound spirals. In all regimes, particularly for the highly unstable transient capture orbits, the framework successfully maintained conserved quantities at the machine-precision limit. This exceptional numerical fidelity enabled the resolution of the infinite hierarchy of higher-order photon sub-rings-a feature typically smeared out or entirely lost by standard numerical methods due to secular drift.

Applying this framework to astrophysical imaging, we successfully reproduced the characteristic asymmetric "D-shaped" shadow of a near-extremal Kerr black hole ($a = 0.998$) induced by frame-dragging. Through rigorous UV mapping calibration using a four-color test image, we ensured physically accurate lensing of realistic backgrounds, such as the Milky Way. Furthermore, by incorporating a geometrically thin accretion disk model with relativistic radiative transfer ($I \propto g^4$), we captured key observable phenomena, including extreme Doppler effect and the complex gravitational lensing of the disk's far side. Beyond accuracy, the

framework is highly efficient; on a standard 16-core laptop, it processes computationally demanding trajectories at an average rate of one million rays per hour, making it a practical tool for large-scale parameter space exploration.

5.2 Future Research Directions

The computational infrastructure developed in this thesis lays the groundwork for several promising research extensions. The metric-agnostic architecture, the machine-precision conservation of invariants, and the high performance of the framework make it a versatile platform that can be expanded in multiple directions relevant to next-generation observational astrophysics.

5.2.1 GPU Acceleration and Real-Time Rendering

While the current CPU multi-threaded implementation achieves a throughput of approximately one million rays per hour on a standard 16-core laptop, the embarrassingly parallel nature of backward ray tracing makes it an ideal candidate for massive parallelization on Graphics Processing Units (GPUs). Each pixel on the observer's screen corresponds to an independent initial value problem, meaning that millions of geodesics can be integrated simultaneously without inter-thread communication overhead.

Porting the inner geodesic and projection kernels to GPUs would reduce the rendering time from hours to seconds, enabling real-time visualization of black hole shadows at high resolution. This capability would be particularly valuable for interactive scientific exploration, educational purposes, and the rapid generation of large datasets required for machine learning applications. Furthermore, GPU acceleration would make feasible the computation of time-dependent images from dynamical spacetimes, such as those arising from binary black hole mergers or accretion disk instabilities.

5.2.2 Extension to Alternative Metrics and Beyond-GR Theories

One of the most significant advantages of the framework is its metric-agnostic nature, stemming from the use of Automatic Differentiation to compute Christoffel symbols. This feature allows the code to be applied to any spacetime geometry without requiring manual algebraic derivation of connection coefficients, making it an ideal tool for testing alternative theories of gravity. By systematically comparing the synthetic images produced by these alternative metrics with observational data from the Event Horizon Telescope (EHT) and its next-generation upgrade (ngEHT), one could place quantitative constraints on deviations from General Relativity in the strong-field regime.

5.2.3 Template Generation and Parameter Estimation

The high fidelity and computational efficiency of the framework make it well-suited for the generation of large libraries of synthetic images (templates) that can be used in Bayesian parameter estimation pipelines. Current EHT analyses rely on comparing observed interferometric visibilities with pre-computed template libraries generated by general relativistic magnetohydrodynamic (GRMHD) simulations coupled with ray-tracing codes.

By integrating the forward ray-tracer with differentiable programming techniques, one could construct a fully differentiable imaging pipeline in which the synthetic image is a differentiable function of the black hole parameters (mass M, spin a, inclination θ_0, accretion rate, etc.).

Such a differentiable pipeline would represent a significant advancement in the analysis of horizon-scale observations, bridging the gap between theoretical predictions and observational data in a computationally efficient and statistically rigorous manner.

Appendix A

Simulation Configurations

The following table details the precise spatial parameters and initial observer configurations utilized in the Julia ray-tracing framework to generate the figures presented in Chapter 4.

Figure	**Radial Dist.** r_0 **(**M**)**	**Inclination** θ_0 **(°)**	**FOV Limits** α, β **(**M**)**
Fig. 4.2 (Scattering Orbit)	50.0	85.0	N/A (Single Ray)
Fig. 4.3 (Bound Spiral Orbit)	2.5	90.0	N/A (Single Ray)
Fig. 4.4 (Direct Capture Orbit)	50.0	75.0	N/A (Single Ray)
Fig. 4.5 (Point Source)	30.0	87.5	$\alpha \in [-20, 20], \beta \in [-20, 20]$
Fig. 4.6 (Calibration)	500.0	87.5	$\alpha \in [-9, 9], \beta \in [-9, 9]$
Fig. 4.7 (MW Shadow)	30.0	87.5	$\alpha \in [-30, 30], \beta \in [-60, 60]$
Fig. 4.8 (Schw. Disk)	500.0	87.5	$\alpha \in [-15, 15], \beta \in [-15, 15]$
Fig. 4.9 (Kerr Disk)	500.0	87.5	$\alpha \in [-15, 15], \beta \in [-15, 15]$
Fig. 4.10 (Full Render)	30.0	87.5	$\alpha \in [-15, 15], \beta \in [-15, 15]$

Table A.1: Detailed configuration matrix outlining the initial observer radial distance, inclination angle, and Field of View (impact parameter bounds) for each corresponding figure.

Bibliography

[1] Jeff Bezanson, Alan Edelman, Stefan Karpinski, and Viral B Shah. Julia: A fresh approach to numerical computing. *SIAM review*, 59(1):65–98, 2017.

[2] Robert H Boyer and Richard W Lindquist. Maximal analytic extension of the kerr metric. *Journal of Mathematical Physics*, 8(2):265–281, 1967.

[3] Subrahmanyan Chandrasekhar. *The mathematical theory of black holes*. Oxford University Press, 1983.

[4] Christopher T Cunningham. The effects of redshifts and focusing on the spectrum of an accretion disk around a kerr black hole. *The Astrophysical Journal*, 202:788–802, 1975.

[5] Event Horizon Telescope Collaboration. First m87 event horizon telescope results. v. physical origin of the asymmetric ring. *The Astrophysical Journal Letters*, 875(1):L5, 2019.

[6] Heino Falcke, Fulvio Melia, and Eric Agol. Viewing the shadow of the black hole at the galactic center. *The Astrophysical Journal Letters*, 528(1):L13, 2000.

[7] Samuel E Gralla, Daniel E Holz, and Robert M Wald. Black hole shadows, photon rings, and lensing rings. *Physical Review D*, 100(2):024018, 2019.

[8] Andreas Griewank and Andrea Walther. *Evaluating derivatives*. Society for Industrial and Applied Mathematics, Cambridge, TAS, Australia, 2 edition, January 2008.

[9] Ernst Hairer, Christian Lubich, and Gerhard Wanner. *Geometric numerical integration*. Springer Series in Computational Mathematics. Springer, Berlin, Germany, 2 edition, February 2006.

[10] Roy P Kerr. Gravitational field of a spinning mass as an example of algebraically special metrics. *Physical Review Letters*, 11(5):237–238, 1963.

[11] Jean-Pierre Luminet. Image of a spherical black hole with thin accretion disk. *Astronomy and Astrophysics*, 75:228–235, 1979.

[12] Charles W Misner, Kip S Thorne, and John Archibald Wheeler. *Gravitation*. W. H. Freeman and Company, San Francisco, 1973.

[13] Jarrett Revels, Miles Lubin, and Theodore Papamarkou. Forward-mode automatic differentiation in julia. *arXiv preprint arXiv:1607.07892*, 2016.

[14] Nikolai I Shakura and Rashid A Sunyaev. Black holes in binary systems. observational appearance. *Astronomy and Astrophysics*, 24:337–355, 1973.

[15] John Lighton Synge. The escape of photons from gravitationally intense stars. *Monthly Notices of the Royal Astronomical Society*, 131(3):463–466, 1966.

[16] Robert M Wald. *General Relativity*. University of Chicago Press, 1984.

[17] R E Wengert. A simple automatic derivative evaluation program. *Commun. ACM*, 7(8):463–464, August 1964.