
Shelter in Place Triggers

Chris Hoover 6/4/2020

Testing



Model

We use a slight tweak to LEMMA to add an explicit deaths compartment in order to fit to deaths data in addition to hospitalizations

$$\dot{S} = -\beta S(I_R + I_H)/N$$

$$\dot{E} = \beta S(I_R + I_H)/N - \sigma E$$


$$\dot{I}_R = \sigma (1 - \alpha)E - \gamma_R I_R$$

$$\dot{I}_H = \sigma \alpha E - \rho I_H$$

$$\dot{H} = \rho I_H - \gamma_H H$$

$$\dot{D} = \gamma_H \mu H$$


$$\dot{R} = \gamma_R I_R \gamma_H (1 - \mu) H$$




Warning: Removed 22 rows containing missing values (geom_point).


Table 1: Parameter values and descriptions used in the model

	Value	Definition
\overline{N}	883305	population size
t.sim	110	time to run simulation
E_0	2	starting number of exposed
β	0.7	baseline transmission rate
c_r	1	Relative contact rate between S and Ir
c_h	1	Relative contact rate between S and Ih
σ	0.333	1/serial interval
α	0.05	proportion severely symptomatic (will be hospitalized)
ρ	0.25	time between symptom onset and hospitalization
γ_r	0.2	1/time to recovery (non-infectiousness) for mildly symptomatic
γ_h	0.083	1/time hospitalized
μ	0.2	proportion of hospitalized cases who die




Warning: Removed 6 rows containing missing values (geom_point).






Warning: Removed 22 rows containing missing values (geom_point).



Warning: Removed 6 rows containing missing values (geom_point).



Warning: Removed 1 rows containing missing values (geom_path).

