

# Shelter in Place Triggers

*Chris Hoover*


*6/4/2020*

## Purpose

Evaluate potential triggers to intervene on COVID transmission (e.g. shelter in place or other method to reduce contact and transmission) from signals in testing data.

## Testing

```
## Warning: Removed 12 rows containing missing values (geom_path).
```



## Model

We use a slight tweak to LEMMA to add an explicit deaths compartment in order to fit to deaths data in addition to hospitalizations


Table 1: Best fit model parameters

	Value	Definition
$N$	883305	population size
t.sim	113	time to run simulation
$E_0$	5.921	starting number of exposed
$c_r$	1	Relative contact rate between S and Ir
$c_h$	1	Relative contact rate between S and Ih
$\sigma$	0.333	1/serial interval
$\alpha$	0.039	proportion severely symptomatic (will be hospitalized)
$\rho$	0.25	time between symptom onset and hospitalization
$\gamma_r$	0.2	1/time to recovery (non-infectiousness) for mildly symptomatic
$\gamma_h$	0.083	1/time hospitalized
$\mu$	0.112	proportion of hospitalized cases who die


$$\begin{aligned}
\dot{S} &= -\beta S(I_R + I_H)/N \\
\dot{E} &= \beta S(I_R + I_H)/N - \sigma E \\
\dot{I}_R &= \sigma(1 - \alpha)E - \gamma_R I_R \\
\dot{I}_H &= \sigma\alpha E - \rho I_H \\
\dot{H} &= \rho I_H - \gamma_H H \\
\dot{D} &= \gamma_H \mu H \\
\dot{R} &= \gamma_R I_R + \gamma_H (1 - \mu)H
\end{aligned}$$

## Model fit


## Warning: Removed 22 rows containing missing values (position\_stack).



## Warning: Removed 6 rows containing missing values (position\_stack).




## Warning: Removed 1 rows containing missing values (geom\_path).




### Comparison to testing data

## Warning: Removed 13 rows containing missing values (geom\_path).

SF COVID testing and modeled cases



## Warning: Removed 16 rows containing missing values (geom\_path).



### Match modeled testing to observed testing

Assume all new hospitalized cases are tested and confirmed positive, then assume remaining tests are allocated to non-hospitalized population. In the model, this is equivalent to a sample of  $S$ ,  $E$ , and  $I_r + I_h$ . In reality, lots of nuance in the  $E$  and  $R$  compartments with regard to testing, but for simplicity, we'll assume positive tests from this sample only come from  $I_r + I_h$  and  $R$ s are not tested. So want to solve for sampling bias,  $\mathcal{B}$ , from:

$$\frac{+Tests - H_{new}}{Tests - H_{new}} = \frac{\mathcal{B}(I_r + I_h)}{S + E + I_r + I_h}$$

## Warning: Removed 8 rows containing missing values (geom\_path).






Figure 1: Distribution of estimated sampling bias since end of April









## Model forecast

$\mathcal{R}_e \rightarrow 1.4$


## Warning: Removed 22 rows containing missing values (position\_stack).

Projected Hospital Census with  
Re reaching 1.4 in 7 days  
and remaining until 2020-08-01






Projected Hospital Admissions with  
Re reaching 1.4 in 7 days  
and remaining until 2020-08-01



### Simulations of potential SiP Triggers

Assuming 1600 tests per day,  $B=45$ ,  $R_e=1.4$  by June 15

Blue—positive tests, Red—Hospital admissions



## Warning: Removed 2 rows containing missing values (geom\_path).

