

Additional exercises for MA505250

Exercise 1. Let X be a Banach space and $\mathcal{J}_1 : X \rightarrow \mathbb{R}$, $\mathcal{J}_2 : X \rightarrow \mathbb{R}$ two functionals. Prove that then

- (a) If \mathcal{J}_1 and \mathcal{J}_2 are weak l.s.c., $\alpha \geq 0$, then $\alpha\mathcal{J}_i$ and $\mathcal{J}_1 + \mathcal{J}_2$ are weak l.s.c.
- (b) If \mathcal{J}_1 is weak l.s.c. and $\varphi : \mathbb{R} \rightarrow \mathbb{R}$ is monotonically increasing and l.s.c., then $\varphi \circ \mathcal{J}_1$ is weak l.s.c.
- (c) If \mathcal{J}_1 is weak l.s.c. then for a Banach space Y and weak sequentially continuous $\Phi : \mathcal{Y} \rightarrow X$ it follows that $\mathcal{J}_1 \circ \Phi$ is weak l.s.c.
- (d) For any (non-empty) family of weak l.s.c. functionals $\mathcal{J}_i : X \rightarrow \mathbb{R}$, $i \in I$ we have that $\sup_{i \in I} \mathcal{J}_i$ is weak l.s.c.
- (e) Let $\varphi : K \rightarrow \mathbb{R}$ be l.s.c., where K is either \mathbb{R} or \mathbb{C} , and $x^* \in X^*$. Then the functional

$$L_{x^*, \varphi} = \varphi \circ \langle x^*, \cdot \rangle_{X^* \times X},$$

is weak l.s.c. in X .

Collect what you have proven (as appropriate) and show that $\varphi(\|u\|_X)$ is weak l.s.c. for any $\varphi : [0, \infty) \rightarrow \mathbb{R}$ that is monotonically increasing and l.s.c.

Exercise 2. For a given noisy image $g \in L^2(\Omega)$ and rectangular image domain $\Omega \subset \mathbb{R}^2$ we consider the following variational problem

$$u_\alpha = \operatorname{argmin}_{u \in L^2(\Omega)} \{ \alpha \|Du\|_2^2 + \|u - g\|_2^2 \},$$

for $\alpha > 0$. Prove that there exists a unique minimiser u_α to the above problem. Show further, that under the additional assumption that $L \leq g \leq R$ a.e. in Ω the minimiser u_α of the above problem also fulfils $L \leq u_\alpha \leq R$ a.e. in Ω .