

SIR Model Stability

Charlotte Maryjean White

December 10, 2025

1 Model

We can begin with a simplified SIR (Susceptible-Infected-Removed) model with no births or deaths.

$$\begin{aligned}\frac{dS}{dt} &= -\frac{\alpha}{N}SI \\ \frac{dI}{dt} &= \frac{\alpha}{N}SI - \gamma I \\ \frac{dR}{dt} &= \gamma I\end{aligned}$$

Where α is the contact rate, and γ is the recovery rate. The initial conditions would be a constant population, N , and a positive, non-zero S_0 and I_0 and non-negative R_0 .

$$S_0 + I_0 + R_0 = N$$

$$S_i + I_i + R_i = N$$

$$S_0 > 0, I_0 > 0, R_0 \geq 0$$

In such a model with a closed population, there will be no endemic equilibrium. The disease would die out as the susceptible population is eventually depleted. Thus a more useful model to examine is the SIR model with births and deaths:

References

- [1] Brauer, F., Castillo-Chavez, C., Feng, Z. (2019). Introduction: A Prelude to Mathematical Epidemiology. In: Mathematical Models in Epidemiology. Texts in Applied Mathematics, vol 69. Springer, New York, NY. https://doi.org/10.1007/978-1-4939-9828-9_1
- [2] Allen, L.J.S. (1994). Some discrete-time SI, SIR, and SIS epidemic models. Mathematical Biosciences 124(1), 83-105. [https://doi.org/10.1016/0025-5564\(94\)90025-6](https://doi.org/10.1016/0025-5564(94)90025-6)

- [3] Strogatz, S. (2001). Nonlinear dynamics and chaos: with applications to physics, biology, chemistry and engineering. Perseus Books Group.