Chapter 1

Milestone 4

1.1 Force-Derivation from the Lenard Jones Potential

$$\vec{f_k} = \sum_i \frac{\partial V}{\partial r_{ik}} \hat{r_{ik}} \tag{1.1}$$

$$V(r) = 4\epsilon \left[\left(\frac{\sigma}{r} \right)^{12} - \left(\frac{\sigma}{r} \right)^{6} \right] \tag{1.2}$$

$$\frac{\partial V}{\partial r_{ik}} = 4\epsilon \left(\frac{6\sigma^6}{r_{ik}^6} - \frac{12\sigma^{12}}{r_{ik}^{13}} \right) \tag{1.3}$$

```
[4]: import sympy as sp
     import warnings
     warnings.filterwarnings('ignore')
     sp.init_printing()
     eps = sp.Symbol("e")
     sig = sp.Symbol("s")
     rad = sp.Symbol("r")
     energyRad = 4 * eps * ((sig/rad)**12 - (sig/rad)**6)
     energyRad.diff(rad)
```

[4]: $4e \left(\dfrac{6s^6}{r^7} - \dfrac{12s^{12}}{r^{13}} \right)$

## 1.2 Different Time Steps

1.3 Simulation Snapshots

Figure 1.1: Simulation with a time step of 0.001

Figure 1.2: Simulation with a time step of 0.005

Figure 1.3: Simulation with a time step of 0.01

Figure 1.4: Simulation with a time step of 0.02

Figure 1.5: Simulation with a time step of 0.03

Figure 1.6: Simulation with a time step of 0.04

Figure 1.7: Simulation

Figure 1.8: Simulation

Figure 1.9: Simulation

Figure 1.10: Simulation

Figure 1.11: Simulation

Chapter 2

Milestone 5

2.1 Berendsen Thermostat Teststrategy

$$\lambda = \sqrt{1 + \left(\frac{T_0}{T} - 1\right)\frac{\Delta t}{\tau}}$$

(2.1)

$$T(t) = T_0 + (T_1 - T_0)e^{-t/\tau}$$

(2.2)

2.2 Simulation Time

Figure 2.1: Simulationtime from 8 to 192 Atoms

Chapter 3

Milestone 6

The Images are outdated!!!!

Figure 3.1: Simulationtime without Neighborlist

New images that are correct.

Figure 3.2: Simulationtime with the old Neighborlist

Figure 3.3: Simulationtime with the new Neighborlist

Figure 3.4: Simulationtime with the new Neighborlist

Chapter 4

Introduction

Chapter 5

Methods

Chapter 6

Implementation

Chapter 7

Results

Chapter 8

Conclusion