

Integrable Liouville Minisuperspace Model

Self-adjointness of the Energy Density and Semi-classical Wave Functions

Alexander A. Andrianov^{1,4} Chen Lan (蘭 晨)² Oleg O. Novikov
Yi-Fan Wang (王 一帆)³

¹Saint-Petersburg State University, Ulyanovskaya str. 1, Petrodvorets, Sankt-Petersburg 198504, 俄羅斯

²ELI-ALPS Research Institute, Budapesti út 5, H-67228 Szeged, 匈牙利

³Institut für Theoretische Physik, Universität zu Köln, Zülpicher Straße 77, D-50937 Köln, 德國

⁴Institut de Ciències del Cosmos, Universitat de Barcelona, Martí i Franquès 1, E-08028 Barcelona, 西班牙

May 26, 2018



Outline

1. Introduction

2. The classical model and the implicit trajectories

Lagrangian formalism and the first integral

3. Dirac quantisation and self-adjointness of the energy density

Dirac quantisation and the quantum-mechanical analogy

Self-adjointness of the energy density

4. The semi-classical wave functions

The Wentzel–Kramers–Brillouin approximation

Inner products and wave packets

5. Conclusions



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and self-adjointness of the energy density
Dirac quantisation and the quantum-mechanical analogy
Self-adjointness of the energy density
4. The semi-classical wave functions
The Wentzel–Kramers–Brillouin approximation
Inner products and wave packets
5. Conclusions



Introduction

The Friedmann–Lemaître model with quintessence and phantom Liouville fields

- Dynamical Dark Energy has been modelled by quintessence¹ and phantom² matter, which can be realised by minimally-coupled real scalar fields with $\ell = \pm 1$ ³

$$\mathcal{S}_L = \int d^4x \sqrt{-g} \left\{ -\ell \frac{g^{\mu\nu}}{2} (\partial_\mu \phi)(\partial_\nu \phi) - \mathcal{V}(\phi) \right\}. \quad (1)$$

- The Liouville field⁴ $\mathcal{V}(\phi) = V e^{\lambda\phi}$ is of interest, where λ, V are real parameters.
- Assume a flat Robertson–Walker metric $g_{\mu\nu} dx^\mu dx^\nu = -N^2(t) dt^2 + \kappa e^{2\alpha(t)} d\Omega_{3F}^2$, where $\kappa := 8\pi G$, $d\Omega_{3F}^2 := d\chi^2 + \chi^2(d\theta^2 + \sin^2 \theta d\varphi^2)$, and N lapse function.
- The total minisuperspace action reads $\mathcal{S} := \mathcal{S}_{EH} + \mathcal{S}_{GHY} + \mathcal{S}_L = \int d\Omega_{3F}^2 \int dt L$, where

$$L := \kappa^{3/2} N e^{3\alpha} \left(-\frac{3}{\kappa} \frac{\dot{\alpha}^2}{N^2} + \ell \frac{\dot{\phi}^2}{2N^2} - V e^{\lambda\phi} \right). \quad (2)$$

¹R. R. Caldwell *et al.*, *Phys. Rev. Lett.* **80**, 1582–1585 (Aug. 7, 1997).

²R. R. Caldwell, *Phys. Lett. B* **545**, 23–29 (Aug. 16, 1999).

³The signature of metric is $(-, +, \dots, +)$.

⁴Y. Nakayama, *Int. J. Mod. Phys. A* **19**, 2771–2930 (Feb. 2, 2004).



Integrability

With the help of an integral of motion, implicit trajectories can be obtained explicitly, and the minisuperspace Wheeler–DeWitt equation can be integrated exactly.

Self-adjointness

Imposing self-adjointness of the energy density leads to removal of the degeneracy and discretisation of the spectrum.

Semi-classical wave functions

Semi-classical wave functions can be constructed by the principle of constructive interference, the Wentzel–Kramers–Brillouin approximation and numerical methods.



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and self-adjointness of the energy density
Dirac quantisation and the quantum-mechanical analogy
Self-adjointness of the energy density
4. The semi-classical wave functions
The Wentzel–Kramers–Brillouin approximation
Inner products and wave packets
5. Conclusions



Decoupling the variables

Via orthogonal transformation

- By rescaling $\bar{N} := N e^{-3\alpha}$ and an orthogonal transformation from (α, ϕ) to (β, χ) , the system can be decoupled

$$L = \varkappa^{3/2} \bar{N} \left(-\varsigma \frac{3}{\varkappa} \frac{\dot{\beta}^2}{\bar{N}^2} + \ell \varsigma \frac{\dot{\chi}^2}{2\bar{N}^2} - V e^{g\chi} \right), \quad (3)$$

where ς is the sign of $(\lambda^2 - 6\ell\varkappa)$, and $g = \varsigma \sqrt{|\lambda^2 - 6\ell\varkappa|}$ takes the place of λ .

- Using the conservation of p_β ⁵, $\bar{N}(t)$ can be eliminated, leading to the exact integrations of the equations of motion as implicit trajectories.
 - $p_\beta \neq 0$: four cases according to $(\ell, \varsigma v)$, v is the sign of V ; see below.
 - $p_\beta = 0$: well-known power-law special solution for $(+, -)$ and $(-, +)$; read literature⁶.

⁵The same first integral has been found in C. Lan, PhD thesis, Saint Petersburg State University, 2016, <https://search.rsl.ru/ru/record/01006663434>, A. A. Andrianov et al., *Theor. Math. Phys.* **184**, 1224–1233 (Mar. 18, 2015), in canonical formalism.


⁶For instance, A. R. Liddle, D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, (Cambridge University, 2000) , ch. 3.



Minisuperspace trajectories for quintessence model (+, +)


sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ

$$e^{g\chi} = \frac{p_\beta^2}{12\kappa^2|V|} \operatorname{sech}^2\left(\sqrt{\frac{3}{2\kappa}} g(\beta - \beta_0)\right); \text{ no power-law special solution.}$$



Trajectories for quintessence model (+, -)


csh, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ



Minisuperspace trajectories for phantom model $(-, +)$

sec, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$

$e^{g\chi} = \frac{p_\beta^2}{12\varkappa^2|V|} \sec^2\left(\sqrt{\frac{3}{2\varkappa}} g(\beta - \beta_0)\right)$, power-law special solution parallel to asymptotes.



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and self-adjointness of the energy density
Dirac quantisation and the quantum-mechanical analogy
Self-adjointness of the energy density
4. The semi-classical wave functions
The Wentzel–Kramers–Brillouin approximation
Inner products and wave packets
5. Conclusions



Dirac quantisation

- The primary Hamiltonian of this constrained, singular system⁷ reads $H^p = \bar{N}H_\perp + v^{\bar{N}}p_{\bar{N}}$, in which the Hamiltonian constraint is

$$H_\perp := -\beta \frac{p_\beta^2}{12\kappa^{1/2}} + \ell\beta \frac{p_\chi^2}{2\kappa^{3/2}} + \kappa^{3/2}V e^{g\chi}. \quad (4)$$

- Applying the Dirac quantisation rules with the Laplace–Beltrami operator ordering⁸, one gets the minisuperspace Wheeler–DeWitt equation with (β, χ)

$$0 = \widehat{H}_\perp \Psi(\beta, \chi) := \left(\beta \frac{\hbar^2}{12\kappa^{1/2}} \partial_\beta^2 - \ell\beta \frac{\hbar^2}{2\kappa^{3/2}} \partial_\chi^2 + \kappa^{3/2}V e^{g\chi} \right) \Psi. \quad (5)$$

- Equation (5) is Klein–Gordon-like, hyperbolic for $\ell = +1$ and **elliptic** for $\ell = -1$.

⁷D. M. Gitman, I. V. Tyutin, *Quantization of Fields with Constraints*, (Springer, 1990), H. J. Rothe, K. D. Rothe, *Classical and Quantum Dynamics of Constrained Hamiltonian Systems*, (World Scientific, Apr. 2010).

⁸C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012), ch. 8.



Integration of the Wheeler–DeWitt equation

The quantum-mechanical analogy

- Fourier transforming β to $k_\beta \in \mathbb{R}$ yields the time-independent Schrödinger equation

$$E\psi(x) := \widehat{H}_{\text{eff}}\psi := \left(-\frac{\hbar^2}{2M_p}\partial_x^2 + V_{\text{eff}}\right)\psi, \quad V_{\text{eff}} := \ell s v \widetilde{V} e^{\tilde{g}x}, \quad (6)$$

in which $E \propto \ell\nu^2$, $M_p := \sqrt{\frac{\hbar}{\pi}}$, $x \propto \chi$, $\widetilde{V} \propto |V| \geq 0$, and $\nu \propto |k_\beta| \geq 0$.

- \widehat{H}_{eff} plays the role of energy density in cosmology.
- The Liouville potential V_{eff} is a special case of the Morse potential⁹.
- The Starobinsky model in the Einstein frame is related to the Morse potential.
- Let $e^{2y} := 8M_p \widetilde{V}/\hbar^2 \tilde{g}^2 \cdot e^{\tilde{g}x}$; there are four cases according to (ℓ, sv) , and the mode functions are modified and unmodified Bessel functions of imaginary and real orders

	$E \geq 0$	$E < 0$
$V_{\text{eff}} > 0$	$(+, +); K_{\pm\nu}(e^y)$	$(-, -); \text{none}$
$V_{\text{eff}} < 0$	$(+, -); F_{\pm\nu}(e^y), G_{\pm\nu}(e^y)$ ¹⁰	$(-, +); J_\nu(e^y)$

⁹P. M. Morse, *Phys. Rev.* **34**, 57–64 (July 1929), D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012).

¹⁰These are linear combinations of $J_{\pm\pm\nu}(e^y)$; see T. M. Dunster, *SIAM J. Math. Anal.* **21**, 995–1018 (July 1990).



Self-adjointness of the energy density

Results for the Liouville potential¹⁴

- k_β being real enforces the reality of E and the self-adjointness of \widehat{H}_{eff} .
- For $V_{\text{eff}} > 0$: \widehat{H}_{eff} is **essentially self-adjoint**¹¹. Nothing special for quintessence (+,+).
- For $V_{\text{eff}} < 0$: \widehat{H}_{eff} admits a **family of self-adjoint extensions**.
- Quintessence (+,-): “eigenfunctions” are linear combinations of F and G; **the degeneracy is removed**
- Phantom (-,+): the spectrum becomes **discrete**

$$\psi_{-, \nu}(e^y) \propto J_\nu(e^y), \quad \nu = 2n + a_<, \quad n \in \mathbb{Z}_\geq, \quad (7)$$

and the corresponding full wave-functions are Bloch-periodic. Imposing (anti-)periodic boundary condition fixes $a_< = 0$ (1).

- The self-adjointness has been discussed in quantum cosmology in e.g.¹² (operator ordering) and more recently in¹³ (Λ as a quantum number).

¹¹See also B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267 , sec. 9.9.

¹²C. R. Almeida et al., *Gravitation Cosmol.* **21**, 191–199 (Jan. 17, 2015).

¹³S. Gryb, K. P. Y. Thébault, arXiv: 1801.05789 (gr-qc) (Jan. 17, 2018).

¹⁴See also D. M. Gitman et al., *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012) , sec. 8.5.



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and self-adjointness of the energy density
Dirac quantisation and the quantum-mechanical analogy
Self-adjointness of the energy density
4. The semi-classical wave functions
The Wentzel–Kramers–Brillouin approximation
Inner products and wave packets
5. Conclusions



Matching quantum number with classical first integral

Apply the principle of constructive interference; take $(+, +)$ as an example

- Write the mode function in the Wentzel–Kramers–Brillouin form
 $\Psi_\nu(\beta, \chi) \sim \sqrt{\rho(\beta, \chi)} \exp\left\{\frac{i}{\hbar} S(\beta, \chi)\right\}$. For $S/\hbar \gg 1$ and $\nu \gg 1$, $S(\beta, \chi)$ becomes the Hamilton principal function in the leading-order approximation, which is stationary with respect to the variation of integral constants $\partial S / \partial \nu = 0$.
- Fixing $\nu/\sigma > 1$,

$$K_{\pm\nu}(\sigma) \sim \frac{\sqrt{2\pi} \cos\left(\sqrt{\nu^2 - \sigma^2} - \nu \arccos \frac{\nu}{\sigma} - \frac{\pi}{4}\right)}{(\nu^2 - \sigma^2)^{1/4} e^{\mp\nu/2}} + O\left(\frac{1}{\sigma}\right). \quad (8)$$

- Applying *the principle* to the phase of $\Psi_\nu(\sigma)$, one has $\sigma/\nu = \operatorname{sech}(\beta_\beta \gamma)$, which matches the trajectory if $\hbar k_\beta \equiv \hbar \sqrt{\frac{3}{2\nu}} g\nu = p_\beta$.
- Fixing $\nu/\sigma < 1$, the asymptotic expansion is not oscillatory, but exponential; it is not within the Wentzel–Kramers–Brillouin regime.
- The conclusions also hold for $F_{\pm\nu}(\sigma)$, $G_{\pm\nu}(\sigma)$ for $(+, -)$, and $J_\nu(\sigma)$ for $(-, +)$.



Inner products for wave functions

Schrödinger and Mostafazadeh inner product

- A common starting point is the Schrödinger product¹⁵

$$(\Psi_1, \Psi_2)_S := \int d\chi \Psi_1^*(\beta, \chi) \Psi_2(\beta, \chi); \quad (9)$$

- $(\Psi, \Psi)_S$ is positive-definite, and the integrand $\rho_S(\beta, \chi)$ is non-negative
- The corresponding Schrödinger current is real but does not satisfy continuity equation
- Mostafazadeh¹⁶: for self-adjoint \mathbb{D} with positive spectrum in $(-\partial_t^2 + \mathbb{D})\Psi = 0$:

$$(\Psi_1, \Psi_2)_M^\kappa := \kappa \left\{ (\Psi_1, \mathbb{D}^{+1/2} \Psi_2)_S + (\partial_\beta \Psi_1, \mathbb{D}^{-1/2} \partial_\beta \Psi_2)_S \right\}, \quad \kappa > 0. \quad (10)$$

- Real power of \mathbb{D} is defined by spectral decomposition
- It can be shown¹⁷ that the density

$$\varrho_M^\kappa := \kappa \left\{ |\mathbb{D}^{+1/4} \Psi|^2 + |\mathbb{D}^{-1/4} \partial_\beta \Psi|^2 \right\} \quad (11)$$

- is equivalent to the integrand ρ_M^κ up to a boundary term $\int d\chi \varrho_M^\kappa = \int d\chi \rho_M^\kappa \equiv (\Psi_1, \Psi_2)_M^\kappa$
- is non-negative
- The corresponding current $\vec{\mathcal{J}}_M^\kappa$ is real but not conserved¹⁸.

¹⁵C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012), ch. 5.

¹⁶A. Mostafazadeh, *Class. Quantum Grav.* **20**, 155–171 (Sept. 6, 2002).

¹⁷A. Mostafazadeh, F. Zamani, *Ann. Phys.* **321**, 2183–2209 (Feb. 17, 2006).

¹⁸B. Rosenstein, L. P. Horwitz, *J. Phys. A: Math. Gen.* **18**, 2115–2121 (Aug. 1985).



Wave packets of Gaussian amplitude for continuous spectrum

Quintessence models

- It is difficult to find an amplitude such that the wave packet is Gaussian
- Instead, one can choose a Gaussian amplitude

$$A(\nu; \bar{\nu}, \sigma) := \left(\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\nu - \bar{\nu})^2}{2\sigma^2}\right) \right)^{1/2} \quad (12)$$

- In¹⁹, $A(\nu; \bar{\nu}, \sigma/\sqrt{2})$ was chosen.


¹⁹M. P. Dąbrowski et al., *Phys. Rev. D* **74**, arXiv: hep-th/0605229 (hep-th) (May 23, 2006).




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\kappa^{1/2}/5$, $V = +\kappa^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\kappa^{1/2}/5$, $V = +\kappa^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger



Mostafazadeh



Wave packets with Poissonian amplitude for discrete spectrum

Discrete phantom model $(-, +)$

- Gaussian distribution works for continuous variable
- For discrete spectrum, one can choose a Poissonian amplitude

$$A_n(\bar{n}) := \left(e^{-\bar{n}} \frac{\bar{n}^n}{n!} \right)^{1/2} \quad (13)$$

- In²⁰, $A_n(\bar{n}/\sqrt{2})$ was chosen.


²⁰C. Kiefer, *Nucl. Phys. B* **341**, 273–293 (Sept. 1990).




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger




Mostafazadeh




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger



Mostafazadeh




Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and self-adjointness of the energy density
Dirac quantisation and the quantum-mechanical analogy
Self-adjointness of the energy density
4. The semi-classical wave functions
The Wentzel–Kramers–Brillouin approximation
Inner products and wave packets
5. Conclusions



Highlights

Revisited



- An **integral of motion** was found for the Liouville cosmological model.
 - Implicit trajectories in minisuperspace were obtained explicitly.
 - The minisuperspace Wheeler–DeWitt equation was integrated exactly.
- The **self-adjointness** of the energy density was found to be non-trivial.
 - The degeneracy of the quintessence model $(+, -)$ was removed.
 - The spectrum of the phantom model $(-, +)$ became discrete.
- The **semi-classical wave functions** were compared with the classical trajectory.
 - Mode functions in the Wentzel–Kramers–Brillouin approximation were matched with the classical Hamilton principal function.
 - Semi-classical wave packets were constructed and compared with the classical trajectory under the Schrödinger and Mostafazadeh inner product.

Outlook

- Beyond isotropy: Bianchi models²¹
- Beyond homogeneity: cosmological perturbations²² (working)
- Beyond single Liouville field: multiple potentials / multiple fields²³
- Beyond classical matter and Schrödinger inner product: *PT*-symmetric fields²⁴

²¹A. Y. Kamenshchik *et al.*, *Phys. Rev. D* **95**, arXiv: 1702.02314 (gr-qc) (Feb. 8, 2017).

²²D. Brizuela *et al.*, *Phys. Rev. D* **93**, arXiv: 1511.05545 (gr-qc) (Nov. 17, 2015), D. Brizuela *et al.*, *Phys. Rev. D* **94**, arXiv: 1611.02932 (gr-qc) (Nov. 9, 2016), O. O. Novikov, arXiv: 1712.03939 (hep-th) (Dec. 11, 2017), A. Y. Kamenshchik *et al.*, *Class. Quantum Grav.* **35**, 015012 (Sept. 29, 2017), C. Kiefer, D. Wichmann, arXiv: 1802.01422 (gr-qc) (Feb. 5, 2018).


²³A. A. Andrianov *et al.*, *Theor. Math. Phys.* **184**, 1224–1233 (Mar. 18, 2015).

²⁴A. A. Andrianov *et al.*, *Int. J. Mod. Phys. D* **19**, 97–111 (Jan. 2010), A. A. Andrianov *et al.*, in *Springer Proceedings in Physics* (Springer International Publishing, 2016), pp. 29–44.



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $\lambda^2 = 3\kappa$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying V




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model $(+, -)$: csch

csch, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for quintessence model (+, -): csch

csch, with $V = \kappa^{-2}$ and $\lambda^2 = \kappa$; varying p_β




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for phantom model $(-, +)$

sec, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β



- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Integration of the transformed first Friedmann equation

General integral for $p_\beta \neq 0$

In order to integrate the equation under the implicit gauge

$$s \frac{p_\beta^2}{12} \left(-\ell \frac{\varkappa^{1/2}}{6} \left(\frac{d\chi}{d\beta} \right)^2 + \varkappa^{-1/2} \right) - \varkappa^{3/2} V e^{g\chi} = 0, \quad (14)$$

one can substitute

$$\gamma := \sqrt{\frac{3}{2\varkappa} g\beta}, \quad \tilde{\sigma}^2 := \frac{p_\beta^2}{12\varkappa^2 |V|} e^{-g\chi}, \quad (15)$$

to get

$$\left(\frac{d\tilde{\sigma}}{d\gamma} \right)^2 + \ell(s\upsilon - \tilde{\sigma}^2) = 0 \quad \Rightarrow \quad \frac{d\gamma}{d\tilde{\sigma}} = \pm \frac{1}{\sqrt{\ell(-s\upsilon + \tilde{\sigma}^2)}}, \quad (16)$$

which is of the standard inverse hyperbolic / trigonometric form **except for $(-, -)$** .



Integration of the separated minisuperspace WDW equation

In order to integrate the separated minisuperspace Wheeler–DeWitt equation

$$\mathbb{D}\psi(\chi) = k_\beta^2 \psi(\chi), \quad \mathbb{D} := -\ell \frac{6}{\varkappa} \partial_\chi^2 + \mathfrak{s} v \frac{12\varkappa^2 |V|}{\hbar^2}, \quad (17)$$

one can transform

$$\nu := \sqrt{\frac{2\varkappa}{3}} \frac{k_\beta}{g}, \quad \sigma^2 := \frac{8\varkappa^3 |V| e^{g\chi}}{\hbar^2 g^2}, \quad (18)$$

to get

$$\sigma^2 \psi''(\sigma) + \sigma \psi'(\sigma) + \ell(\nu^2 - \mathfrak{s} v \sigma^2) \psi(\sigma) = 0, \quad (19)$$

which is of the standard Besselian form.



Self-adjointness of unbounded operators

General theory

- Even in quantum geometrodynamics, the self-adjointness of the energy density is desirable, which is typically an unbounded operator in the Hilbert space \mathbf{F} endowed with the Schrödinger inner product $(\cdot, \cdot)_S$.
- Mathematically, an *unbounded* operator H is characterised not only by its action on a vector, but also by its domain $\text{Dom}(H) \subsetneq \mathbf{F}^{25}$.
- In addition to the *symmetry* $(H^\dagger \phi_1, \phi_2) \equiv (\phi_1, H\phi_2)$, the self-adjointness of an unbounded operator also requires $\text{Dom}(H^\dagger) = \text{Dom}(H)$.
- For quantum systems not bounded below, not only the self-adjointness, but also the symmetry of the energy density is not guaranteed automatically.
- Even when one could find a $\text{Dom}_0(H)$ such that H is symmetric, one would still be left with $\text{Dom}_0(H^\dagger) \supsetneq \text{Dom}_0(H)$ in general.
- Sloppily speaking, the process of extending $\text{Dom}(H)$ such that $\text{Dom}(H^\dagger) = \text{Dom}(H)$ is called **self-adjoint extension**²⁶; if the extension is unique, the operator is called *essentially self-adjoint*.

²⁵B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267 , ch. 9.

²⁶D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012) , G. Bonneau *et al.*, *Am. J. Phys* **69**, 322–331 (Mar. 28, 2001), V. S. Araujo *et al.*, *Am. J. Phys* **72**, 203–213 (Feb. 2004), A. M. Essin, D. J. Griffiths, *Am. J. Phys* **74**, 109–117 (Feb. 2006).

