

Integrable Liouville Cosmological Models

The self-adjointness of Hamiltonian and the Semi-classical Wave Functions

Alexander A. Andrianov^{1,4} Chen Lan² Oleg O. Novikov¹ Yi-Fan Wang³

¹Saint-Petersburg State University, Ulyanovskaya str. 1, Petrodvorets, Sankt-Petersburg 198504, Russland

²ELI-ALPS Research Institute, Budapesti út 5, H-67228 Szeged, Ungarn

³Institut für Theoretische Physik, Universität zu Köln, Zülpicher Straße 77, D-50937 Köln, Deutschland

⁴Institut de Ciències del Cosmos, Universitat de Barcelona, Martí i Franquès 1, E-08028 Barcelona, Spanien

March 23, 2018



Outline

1. Introduction

2. The classical model and the implicit trajectories

Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian

Dirac quantisation and the quantum-mechanical analogy

The self-adjointness of Hamiltonian

4. The semi-classical wave packets

The Wentzel–Kramers–Brillouin approximation

The inner product and wave packet

5. Conclusions



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
The Wentzel–Kramers–Brillouin approximation
The inner product and wave packet
5. Conclusions



Introduction

The Friedmann–Lemaître model with quintessence and phantom Liouville fields

- Dynamical Dark Energy has been modelled by quintessence¹ and phantom² matter, which can be realised by minimally-coupled real scalar fields with $\ell = \pm 1$ ³

$$\mathcal{S}_L = \int d^4x \sqrt{-g} \left\{ -\ell \frac{g^{\mu\nu}}{2} (\partial_\mu \phi)(\partial_\nu \phi) - \mathcal{V}(\phi) \right\}. \quad (1)$$

- The Liouville field⁴ $\mathcal{V}(\phi) = V e^{\lambda\phi}$ is of interest, where $\lambda, V \in \mathbb{R}$.
- Assume a flat Robertson–Walker metric $g_{\mu\nu} dx^\mu dx^\nu = -N^2(t) dt^2 + \kappa e^{2\alpha(t)} d\Omega_{3F}^2$, where $\kappa := 8\pi G$, $d\Omega_{3F}^2 := d\chi^2 + \chi^2(d\theta^2 + \sin^2 \theta d\varphi^2)$, and N lapse function.
- The total action reads $\mathcal{S} := \mathcal{S}_{EH} + \mathcal{S}_{GHY} + \mathcal{S}_L = \int d\Omega_{3F}^2 \int dt L$, where

$$L := \kappa^{3/2} N e^{3\alpha} \left(-\frac{3}{\kappa} \frac{\dot{\alpha}^2}{N^2} + \ell \frac{\dot{\phi}^2}{2N^2} - V e^{\lambda\phi} \right). \quad (2)$$

¹R. R. Caldwell *et al.*, *Phys. Rev. Lett.* **80**, 1582–1585 (Feb. 1998).

²R. R. Caldwell, *Phys. Lett. B* **545**, 23–29 (Oct. 2002).

³The signature of metric is $(-, +, \dots, +)$.

⁴Y. Nakayama, *Int. J. Mod. Phys. A* **19**, 2771–2930 (July 2004).



Introduction

Highlights

Integrability

Implicit trajectories can be obtained explicitly; the minisuperspace Wheeler–DeWitt equation can be integrated exactly.

Self-adjointness

The Hamiltonian can be asymmetric; imposing self-adjointness leads to removal of degeneracy and discretisation of levels.

Semi-classical wave functions

Semi-classical wave functions can be constructed by the principle of constructive interference, the Wentzel–Kramers–Brillouin approximation and numerical methods.



Outline

1. Introduction

2. The classical model and the implicit trajectories

Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian

Dirac quantisation and the quantum-mechanical analogy

The self-adjointness of Hamiltonian

4. The semi-classical wave packets

The Wentzel–Kramers–Brillouin approximation

The inner product and wave packet

5. Conclusions



Decoupling the variables

Via orthogonal transformation

- By rescaling $\bar{N} := N e^{-3\alpha}$ and the orthogonal transformation

$$\begin{pmatrix} \alpha(t) \\ \phi(t) \end{pmatrix} = \frac{s}{g} \begin{pmatrix} \lambda & -\ell \nu \\ -6 & \lambda \end{pmatrix} \begin{pmatrix} \beta(t) \\ \chi(t) \end{pmatrix} \quad (3)$$

the system can be decoupled ($\Delta := \lambda^2 - 6\ell\nu$, $s := \operatorname{sgn} \Delta$ and $g := s\sqrt{|\Delta|} \equiv s\sqrt{s\Delta}$.)

$$L = \nu^{3/2} \bar{N} \left(-s \frac{3}{\pi} \frac{\dot{\beta}^2}{\bar{N}^2} + \ell s \frac{\dot{\chi}^2}{2\bar{N}^2} - V e^{g\chi} \right), \quad (4)$$

- Using the conservation of p_β ⁵, $\bar{N}(t)$ can be eliminated, leading to the exact integrations of the equations of motion as implicit trajectories.
 - $p_\beta \neq 0$: four cases according to $(\ell, s\nu)$, $v = \operatorname{sgn} V$
 - $p_\beta = 0$: well-known power-law special solution for $(+, -)$ and $(-, +)$.


⁵The same first integral has been found in C. Lan, PhD thesis, Saint Petersburg State University, 2016, <https://search.rsl.ru/ru/record/01006663434>, A. A. Andrianov et al., *Theor. Math. Phys.* **184**, 1224–1233 (Sept. 2015), in canonical formalism.



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ


$$e^{g\chi} = \frac{p_\beta^2}{12\kappa^2|V|} \operatorname{sech}^2\left(\sqrt{\frac{3}{2\kappa}} g(\beta - \beta_0)\right) \quad (5)$$



Trajectories for quintessence model $(+, -)$

csch , with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ


$$e^{g\chi} = \frac{p_\beta^2}{12\kappa^2|V|} \text{csch}^2\left(\sqrt{\frac{3}{2\kappa}} g(\beta - \beta_0)\right) \quad (6)$$



Trajectories for phantom model $(-, +)$

sec, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$

$$\mathbb{E}^{g\chi} = \frac{p_\beta^2}{12\varkappa^2|V|} \sec^2\left(\sqrt{\frac{3}{2\varkappa}} g(\beta - \beta_0)\right) \quad (7)$$



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
The Wentzel–Kramers–Brillouin approximation
The inner product and wave packet
5. Conclusions



Dirac quantisation

- The primary Hamiltonian and the Hamiltonian constraint⁶ reads

$$H^p = \bar{N} H_{\perp} + v^{\bar{N}} p_{\bar{N}}, \quad (8)$$

$$H_{\perp} := -\beta \frac{p_{\beta}^2}{12\nu^{1/2}} + \ell \beta \frac{p_{\chi}^2}{2\nu^{3/2}} + \nu^{3/2} V e^{g\chi}. \quad (9)$$

- Applying the Dirac quantisation rules with the Laplace–Beltrami operator ordering⁷, one gets the minisuperspace Wheeler–DeWitt equation with (β, χ)

$$0 = \widehat{H}_{\perp} \Psi(\beta, \chi) := \left(\beta \frac{\hbar^2}{12\nu^{1/2}} \partial_{\beta}^2 - \ell \beta \frac{\hbar^2}{2\nu^{3/2}} \partial_{\chi}^2 + \nu^{3/2} V e^{g\chi} \right) \Psi. \quad (10)$$

- Equation (10) is Klein–Gordon-like, hyperbolic for $\ell = +1$ and *elliptic* for $\ell = -1$.

⁶D. M. Gitman, I. V. Tyutin, *Quantization of Fields with Constraints*, (Springer, 1990), H. J. Rothe, K. D. Rothe, *Classical and Quantum Dynamics of Constrained Hamiltonian Systems*, (World Scientific, Apr. 2010).

⁷C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012), ch. 8.



Integration of the Wheeler–DeWitt equation

The quantum-mechanical analogy

- Fourier transforming β to $k_\beta \in \mathbb{R}$ yields the time-independent Schrödinger equation

$$E\psi(x) := \ell \frac{\hbar^2 \tilde{g}^2 \nu^2}{8M_P} \psi = \widehat{H}_{\text{eff}} \psi := \left(-\frac{\hbar^2}{2M_P} \partial_x^2 + V_{\text{eff}} \right) \psi, \quad V_{\text{eff}} := \ell s v \widetilde{V} e^{\tilde{g}x}, \quad (11)$$

in which $M_P := \sqrt{\frac{\hbar}{\nu}}$, $x := \sqrt{\hbar\nu} \cdot \chi$, $\widetilde{V} := M_P^{-3} |V| \geq 0$, and $\nu = \sqrt{\frac{2\nu}{3}} \frac{|k_\beta|}{\tilde{g}} \geq 0$.

- V_{eff} is a special case of the Morse potential⁸.
- Let $e^{2y} := 8M_P \widetilde{V} / \hbar^2 \tilde{g}^2 \cdot e^{\tilde{g}x}$

	$E \geq 0$	$E < 0$
$V_{\text{eff}} > 0$	$(+, +); K_{i\nu}(e^y)$	$(-, -); \text{none}$
$V_{\text{eff}} < 0$	$(+, -); F_{i\nu}(e^y), G_{i\nu}(e^y)$	$(-, +); J_\nu(e^y)$

⁸P. M. Morse, *Phys. Rev.* **34**, 57–64 (July 1929), D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012).



Self-adjointness of the Hamiltonian

Results for the Liouville potential¹²

- $k_\beta \in \mathbb{R}$, enforcing the self-adjointness of \widehat{H}_{eff} .
- For $V_{\text{eff}} > 0$: \widehat{H}_{eff} is **essentially self-adjoint**⁹.
 - Quintessence (+, +): nothing special.
- For $V_{\text{eff}} < 0$: \widehat{H}_{eff} admits a **family of self-adjoint extensions**.
 - Quintessence (+, -): "eigenfunctions" are lin. combi. of F and G; **the degeneracy is removed**
 - Phantom (-, +): the eigenfunctions are **discretised**

$$\psi_{-, \nu}(e^y) \propto J_\nu(e^y), \quad \nu = 2n + a_<, \quad n \in \mathbb{Z}_+, \quad (12)$$

and the corresponding full wave-functions are Bloch-periodic Imposing (anti-)periodic boundary condition fixes $a_< = 0$ (1).

- The self-adjointness has been discussed in quantum cosmology in e.g.¹⁰ (in the operator ordering scenario) and more recently in¹¹.

⁹See also B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267 , sec. 9.9.

¹⁰C. R. Almeida et al., *Gravitation Cosmol.* **21**, 191–199 (July 2015).

¹¹S. Gryb, K. P. Y. Thébault, arXiv: 1801.05789 (gr-qc) (Jan. 17, 2018).

¹²See also D. M. Gitman et al., *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012) , sec. 8.5.



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian

4. The semi-classical wave packets
The Wentzel–Kramers–Brillouin approximation
The inner product and wave packet

5. Conclusions



Matching quantum number with classical first integral

Principle of constructive interference; $(+, +)$ as the example

- Write the mode function in the Wentzel–Kramers–Brillouin form
 $\Psi_\nu(\beta, \chi) \sim \sqrt{\rho(\beta, \chi)} \exp\left\{\frac{i}{\hbar} S(\beta, \chi)\right\}$. For $S/\hbar \gg 1$ and $\nu \gg 1$, $S(\beta, \chi)$ becomes the Hamilton principal function in the leading-order approximation, which is stationary with respect to the variation of integral constants $\partial S / \partial \nu = 0$.
- Fixing $\nu/\sigma > 1$,

$$K_{\pm\nu}(\sigma) \sim \frac{\sqrt{2\pi} \cos\left(\sqrt{\nu^2 - \sigma^2} - \nu \arccos \frac{\nu}{\sigma} - \frac{\pi}{4}\right)}{(\nu^2 - \sigma^2)^{1/4} e^{\mp\nu/2}} + O\left(\frac{1}{\sigma}\right). \quad (13)$$

- Applying *the principle* to the phase of $\Psi_\nu(\sigma)$, one has $\sigma/\nu = \operatorname{sech}(\beta_\beta \gamma)$, which matches the trajectory if $\hbar k_\beta \equiv \hbar \sqrt{\frac{3}{2\nu}} g\nu = p_\beta$.
- Fixing $\nu/\sigma < 1$, the asymptotic expansion is not oscillatory, but exponential; it is not within the Wentzel–Kramers–Brillouin regime.
- The conclusions also hold for $F_{\pm\nu}(\sigma)$, $G_{\pm\nu}(\sigma)$ for $(+, -)$, and $J_\nu(\sigma)$ for $(-, +)$.



Inner products for wave functions

Schrödinger and Mostafazadeh inner product

- A common starting point is the Schrödinger product¹³

$$(\Psi_1, \Psi_2)_S := \int d\chi \Psi_1^*(\beta, \chi) \Psi_2(\beta, \chi); \quad (14)$$

- $(\Psi, \Psi)_S$ is positive-definite, and the integrand $\rho_S(\beta, \chi)$ is non-negative
- The corresponding Schrödinger current is real but does not satisfy continuity equation
- Mostafazadeh¹⁴ found an inner product for self-adjoint \mathbb{D} with positive spectrum:

$$(\Psi_1, \Psi_2)_M^\kappa := \kappa \left\{ (\Psi_1, \mathbb{D}^{+1/2} \Psi_2)_S + (\partial_\beta \Psi_1, \mathbb{D}^{-1/2} \partial_\beta \Psi_2)_S \right\}, \quad \kappa > 0. \quad (15)$$

- Real power of \mathbb{D} is defined by spectral decomposition
- It can be shown¹⁵ that the density

$$\varrho_M^\kappa := \kappa \left\{ |\mathbb{D}^{+1/4} \Psi|^2 + |\mathbb{D}^{-1/4} \partial_\beta \Psi|^2 \right\} \quad (16)$$

- is equivalent to the integrand ρ_M^κ up to a boundary term $\int d\chi \varrho_M^\kappa = \int d\chi \rho_M^\kappa \equiv (\Psi_1, \Psi_2)_M^\kappa$
- is non-negative
- The corresponding current $\vec{\mathcal{J}}_M^\kappa$ is real but not conserved¹⁶.

¹³C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012), ch. 5.

¹⁴A. Mostafazadeh, *Class. Quantum Grav.* **20**, 155–171 (Dec. 2002).

¹⁵A. Mostafazadeh, F. Zamani, *Ann. Phys.* **321**, 2183–2209 (Sept. 2006).

¹⁶B. Rosenstein, L. P. Horwitz, *J. Phys. A: Math. Gen.* **18**, 2115–2121 (Aug. 1985).



Wave packets of Gaussian amplitude for continuous spectrum

Quintessence models

- It is difficult to find an amplitude such that the wave packet is Gaussian
- Instead, one can choose a Gaussian amplitude

$$A(\nu; \bar{\nu}, \sigma) := \left(\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\nu - \bar{\nu})^2}{2\sigma^2}\right) \right)^{1/2} \quad (17)$$

- In¹⁷, $A(\nu; \bar{\nu}, \sigma/\sqrt{2})$ was chosen.


¹⁷ M. P. Dąbrowski et al., *Phys. Rev. D* **74**, arXiv: hep-th/0605229 (hep-th) (Aug. 2006).




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\kappa^{1/2}/5$, $V = +\kappa^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger



Mostafazadeh



Wave packets with Poissonian amplitude for discrete spectrum

Discrete phantom model $(-, +)$

- Gaussian distribution works for continuous variable
- For discrete spectrum, one can choose a Poissonian amplitude

$$A_n(\bar{n}) := \left(e^{-\bar{n}} \frac{\bar{n}^n}{n!} \right)^{1/2} \quad (18)$$

- In¹⁸C, $A_n(\bar{n}/\sqrt{2})$ was chosen.


¹⁸C. Kiefer, *Nucl. Phys. B* **341**, 273–293 (Sept. 1990).




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger




Mostafazadeh




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger



Mostafazadeh



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
The Wentzel–Kramers–Brillouin approximation
The inner product and wave packet
5. Conclusions



Highlights

Revisited

- An **integral of motion** was found for the Liouville cosmological model.
 - Implicit trajectories in minisuperspace were obtained explicitly.
 - The minisuperspace Wheeler–Dewitt equation was integrated exactly.
- The **self-adjointness** of matter Hamiltonian was found to be non-trivial.
 - The degeneracy of the quintessence model $(+, -)$ was removed.
 - The levels of the phantom model $(-, +)$ were discretised.
- The **semi-classical wave functions** were compared with the classical trajectory.
 - Mode functions in the Wentzel–Kramers–Brillouin approximation were matched with the classical Hamilton principal function.
 - Semi-classical wave packets were constructed and compared with the classical trajectory under the Schrödinger and Mostafazadeh inner product.




Outlook

- Beyond isotropy: Bianchi models
- Beyond homogeneity: cosmological perturbations
- Beyond single Liouville field: multiple potentials / multiple fields
- Beyond classical matter and Schrödinger inner product: PT -symmetric field



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $\lambda^2 = 3\kappa$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying V




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model $(+, -)$: csch

csch, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for quintessence model (+, -): csch

csch, with $V = \kappa^{-2}$ and $\lambda^2 = \kappa$; varying p_β




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for phantom model $(-, +)$

sec, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β



- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Integration of the transformed first Friedmann equation

General integral for $p_\beta \neq 0$

In order to integrate the equation under the implicit gauge

$$s \frac{p_\beta^2}{12} \left(-\ell \frac{\varkappa^{1/2}}{6} \left(\frac{d\chi}{d\beta} \right)^2 + \varkappa^{-1/2} \right) - \varkappa^{3/2} V e^{g\chi} = 0, \quad (19)$$

one can substitute

$$\gamma := \sqrt{\frac{3}{2\varkappa} g\beta}, \quad \tilde{\sigma}^2 := \frac{p_\beta^2}{12\varkappa^2 |V|} e^{-g\chi}, \quad (20)$$

to get

$$\left(\frac{d\tilde{\sigma}}{d\gamma} \right)^2 + \ell(s\upsilon - \tilde{\sigma}^2) = 0 \quad \Rightarrow \quad \frac{d\gamma}{d\tilde{\sigma}} = \pm \frac{1}{\sqrt{\ell(-s\upsilon + \tilde{\sigma}^2)}}, \quad (21)$$

which is of the standard inverse hyperbolic / trigonometric form **except for $(-, -)$** .



Integration of the separated minisuperspace WDW equation

In order to integrate the separated minisuperspace Wheeler–DeWitt equation

$$\mathbb{D}\psi(\chi) = k_\beta^2 \psi(\chi), \quad \mathbb{D} := -\ell \frac{6}{\varkappa} \partial_\chi^2 + \mathfrak{s} v \frac{12\varkappa^2 |V|}{\hbar^2}, \quad (22)$$

one can transform

$$\nu := \sqrt{\frac{2\varkappa}{3}} \frac{k_\beta}{g}, \quad \sigma^2 := \frac{8\varkappa^3 |V| e^{g\chi}}{\hbar^2 g^2}, \quad (23)$$

to get

$$\sigma^2 \psi''(\sigma) + \sigma \psi'(\sigma) + \ell(\nu^2 - \mathfrak{s} v \sigma^2) \psi(\sigma) = 0, \quad (24)$$

which is of the standard Besselian form.



Self-adjointness of unbounded operators

General theory

- Even in quantum geometrodynamics, the self-adjointness of the matter Hamiltonian is desirable, which is typically an unbounded operator in the Hilbert space \mathbf{F} endowed with the Schrödinger inner product $(\cdot, \cdot)_S$.
- Mathematically, an *unbounded* operator H is characterised not only by its action on a vector, but also by its domain $\text{Dom}(H) \subsetneq \mathbf{F}^{19}$.
- In addition to the *symmetry* $(H^\dagger \phi_1, \phi_2) \equiv (\phi_1, H\phi_2)$, the self-adjointness of an unbounded operator also requires $\text{Dom}(H^\dagger) = \text{Dom}(H)$.
- For quantum systems not bounded below, not only the self-adjointness, but also the symmetry of the Hamiltonian is not guaranteed automatically.
- Even when one could find a $\text{Dom}_0(H)$ such that H is symmetric, one would still be left with $\text{Dom}_0(H^\dagger) \supsetneq \text{Dom}_0(H)$ in general.
- Sloppily speaking, the process of extending $\text{Dom}(H)$ such that $\text{Dom}(H^\dagger) = \text{Dom}(H)$ is called **self-adjoint extension**²⁰; if the extension is unique, the operator is called *essentially self-adjoint*.

¹⁹B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267 , ch. 9.

²⁰D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012) , G. Bonneau *et al.*, *Am. J. Phys* **69**, 322–331 (Mar. 28, 2001), V. S. Araujo *et al.*, *Am. J. Phys* **72**, 203–213 (Feb. 2004), A. M. Essin, D. J. Griffiths, *Am. J. Phys* **74**, 109–117 (Feb. 2006).

