

Integrable Liouville Cosmological Models

The self-adjointness of Hamiltonian and the Semi-classical Wave Functions

Alexander A. Andrianov^{1,4} Chen Lan² Oleg O. Novikov¹ Yi-Fan Wang³

¹Saint-Petersburg State University, Ulyanovskaya str. 1, Petrodvorets, Sankt-Petersburg 198504, Russland

²ELI-ALPS Research Institute, Budapesti út 5, H-67228 Szeged, Ungarn

³Institut für Theoretische Physik, Universität zu Köln, Zülpicher Straße 77, D-50937 Köln, Deutschland

⁴Institut de Ciències del Cosmos, Universitat de Barcelona, Martí i Franquès 1, E-08028 Barcelona, Spanien

March 23, 2018



Outline

1. Introduction

2. The classical model and the implicit trajectories

Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian

Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian

4. The semi-classical wave packets

Inner product and wave packet

5. Conclusions



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
Inner product and wave packet
5. Conclusions



Introduction

The Friedmann–Lemaître model with quintessence and phantom Liouville fields

- Dynamical Dark Energy has been modelled by quintessence¹ and phantom² matter, which can be realised by minimally-coupled real scalar fields with $\ell = \pm 1$ ³

$$\mathcal{S}_L = \int d^4x \sqrt{-g} \left\{ -\ell \frac{g^{\mu\nu}}{2} (\partial_\mu \phi)(\partial_\nu \phi) - \mathcal{V}(\phi) \right\}. \quad (1)$$

- The Liouville field⁴ $\mathcal{V}(\phi) = V e^{\lambda\phi}$ is of interest, where $\lambda, V \in \mathbb{R}$.
- Assume a flat Robertson–Walker metric $g_{\mu\nu} dx^\mu dx^\nu = -N^2(t) dt^2 + \varkappa e^{2\alpha(t)} d\Omega_{3F}^2$, where $\varkappa := 8\pi G$, $d\Omega_{3F}^2 := d\chi^2 + \chi^2(d\theta^2 + \sin^2 \theta d\varphi^2)$, and N lapse function.
- The total action reads $\mathcal{S} := \mathcal{S}_{EH} + \mathcal{S}_{GHY} + \mathcal{S}_L = \int d\Omega_{3F}^2 \int dt L$, where

$$L := \varkappa^{3/2} N e^{3\alpha} \left(-\frac{3}{\varkappa} \frac{\dot{\alpha}^2}{N^2} + \ell \frac{\dot{\phi}^2}{2N^2} - V e^{\lambda\phi} \right). \quad (2)$$

¹R. R. Caldwell *et al.*, *Phys. Rev. Lett.* **80**, 1582–1585 (Feb. 1998).

²R. R. Caldwell, *Phys. Lett. B* **545**, 23–29 (Oct. 2002).

³The signature of metric is mostly positive.

⁴Y. Nakayama, *Int. J. Mod. Phys. A* **19**, 2771–2930 (July 2004).



Introduction

Highlights

Integrability

Implicit trajectories can be obtained explicitly; the minisuperspace Wheeler–DeWitt equation can be integrated exactly.

Self-adjointness

The Hamiltonian is not even naturally symmetric; imposing self-adjointness leads to non-trivial physical results.

Semi-classical wave functions

Semi-classical wave functions can be constructed by the principle of constructive interference, the JWKB approximation and numerical methods.



Outline

1. Introduction

2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian

4. The semi-classical wave packets
Inner product and wave packet

5. Conclusions



Decoupling the variables

Via orthogonal transformation

- By rescaling $\bar{N} := N e^{-3\alpha}$ and the orthogonal transformation

$$\begin{pmatrix} \alpha(t) \\ \phi(t) \end{pmatrix} = \frac{\varsigma}{g} \begin{pmatrix} \lambda & -\ell \varkappa \\ -6 & \lambda \end{pmatrix} \begin{pmatrix} \varsigma_\beta \cdot \beta(t) \\ \varsigma_\chi \cdot \chi(t) \end{pmatrix} \quad \text{where } \varsigma_\beta, \varsigma_\chi = \pm 1, \quad (3)$$

the effective Lagrangian eq. (2) can be decoupled ($\varsigma_\beta = \varsigma_\chi = +1$ from now on)

$$L = \varkappa^{3/2} \bar{N} \left(-\varsigma \frac{3}{\varkappa} \frac{\dot{\beta}^2}{\bar{N}^2} + \ell \varsigma \frac{\dot{\chi}^2}{2\bar{N}^2} - V e^{g\chi} \right), \quad (4)$$

where $\Delta := \lambda^2 - 6\ell\varkappa$, $\varsigma := \operatorname{sgn} \Delta$ and $g := \varsigma \sqrt{|\Delta|} \equiv \varsigma \sqrt{3\Delta}$.

- The Euler–Lagrange equations with respect to \bar{N} , β and χ will be called the modified first, second Friedmann equations and the Klein–Gordon equation, respectively.
- Since $\beta(t)$ is cyclic, its conjugate momentum p_β is conserved⁵, and the modified second Friedmann equation can be readily integrated.

⁵The same first integral has been found in C. Lan, PhD thesis, Saint Petersburg State University, 2016, <https://search.rsl.ru/ru/record/01006663434>, A. A. Andrianov et al., *Theor. Math. Phys.* **184**, 1224–1233 (Sept. 2015), in canonical formalism.



Integration and the implicit trajectories

- For $p_\beta \neq 0$, fixing the *implicit gauge* $\bar{N} = -6\mathfrak{s}\sqrt{\nu}\dot{\beta}/p_\beta$, the modified first Friedmann equation can be integrated, yielding the **implicit** trajectory

$$e^{g\chi} = \frac{p_\beta^2}{12\nu^2|V|} s^2 \left(\sqrt{\frac{3}{2\nu}} g(\beta - \beta_0) \right), \quad (5)$$

in which $v := \operatorname{sgn} V$, $(\operatorname{sgn}, \operatorname{sgn})$ means $(\ell, \mathfrak{s}v)$, and

$$\begin{aligned} (+, +)s(\gamma) &:= \operatorname{sech}(\gamma), & (+, -)s(\gamma) &:= \operatorname{csch}(\gamma), \\ (-, +)s(\gamma) &:= \sec(\gamma), & (-, -)s(\gamma) &:= \mathbb{i} \csc(\gamma). \end{aligned} \quad (6)$$

- The integral for the phantom model with $(-, +)$ is periodic for β with $T_\beta = \sqrt{\frac{2\nu}{3}} \frac{2\pi}{g}$.
- The integral for the phantom model with $(-, -)$ is not real.
- The trajectories can be parametrised by β , inspiring recognising β as a ‘time variable’.
- For $p_\beta = 0$, integrating the modified second Friedmann equation yields $\beta \equiv \beta_0$ or $\phi - \phi_0 = -\ell \lambda \alpha / \nu^6$, which is the well-known power-law special solution⁷.

⁶It only works for $(+, -)$ or $(-, +)$ if one checks consistency with the modified first Friedmann equation.

⁷For instance, A. R. Liddle, D. H. Lyth, *Cosmological Inflation and Large-Scale Structure*, (Cambridge University, 2000) , ch. 3.



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral

3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian

4. The semi-classical wave packets
Inner product and wave packet

5. Conclusions



Dirac quantisation

- The primary Hamiltonian and the Hamiltonian constraint⁸ reads

$$H^p = \bar{N} H_{\perp} + v^{\bar{N}} p_{\bar{N}}, \quad (7)$$

$$H_{\perp} := -\beta \frac{p_{\beta}^2}{12\nu^{1/2}} + \ell \beta \frac{p_{\chi}^2}{2\nu^{3/2}} + \nu^{3/2} V e^{g\chi}. \quad (8)$$

- Applying the Dirac quantisation rules with the Laplace–Beltrami operator ordering⁹, one gets the minisuperspace Wheeler–DeWitt equation with (β, χ)

$$0 = \widehat{H}_{\perp} \Psi(\beta, \chi) := \left(\beta \frac{\hbar^2}{12\nu^{1/2}} \partial_{\beta}^2 - \ell \beta \frac{\hbar^2}{2\nu^{3/2}} \partial_{\chi}^2 + \nu^{3/2} V e^{g\chi} \right) \Psi. \quad (9)$$

- Equation (9) is Klein–Gordon-like, hyperbolic for $\ell = +1$ and *elliptic* for $\ell = -1$.

⁸D. M. Gitman, I. V. Tyutin, *Quantization of Fields with Constraints*, (Springer, 1990), H. J. Rothe, K. D. Rothe, *Classical and Quantum Dynamics of Constrained Hamiltonian Systems*, (World Scientific, Apr. 2010).

⁹C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012), ch. 8.



Integration of the Wheeler–DeWitt equation

The quantum-mechanical analogy

- Fourier transforming β to $k_\beta \in \mathbb{R}$ yields the time-independent Schrödinger equation

$$\ell \frac{\hbar^2 \tilde{g}^2 \nu^2}{8M_P} \psi(x) = \widehat{H}_{\text{eff}} \psi(x) := \left(-\frac{\hbar^2}{2M_P} \partial_x^2 + \widehat{V}_{\text{eff}}(x) \right) \psi(x), \quad V_{\text{eff}} = \ell s v \widetilde{V} e^{\tilde{g} x}, \quad (10)$$

in which $M_P := \sqrt{\frac{\hbar}{\kappa}}$, $x := \sqrt{\hbar \kappa} \cdot \chi$, $\widetilde{V} := M_P^{-3} |V| \geq 0$, and $\nu = \sqrt{\frac{2\kappa}{3}} \frac{|k_\beta|}{\tilde{g}} \geq 0$.

- V_{eff} is a special case of the Morse potential¹⁰.
- For the quintessence model $(+, +)$: V_{eff} is bounded **below**; generalised eigenfunctions are the Besselian $K_{\pm\nu}(e^{\tilde{x}})$'s, $e^{2\tilde{x}} = 8M_P \widetilde{V}/\hbar^2 \tilde{g}^2 \cdot e^{\tilde{g} x}$.
- For the quintessence model $(+, -)$: V_{eff} is bounded **above**; generalised eigenfunctions for positive spectrum (scattering states) are the Besselian $F_{\pm\nu}(e^{\tilde{x}})$'s and $G_{\pm\nu}(e^{\tilde{x}})$ 's¹¹.
- For the phantom model $(-, +)$: V_{eff} is bounded **above**; eigenfunctions for negative eigenvalues (bound states) are the Besselian $J_\nu(e^{\tilde{x}})$'s.

¹⁰P. M. Morse, *Phys. Rev.* **34**, 57–64 (July 1929), D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012).

¹¹These are linear combinations of $J_{\pm\pm\nu}(e^{\tilde{x}})$; see T. M. Dunster, *SIAM J. Math. Anal.* **21**, 995–1018 (July 1990).



The self-adjointness of unbounded operators

The general theory

- Even in quantum geometrodynamics, the self-adjointness of the matter Hamiltonian is desirable, which is typically an unbounded operator in the Hilbert space \mathbf{F} endowed with the Schrödinger inner product $(\cdot, \cdot)_S$.
- Mathematically, an *unbounded* operator H is characterised not only by its action on a vector, but also by its domain $\text{Dom}(H) \subsetneq \mathbf{F}^{12}$.
- In addition to the *symmetry* $(\phi_1, H^\dagger \phi_2) \equiv (H\phi_1, \phi_2)$, the self-adjointness of an unbounded operator also requires $\text{Dom}(H^\dagger) = \text{Dom}(H)$.
- For quantum systems not bounded below, not only the self-adjointness, but also the symmetry of the Hamiltonian is not guaranteed automatically.
- Even when one could find a $\text{Dom}_0(H)$ such that H is symmetric, one would still be left with $\text{Dom}_0(H^\dagger) \supsetneq \text{Dom}_0(H)$ in general.
- Sloppily speaking, the process of extending $\text{Dom}(H)$ such that $\text{Dom}(H^\dagger) = \text{Dom}(H)$ is called **self-adjoint extension**¹³; if the extension is unique, the operator is called *essentially self-adjoint*.

¹²B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267 , ch. 9.

¹³D. M. Gitman *et al.*, *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012) , G. Bonneau *et al.*, *Am. J. Phys* **69**, 322–331 (Mar. 28, 2001), V. S. Araujo *et al.*, *Am. J. Phys* **72**, 203–213 (Feb. 2004), A. M. Essin, D. J. Griffiths, *Am. J. Phys* **74**, 109–117 (Feb. 2006).



The self-adjointness of the Hamiltonian

Results for the special Morse potential¹⁵

- For the quintessence model $(+, +)$: \widehat{H}_{eff} is essentially self-adjoint on $C_c^\infty(\mathbb{R})$ ¹⁴.
- For $(+, -)$ and $(-, +)$: \widehat{H}_{eff} is not self-adjoint but admits a family of self-adjoint extensions parametrised by $a \in [0, 2)$.
- For the quintessence model $(+, -)$, the generalised eigenfunctions are

$$_{(+,-)}\psi_\nu \propto F_{\mathbb{I}\nu} \cos \frac{\mathbb{I}a_{\geq}}{2} + G_{\mathbb{I}\nu} \sin \frac{\mathbb{I}a_{\geq}}{2} \quad (11)$$

with asymptotic behaviour $_{(+,-)}\psi_\nu(e^{\tilde{x}}) \sim e^{-\tilde{x}/2} \cos(e^{\tilde{x}} - \frac{\mathbb{I}a}{2} - \frac{\mathbb{I}}{4})$ as $x \rightarrow +\infty$.

- For the phantom model $(-, +)$, the eigenfunctions are

$$_{(-,+)}\psi_\nu \propto J_\nu, \quad \nu = 2n + a_<, n \in \mathbb{Z}_{\geq}, \quad (12)$$

and the corresponding full wave-functions are Bloch-periodic

$$_{(-,+)}\Psi_n(\beta, \chi) = e^{\mathbb{I}\pi a_<} {}_{(-,+)}\Psi_n(\beta + T_\beta, \chi).$$

- Equation (12) can also be shown by noting $(J_\mu, \widehat{H}_{\text{eff}} J_\nu)_S - (\widehat{H}_{\text{eff}} J_\mu, J_\nu)_S \propto \sin \frac{\mathbb{I}(\mu - \nu)}{2}$ and then imposing symmetry.
- Imposing (anti-)periodic boundary condition would fix $a_< = 0$ (1).

¹⁴ See also B. C. Hall, *Quantum Theory for Mathematicians*, (Springer, 2013), vol. 267, sec. 9.9.

¹⁵ D. M. Gitman et al., *Self-adjoint Extensions in Quantum Mechanics*, (Birkhäuser, 2012), sec. 8.5.



The self-adjointness of the Hamiltonian

Discussions

- The self-adjointness of \widehat{H}_{eff} is non-trivial and not unique for $(+, -)$ and $(-, +)$.
- For the phantom model $(-, +)$, it **discretises** the spectrum.
- The self-adjointness has been discussed in quantum cosmology in e.g.¹⁶ (in the operator ordering scenario) and more recently in¹⁷.
- The self-adjoint extension is also non-trivial for the x^{-2} potential¹⁸, which is of cosmological relevance as well¹⁹.

¹⁶C. R. Almeida *et al.*, *Gravitation Cosmol.* **21**, 191–199 (July 2015).

¹⁷S. Gryb, K. P. Y. Thébault, arXiv: 1801.05789 (gr-qc) (Jan. 17, 2018).

¹⁸A. M. Essin, D. J. Griffiths, *Am. J. Phys.* **74**, 109–117 (Feb. 2006).

¹⁹M. Bouhmadi-López *et al.*, *Phys. Rev. D* **79**, arXiv: 0905.2421 (gr-qc) (June 2009).



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
Inner product and wave packet
5. Conclusions



Inner product for wave functions

Schrödinger product

- To make sense of amplitude and wave packet, an inner product is necessary
- In terms of a Klein–Gordon equation, call β the “temporal” variable, and χ the “spacial” variable.
- A common starting point is the *Schrödinger product*²⁰

$$(\Psi_1, \Psi_2)_S := \int d\chi \Psi_1^*(\beta, \chi) \Psi_2(\beta, \chi); \quad (13)$$

- $(\Psi, \Psi)_S$ is **positive-definite**, and the integrand $\rho_S(\beta, \chi)$ is **non-negative**
- The corresponding Schrödinger current is **real** but **does not satisfy continuity equation** $\dot{\rho}_S + \nabla \cdot \vec{j}_S = 0$, because eq. (9) is KG-like.
- $K_{\parallel\nu}^{21}$ for $(+, +)$, $F_{\parallel\nu}$ and $G_{\parallel\nu}$ for $(+, -)$ can be proved to be **orthogonal** and **complete** among themselves, as well as **can be normalised**.
 - $J_{\parallel\nu}$'s for $(+, -)$ are **not orthogonal**

²⁰C. Kiefer, *Quantum Gravity*, (Oxford University, ed. 3, Apr. 2012) , ch. 5.

²¹S. B. Yakubovich, *Opuscula Math.* **26**, 161–172 (2006), A. Passian *et al.*, *J. Math. Anal. Appl.* **360**, 380–390 (Dec. 2009), R. Szmytkowski, S. Bielski, *J. Math. Anal. Appl.* **365**, 195–197 (May 2010).



Further inner products for wave functions

Klein–Gordon and Mostafazadeh product

- Since eq. (9) is KG-like, another popular choice is the KG product

$$(\Psi_1, \Psi_2)_{\text{KG}}^g := \mathbb{i}g \left\{ (\Psi_1, \partial_\beta \Psi_2)_S - (\partial_\beta \Psi_1, \Psi_2)_S \right\}, \quad g > 0. \quad (14)$$

- $(\Psi, \Psi)_{\text{KG}}^g$ is **real** but **not positive-definite**, so does the integrand ρ_{KG} ;
- The corresponding \vec{J}_{KG} is **conserved** $\dot{\rho}_{\text{KG}} + \nabla \cdot \vec{J}_{\text{KG}} = 0$ and **real**.
- Mostafazadeh²² found a product *for Hermitian \mathbb{D} with positive spectrum*:

$$(\Psi_1, \Psi_2)_M^\kappa := \kappa \left\{ (\Psi_1, \mathbb{D}^{+1/2} \Psi_2)_S + (\partial_\beta \Psi_1, \mathbb{D}^{-1/2} \partial_\beta \Psi_2)_S \right\}, \quad \kappa > 0. \quad (15)$$

- $(\Psi, \Psi)_M^\kappa$ is **positive-definite**, but the integrand ρ_M^κ is **complex**
- The corresponding \vec{J}_M^κ is **conserved** $\dot{\rho}_M^\kappa + \nabla \cdot \vec{J}_M^\kappa = 0$ but also **complex**.

²²A. Mostafazadeh, *Class. Quantum Grav.* **20**, 155–171 (Dec. 2002).



Mostafazadeh inner product and the corresponding density

- Real power of \mathbb{D} is defined by spectral decomposition $\mathbb{D}^\gamma := \sum_\nu \nu^{2\gamma} \mathbf{P}_\nu$,
 $\mathbf{P}_\nu \Psi := \Psi_\nu (\Psi_\nu, \Psi)_S$.
- It can be shown²³ that *the density*

$$\varrho_M^\kappa := \kappa \left\{ |\mathbb{D}^{+1/4} \Psi|^2 + |\mathbb{D}^{-1/4} \partial_\beta \Psi|^2 \right\} \quad (16)$$

- is equivalent to ρ_M^κ up to a boundary term

$$\int d\chi \varrho_M^\kappa = \int d\chi \rho_M^\kappa \equiv (\Psi_1, \Psi_2)_M^\kappa; \quad (17)$$

- is non-negative.
- The corresponding current $\vec{\mathcal{J}}_M^\kappa$ is real but not conserved²⁴.

²³A. Mostafazadeh, F. Zamani, *Ann. Phys.* **321**, 2183–2209 (Sept. 2006).

²⁴B. Rosenstein, L. P. Horwitz, *J. Phys. A: Math. Gen.* **18**, 2115–2121 (Aug. 1985).



Wave packets of Gaussian amplitude for continuous spectrum

Quintessence models

- It is difficult to find an amplitude such that the wave packet is Gaussian
- Instead, one can choose a Gaussian amplitude

$$A(\nu; \bar{\nu}, \sigma) := \left(\frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\nu - \bar{\nu})^2}{2\sigma^2}\right) \right)^{1/2} \quad (18)$$

- In²⁵, $A(\nu; \bar{\nu}, \sigma/\sqrt{2})$ was chosen.


²⁵M. P. Dąbrowski et al., *Phys. Rev. D* **74**, arXiv: hep-th/0605229 (hep-th) (Aug. 2006).




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\ell\nu}$, with $\lambda = \varkappa^{1/2}/2$, $V = -\varkappa^{-2}$, $\bar{k}_\beta = -2$ and $\sigma_\beta = 5/4$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\nu^{1/2}/5$, $V = +\nu^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\parallel\nu}$, with $\lambda = 4\kappa^{1/2}/5$, $V = +\kappa^{-2}$, $\bar{k}_\beta = -7/2$ and $\sigma_\beta = 7/5$

Schrödinger



Mostafazadeh



Wave packets with Poissonian amplitude for discrete spectrum

Discrete phantom model $(-, +)$

- Gaussian distribution works for continuous variable
- For discrete spectrum, one can choose a Poissonian amplitude

$$A_n(\bar{n}) := \left(e^{-\bar{n}} \frac{\bar{n}^n}{n!} \right)^{1/2} \quad (19)$$

- In²⁶, $A_n(\bar{n}/\sqrt{2})$ was chosen.


²⁶C. Kiefer, *Nucl. Phys. B* **341**, 273–293 (Sept. 1990).




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger




Mostafazadeh




Wave packets of Poissonian amplitude for phantom model

J_{2n+1} , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$ and $\bar{k}_\beta = 8$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for phantom model

J_ν , with $\lambda = 2\kappa^{1/2}$, $V = +\kappa^{-2}$, $\bar{k}_\beta = 8$ and $\sigma_\beta = 11/2$

Schrödinger




Mostafazadeh




Wave packets of Gaussian amplitude for phantom model

J_ν , with $\lambda = 2\nu^{1/2}$, $V = +\nu^{-2}$, $\bar{k}_\beta = 8$ and $\sigma_\beta = 11/2$

Schrödinger



Mostafazadeh



Outline

1. Introduction
2. The classical model and the implicit trajectories
Lagrangian formalism and the first integral
3. Dirac quantisation and the self-adjointness of Hamiltonian
Dirac quantisation and the quantum-mechanical analogy
The self-adjointness of Hamiltonian
4. The semi-classical wave packets
Inner product and wave packet
5. Conclusions



Highlights

- An **integral of motion** was found for Liouville cosmological models.
- **Implicit trajectories** in minisuperspace were obtained.
- The levels of phantom Liouville model were found to be **discrete** due to the Hermiticity requirement of observables.
- The semi-classical quantum mode functions and wave packets was compared directly with the classical trajectory.



Issues

- In $(+, -)$ and $(-, +)$, wave packets contain multiple branches; however, the classical universe runs only on one trajectory.
- Quantum-corrected \bar{k}_β is to be understood.
- Wave packets with Gaussian profile are to be constructed, instead of inserting Gaussian / Poissonian amplitude artificially.
- A normalising κ for $(\cdot, \cdot)_M^\kappa$ is to be evaluated, otherwise a quantitative comparison of $(\cdot, \cdot)_S$ and $(\cdot, \cdot)_M^\kappa$ is not possible.



Outlook

- Beyond isotropy: generalise to Bianchi models
 - Bianchi Type-I: a natural extension, **under investigation**
- Beyond homogeneity: cosmological perturbation, **under investigation**
- Beyond single field
 - Two exponential potentials: $|V_1| = |V_2|$ and special λ_i
 - Multiple Liouville fields: mixing kinetic terms needed²⁷
- Beyond classic matter
 - PT -symmetric Liouville field²⁸: may cross the phantom divide $w = -1$.


²⁷A. A. Andrianov *et al.*, *Theor. Math. Phys.* **184**, 1224–1233 (Sept. 2015).

²⁸A. A. Andrianov *et al.*, *Int. J. Mod. Phys. D* **19**, 97–111 (Jan. 2010), A. A. Andrianov *et al.*, in *Springer Proceedings in Physics* (Springer International Publishing, 2016), pp. 29–44.



Trajectories for quintessence model $(+, +)$

sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model (+, +)

sech, with $\beta_0 = 0$, $\lambda^2 = 3\kappa$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying V




- has two asymptotes $\chi \propto \pm \beta$



Trajectories for quintessence model (+, -)

csch, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for quintessence model $(+, -)$: csch

csch, with $\beta_0 = 0$, $|V| = \kappa^{-2}$ and $p_\beta^2 = \kappa^2 \sqrt{|V|}$; varying λ




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for quintessence model (+, -): csch

csch, with $V = \kappa^{-2}$ and $\lambda^2 = \kappa$; varying p_β




- contains two distinct solutions, separated by $\beta = 0$
- has three asymptotes $\chi \propto \pm\beta$ and $\beta = 0$



Trajectories for phantom model $(-, +)$

csc, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \varkappa^{-2}$ and $p_\beta^2 = 3\varkappa^2 \sqrt{|V|}$




- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Trajectories for phantom model $(-, +)$

csc, with $V = \kappa^{-2}$ and $\lambda^2 = 3\kappa$; varying p_β



- contains infinite distinct solutions
- has infinite parallel asymptotes $\beta \propto (n + 1/2)\pi$



Integration of the transformed first Friedmann equation

General integral for $p_\beta \neq 0$

In order to integrate the equation under the implicit gauge

$$s \frac{p_\beta^2}{12} \left(-\ell \frac{\varkappa^{1/2}}{6} \left(\frac{d\chi}{d\beta} \right)^2 + \varkappa^{-1/2} \right) - \varkappa^{3/2} V e^{g\chi} = 0, \quad (20)$$

one can substitute

$$\gamma := \sqrt{\frac{3}{2\varkappa} g\beta}, \quad \tilde{\sigma}^2 := \frac{p_\beta^2}{12\varkappa^2 |V|} e^{-g\chi}, \quad (21)$$

to get

$$\left(\frac{d\tilde{\sigma}}{d\gamma} \right)^2 + \ell(s\upsilon - \tilde{\sigma}^2) = 0 \quad \Rightarrow \quad \frac{d\gamma}{d\tilde{\sigma}} = \pm \frac{1}{\sqrt{\ell(-s\upsilon + \tilde{\sigma}^2)}}, \quad (22)$$

which is of the standard inverse hyperbolic / trigonometric form **except for $(-, -)$.**



Integration of the separated minisuperspace WDW equation

In order to integrate the separated minisuperspace Wheeler–DeWitt equation

$$\mathbb{D}\psi(\chi) = k_\beta^2 \psi(\chi), \quad \mathbb{D} := -\ell \frac{6}{\varkappa} \partial_\chi^2 + \mathfrak{s} v \frac{12\varkappa^2 |V|}{\hbar^2}, \quad (\text{?? rev.})$$

one can transform

$$\nu := \sqrt{\frac{2\varkappa}{3}} \frac{k_\beta}{g}, \quad \sigma^2 := \frac{8\varkappa^3 |V| e^{g\chi}}{\hbar^2 g^2}, \quad (\text{?? rev.})$$

to get

$$\sigma^2 \psi''(\sigma) + \sigma \psi'(\sigma) + \ell(\nu^2 - \mathfrak{s} v \sigma^2) \psi(\sigma) = 0, \quad (23)$$

which is of the standard Besselian form.



