

# Integrable Cosmological Models with Liouville Fields

Alexander A. Andrianov<sup>1,4</sup> Chen Lan<sup>2</sup> Oleg O. Novikov<sup>1</sup>  
Yi-Fan Wang<sup>3</sup>

<sup>1</sup>Saint-Petersburg State University, St. Petersburg 198504, Russia

<sup>2</sup>ELI-ALPS, ELI-Hu NKft, Dugonics tér 13, Szeged 6720, Hungary

<sup>3</sup>Institut für Theoretische Physik, Universität zu Köln, Zülpicher Straße 77, 50937 Köln, Germany

<sup>4</sup>Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona, Spain

December 10, 2017



# Outline

## 1. Introduction

## 2. Classical model and the implicit trajectories

Lagrangian formalism

## 3. Quantised model and the wave packets

Dirac quantisation  
Semi-classical approximation  
Inner product and wave packet

## 4. Conclusions



# Outline

## 1. Introduction

## 2. Classical model and the implicit trajectories

Lagrangian formalism

## 3. Quantised model and the wave packets

Dirac quantisation  
Semi-classical approximation  
Inner product and wave packet

## 4. Conclusions



# Introduction

## Introduction

- 



# Outline

1. Introduction

2. Classical model and the implicit trajectories  
Lagrangian formalism

3. Quantised model and the wave packets  
Dirac quantisation  
Semi-classical approximation  
Inner product and wave packet

4. Conclusions



# The Friedmann–Lemaître model

123

- Flat Robertson–Walker metric  $\text{d}s^2 = -N^2(t) \text{d}t^2 + \varkappa^{-1} e^{2\alpha(t)} \text{d}\Omega_3^2$ 
  - $\varkappa := 8\pi G$ ;  $\text{d}\Omega_3^2$  dimensionless spacial metric
- Real Klein–Gordon field with potential  $V e^{\lambda\phi}$  (Liouville), where  $\lambda, V \in \mathbb{R}$ , and kinetic term with sign  $\ell = \pm 1$  (quintessence / phantom model).
- Total action  $\mathcal{S} := S_{\text{EH}} + S_{\text{GHY}} + S_L = \int \text{d}\Omega_3^2 \int \text{d}t L$ ,

$$L := \varkappa^{3/2} N e^{3\alpha} \left( -\frac{3}{\varkappa} \frac{\dot{\alpha}^2}{N^2} + \ell \frac{\dot{\phi}^2}{2N^2} - V e^{\lambda\phi} \right), \quad (1)$$

in which dot means  $\text{d}/\text{d}t$  and  $\ell = \pm 1$ .



# Decoupling the variables

Via *rescaled special orthogonal transformation*

- Setting  $\bar{N} := N e^{-3\alpha}$ , eq. (1) transforms to

$$L = \varkappa^{3/2} \bar{N} \left( -\frac{3}{\varkappa} \frac{\dot{\alpha}^2}{\bar{N}^2} + \ell \frac{\dot{\phi}^2}{2\bar{N}^2} - V e^{\lambda\phi+6\alpha} \right) \quad (2)$$

- Defining  $\Delta := \lambda^2 - 6\ell\varkappa$ ,  $s := \operatorname{sgn} \Delta$  and  $g := s\sqrt{|\Delta|} \equiv s\sqrt{s\Delta}$ , the *rescaled special orthogonal transformation*

$$\begin{pmatrix} \alpha \\ \phi \end{pmatrix} = \frac{s}{g} \begin{pmatrix} \lambda & -\ell\varkappa \\ -6 & \lambda \end{pmatrix} \begin{pmatrix} s_\beta \beta \\ s_\chi \chi \end{pmatrix} \quad \text{where } s_\beta, s_\chi = \pm 1 \quad (3)$$

gives the decoupled Lagrangian

$$L = \varkappa^{3/2} \bar{N} \left( -s \frac{3}{\varkappa} \frac{\dot{\beta}^2}{\bar{N}^2} + \ell s \frac{\dot{\chi}^2}{2\bar{N}^2} - V e^{s_\chi g \chi} \right). \quad (4)$$

- The Euler–Lagrange equations w.r.t.  $\bar{N}$ ,  $\beta$  and  $\chi$  will be called the trsfed. 1st, 2nd Friedmann eqs. and the Klein–Gordon eq., respectively.



# Implicit integration

General integral for  $p_\beta \neq 0$

- Since  $\beta$  is cyclic in eq. (4), the trsfed. 2nd Friedmann eq. can be integrated<sup>1</sup>

$$\text{const.} \equiv p_\beta := \frac{\partial L}{\partial \dot{\beta}} = -6\mathfrak{s}\nu^{1/2} \frac{\dot{\beta}}{\bar{N}} \equiv -6\mathfrak{s}\mathfrak{s}_\beta \frac{\nu^{1/2}}{g} \frac{\lambda\dot{\alpha} + \ell\nu\dot{\phi}}{\bar{N}}. \quad (5)$$

- For  $p_\beta \neq 0$ , fixing the *implicit gauge*  $\bar{N} = -6\mathfrak{s}\sqrt{\nu}\dot{\beta}/p_\beta$ , the trsfed. 1st Friedmann equation **can be integrated to get the trajectory**

$$e^{\mathfrak{s}_X g\chi} = \frac{p_\beta^2}{12\nu^2|V|} S^2 \left( \mathfrak{s}_\beta \sqrt{\frac{3}{2\nu}} g\beta + C \right), \quad (6)$$

in which  $\nu := \operatorname{sgn} V$ ,  $(\operatorname{sgn}, \operatorname{sgn})$  means  $(\ell, \mathfrak{s}\nu)$ , and

$$\begin{array}{ll} {}_{(+,+)}S(\gamma) := \operatorname{sech}(\gamma), & {}_{(+,-)}S(\gamma) := \operatorname{csch}(\gamma), \\ {}_{(-,+)}S(\gamma) := \operatorname{sec}(\gamma), & {}_{(-,-)}S(\gamma) := i \operatorname{csc}(\gamma). \end{array} \quad (7)$$

---


<sup>1</sup>The same first integral in canonical formalism has been found in C. Lan. PhD thesis. Saint Petersburg State University, 2016, A. A. Andrianov, O. O. Novikov, and C. Lan. In: *Theoretical and Mathematical Physics* 184.3 (2015), pp. 1224–1233.



# Implicit integration

Specific integral for  $p_\beta = 0$

- For  $p_\beta = 0$ , one has  $\beta \equiv \beta_0$  or  $\phi - \phi_0 = -\ell \lambda \alpha / \nu$ , which is the familiar power-law special solution<sup>2</sup>.
- Further integrating the first Friedmann equation demands  $(+, -)$  or  $(-, +)$  to guarantee  $\bar{N} > 0$ , and the result is automatically consistent with the trsfed. Klein–Gordon equation.




---

<sup>2</sup> M. P. Dąbrowski, C. Kiefer, and B. Sandhöfer. In: *Physical Review D* 74.4 (2006).



# Trajectories for quintessence model $(+, +)$

$\text{sech}$ , with  $C = 0$ ,  $|V| = \kappa^{-2}$  and  $p_\beta^2 = \kappa^2 \sqrt{|V|}$ ; varying  $\lambda$




- has two asymptotes  $\chi \propto \pm \beta$



# Trajectories for quintessence model $(+, +)$

$\text{sech}$ , with  $C = 0$ ,  $|V| = \nu^{-2}$  and  $\lambda^2 = 3\nu$ ; varying  $p_\beta$




- has two asymptotes  $\chi \propto \pm\beta$



# Trajectories for quintessence model $(+, +)$

sech, with  $C = 0$ ,  $\lambda^2 = 3\nu$  and  $p_\beta^2 = \nu^2 \sqrt{|V|}$ ; varying  $V$




- has two asymptotes  $\chi \propto \pm \beta$



# Trajectories for quintessence model $(+, +)$

sech, with  $C = 0$ ,  $\lambda^2 = 3\nu$  and  $p_\beta^2 = \nu^2 \sqrt{|V|}$ ; varying  $\nu$




- has two asymptotes  $\chi \propto \pm \beta$



# Trajectories for quintessence model $(+, -)$

$\text{csch}$ , with  $C = 0$ ,  $|V| = \kappa^{-2}$  and  $p_\beta^2 = \kappa^2 \sqrt{|V|}$ ; varying  $\lambda$




- contains two distinct solutions, separated by  $\beta = 0$
- has three asymptotes  $\chi \propto \pm\beta$  and  $\beta = 0$



# Trajectories for quintessence model $(+, -)$ : csch

csch, with  $C = 0$ ,  $|V| = \kappa^{-2}$  and  $p_\beta^2 = \kappa^2 \sqrt{|V|}$ ; varying  $\lambda$




- contains two distinct solutions, separated by  $\beta = 0$
- has three asymptotes  $\chi \propto \pm\beta$  and  $\beta = 0$



# Trajectories for quintessence model (+, -): csch

csch, with  $V = \kappa^{-2}$  and  $\lambda^2 = \kappa$ ; varying  $p_\beta$




- contains two distinct solutions, separated by  $\beta = 0$
- has three asymptotes  $\chi \propto \pm\beta$  and  $\beta = 0$



## Trajectories for phantom model $(-, +)$

csc, with  $C = 0$ ,  $V = \kappa^{-2}$  and  $p_\beta^2 = 3\kappa^2 \sqrt{|V|}$ ; varying  $\lambda$




- contains infinite distinct solutions
- has infinite parallel asymptotes  $\beta \propto n\pi$
- is  $\beta$ -even for  $C = 0$



## Trajectories for phantom model $(-, +)$

csc, with  $C = 0$ ,  $V = \kappa^{-2}$  and  $p_\beta^2 = 3\kappa^2 \sqrt{|V|}$ ; varying  $\lambda$




- contains infinite distinct solutions
- has infinite parallel asymptotes  $\beta \propto n\pi$
- is  $\beta$ -even for  $C = 0$



# Trajectories for phantom model $(-, +)$

csc, with  $C = 0$ ,  $|V| = \kappa^{-2}$  and  $\lambda^2 = 3\kappa$ ; varying  $p_\beta$



- contains infinite distinct solutions
- has infinite parallel asymptotes  $\beta \propto n\pi$
- is  $\beta$ -even for  $C = 0$



# Integration

## Further discussions



- The integrals are consistent with the trsfed. Klein–Gordon equation.
- The integral for  $(-, -)$  is not real.
- The implicit integration enables one to compare trajectories with wave functions, see below.

# Outline

## 1. Introduction

## 2. Classical model and the implicit trajectories Lagrangian formalism

## 3. Quantised model and the wave packets

- Dirac quantisation
- Semi-classical approximation
- Inner product and wave packet

## 4. Conclusions



# Dirac quantisation and the mode functions

- The primary Hamiltonian and the Hamiltonian constraint reads

$$H^P = \bar{N} H_{\perp} + v^{\bar{N}} p_{\bar{N}}, \quad (8)$$

$$H_{\perp} := -s \frac{p_{\beta}^2}{12\nu^{1/2}} + \ell s \frac{p_{\chi}^2}{2\nu^{3/2}} + \nu^{3/2} V e^{g_s \chi} \chi. \quad (9)$$

- Applying the Dirac quantisation rules with Laplace–Beltrami operator, one gets the mss. Wheeler–DeWitt eq. with  $(\beta, \chi)$

$$0 = \widehat{H}_{\perp} \Psi(\beta, \chi) := \left( s \frac{\hbar^2}{12\nu^{1/2}} \partial_{\beta}^2 - \ell s \frac{\hbar^2}{2\nu^{3/2}} \partial_{\chi}^2 + \nu^{3/2} V e^{g_s \chi} \chi \right) \Psi. \quad (10)$$

- Equation (10) is KG-like, hyperbolic for  $\ell = +1$  and *elliptic* for  $\ell = -1$ .



## Separation of the variables and mode functions

- Writing  $\Psi(\beta, \chi) = \varphi(\beta)\psi(\chi)$ , eq. (10) can be separated into

$$\ddot{\varphi}(\beta) := \partial_\beta^2 \varphi(\beta) = k_\beta^2 \varphi(\beta); \quad (11)$$

$$\mathbb{D}\psi(\chi) = k_\beta^2 \psi(\chi), \quad \mathbb{D} := -\ell \frac{6}{\varkappa} \partial_\chi^2 + s\nu \frac{12\varkappa^2 |V|}{\hbar^2}. \quad (12)$$

- Equation (12) turns out to be Besselian, and the mode functions are

$$\Psi_\nu(\beta, \chi) := \sum_{i=1}^2 c_i \varphi_\nu^{(i)}(\gamma) \sum_{j=1}^2 a_j B_\nu^{(j)}(\sigma), \quad \nu \geq 0; \quad (13)$$

$$\nu := \sqrt{\frac{2\varkappa}{3}} \frac{1}{g} k_\beta, \quad \gamma := \sqrt{\frac{3}{2\varkappa}} \frac{g}{1} \beta, \quad \sigma^2 := \frac{8\varkappa^3 |V| e^{g\beta_X \chi}}{\hbar^2 g^2}, \quad (14)$$

$${}_{(+,+)} B_\nu^{(i)}(\sigma) := K \text{ and } I_{i\nu}(\sigma), \quad {}_{(+,-)} B_\nu^{(i)}(\sigma) := F \text{ and } G_{i\nu}(\sigma),$$

$${}_{(-,+)} B_\nu^{(i)}(\sigma) := J \text{ and } Y_\nu(\sigma), \quad {}_{(-,-)} B_\nu^{(i)}(\sigma) := K \text{ and } I_\nu(\sigma).$$

- Adapted to imaginary order,  $F_\nu(\sigma)$  and  $G_\nu(\sigma)$  are defined in<sup>3</sup>.

<sup>3</sup> T. M. Dunster. In: *SIAM Journal on Mathematical Analysis* 21.4 (1990), pp. 995–1018.



# Physical mode functions

- Physical mode functions are expected to be regular on the boundary.
- $(+, +)$ :  $|I_{i\nu}(\sigma)| \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$
- $(-, +)$ :
  - $\forall n \in \mathbb{N}$ ,  $|Y_n(\sigma)| \rightarrow +\infty$  as  $\alpha \rightarrow -\infty$ .
  - $\forall \nu \in \mathbb{R}^+ \setminus \mathbb{N}$ , choose  $J_{-\nu}$  instead of  $Y_\nu$ , since  $J_{\pm\nu}$  are also linearly independent.
  - $\forall \nu \in \mathbb{R}^+ \setminus \mathbb{N}$ ,  $|J_{-\nu}(\sigma)| \rightarrow +\infty$  as  $\alpha \rightarrow -\infty$ .
- $(-, -)$ :  $|K_\nu(\sigma)| \rightarrow +\infty$  as  $\alpha \rightarrow -\infty$ ;  $|I_\nu(\sigma)| \rightarrow +\infty$  as  $\alpha \rightarrow +\infty$
- These are not to be included in the space of physical wave functions.  
 $\forall \nu \geq 0$ ,
  - $(+, +)$ :  $K_{i\nu}(\sigma)$  survives
  - $(+, -)$ :  $F_{i\nu}(\sigma)$  and  $G_{i\nu}(\sigma)$  survives
  - $(-, +)$ :  $J_\nu(\sigma)$  survives
  - $(-, -)$ : drops out



# Matching quantum number with classical first integral

## Principle of constructive interference

- Baustelle
- In order to match the quantum number  $k_\beta$  (or *linearly*,  $\nu$ ) with the classical first integral  $p_\beta$ , one may apply the *principle of constructive interference*<sup>4</sup>.
- Writing the mode function in the WKB form

$$\Psi_{k_\beta}(\beta, \chi) \sim \sqrt{\rho} e^{\frac{iS}{\hbar}}, \quad S/\hbar \gg 1 \text{ and } k_\beta \gg 1, \quad (15)$$

the principle demands that  $\partial S / \partial k_\beta = 0$  be equivalent to the classical trajectory.

<sup>4</sup> C. Kiefer. In: *Nuclear Physics B* 341.1 (1990), pp. 273–293.



# Matching quantum number with classical first integral

(+, +) as exemplar

- Fixing  $\nu/\sigma > 1$ , the asymptotic expansion reads

$$K_{\pm\nu}(\sigma) \sim \frac{\sqrt{2\pi} \cos\left(\sqrt{\nu^2 - \sigma^2} - \nu \arccos \frac{\nu}{\sigma} - \frac{\pi}{4}\right)}{(\nu^2 - \sigma^2)^{1/4} e^{\mp\nu/2}} + O\left(\frac{1}{\sigma}\right). \quad (16)$$

- There are two phases with opposite signs. Assuming  $c_i, a_j$ 's are real and applying *the principle* to  $\Psi_\nu(\sigma)$ , one has  $\sigma/\nu = \operatorname{sech}(\beta_\beta \gamma)$ , which matches the trajectory with  $C = 0$  if

$$\hbar k_\beta \equiv \hbar \sqrt{\frac{3}{2\nu}} g \hbar \nu = p_\beta, \quad (17)$$

- Non-vanishing  $C$  can be compensated by the phase of  $c_i$  and  $a_j$ 's.
- Fixing  $\nu/\sigma < 1$ , the asymptotic expansion is not oscillatory, but exponential; it is not within the WKB regime.
- The conclusions also hold for  $F_{\pm\nu}(\sigma), G_{\pm\nu}(\sigma)$  for  $(+,-)$ , and  $J_\nu(\sigma)$  for  $(-,+)$ .



# Inner product for wave functions

## Schrödinger product

- To make sense of amplitude and wave packet, an inner product is necessary
- A common starting point is the *Schrödinger product*

$$(\Psi_1, \Psi_2)_S := \int d\chi \Psi_1^*(\beta, \chi) \Psi_2(\beta, \chi); \quad (18)$$

- $(\Psi, \Psi)_S$  is **positive-definite**, and the integrand  $\rho_S(\beta, \chi)$  is **non-negative**
- The corresponding Schrödinger current is **real** but **does not satisfy continuity equation**  $\dot{\rho}_S + \nabla \cdot \vec{j}_S = 0$ , because eq. (10) is KG-like.
- $K_{i\nu}^5$  for  $(+, +)$ ,  $F_{i\nu}$  and  $G_{i\nu}$  for  $(+, -)$  can be proved to be **orthogonal** and **complete** among themselves, as well as **can be normalised**.
  - $J_{i\nu}$ 's for  $(+, -)$  are **not orthogonal**

---

<sup>5</sup> S. B. Yakubovich. In: *Opuscula Math.* 26.1 (2006), pp. 161–172, A. Passian et al. In: *Journal of Mathematical Analysis and Applications* 360.2 (2009), pp. 380–390, R. Szmytkowski and S. Bielski. In: *Journal of Mathematical Analysis and Applications* 365.1 (2010), pp. 195–197.



# Peculiarity of the phantom model $(-, +)$

Orthogonality for mode functions; Hermiticity for operators

- $J_\nu(\sigma)$ 's are not orthogonal under the Schrödinger product

$$(J_\nu, J_{\tilde{\nu}})_S \propto \int_{-\infty}^{+\infty} dx J_\nu^*(e^x) J_{\tilde{\nu}}(e^x) = \frac{2\sin(\pi(\nu - \tilde{\nu})/2)}{\pi(\nu^2 - \tilde{\nu}^2)}, \quad (19)$$

therefore  $\mathbb{D}$  in eq. (12) is not Hermitian (though we do not need it so far)

- $\hat{p}_\chi^2$  is not Hermitian for  $\{J_\nu(\sigma)\}$  under the Schrödinger product

$$\int_{-\infty}^{+\infty} dx J_\nu^*(-\partial_x^2 J_{\tilde{\nu}}) - \int_{-\infty}^{+\infty} dx (-\partial_x^2 J_\nu)^* J_{\tilde{\nu}} = \frac{2}{\pi} \sin \frac{\pi(\nu - \tilde{\nu})}{2}. \quad (20)$$

- In order to save Hermiticity for  $p_\chi^2$  and  $\mathbb{D}$  and orthogonality of the modes under Schrödinger product, one can restrict

$$\nu = 2n + \nu_0, \quad n \in \mathbb{N}, \quad \nu_0 \in [0, 2]. \quad (21)$$

- The classical trajectory is  $\beta$ -even; imposing the same condition fixes  $\nu_0 = 1$ .



# Discretisation of the phantom model $(-, +)$

Levels of the phantom model are discretised

...if one imposes Hermiticity of squared momenta under the Schrödinger product

- This kind of subtlety on Hermiticity is named as *self-adjoint extension*, which arises already for infinite square well<sup>6</sup>.
- It also applies to  $x^{-2}$  potentials<sup>7</sup>, which is of cosmological relevance<sup>8</sup>.

---

<sup>6</sup> G. Bonneau, J. Faraut, and G. Valent. In: *American Journal of Physics* 69.3 (2001), pp. 322–331.

<sup>7</sup> A. M. Essin and D. J. Griffiths. In: *American Journal of Physics* 74.2 (2006), pp. 109–117,  
V. S. Araujo, F. A. B. Coutinho, and J. F. Perez. In: *American Journal of Physics* 72.2 (2004),  
pp. 203–213.

<sup>8</sup> M. Bouhmadi-López et al. In: *Physical Review D* 79.12 (2009).



# Further inner products for wave functions

## Klein–Gordon and Mostafazadeh product

- Since eq. (10) is KG-like, another popular choice is the KG product

$$(\Psi_1, \Psi_2)^g_{\text{KG}} := \mathrm{i}g \left\{ (\Psi_1, \dot{\Psi}_2)_S - (\dot{\Psi}_1, \Psi_2)_S \right\}, \quad g > 0. \quad (22)$$

- $(\Psi, \Psi)^g_{\text{KG}}$  is **real** but **not positive-definite**, so does the integrand  $\rho_{\text{KG}}$ ;
- The corresponding  $\vec{J}_{\text{KG}}$  is **conserved**  $\dot{\rho}_{\text{KG}} + \nabla \cdot \vec{J}_{\text{KG}} = 0$  and **real**.
- Mostafazadeh<sup>9</sup> found a product *for Hermitian  $\mathbb{D}$  with positive spectrum*:

$$(\Psi_1, \Psi_2)^\kappa_M := \kappa \left\{ (\Psi_1, \mathbb{D}^{+1/2}\Psi_2)_S + (\dot{\Psi}_1, \mathbb{D}^{-1/2}\dot{\Psi}_2)_S \right\}, \quad \kappa > 0. \quad (23)$$

- $(\Psi, \Psi)^\kappa_M$  is **positive-definite**, but the integrand  $\rho_M^\kappa$  is **complex**
- The corresponding  $\vec{J}_M^\kappa$  is **conserved**  $\dot{\rho}_M^\kappa + \nabla \cdot \vec{J}_M^\kappa = 0$  but also **complex**.

<sup>9</sup> A. Mostafazadeh. In: *Classical and Quantum Gravity* 20.1 (2002), pp. 155–171.



# Mostafazadeh inner product and the corresponding density

- The real power of  $\mathbb{D}$  is defined by spectral decomposition  $\mathbb{D}^\gamma := \sum_\nu \nu^{2\gamma} \mathbf{P}_\nu$ ,  
 $\mathbf{P}_\nu \Psi := \Psi_\nu (\Psi_\nu, \Psi)_S$ .
- It can be shown<sup>10</sup> that *the density*

$$\varrho_M^\kappa := \kappa \left\{ |\mathbb{D}^{+1/4} \Psi|^2 + |\mathbb{D}^{-1/4} \dot{\Psi}|^2 \right\} \quad (24)$$

- is equivalent to  $\rho_M^\kappa$  up to a boundary term  $\int d\chi \varrho_M^\kappa = \int d\chi \rho_M^\kappa \equiv (\Psi_1, \Psi_2)_M^\kappa$ ;
- is non-negative.
- The corresponding current  $\vec{J}_M^\kappa$  is real but not conserved<sup>11</sup>.

<sup>10</sup> A. Mostafazadeh and F. Zamani. In: *Annals of Physics* 321.9 (2006), pp. 2183–2209.

<sup>11</sup> B. Rosenstein and L. P. Horwitz. In: *Journal of Physics A: Mathematical and General* 18.11 (1985), pp. 2115–2121.



# Wave packets of Gaussian amplitude for continuous spectrum

## Quintessence models

- It is difficult to find an amplitude such that the wave packet is Gaussian
- Instead, one can choose a Gaussian amplitude

$$A(\nu; \bar{\nu}, \sigma) := \left( \frac{1}{\sqrt{2\pi}\sigma} \exp\left(-\frac{(\nu - \bar{\nu})^2}{2\sigma^2}\right) \right)^{1/2} \quad (25)$$

- In<sup>12</sup>,  $A(\nu; \bar{\nu}, \sigma/\sqrt{2})$  was chosen.

---


<sup>12</sup> M. P. Dąbrowski, C. Kiefer, and B. Sandhöfer. In: *Physical Review D* 74.4 (2006).




# Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\parallel\nu}$ , with  $\lambda = \kappa^{1/2}/2$ ,  $V = -\kappa^{-2}$ ,  $\bar{k}_\beta = -2$  and  $\sigma_\beta = 5/4$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for quintessence model (+, +)

$K_{\parallel\nu}$ , with  $\lambda = \kappa^{1/2}/2$ ,  $V = -\kappa^{-2}$ ,  $\bar{k}_\beta = -2$  and  $\sigma_\beta = 5/4$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\text{I}\nu}$ , with  $\lambda = 4\kappa^{1/2}/5$ ,  $V = +\kappa^{-2}$ ,  $\bar{k}_\beta = -7/2$  and  $\sigma_\beta = 7/5$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for quintessence model (+, -)

$F_{\text{I}\nu}$ , with  $\lambda = 4\kappa^{1/2}/5$ ,  $V = +\kappa^{-2}$ ,  $\bar{k}_\beta = -7/2$  and  $\sigma_\beta = 7/5$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\text{I}\nu}$ , with  $\lambda = 4\kappa^{1/2}/5$ ,  $V = +\kappa^{-2}$ ,  $\bar{k}_\beta = -7/2$  and  $\sigma_\beta = 7/5$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for quintessence model (+, -)

$G_{\mu\nu}$ , with  $\lambda = 4\kappa^{1/2}/5$ ,  $V = +\kappa^{-2}$ ,  $\bar{k}_\beta = -7/2$  and  $\sigma_\beta = 7/5$

Schödinger



Mostafazadeh



# Wave packets with Poissonian amplitude for discrete spectrum

Discrete phantom model  $(-, +)$

- Gaussian distribution works for continuous variable
- For discrete spectrum, one can choose a Poissonian amplitude

$$A_n(\bar{n}) := \left( e^{-\bar{n}} \frac{\bar{n}^n}{n!} \right)^{1/2} \quad (26)$$

- In<sup>13</sup>,  $A_n(\bar{n}/\sqrt{2})$  was chosen.

---


<sup>13</sup> C. Kiefer. In: *Nuclear Physics B* 341.1 (1990), pp. 273–293.




# Wave packets of Poissonian amplitude for phantom model

$J_{2n+1}$ , with  $\lambda = 2\kappa^{1/2}$ ,  $V = +\kappa^{-2}$  and  $\bar{k}_\beta = 8$

Schödinger




Mostafazadeh




# Wave packets of Poissonian amplitude for phantom model

$J_{2n+1}$ , with  $\lambda = 2\kappa^{1/2}$ ,  $V = +\kappa^{-2}$  and  $\bar{k}_\beta = 8$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for phantom model

$J_\nu$ , with  $\lambda = 2\nu^{1/2}$ ,  $V = +\nu^{-2}$ ,  $\bar{k}_\beta = 8$  and  $\sigma_\beta = 11/2$

Schödinger




Mostafazadeh




# Wave packets of Gaussian amplitude for phantom model

$J_\nu$ , with  $\lambda = 2\nu^{1/2}$ ,  $V = +\nu^{-2}$ ,  $\bar{k}_\beta = 8$  and  $\sigma_\beta = 11/2$

Schödinger



Mostafazadeh



# Outline

## 1. Introduction

## 2. Classical model and the implicit trajectories Lagrangian formalism

## 3. Quantised model and the wave packets Dirac quantisation Semi-classical approximation Inner product and wave packet

## 4. Conclusions



## Issues

- Wave packets with multiple branches in  $(+, -)$  and  $(-, +)$  is to be understood
- Quantum-corrected  $\bar{k}_\beta$  is to be understood
- Wave packets with Gaussian profile are to be constructed, instead of inserting Gaussian / Poissonian amplitude artificially.
- A normalising  $\kappa$  for  $(\cdot, \cdot)_M^\kappa$  is to be evaluated, otherwise a quantitative comparison of  $(\cdot, \cdot)_S$  and  $(\cdot, \cdot)_M^\kappa$  is not possible.



# Outlook

- Beyond isotropy: generalise to Bianchi models
  - Bianchi Type-I: natural extension; **under investigation**
- Beyond homogeneity: cosmological perturbation, **under investigation**
- Beyond single field
  - Two exponential potentials:  $V_1 = V_2$  and special  $\lambda_i$
  - Multiple Liouville fields: mixing kinetic terms needed<sup>14</sup>
- Beyond classic matter
  - Non-Hermitian,  $PT$ -symmetric Liouville fields<sup>15</sup>: may avoid big-rip etc.

---

<sup>14</sup> A. A. Andrianov, O. O. Novikov, and C. Lan. In: *Theoretical and Mathematical Physics* 184.3 (2015), pp. 1224–1233.

<sup>15</sup> A. A. Andrianov et al. In: *International Journal of Modern Physics D* 19.01 (2010), pp. 97–111, A. A. Andrianov, C. Lan, and O. O. Novikov. In: *Springer Proceedings in Physics*. Springer International Publishing, 2016, pp. 29–44.



# Integration of the transformed first Friedmann equation

General integral for  $p_\beta \neq 0$

In order to integrate the equation under the implicit gauge

$$\beta \frac{p_\beta^2}{12} \left( -\ell \frac{\nu^{1/2}}{6} \left( \frac{d\chi}{d\beta} \right)^2 + \nu^{-1/2} \right) - \nu^{3/2} V e^{g_\beta \chi} = 0, \quad (27)$$

one can substitute

$$\gamma := \sqrt{\frac{3}{2\nu}} g\beta, \quad \tilde{\sigma}^2 := \frac{p_\beta^2}{12\nu^2 |V|} e^{-g_\beta \chi}, \quad (28)$$

to get

$$\left( \frac{d\tilde{\sigma}}{d\gamma} \right)^2 + \ell(\beta v - \tilde{\sigma}^2) = 0 \quad \Rightarrow \quad \frac{d\gamma}{d\tilde{\sigma}} = \pm \frac{1}{\sqrt{\ell(-\beta v + \tilde{\sigma}^2)}}, \quad (29)$$

which is of the standard inverse hyperbolic / trigonometric form **except for  $(-, -)$ .**



# Integration of the separated minisuperspace WDW equation

233

In order to integrate the separated minisuperspace Wheeler–DeWitt equation

$$\mathbb{D}\psi(\chi) = k_\beta^2 \psi(\chi), \quad \mathbb{D} := -\ell \frac{6}{\nu} \partial_\chi^2 + \varsigma v \frac{12\nu^2 |V|}{\hbar^2}, \quad (12 \text{ rev.})$$

one can transform

$$\nu := \sqrt{\frac{2\nu}{3}} \frac{k_\beta}{g}, \quad \sigma^2 := \frac{8\nu^3 |V| e^{g\beta_X \chi}}{\hbar^2 g^2}, \quad (14 \text{ rev.})$$

to get

$$\sigma^2 \psi''(\sigma) + \sigma \psi'(\sigma) + \ell(\nu^2 - \varsigma v \sigma^2) \psi(\sigma) = 0, \quad (30)$$

which is of the standard Besselian form.

