

Notes on Canonical Singular Dynamics

Yi-Fan Wang (王一帆)

April 10, 2017

1 Canonical formalism

Lagrangian with velocity

$$L^\vee := L|_{\dot{q}=v} \quad (1.1)$$

Equations of motion

$$\sum_j M_{ij} \dot{v}_j = K_i^\vee, \quad \dot{q}_i = v_i. \quad (1.2)$$

where

$$M_{ij}(q, v) := \frac{\partial^2 L^\vee}{\partial v_i \partial v_j}. \quad (1.3)$$

Adding

$$p_i := \frac{\partial L^\vee}{\partial v_i}. \quad (1.4)$$

Variation of

$$S[q, p; v] := \int dt \left[L^\vee + \sum_i p_i (\dot{q}_i - v_i) \right]. \quad (1.5)$$

gives the *extended Euler–Lagrange equations*

$$\dot{q}_i = v_i, \quad \dot{p}_i = \frac{\partial L^\vee}{\partial q_i}, \quad p_i = \frac{\partial L^\vee}{\partial v_i}. \quad (1.6)$$

Extended Hamiltonian

$$H^\vee(q, p; v) := \sum_i p_i v_i - L^\vee. \quad (1.7)$$

Identities

$$\frac{\partial H^\vee}{\partial q_i} \equiv -\frac{\partial L^\vee}{\partial q_i}, \quad \frac{\partial H^\vee}{\partial p_i} \equiv v_i, \quad \frac{\partial H^\vee}{\partial v_i} \equiv p_i - \frac{\partial L^\vee}{\partial v_i}. \quad (1.8)$$

Variation of

$$S[q, p; v] := \int dt \left[\sum_i p_i \dot{q}_i - H^\vee \right] \quad (1.9)$$

gives the *extended canonical equations*

$$\dot{q}_i = [q_i, H^\vee]_P, \quad \dot{p}_i = [p_i, H^\vee]_P, \quad \frac{\partial H^\vee}{\partial v_i} = 0, \quad (1.10)$$

where the Poisson bracket is defined as

$$[f^\vee, g^\vee]_P := \sum_i \left(\frac{\partial f^\vee}{\partial q_i} \frac{\partial g^\vee}{\partial p_i} - \frac{\partial f^\vee}{\partial p_i} \frac{\partial g^\vee}{\partial q_i} \right). \quad (1.11)$$

$v_a = \bar{v}_a(q, p)$ can be solved, $a = 1, 2, \dots, r_M$; v^α cannot be solved, $\alpha = r_M + 1, \dots, n$, where $r_M = \text{rank } M$.

(need to show $v_a = \bar{v}_a(q, p_a)$)

Primary constraints in the standard form

$$\phi_\alpha^{(0)}(q, p) := \left. \frac{\partial H^\vee}{\partial v_\alpha} \right|_{\{v_\alpha = \bar{v}_\alpha\}} \equiv p_\alpha - \bar{p}_\alpha(q, \{p_a\}), \quad (1.12)$$

where

$$\bar{p}_\alpha(q, \{p_a\}) := \left. \frac{\partial L^\vee}{\partial v_\alpha} \right|_{\{v_a = \bar{v}_a\}}. \quad (1.13)$$

Total Hamiltonian

$$H^t := H^\vee|_{\{v_a = \bar{v}_a\}} \equiv H^\vee(q, p; \{\bar{v}^a(q, p_a), v_\alpha\}). \quad (1.14)$$

Subspace of primary constraints

$$\Gamma_P = \{(q, p) \mid \phi_\alpha^0(q, p) = 0, \forall \alpha\} \quad (1.15)$$

Since

$$\frac{\partial H^t}{\partial v_\alpha} = \left. \frac{\partial H^\vee}{\partial v_\alpha} \right|_{\{v_a = \bar{v}_a\}} = \phi_\alpha^{(0)} \equiv p_\alpha - \bar{p}_\alpha(q, \{p_a\}), \quad (1.16)$$

H^t is linear in v_α . One writes

$$H^t(q, \{p_a\}; \{p_\alpha\}, \{v_\alpha\}) = H(q, \{p_a\}) + \sum_\alpha v_\alpha \phi_\alpha^{(0)}, \quad (1.17)$$

where H is the *canonical Hamiltonian* or simply *Hamiltonian*.

Proposition show H is independent of $\{p_\alpha\}$.

Proposition Canonical equations with primary constraints

$$\dot{q}_i = [q_i, H]_p + \sum_{\beta} v_{\beta} [q_i, \phi_{\beta}]_p, \quad (1.18)$$

$$\dot{p}_i = [p_i, H]_p + \sum_{\beta} v_{\beta} [p_i, \phi_{\beta}]_p, \quad (1.19)$$

$$\phi_{\alpha}^{(0)}(q, p) = 0, \quad (1.20)$$

where v_{β} 's are undetermined. Note that eq. (1.18) for $i = \alpha$ holds identically: $\dot{q}_{\alpha} = \dot{q}_{\alpha}$.

Weak equality: $f_1 \approx f_2$ iff $f_1|_{\Gamma_p} = f_2|_{\Gamma_p}$.

Proposition if f and g are two functions over the phase space Γ , and $f \approx h$, then

$$\frac{\partial}{\partial q_i} \left(f - \sum_{\beta} \phi_{\beta} \frac{\partial f}{\partial p_{\beta}} \right) \approx \frac{\partial}{\partial q_i} \left(h - \sum_{\beta} \phi_{\beta} \frac{\partial h}{\partial p_{\beta}} \right), \quad (1.21)$$

$$\frac{\partial}{\partial p_i} \left(f - \sum_{\beta} \phi_{\beta} \frac{\partial f}{\partial p_{\beta}} \right) \approx \frac{\partial}{\partial p_i} \left(h - \sum_{\beta} \phi_{\beta} \frac{\partial h}{\partial p_{\beta}} \right). \quad (1.22)$$

Corollary $\forall H_1 \approx H$,

$$\dot{q}_i \approx [q_i, H]_p, \quad \dot{p}_i \approx [p_i, H]_p. \quad (1.23)$$

2 Examples

2.1 Toy examples

Example 0

Gitman and Tyutin 1990, sec. 1.2

$$L = \frac{1}{2}(\dot{x} - y)^2 \quad (2.1)$$

Example 1

$$L = \frac{1}{2}\dot{x}^2 + \dot{x}y - \frac{1}{2}(x - y)^2. \quad (2.2)$$

One has

$$L^v = \frac{1}{2}v_x^2 + v_x y - \frac{1}{2}(x - y)^2, \quad (2.3)$$

so that

$$p_x = \frac{\partial L^v}{\partial v_x} = v_x + y, \quad p_y = 0, \quad (2.4)$$

thus

$$\bar{v}_x = p_x - y. \quad (2.5)$$

So that v_y is the primary inexpressible velocity.

The extended Hamiltonian reads

$$H^v(q, p; v) = v_x p_x + v_y p_y - \frac{1}{2} v_x^2 - v_x y + \frac{1}{2} (x - y)^2, \quad (2.6)$$

whilst the total Hamiltonian is

$$H^t(q, p; \bar{v}_x, v_y) = \frac{1}{2} (p_x - y)^2 + \frac{1}{2} (x - y)^2 + v_y p_y. \quad (2.7)$$

Example 2

$$L = \frac{1}{2} \dot{x}^2 + \dot{x}y + \frac{1}{2} (x - y)^2 \quad (2.8)$$

Primary constraint

$$p_y = 0; \quad (2.9)$$

total Hamiltonian

$$H^t = \frac{1}{2} p_x^2 - p_x y - \frac{1}{2} x^2 + xy + v_y p_y. \quad (2.10)$$

Example 3

$$L = \frac{1}{2} (\dot{q}_2 - e^{q_1})^2 + \frac{1}{2} (\dot{q}_3 - q_2)^2. \quad (2.11)$$

2.2 Parametrised systems

Non-relativistic point particle

Relativistic charged point particle

Landau and Lifshitz 1975, sec. 15

$$S := \int -m \, ds + q A_\mu \, dx^\mu =: \int dt \, L, \quad (2.12)$$

where the Lagrangian reads

$$L = -m \sqrt{-\dot{x}^\mu \dot{x}_\mu} + q \dot{x}^\mu A_\mu. \quad (2.13)$$

Blumenhagen, Lüst, and Theisen 2013, sec. 2.1

$$L := \frac{1}{2} (e^{-1} \dot{x}^\mu \dot{x}_\mu - m^2 e) \quad (2.14)$$

2.3 Maxwell–Proca theory

$$\mathcal{L} = -\frac{1}{4}F_{\mu\nu}F^{\mu\nu} - \frac{1}{2}m^2 A_\mu A^\mu + \frac{1}{2}F_{\mu\nu}S^{\mu\nu} + A_\mu J^\mu_{\text{f}}, \quad (2.15)$$

where $m > 0$ corresponds to the Proca theory Gitman and Tyutin 1990, sec. 2.3, and $m = 0$ the Maxwell theory H. J. Rothe and K. D. Rothe 2010, sec. 3.3.3 Gitman and Tyutin 1990, sec. 2.4.

2.4 String theories

Nambu–Goto action

Generalising the kinetic part of (2.12), one has

$$S_{\text{NG}} := -T \int_{\Sigma} \mathrm{d}A =: -T \int_{\Sigma} \mathrm{d}^2\sigma \mathcal{L}, \quad (2.16)$$

where the Lagrangian density

$$\mathcal{L} = \sqrt{-\Gamma}, \quad \Gamma := \det \Gamma_{\alpha\beta}, \quad \Gamma_{\alpha\beta} := \frac{\partial X^\nu}{\partial \sigma^\alpha} \frac{\partial X_\nu}{\partial \sigma^\beta}. \quad (2.17)$$

Historically Gotō 1971; Nambu 1970; Reference e.g. Blumenhagen, Lüst, and Theisen 2013; Kiefer 2012

Polyakov action

Generalising (2.14)

$$S_{\text{p}}[X^\mu, h_{\alpha\beta}] = -\frac{T}{2} \int_{\Sigma} \mathcal{L}, \quad (2.18)$$

where

$$\mathcal{L} := \sqrt{-h} h^{\alpha\beta} \Gamma_{\alpha\beta}. \quad (2.19)$$

Historically Brink, Di Vecchia, and Howe 1976; Deser and Zumino 1976; Polyakov 1981

References

- Blumenhagen, Ralph, Dieter Lüst, and Stefan Theisen (2013). *Basic Concepts of String Theory*. Theoretical and Mathematical Physics. Springer. ISBN: <http://id.crossref.org/isbn/978-3-642-29497-6>. DOI: 10.1007/978-3-642-29497-6. URL: <http://dx.doi.org/10.1007/978-3-642-29497-6>.
- Brink, L., P. Di Vecchia, and P. Howe (1976). “A locally supersymmetric and reparametrization invariant action for the spinning string”. In: *Physics Letters B* 65.5, pp. 471–474. ISSN: 0370-2693. DOI: 10.1016/0370-2693(76)90445-7. URL: [http://dx.doi.org/10.1016/0370-2693\(76\)90445-7](http://dx.doi.org/10.1016/0370-2693(76)90445-7).

- Deser, S. and B. Zumino (1976). "A complete action for the spinning string". In: *Physics Letters B* 65.4, pp. 369–373. ISSN: 0370-2693. DOI: 10.1016/0370-2693(76)90245-8. URL: [http://dx.doi.org/10.1016/0370-2693\(76\)90245-8](http://dx.doi.org/10.1016/0370-2693(76)90245-8).
- Gitman, Dmitriy M. and Igor V. Tyutin (1990). *Quantization of Fields with Constraints*. Springer Series in Nuclear and Particle Physics. Springer Berlin Heidelberg. ISBN: <http://id.crossref.org/isbn/978-3-642-83938-2>. DOI: 10.1007/978-3-642-83938-2. URL: <http://dx.doi.org/10.1007/978-3-642-83938-2>. Дмитрий Максимович Гитман and Игорь Викторович Тютин. *Каноническое квантование полей со связями*. 030077 г. Новосибирск-77, Станиславского, 25: Наука, 1986.
- Gotō, Tetsuo (1971). "Relativistic Quantum Mechanics of One-Dimensional Mechanical Continuum and Subsidiary Condition of Dual Resonance Model". In: *Progress of Theoretical Physics* 46.5, pp. 1560–1569. ISSN: 0033-068X. DOI: 10.1143/ptp.46.1560. URL: <http://dx.doi.org/10.1143/PTP.46.1560>.
- Kiefer, Claus (2012). *Quantum Gravity*. 3rd. International Series of Monographs on Physics. Oxford University Press. ISBN: <http://id.crossref.org/isbn/9780199585205>. DOI: 10.1093/acprof:oso/9780199585205.001.0001. URL: <http://dx.doi.org/10.1093/acprof:oso/9780199585205.001.0001>.
- Landau, Lev Davidovich and Evgeny Mikhailovich Lifshitz (1975). *The Classical Theory of Fields*. 4th. Vol. 2. Course of Theoretical Physics. Butterworth-Heinemann. ISBN: <http://id.crossref.org/isbn/9780080250724>. DOI: 10.1016/c2009-0-14608-1. URL: <http://dx.doi.org/10.1016/c2009-0-14608-1>.
- Nambu, Yoichiro (1970). "Quark model and the factorization of the Veneziano amplitude". In: *Symmetries and Quark Models*. Ed. by R. Chand. International Conference on Symmetries and Quark Models, Wayne State U., Detroit Detroit, Mich., USA, June 18-20, 1969, pp. 269–278.
- Polyakov, A.M. (1981). "Quantum geometry of bosonic strings". In: *Physics Letters B* 103.3, pp. 207–210. ISSN: 0370-2693. DOI: 10.1016/0370-2693(81)90743-7. URL: [http://dx.doi.org/10.1016/0370-2693\(81\)90743-7](http://dx.doi.org/10.1016/0370-2693(81)90743-7).
- Rothe, Heinz J and Klaus D Rothe (2010). *Classical and Quantum Dynamics of Constrained Hamiltonian Systems*. World Scientific Lecture Notes in Physics. World Scientific. ISBN: <http://id.crossref.org/isbn/978-981-4299-65-7>. DOI: 10.1142/7689. URL: <http://dx.doi.org/10.1142/7689>.