

Integrable Cosmological Models with Liouville Fields

Alexander A. Andrianov^{1,4} Chen Lan² Oleg O. Novikov¹
Yi-Fan Wang³

¹ Saint-Petersburg State University, St. Petersburg 198504, Russia

² ELI-ALPS, ELI-Hu NKft, Dugonics tér 13, Szeged 6720, Hungary

³ Institut für Theoretische Physik, Universität zu Köln, Zùlpicher StraÙe 77, 50937 Köln, Germany

⁴ Institut de Ciències del Cosmos (ICCUB), Universitat de Barcelona, Spain

December 7, 2017



Outline

1. Introduction

2. Classical model and the implicitised trajectories

3. Dirac quantisation and the wave functions



Introduction

Introduction



The Friedmann-Lemaître model

123

- Flat Robertson–Walker metric $ds^2 = -N^2(t) dt^2 + \varkappa^{-1} e^{2\alpha(t)} d\Omega_3^2$, where $\varkappa := 8\pi G$, $d\Omega_3^2$ dimensionless spacial metric.
- Homogeneous real Klein–Gordon field with potential $V e^{\lambda\phi}$ (Liouville), where $\lambda, V \in \mathbb{R}$, and kinetic term with sign $\ell = \pm 1$ (quintessence / phantom model).
- Total action $\mathcal{S} := S_{\text{EH}} + S_{\text{GHY}} + S_{\text{L}} = \int d\Omega_3^2 \int dt L$, in which the effective Lagrangian reads

$$L := \varkappa^{3/2} N e^{3\alpha} \left(-\frac{3}{\varkappa} \frac{\dot{\alpha}^2}{N^2} + \ell \frac{\dot{\phi}^2}{2N^2} - V e^{\lambda\phi} \right), \quad (1)$$

in which dot means d/dt and $\ell = \pm 1$.



Decoupling the variables

Via *rescaled* special orthogonal transformation

- Setting $\overline{N} := N e^{-3\alpha}$, eq. (1) transforms to

$$L = \kappa^{3/2} \overline{N} \left(-\frac{3}{\kappa} \frac{\dot{\alpha}^2}{\overline{N}^2} + \ell \frac{\dot{\phi}^2}{2\overline{N}^2} - V e^{\lambda\phi+6\alpha} \right) \quad (2)$$

- Defining $\Delta := \lambda^2 - 6\ell\kappa$, $\jmath := \text{sgn } \Delta$ and $g := \jmath\sqrt{|\Delta|} \equiv \jmath\sqrt{\jmath\Delta}$, the *rescaled* special orthogonal transformation

$$\begin{pmatrix} \alpha \\ \phi \end{pmatrix} = \frac{\jmath}{g} \begin{pmatrix} \lambda & -\ell\kappa \\ -6 & \lambda \end{pmatrix} \begin{pmatrix} \jmath_\beta \beta \\ \jmath_\chi \chi \end{pmatrix} \quad \text{where } \jmath_\beta, \jmath_\chi = \pm 1 \quad (3)$$

gives the decoupled Lagrangian

$$L = \kappa^{3/2} \overline{N} \left(-\jmath \frac{3}{\kappa} \frac{\dot{\beta}^2}{\overline{N}^2} + \ell \jmath \frac{\dot{\chi}^2}{2\overline{N}^2} - V e^{\jmath_\chi g \chi} \right). \quad (4)$$

- The Euler–Lagrange equations w.r.t. \overline{N} , β and χ will be called the trsfed. 1st, 2nd Friedmann eqs. and the Klein–Gordon eq., respectively.



Implicitised integration

$p_\beta \neq 0$

- Since β is cyclic in eq. (4), the trsfed. 2nd Friedmann eq. can be integrated¹

$$\text{const.} \equiv p_\beta := \frac{\partial L}{\partial \dot{\beta}} = -6\mathfrak{s}\kappa^{1/2} \frac{\dot{\beta}}{\bar{N}} \equiv -6\mathfrak{s}\mathfrak{s}_\beta \frac{\kappa^{1/2}}{g} \frac{\lambda\dot{\alpha} + \ell\kappa\dot{\phi}}{\bar{N}}. \quad (5)$$

- For $p_\beta \neq 0$, fixing the **implicitising gauge** $\bar{N} = -6\mathfrak{s}\sqrt{\kappa}\dot{\beta}/p_\beta$, the trsfed. 1st Friedmann equation can be integrated

$$\mathfrak{e}^{\mathfrak{s}_\chi g\chi} = \frac{p_\beta^2}{12\kappa^2|V|} S^2 \left(\mathfrak{s}_\beta \sqrt{\frac{3}{2\kappa}} g\beta \right), \quad (6)$$

in which $\mathfrak{v} := \text{sgn } V$, and

$$S(\gamma) := \begin{cases} \text{sech}(\gamma + C_{++}) & (\ell, \mathfrak{s}\mathfrak{v}) = (+, +), \\ \text{csch}(\gamma + C_{+-}) & (\ell, \mathfrak{s}\mathfrak{v}) = (+, -), \\ \text{sec}(\gamma + C_{-+}) & (\ell, \mathfrak{s}\mathfrak{v}) = (-, +), \\ \text{icsc}(\gamma + C_{--}) & (\ell, \mathfrak{s}\mathfrak{v}) = (-, -). \end{cases} \quad (7)$$

¹Chen Lan. PhD thesis. Saint Petersburg State University, 2016. URL: <https://search.rsl.ru/ru/record/01006663434>.



Integration

Discussions

- The integrals are consistent with the trsfed. Klein–Gordon equation.
- The integral for $(+, +)$
 - has two asymptotes
- The implicitised integral for $(+, -)$
 - contains two distinct solutions
 - has three asymptotes
- The implicitised integral for $(-, +)$
 - contains infinite distinct solutions
 - has infinite asymptotes, which are pairwise parallel
- The integral for $(-, -)$
 - is not real
- The implicitised integral enables one to compare trajectories with wave functions, see below.



Implicitised integration

$$p_\beta = 0$$

- For $p_\beta = 0$, one has $\beta \equiv \beta_0$ or $\phi - \phi_0 = -\ell\lambda\alpha/\kappa$, which is the familiar power-law special solution².
- Further integrating the first Friedmann equation demands $(+, -)$ or $(-, +)$ to guarantee $\bar{N} > 0$, and the result is automatically consistent with the trsfed. Klein–Gordon equation.
- Fixing $\bar{N} = (2\kappa^2|V|)^{-1/2}$ yields

$$e^{g\beta_X X} = \left(\frac{2\kappa}{g(t-t_0)} \right)^2. \quad (8)$$

²Mariusz P. Dąbrowski, Claus Kiefer, and Barbara Sandhöfer. In: *Physical Review D* 74.4 (Aug. 2006).



Introduction

Introduction



- The primary Hamiltonian and the Hamiltonian constraint reads

$$H^p = \overline{N} H_{\perp} + p_{\overline{N}} v^{\overline{N}}, \quad (9)$$

$$H_{\perp} = -\mathfrak{J} \frac{p_{\beta}^2}{12\kappa^{1/2}} + \ell \mathfrak{J} \frac{p_{\chi}^2}{2\kappa^{3/2}} + \kappa^{3/2} V e^{g_{\mathfrak{J}} \chi}. \quad (10)$$

- Applying the Dirac quantisation rules with the Laplace–Beltrami operator, one gets the mss. Wheeler–DeWitt eq. with (β, χ) $\widehat{H}_{\perp} \Psi(\beta, \chi) = 0$,

$$-\mathfrak{J} \frac{12\kappa^{1/2}}{\hbar^2} \widehat{H}_{\perp} \Psi =: -\partial_{\beta}^2 \Psi + \mathbb{D} \Psi, \quad \mathfrak{d} = \ell \frac{6}{\kappa} \partial_{\chi}^2 - \mathfrak{J} \frac{12\kappa^2 V e^{g_{\mathfrak{J}} \chi}}{\hbar^2}. \quad (11)$$

- Equation (11) is KG-like, hyperbolic for $\ell = +1$ and **elliptic** for $\ell = -1$.

- Inserting the separating Ansatz $\Psi(\beta, \chi) = e^{-ik_\beta \beta} \psi(\chi)$, the remaining equation turns out to be Besselian, and the mode functions are

$$\Psi_\nu(\beta, \chi) := e^{-i\nu\gamma} \psi_\nu(\chi) := e^{-i\nu\gamma} (C_1 B_\nu^{(1)}(\sigma) + C_2 B_\nu^{(2)}(\sigma)), \quad (12)$$

in which

$$\nu := \sqrt{\frac{2\kappa}{3}} \frac{k_\beta}{g}, \quad \gamma := \sqrt{\frac{3}{2\kappa}} g\beta, \quad \sigma^2 := \frac{8\kappa^3 |V| e^{g\beta} \chi}{\hbar^2 g^2}, \quad (13)$$

$$B_\nu^{(i)}(\sigma) := \begin{cases} K \text{ or } I_\nu(\sigma) & (\ell, \mathfrak{s}\nu) = (+, +), \\ F \text{ or } G_\nu(\sigma) & (\ell, \mathfrak{s}\nu) = (+, -), \\ J \text{ or } Y_\nu(\sigma) & (\ell, \mathfrak{s}\nu) = (-, +), \\ K \text{ or } I_\nu(\sigma) & (\ell, \mathfrak{s}\nu) = (-, -). \end{cases} \quad (14)$$

- Adapted to imaginary order, $F_\nu(\sigma)$ and $G_\nu(\sigma)$ are defined in³.

³T. Mark Dunster. In: *SIAM Journal on Mathematical Analysis* 21.4 (July 1990), pp. 995–1018.



- Physical mode functions are expected to be regular on the boundary.
- $(+, +)$: $|I_{i\nu}(\sigma)| \rightarrow +\infty$ as $\alpha \rightarrow +\infty$
- $(-, +)$:
 - $\forall n \in \mathbb{Z}$, $|Y_n(\sigma)| \rightarrow +\infty$ as $\alpha \rightarrow -\infty$.
 - $\forall \nu \in \mathbb{R} \setminus \mathbb{Z}$, choose $J_{-\nu}$ instead of $Y_{+\nu}$, since $J_{\pm\nu}$ are also linearly independent.
 - $\forall \nu \in \mathbb{R}^+ \setminus \mathbb{Z}$, $|J_{-\nu}(\sigma)| \rightarrow +\infty$ as $\alpha \rightarrow -\infty$.
- $(-, -)$: $|K_\nu(\sigma)| \rightarrow +\infty$ as $\alpha \rightarrow -\infty$; $|I_\nu(\sigma)| \rightarrow +\infty$ as $\alpha \rightarrow +\infty$
- These are not to included in the space of physical wave functions.
- For $(-, +)$, **only J_ν with $\nu \geq 0$ survives.**



Matching quantum number with classical first integral

233

- Baustelle
- In order to match the quantum number k_β (or linearly, ν) with the classical first integral p_β , one may apply the principle of constructive interference.
- Writing the mode function in the WKB form

$$\Psi_{k_\beta}(\beta, \chi) \sim \sqrt{\rho} e^{iS/\hbar}, \quad S/\hbar \gg 1 \text{ and } k_\beta \gg 1, \quad (15)$$

the principle demands that $\partial S / \partial k_\beta = 0$ be equivalent to the classical trajectory.



Matching quantum number with classical first integral

(+, +) as exemplar

- Fixing $\nu/\sigma > 1$, the asymptotic expansion reads

$$K_{i\nu}(\sigma) \sim \frac{\sqrt{2\pi} \cos\left(\sqrt{\nu^2 - \sigma^2} - \nu \arccos \frac{\nu}{\sigma} - \frac{\pi}{4}\right)}{(\nu^2 - \sigma^2)^{1/4} e^{\pi\nu/2}} + O\left(\frac{1}{\sigma}\right). \quad (16)$$

There are two phases with opposite signs. Applying the principle to the full mode function Ψ_ν , one has $\sigma/\nu = \text{sech}(\beta\gamma)$, which match the trajectory if

$$\hbar k_\beta \equiv \hbar \sqrt{\frac{3}{2\kappa}} g \hbar \nu = p_\beta, \quad (17)$$

as expected.

- Fixing $\nu/\sigma < 1$, the asymptotic expansion is not oscillatory, but exponential; hence no WKB approximation works.
- The conclusions also hold for $F_{i\nu}(\sigma)$ and $G_{i\nu}(\sigma)$ for $(+, -)$, as well as $J_\nu(\sigma)$ for $(-, +)$.



Inner product for wave functions

Schrödinger product

- To make sense of amplitude and wave packet, an inner product is necessary
- A common starting point of most approaches is the **Schrödinger product**

$$(\Psi_1, \Psi_2)_S := \int d\chi \Psi_1^*(\beta, \chi) \Psi_2(\beta, \chi); \quad (18)$$

- In terms of a norm, $(\Psi, \Psi)_S \equiv \int d\chi \rho_S(\beta, \chi)$, in which $\rho_S := \Psi^* \Psi$. Manifestly $\rho_S \geq 0$; one has $(\Psi, \Psi)_S > 0$.
- The corresponding Schrödinger current does not satisfy continuity equation $\dot{\rho}_S + \nabla \cdot \vec{j}_S = 0$, because eq. (11) is KG-like.
- $K_{i\nu}$ ⁴ for $(+, +)$, $F_{i\nu}$ and $G_{i\nu}$ (not $J_{i\nu}$!) for $+, -$ can be proved to be orthogonal and complete individually, as well as normalised.

⁴Semyon B. Yakubovich. In: *Opuscula Math.* 26.1 (2006), pp. 161–172. URL: http://www.opuscula.agh.edu.pl/vol26/1/art/opuscula_math_2612.pdf, A. Passian et al. In: *Journal of Mathematical Analysis and Applications* 360.2 (Dec. 2009), pp. 380–390, Radosław Szmytkowski and Sebastian Bielski. In: *Journal of Mathematical Analysis and Applications* 365.1 (May 2010), pp. 195–197.



Peculiarity for the phantom model

Orthonormality and completeness for mode functions; self-adjointness for operators

- $\{J_\nu(\sigma)\}$'s are not orthogonal

$$\int_0^{+\infty} \frac{d\sigma}{\sigma} J_\nu(\sigma) J_{\tilde{\nu}}(\sigma) = \frac{2\sin(\pi(-\tilde{\nu})/2)}{\pi(\nu^2 - \tilde{\nu}^2)}. \quad (19)$$

- Imposing self-adjointness for \hat{p}_χ^2 , one finds that



Integration of the transformed first Friedmann equation

$$p_\beta \neq 0$$

In order to integrate the equation under the implicitising gauge

$$\mathcal{J} \frac{p_\beta^2}{12} \left(-\ell \frac{\kappa^{1/2}}{6} \left(\frac{d\chi}{d\beta} \right)^2 + \kappa^{-1/2} \right) - \kappa^{3/2} V e^{g\mathcal{J}_\chi \chi} = 0, \quad (20)$$

one can substitute

$$\gamma := \sqrt{\frac{3}{2\kappa}} g\beta, \quad \tilde{\sigma}^2 := \frac{p_\beta^2}{12\kappa^2 |V|} e^{-g\mathcal{J}_\chi \chi}, \quad (21)$$

to get

$$\left(\frac{d\tilde{\sigma}}{d\gamma} \right)^2 + \ell(\mathcal{J}v - \tilde{\sigma}^2) = 0 \quad \Rightarrow \quad \frac{d\gamma}{d\tilde{\sigma}} = \pm \frac{1}{\sqrt{\ell(-\mathcal{J}v + \tilde{\sigma}^2)}}, \quad (22)$$

which is of the standard inverse hyperbolic / trigonometric form for $(+, +)$, $(+, -)$ and $(-, +)$.



Integration of the separated mss. Wheeler–DeWitt equation

233

In order to integrate the separated minisuperspace Wheeler–DeWitt equation

$$e^{-ik_{\beta} s_{\beta} \beta} \left(-\ell s \frac{\hbar^2}{2\kappa^{3/2}} \psi''(\chi) - s \frac{\hbar^2 k_{\beta}^2}{12\kappa^{1/2}} \psi(\chi) + \kappa^{3/2} V e^{g s_{\chi} \chi} \psi(\chi) \right) = 0, \quad (23)$$

one can transform

$$\nu := \sqrt{\frac{2\kappa}{3}} \frac{k_{\beta}}{g}, \quad \sigma^2 := \frac{8\kappa^3 |V| e^{g s_{\chi} \chi}}{\hbar^2 g^2}, \quad (14 \text{ rev.})$$

to get

$$\sigma^2 \psi''(\sigma) + \sigma \psi'(\sigma) + \ell(\nu^2 - s \nu \sigma^2) \psi(\sigma) = 0, \quad (24)$$

which is of the standard Besselian form.





- Mit diesem *beamer theme* ist es möglich, Präsentationen in \LaTeX mit der Beamer-Klasse zu erstellen, die dem Corporate Design der Universität zu Köln entsprechen
- Auf die Beamer-Klasse wird in diesem Dokument nicht näher eingegangen, nähere Informationen finden Sie unter <http://latex-beamer.sourceforge.net/>



Das Theme kann mit den folgenden Optionen geladen werden

```
\usepackage[%  
% uk,      %% Farben aller Fakultäten  
wiso,      %% Wiso-Fakultät  
% jura,    %% Rechtswissenschaftliche Fakultät  
% medizin, %% Medizinische Fakultät  
% philo,   %% Philosophische Fakultät  
% matnat,  %% Mathematisch-Naturwissenschaftliche Fakultät  
% human,   %% Humanwissenschaftliche Fakultät  
% verw,    %% Universitätsverwaltung  
{UzK}
```



- Es stehen verschiedene Fußzeilen zur Auswahl, die als Option beim Laden des *themes* übergeben werden:
 - Balken mit allen Fakultätsfarben (Option uk)
 - Balken in jeweils einer Fakultätsfarbe (Optionen wiso, jura, medizin, philo, matnat, human, verw)⁵
- "'Universität zu Köln"' sowie der Name der Fakultät sind im Theme definiert, das Institut oder Seminar kann mit dem Befehl `\institute{}` festgelegt werden
- Die Optionen sind im Quellcode dieser Präsentation dokumentiert

⁵Es werden die offiziellen RGB-Werte aus dem 2-D Handbuch Corporate Design verwendet.



- Der Universitäts- sowie die Fakultätsnamen werden standardmäßig auf Deutsch angezeigt.
- Übergeben Sie dem Paket `babel` die Option `english`, so werden diese Namen entsprechen angepasst.
- Die Übersetzungen können in der Theme-Datei `beamerthemeUzK.sty` geändert werden



block-Umgebungen

Standard (block)

Verwendet die Farbe "Blaugrau Mittel" als Blocktitel-Hintergrund

exampleblock

Bei Verwendung der Fußzeile mit allen Fakultätsfarben Titelhintergrund in Wiso-Grün, sonst in der jeweiligen Fakultätsfarbe

alertblock

Verwendet das Rot der Folientitel



Installation

- Das Theme besteht aus den Dateien `beamerthemeUzK.sty` und `beamercolorthemeUzK.sty` sowie den Grafikdateien `logo.pdf` und `logo-small.pdf`.
- Das Theme kann auf zwei Arten verwendet werden:
 1. Die vier Dateien werden in den selben Ordner wie die zu erstellende Präsentation gelegt
 2. Die vier Dateien werden im lokalen *texmf*-Baum abgelegt
- Die zweite Variante ist der ersten vorzuziehen, da das Theme so an einem zentralen Ort vorliegt



Was noch zu tun ist...

- Erstellen einer eigenen Titelseite
- ...