

Transient Mapping

Daichi Hiramatsu¹ and Corey Mutnik^{1,*}

¹*Department of Physics & Astronomy,
University of Hawaii at Manoa,
2505 Correa Rd, Honolulu, HI, 96822, USA[†]*

DATA COLLECTION

$$l^{II} = 202^\circ \dots b^{II} = \pm 5$$

	$b = (0)$ min	$b = (0)$ max	$b = (1)$ min	$b = (1)$ max
RA (no offset)	93°	+108°	+102°	+117°
Dec (no offset)	-20°	+8°	-15°	+12°
RA (offset)	+93°	+108°	+102°	+117°
Dec (offset)	-20°	+8°	-15°	+12°

DATA REDUCTION

- Sorted data by going through 1deg x 1deg FOV
- Identify stars as most variable
- Run LS
- Discuss how we established uncertainty in period - how this propagates to distance calculations
- How are we going to determine distance - discuss PL-relation

ANALYSIS

Pan-STARRS Comparison

- download Pan-STARRS data (finished)
- compare generated variable star list to PS RA and Dec
- validate observed variable stars
- Determine if PS parameters are worth anything (are candidates actually RR Lyraes)





FIG. 1: Aitoff projection of observed and PS RR Lyrae candidates. Blue are candidates from PS that $\bar{z} = 0.05$, green are PS candidates that $\bar{z} = 0.2$, observed data in red.

Simbad Completeness

- Pull established RR list from Simbad
- Pull other variable data from simbad, too
- Compare list of observed RR to catalogs
- Is anyone actually reading this outline, this bullet point serves no purpose
- Wow, its sad how little Jeff did since class began (especially after JT gave him the code to do it a month ago) - 6 obs x 4 nights = January-April work period haha
- Establish completeness with Simbad


3D map galaxy - var. stars

- Use gri data to identify variable stars
- Use Period-Luminosity relationship to get distance
- Map 3D spatial distribution



(a) *Aitoff map.*

FIG. 2: *Aitoff projection of observed and PS RR Lyrae candidates. Blue are candidates from PS that $\bar{z} = 0.05$, yellow are PS candidates that $\bar{z} = 0.2$, observed data in red, simbad in black.*



(b) *Aitoff projection*

- Determine deviation of variable stars from model

- Variations arise from non-gravitational effects

- Figure out dark matter distribution

CONFIRM ACCELERATED EXPANSION - SUPER NOVEA

- use PanStarrs data to identify supernova locations

* dhiramat@hawaii.edu; cmutnik@hawaii.edu
† Observational Astronomy 301