

UNIVERSITY OF HAWAII • INSTITUTE FOR ASTRONOMY
RESEARCH PROPOSAL – OBSERVING TIME REQUEST

Author* : Daichi Hiramatsu

Author[†] : Corey Mutnik

E-mail* : dhiramat@hawaii.edu

E-mail[†] : cmutnik@hawaii.edu

Institution/Dept^{*,†} : UH

Program Title

A. Measuring the Milky Way

B.

C.

Abstract

Using various data reduction techniques, we will measure the size, shape, and age of the Milky Way galaxy's spiral arms.

TELESCOPE TIME REQUESTED

| Run | Program | Telescope | Instrument | Nights (n) or hours (h) | Observers' initials |
|-----|---------|------------|------------|-------------------------|---------------------|
| 1 | A | Pathfinder | | 22.5 h | DH & CM |

COLLABORATORS

| Name | Institution | E-mail | Program(s) |
|--------------------|-------------|-------------------------|------------|
| John L. Tonry | IfA | jt@ifa.hawaii.edu | A |
| Conor McPartland | IfA | conormcp@ifa.hawaii.edu | A |
| Marielle Dela Cruz | UH | msdelacr@hawaii.edu | A |
| Jeff Kleyner | UH | jeff2012@hawaii.edu | A |
| | | | |
| | | | |
| | | | |
| | | | |

1 SCIENTIFIC JUSTIFICATION

1.1 Immediate Objective

To determine the size and shape of particular spiral arms, variable star distributions must be spatially mapped. In observed regions, the density of variable stars will give insight into the age (*and star formation epoch?*) of our galaxy. The distance to each star will be calculated using the distance modulus,

$$d = 10^{(m-M+5)/5},$$

where d is the distance in *parsecs*, m is the apparent magnitude, and M is the absolute magnitude.

Catalogs such as the Sloan Digital Sky Survey (SDSS) Abazajian et al. (2009), will act as sources of data. Using the search capabilities of VizieR Ochtenbein et al. (2000), variable star locations and magnitudes are easily accessible. From available catalogs, variable star locations within our galaxy will be mapped. Existing catalogs only allow for the distribution of known variable stars to be determined.

Data collected by the gri project will be used to identify new variable stars. Analyzing the gri data, in its entirety, is a cumbersome task. Subtraction of data collected by the gri project will cause transient objects to emerge. Light curves will be used in identification and categorization of desired variable stars. Variable stars comprising less than 1% of all observable stars Allen et al. (2016), significantly reduces the amount of data to analyze. Required computing time also decreases substantially.

1.2 Scientific Rationale

Of the different types of variable stars, we will focus on RR Lyrae, Type 1 Cepheid, and Type 2 Cepheid. These pulsating variables have well established absolute magnitudes B. et al. (2012). From this the luminosity is known, permitting the distance to each star to be calculated using Period-Luminosity (PL) relationship.

1.2.1 RR Lyrae

RR Lyrae have short periods, 1.5 – 24 hours, and are generally classified as stars with spectral type A. On average, absolute magnitudes of RR Lyrae stars fall between 0.6-0.7 Tsujimoto et al. (1998). Using the distance modulus assuming no ISM extinction yields the upper limit on RR Lyrae distance measurements of 7.9 kpc with $m = 15$ (photometric accuracy of 4%). Figure 1 shows the PL relationship for variable stars classified as RR Lyrae Ngeow et al. (1998) (*how do we calibrate different band passes?*). Since the typical age of RR Lyrae is 10 Gyr, it can be used as the lower limit of the age of spiral arms.






Figure 1: Period-Luminosity relationship of RR Lyrae variable stars.

1.2.2 Cepheid

Other variable stars that will be identified are Cepheid Type 1 and Cepheid Type 2. A typical pulsation period of Cepheid star is 1 – 50 days Ryden et al. (2010). These sugergiants span the F, G, and K spectral classes, with average absolute magnitudes between $M = -0.5$ and $M = -6$ Ryden et al. (2010). The PL relation of Cepheid Type 1 and Cepheid Type 2 is shown in Figure 2a. From the PL relation we can extract distances Ngeow et al. (2013). Using $m = 15, 30$ s exposures will allow for photometric accuracy of 4%. A Hertzsprung-Russell diagram of pulsating variable stars is shown by Figure 2b Turner et al. (2012). *Relatively young, star formation epoch?*



(a) Period-Luminosity relationship.



(b) HR-Diagram of pulsating variable stars.

Figure 2: Cepheid Type 1 and Cepheid Type 2 stars.

Near the galactic center, ISM is dense, and the number density of stars is high, which makes optical investigations

quite difficult. Figure 3 shows the distribution of gasses in the Milky Way (Nakanishi and Sofue 2015). In order to determine the spiral arm structure of the Milky Way galaxy, we will avoid the galactic center. To take into account for the ISM extinction, we will use the ratio of total to selective extinction $R = \frac{1}{(\tau_1/\tau_2)-1} = \frac{1}{(\lambda_{eff,2}/\lambda_{eff,1})-1}$ assuming the optical depth $\tau \propto \lambda^{-1}$ according to the Mie scattering for $\lambda < D$ where the D is the dimension of a dust grain (Ryden et al. 2010 & Carroll et al. 2007).




Figure 3: Column density distribution of the sum of HI and H₂ gases.


References

- Abazajian, K. N., Adelman-McCarthy, J. K., Agu eros, M. A., et al. 2009, ApJS, 182, 543
- Allen, S. et al., 2016. (n.d.). The Classification of Stellar Spectra. Retrieved February 15, 2016, from http://www.star.ucl.ac.uk/~pac/spectral_classification.html
- B. et al., 2012. Types of Variables — AAVSO. Retrieved February 16, 2016, from <https://www.aavso.org/types-variables>
- Nakanishi, H. and Sofue, Y., 2015. Three-Dimensional Distribution of the ISM in the Milky Way Galaxy: III. The Total Neutral Gas Disk. *Publ. Astron. Soc. Japan (2014) 00(0), 114*. Retrieved February 15, 2016.
- Ngeow, C. et al., 2013. "Distance Determination From The Cepheid And RR Lyrae Period-Luminosity Relations". *Proc. IAU 9.S301*: 123-128. Web. 18 Feb. 2016.
- Ochsenbein, F., Bauer, P., Marcout, J., et al. 2000, A&AS 143, 221
- Ryden, B. et al., 2010. Foundations of astrophysics. San Francisco: Addison-Wesley.
- Tsujimoto et al., 1998. The Absolute Magnitude of RR Lyrae Stars Derived from the [ITAL]Hipparcos[/ITAL] Catalogue. *The Astrophysical Journal*, 492(1). Retrieved February 15, 2016.
- Turner, R. et al., 2012. H-R Diagram Education Materials — AAVSO. Retrieved February 14, 2016, from <https://www.aavso.org/hr-diagram-education-materials>
- Carroll, B. et al., 2007. An Introduction to Modern Astrophysics. San Francisco: Pearson Addison-Wesley.
- Molnr, L. et al., 2015. Pushing the limits, episode 2: K2 observations of extragalactic RR Lyrae stars in the dwarf galaxy Leo IV ApJ, 812, 2, 2015, arXiv:1508.05587


TECHNICAL JUSTIFICATION

We request 22.5 hours using Pathfinder to obtain g, r, and i imaging for variable stars in the galactic plane.


To calculate an appropriate sampling frequency, a typical RR Lyrae star, EPIC ID 210282474, was used (Molnr et al. 2015) with artificially introduced gaps following the Gaussian distribution. Since sampling in astronomy is always uneven, we used the Lomb-Scargle analysis. The light curve of 210282474 is shown in Figure 4a, and Lomb-Scargle power plots with various sampling conditions are shown in Figures 4b, 4c, and 4d. We want a distinguishable and narrow peak in power plots. To determine distances from the PL relations with $\simeq 5\%$ accuracy, we need peaks with FWHM $\simeq 0.01$ days. From this condition, it is reasonable to choose 15 nights with 10 observations/night as it can be seen in Figures ???. The uncertainty in magnitude was taken to be 10%, and the changes, 1% – 10%, in the uncertainty did not affect the period determinations significantly; we only need 10% photometry.




(a) Light curve of 210282474.



(b) Power plot with 15 nights with 10 observations/night.



(c) Power plot with 10 nights with 10 observations/night.



(d) Power plot with 15 nights with 5 observations/night.

Figure 4: EPIC ID 210282474.

For $m_g = m_r = 16$ and $m_i = 15.7$, an exposure time of 20 sec yields 10% photometry. We want to cover the galactic plane, $90 \text{ deg} < l < 330 \text{ deg}$ and $-2 \text{ deg} < b < 2 \text{ deg}$ avoiding the galactic center and the sun. With the magnitudes, exposure time, and size of the field, we request 1.5 hr/night for 15 nights, resulting the total observational time of 22.5 hr.

Since typical pulsation periods of Cepheid variables are longer than that of RR Lyrae stars, we would be able to sample Cepheid light curves as well by using the sampling rate discussed above.

Things to Discuss Further

- Determine deviation of variable stars from model
- Variations arise from gravitational effects from unknown sources
- Figure out dark matter distribution
- include - Plot of known var star distributions in spiral arms
- comment on knowing m from tying into pan-stars
- difference in 3 variable stars light curves - show examples
- Use catalogs to map known variable star locations
- Cepheid or RR Lyrae? - Focus on RR Lyrae
- gri data will be supplemental - identifying previously unknown variable stars
- More specificity
- Refine overall question
- What are Apparent and Absolute magnitudes
- Sampling necessary to properly describe lightcurve (how often, how long)