

# Transient Mapping

Daichi Hiramatsu<sup>1</sup> and Corey Mutnik<sup>1,\*</sup>

<sup>1</sup>*Department of Physics & Astronomy,  
University of Hawaii at Manoa,  
2505 Correa Rd, Honolulu, HI, 96822, USA<sup>†</sup>*

## DATA COLLECTION

$$l^{II} = 202^\circ \dots b^{II} = \pm 5$$

	$b = (0)$ min	$b = (0)$ max	$b = (1)$ min	$b = (1)$ max
RA (no offset)	93°	+108°	+102°	+117°
Dec (no offset)	-20°	+8°	-15°	+12°
RA (offset)	+93°	+108°	+102°	+117°
Dec (offset)	-20°	+8°	-15°	+12°

## 3D MAP GALAXY - VAR. STARS

- Use gri data to identify variable stars
- Use Period-Luminosity relationship to get distance
- Map 3D spatial distribution
- Determine deviation of variable stars from model

- Variations arise from non-gravitational effects
- Figure out dark matter distribution

## PAN-STARRS COMPARISON

- download Pan-STARRS data (running)
- compare generated variable star list to PS RA and Dec
- validate observed variable stars

## CONFIRM ACCELERATED EXPANSION - SUPER NOVEA

- use PanStarrs data to identify supernova locations

---

\* dhiramat@hawaii.edu; cmutnik@hawaii.edu  
† Observational Astronomy 301