


Solutions to Homework 13

1. Let the state variable represent the number of customers in the system.


(a) The transition diagram is as below:



Solving for π using the “cuts,” we get $\pi = (1/15, 2/15, 4/15, 8/15)$. The throughput is $\lambda_{eff} = 1 \times (1 - \pi_3) = 7/15$.

The average queue length is $L = \pi_2 \times 1 + \pi_3 \times 2 = 4/3$. By Little’s Law, $W = L/\lambda_{eff} = 20/7$.


(b) The transition diagram is as below:



From this diagram, $\pi = (1/7, 2/7, 2/7, 2/7)$. The throughput is $\lambda_{eff} = 1 \times (1 - \pi_3) = 5/7$.

The average queue length is $L = \pi_3 \times 1 = 2/7$. By Little’s Law, $W = L/\lambda_{eff} = 2/5$.


(c) The transition diagram is as below:



$\pi = (3/19, 6/19, 6/19, 4/19)$. The throughput is $\lambda_{eff} = 1 \times (1 - \pi_3) = 15/19$.

The average queue length is $L = 0$. So the average waiting time is 0.

2. Let the state variable represent the number of customers in the system. The transtion diagram is as below:




Using the cuts, we know: $\pi_i \cdot \frac{1}{1+i} \lambda = \pi_{i+1} \mu$ for $i = 0, 1, \dots$. We also know, $\sum_{i=0}^{\infty} \pi_i = 1$. From these equations, we can obtain: $\pi_0 = \frac{1}{\sum_{i=0}^{\infty} \frac{1}{i!} (\frac{\lambda}{\mu})^i} = e^{-\lambda/\mu}$. The last equation comes from the fact $\sum_{i=0}^{\infty} \frac{x^i}{i!} = e^x$. For $i > 0$, $\pi_i = \frac{1}{i!} (\frac{\lambda}{\mu})^i \pi_0$.

The expected number of customers in the system is:

$$\sum_{i=1}^{\infty} i \pi_i = \sum_{i=1}^{\infty} \frac{1}{(i-1)!} (\frac{\lambda}{\mu})^i \pi_0 = \sum_{i=0}^{\infty} \frac{1}{i!} (\frac{\lambda}{\mu})^i \frac{\lambda}{\mu} \pi_0 = \frac{\lambda}{\mu}$$

3. The state represents the number of calls in the system. The transition diagram is as below:



- (a) The generator is as below:

$$\begin{bmatrix} 5 & -5 & 0 & 0 & 0 \\ 4 & -9 & 5 & 0 & 0 \\ 0 & 8 & -13 & 5 & 0 \\ 0 & 0 & -13 & 5 & -8 \\ 0 & 0 & 0 & 8 & -8 \end{bmatrix}$$

(b) Using the cuts method, we know

$$\begin{aligned}\pi_0 \times 5 &= \pi_1 \times 4 \\ \pi_1 \times 5 &= \pi_2 \times 8 \\ \pi_2 \times 5 &= \pi_3 \times 8 \\ \pi_3 \times 5 &= \pi_4 \times 8 \\ 1 &= \pi_0 + \pi_1 + \pi_2 + \pi_3 + \pi_4\end{aligned}$$

The stationary distribution is $\pi = (.261, .327, .204, .128, .080)$.

- (c) An arrival will be accepted if the system is not full (state 4). $\mathbb{P}(\text{accept}) = 1 - \pi_4 = .920$.
- (d) An ambulance is dispatched directly if there is no call or one call in the system. $\mathbb{P}(\text{dispatch directly}) = \pi_0 + \pi_1 = .588$.
- (e) There are 1 and 2 calls waiting in the system when the state is 3 and 4, respectively. No waiting in state 0, 1, or 2. Thus, the average waiting is: $1 \times \pi_3 + 2 \times \pi_4 = .288$.
- (f) As in part (b), a call will be accepted with probability .920. There are three cases when a call being accepted:
 - i. When the call arrives and finds there are no call or one call in the system, his or her waiting time is zero.
 - ii. When the call arrives and finds there are 2 callers in the system, he or she needs to wait until 1 of the 2 finishes the service, which is the minimum of two exponential random variables. In this case, the average waiting time is $15/2$. This happens with probability $\pi_2 = .204$.
 - iii. When the call arrives and finds there are 3 callers in the system, he or she needs to wait until 2 of the 3 finish the service, which is two times the minimum of two exponential random variables. The average waiting time is $2 \times 15/2 = 15$. This happens with probability $\pi_3 = .128$.

For **those accepted to the system**, the average waiting time is $\frac{.204 \times 15/2 + .128 \times 15}{.92} = 3.75$

- (g) As argued in (f), the waiting time the sum of two $\exp(1/7.5)$ random variables.

$$\mathbb{P}(W < 20) = \int_0^{20} 1/7.5 e^{-1/7.5x} \int_0^{20-x} 1/7.5 e^{-1/7.5y} dy dx = .191$$