

Math 2401 - Exam 1

First Name (Print): _____ Last Name (Print): _____ Signature: _____

- There are **5** questions on **5** pages. The exam is worth 50 points in total.
- Answer the questions clearly and completely. You must provide work clearly justifying your solution.
- You can NOT write your work on the back of the page. Use it for scratch work if needed.
- You have 50 minutes to finish your work.

1. (6 points) Find an equation for the plane that passes through $P(1, 1, -1)$, $Q(2, 0, 2)$ and $S(0, -2, 1)$.

Ch. 12.5: # 23 on page 713

Solution.

$\overrightarrow{PQ} = \vec{i} - \vec{j} + 3\vec{k}$, $\overrightarrow{PS} = -\vec{i} - 3\vec{j} + 2\vec{k}$, and $\overrightarrow{PQ} \times \overrightarrow{PS}$ is normal to the plane.

$$\overrightarrow{PQ} \times \overrightarrow{PS} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ 1 & -1 & 3 \\ -1 & -3 & 2 \end{vmatrix} = 7\vec{i} - 5\vec{j} - 4\vec{k}.$$

The equation for the plane is

$$7(x - 1) - 5(y - 1) - 4(z + 1) = 0,$$

or

$$7x - 5y - 4z = 6.$$

■

2. (13 points) The parametrization of a smooth curve is given by

$$\vec{r}(t) = (e^{-t})\vec{i} + (\cos 2t)\vec{j} + (\sin 2t)\vec{k}.$$

Find the angle θ between the velocity vector and acceleration vector at time $t = 0$.

[Leaving θ as an inverse trigonometric function is acceptable.]

Ch. 13.1: # 14 on page 732(even simpler)

Solution.

$$\vec{v}(t) = \frac{d\vec{r}(t)}{dt} = (-e^{-t})\vec{i} + (-2\sin 2t)\vec{j} + (2\cos 2t)\vec{k},$$

$$\vec{a}(t) = \frac{d\vec{v}(t)}{dt} = (e^{-t})\vec{i} + (-4\cos 2t)\vec{j} + (-4\sin 2t)\vec{k},$$

$$\Rightarrow \vec{v}(0) = -\vec{i} + 0\vec{j} + 2\vec{k}, \quad \vec{a}(0) = \vec{i} - 4\vec{j} + 0\vec{k}.$$

Therefore,

$$\vec{v}(0) \cdot \vec{a}(0) = (-1)(1) + (0)(-4) + (2)(0) = -1,$$

$$|\vec{v}(0)| = \sqrt{(-1)^2 + 0^2 + 2^2} = \sqrt{5},$$

$$|\vec{a}(0)| = \sqrt{1^2 + (-4)^2 + 0^2} = \sqrt{17}.$$

Hence,

$$\theta = \cos^{-1} \left(\frac{\vec{v}(0) \cdot \vec{a}(0)}{|\vec{v}(0)||\vec{a}(0)|} \right) = \cos^{-1} \left(-\frac{1}{\sqrt{85}} \right).$$

■

3. (9 points) Solve the initial value problem for $\vec{r}(t)$ as a vector function of t ,

$$\begin{cases} \frac{d\vec{r}}{dt} = \frac{3}{2}(t+1)^{\frac{1}{2}}\vec{i} + e^{-t}\vec{j} + \frac{1}{t+1}\vec{k}, \\ \vec{r}(0) = \vec{k}. \end{cases}$$

Ch. 13.2: # 13 on page 739

Solution.

Integrating both sides of the differential equation with respect to t gives

$$\begin{aligned} \vec{r}(t) &= \left(\int \frac{3}{2}(t+1)^{\frac{1}{2}} dt \right) \vec{i} + \left(\int e^{-t} dt \right) \vec{j} + \left(\int \frac{1}{t+1} dt \right) \vec{k} \\ &= [(t+1)^{\frac{3}{2}}] \vec{i} + (-e^{-t}) \vec{j} + [\ln(t+1)] \vec{k} + \vec{C}, \end{aligned} \tag{0.1}$$

where \vec{C} is a constant vector.

We determine the constant vector \vec{C} by the initial condition. From (0.1), we have

$$\vec{r}(0) = \vec{i} - \vec{j} + \vec{C}.$$

While $\vec{r}(0) = \vec{k}$, so $\vec{C} = -\vec{i} + \vec{j} + \vec{k}$. Therefore,

$$\vec{r}(t) = [(t+1)^{\frac{3}{2}} - 1] \vec{i} + (1 - e^{-t}) \vec{j} + [1 + \ln(t+1)] \vec{k}.$$

■

4. (10 points) The parametrization of a smooth curve is given by

$$\vec{r}(t) = (t \sin t + \cos t)\vec{i} + (t \cos t - \sin t)\vec{j} + \frac{\sqrt{3}}{2}t^2\vec{k}, \quad t \in \mathbb{R}.$$

Ch. 13.3: # 8 and # 12 on page 745 (slightly changed)

(a) Find the arc length of the curve for $-2 \leq t \leq 0$.

Solution.

$$\begin{aligned}\vec{v}(t) &= \vec{r}'(t) = (t \cos t)\vec{i} - (t \sin t)\vec{j} + (\sqrt{3}t)\vec{k} \\ \Rightarrow |\vec{v}(t)| &= \sqrt{(t \cos t)^2 + (-t \sin t)^2 + (\sqrt{3}t)^2} = 2|t|. \\ L &= \int_{-2}^0 |\vec{v}(t)| dt = \int_{-2}^0 2|t| dt = \underbrace{\int_{-2}^0 (-2t) dt}_{\text{since } t \in [-2, 0]} = 4.\end{aligned}$$

■

(b) Find the arc length parameter s along the curve from the base point where $t_0 = 0$.

Solution.

$$s = s(t) = \int_0^t |\vec{v}(\tau)| d\tau = 2 \int_0^t |\tau| d\tau = \begin{cases} t^2, & \text{if } t \geq 0, \\ -t^2, & \text{if } t < 0. \end{cases}$$

■

5. (12 points) The parametrization of a helix is given by

$$\vec{r}(t) = (a \cos t)\vec{i} + (a \sin t)\vec{j} + bt\vec{k}, \quad a, b \geq 0, \quad a^2 + b^2 \neq 0.$$

(a) Find the curvature κ of the helix.

Ch. 13.4: an example given in the lecture!(example 5 on page 750)

Solution.

$$\begin{aligned}\vec{v}(t) &= \frac{d\vec{r}(t)}{dt} = (-a \sin t)\vec{i} + (a \cos t)\vec{j} + b\vec{k}, \\ \vec{a}(t) &= \frac{d\vec{v}(t)}{dt} = (-a \cos t)\vec{i} + (-a \sin t)\vec{j} + 0\vec{k},\end{aligned}$$

$$\vec{v} \times \vec{a} = \begin{vmatrix} \vec{i} & \vec{j} & \vec{k} \\ -a \sin t & a \cos t & b \\ -a \cos t & -a \sin t & 0 \end{vmatrix} = (ab \sin t)\vec{i} + (-ab \cos t)\vec{j} + a^2\vec{k}.$$

$$\begin{aligned}|\vec{v}| &= \sqrt{(-a \sin t)^2 + (a \cos t)^2 + b^2} = \sqrt{a^2 + b^2} \\ |\vec{v} \times \vec{a}| &= \sqrt{(ab \sin t)^2 + (-ab \cos t)^2 + (a^2)^2} = a\sqrt{a^2 + b^2}\end{aligned}$$

$$\kappa = \frac{|\vec{v} \times \vec{a}|}{|\vec{v}|^3} = \frac{a}{a^2 + b^2}.$$

■

(b) Find the torsion τ of the helix

Ch. 13.5: an example given in the lecture!

Solution.

$$\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \dddot{y} & \dddot{z} \end{vmatrix} = \begin{vmatrix} -a \sin t & a \cos t & b \\ -a \cos t & -a \sin t & 0 \\ a \sin t & -a \cos t & 0 \end{vmatrix} = a^2 b.$$

$$\tau = \frac{\begin{vmatrix} \dot{x} & \dot{y} & \dot{z} \\ \ddot{x} & \ddot{y} & \ddot{z} \\ \dddot{x} & \dddot{y} & \dddot{z} \end{vmatrix}}{|\vec{v} \times \vec{a}|^2} = \frac{b}{a^2 + b^2}$$

■