

Quiz 4 Solution

1. Find the equation for the plane containing the intersecting lines:

$$\ell_1(t) = (-1, 2, 1) + t(1, 1, -1), \quad t \in \mathbb{R}; \quad \ell_2(s) = (1, 1, 2) + s(-4, 2, -2), \quad s \in \mathbb{R}.$$

Solution: The normal vector of the plane, is perpendicular to the two lines,

$$\vec{n} = (1, 1, -1) \times (-4, 2, -2) = (0, 6, 6)$$

Any point on the lines is also on the plane, for example $(-1, 2, 1)$ (corresponding to $t = 0$ in the first line). The equation of the plane is

$$(0, 6, 6) \cdot (x, y, z) = (0, 6, 6) \cdot (-1, 2, 1) \implies 6y + 6z = 18$$

2. Using row reduction, find the solution for the following system of equations

$$\begin{cases} x_1 & -3x_3 & = 8 \\ 2x_1 + 2x_2 + 9x_3 & = 7 \\ & x_2 + 5x_3 & = -2 \end{cases}$$

Solution: Row reducing the matrix of the system we get

$$\begin{aligned} \begin{pmatrix} 1 & 0 & -3 & 8 \\ 2 & 2 & 9 & 7 \\ 0 & 1 & 5 & -2 \end{pmatrix} &\rightarrow \begin{pmatrix} 1 & 0 & -3 & 8 \\ 0 & 2 & 15 & -9 \\ 0 & 1 & 5 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -3 & 8 \\ 0 & 0 & 5 & -5 \\ 0 & 1 & 5 & -2 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & -3 & 8 \\ 0 & 1 & 5 & -2 \\ 0 & 0 & 5 & -5 \end{pmatrix} \\ &\rightarrow \begin{pmatrix} 1 & 0 & -3 & 8 \\ 0 & 1 & 5 & -2 \\ 0 & 0 & 1 & -1 \end{pmatrix} \rightarrow \begin{pmatrix} 1 & 0 & 0 & 5 \\ 0 & 1 & 0 & 3 \\ 0 & 0 & 1 & -1 \end{pmatrix} \end{aligned}$$

Therefore the solution is $x_1 = 5$, $x_2 = 3$, $x_3 = -1$.

3. Determine the value(s) of h such that the following matrix is the augmented matrix of a consistent system. Justify your answer. (6 points)

$$\begin{pmatrix} 1 & -4 & 7 & h \\ 0 & 3 & -5 & 2 \\ -2 & 5 & 9 & 6 \end{pmatrix}$$

Solution: Using row reduction

$$\begin{pmatrix} 1 & -4 & 7 & h \\ 0 & 3 & -5 & 2 \\ -2 & 5 & 9 & 6 \end{pmatrix} \xrightarrow{r_3 \rightarrow r_3 + 2r_1} \begin{pmatrix} 1 & -4 & 7 & h \\ 0 & 3 & -5 & 2 \\ 0 & -3 & 23 & 6 + 2h \end{pmatrix} \xrightarrow{r_3 \rightarrow r_3 + r_2} \begin{pmatrix} 1 & -4 & 7 & h \\ 0 & 3 & -5 & 2 \\ 0 & 0 & 18 & 8 + 2h \end{pmatrix}$$

Note that the range of the augmented matrix is the same as the range of the non-augmented matrix, regardless of the value of h . Therefore, the system is consistent for any value of h .