

Full name:

Math 2551, Section B____

Quiz 3 — §13.5, 14.1

Please **clearly** show all work. Scientific calculators are allowed, but no graphing calculators!

(1) Let

$$\mathbf{r}(t) = (\sin t - t \cos t)\mathbf{i} + (\cos t + t \sin t)\mathbf{j}.$$

Find the curvature κ of the curve parameterized by $\mathbf{r}(t)$ as a function of $t > 0$. [8 points]

The velocity of the parameterization is

$$\mathbf{v}(t) = t \sin t \mathbf{i} + t \cos t \mathbf{j} \quad \implies \quad |\mathbf{v}(t)| = |t| = t,$$

the last equality being because we are assuming $t > 0$. The unit tangent vector is therefore

$$\mathbf{T} = \frac{\mathbf{v}}{|\mathbf{v}|} = \sin t \mathbf{i} + \cos t \mathbf{j}.$$

Its derivative is then

$$\frac{d\mathbf{T}}{dt} = \cos t \mathbf{i} - \sin t \mathbf{j} \quad \implies \quad \left| \frac{d\mathbf{T}}{dt} \right| = 1.$$


We can now compute the curvature κ :

$$\kappa = \frac{1}{|\mathbf{v}|} \left| \frac{d\mathbf{T}}{dt} \right| = \boxed{\frac{1}{t}}$$

(2) Let $f(x, y) = \sqrt{2 + x + y}$.

(a) What is the domain of f ? Sketch it. [4 points]

The domain is going to consist of all points (x, y) such that $2 + x + y \geq 0$, or equivalently all points (x, y) such that $y \geq -2 - x$.



(b) What is the range of f ? [4 points]

The range of f is the interval $[0, \infty)$. It is impossible for f to take any values less than 0 since we are (implicitly) taking the positive square root in the definition of f . On the other hand, every value ≥ 0 can be attained by choosing x and y appropriately.

(c) Describe the level curves of f . [4 points]

The level curve $f^{-1}(c)$ has equation $\sqrt{2 + x + y} = c$, which implies that $2 + x + y = c^2$. This is the equation for a line, written more traditionally as $y = c^2 - 2 - x$. The level curves are therefore lines with slope -1 whose y -intercepts are ≥ -2 .