

Stochastic predator-prey models: Population oscillations, spatial correlations, and the effect of randomized rates

ISC-PIF Paris, 7 November 2012

Uwe C. Täuber, Department of Physics, Virginia Tech

Mauro Mobilia, School of Mathematics, University of Leeds

Ivan T. Georgiev, Morgan Stanley, London

Mark J. Washenberger, Webmail.us, Blacksburg, Virginia

Qian He, Citibank, Shanghai

Ulrich Dobramysl, Department of Physics, Virginia Tech

Initial research support through NSF-DMR 0308548, IAESTE



Outline

- Introductory remarks
- Lotka-Volterra predator-prey interaction
- Locally limited resources; predator extinction threshold
- Predator-prey coexistence and oscillations
- Correlations and fluctuation effects; field theory
- Variants: one dimension, triplet model, cyclic predation
- Spatially varying rates and fitness enhancement
- Environmental vs. demographic variability; inheritance
- Summary and conclusions

Fluctuations and correlations in biological systems: fertile ground for statistical physics

- finite number of degrees of freedom: $N^{1/2} / N \sim 1$
 - ⇒ thermodynamic limit need not apply
- complex cooperative, non-equilibrium phenomena:
 - non-random structures: functionally optimized
 - correlations crucial for dynamical processes, e.g., diffusion-limited reactions
 - history dependence, evolving systems with feedback
 - spatial fluctuations non-negligible; prominent both in non-equilibrium steady states and transient features

Physics approach: study simplified models

Lotka-Volterra predator-prey interaction

- **predators:** $A \rightarrow 0$ death, rate μ
- **prey:** $B \rightarrow B+B$ birth, rate σ
- **predation:** $A+B \rightarrow A+A$, rate λ



mean-field factorization \rightarrow rate equations for uniform densities:

$$da(t) / dt = -\mu a(t) + \lambda a(t) b(t)$$

$$db(t) / dt = \sigma b(t) - \lambda a(t) b(t)$$

$$\Rightarrow a^* = \sigma / \lambda, \quad b^* = \mu / \lambda$$

$K = \lambda (a + b) - \sigma \ln a - \mu \ln b$
 conserved \rightarrow neutral cycles,
 population oscillations with
 (linear) frequency $\omega = \sqrt{\sigma\mu}$



(A.J. Lotka, 1920; V. Volterra, 1926)

Model with site restrictions (limited resources)


→ modified rate equations for total carrying capacity $\rho = 1$

$$\begin{aligned} da / dt &= -\mu a(t) + \lambda a(t) b(t) \\ db / dt &= \sigma [1 - a(t) - b(t)] b(t) - \lambda a(t) b(t) \end{aligned}$$

- $\lambda < \mu$: $a \rightarrow 0$, $b \rightarrow 1$;
inactive, **absorbing** state
- active phase: **A** / **B** coexist,
fixed point **node** or **focus** →
transient erratic oscillations
- **active to absorbing transition**:
prey extinction threshold
expect **directed percolation**
(DP) universality class

“individual-based” lattice
Monte Carlo simulations:

at most single particle per site;
 $\sigma = 4.0$, $\mu = 0.1$, 200×200 sites





Note: finite system always reaches
absorbing state, but survival times
 $\sim \exp(c N)$ huge for large N
(see A. Dobrinevski, E. Frey, 2012)

Predator extinction threshold: critical properties


Effective processes for small λ ($b \sim 1$): $A \rightarrow 0$, $A \leftrightarrow A+A$
 \Rightarrow expect DP (*A. Lipowski 1999; T. Antal, M. Droz 2001*)

- **field theory** representation (*M. Doi 1976; L. Peliti 1985*) of master equation for Lotka-Volterra reactions with site restrictions (*F. van Wijland 2001*) \rightarrow **Reggeon effective action**
- measure **critical exponents** in Monte Carlo simulations:
survival probability $P(t) \sim t^{-\delta'}$, $\delta' \approx 0.451$
number of active sites $N(t) \sim t^\theta$, $\theta \approx 0.230$ } 2d **DP** values




Predator / prey coexistence:

Near extinction threshold: stable fixed point is a **node**





observe local predator clusters
(DP clusters in space-time)



Predator / prey coexistence:

Deep in coexistence phase: stable fixed point is a focus





Oscillations near focus:
resonant amplification of
stochastic fluctuations
(A.J. McKane, T.J. Newman, 2005)

Population oscillations in finite systems



$$\sigma = 4.0 ; \mu = 0.1 ; \lambda = 1.0$$

(32x32)





(A. Provata, G. Nicolis, F. Baras, 1999)




$$\lambda = 1.0 ; \sigma = 4.0 ; \mu = 0.1$$








oscillations for **large** system: compare with **mean-field** prediction:



Correlations in the active coexistence phase



abandon site occupation restrictions :



Doi-Peliti field theory: renormalized parameters

Master equation, contribution from predation reaction:

$$\frac{\partial P(n_i, m_i; t)}{\partial t} = \lambda' [(n_i - 1)(m_i + 1) P(n_i - 1, m_i + 1; t) - n_i m_i P(n_i, m_i; t)]$$

Poisson initial distribution: $P(n_i, m_i; 0) = \frac{n_0^{n_i} m_0^{m_i}}{n_i! m_i!} e^{-\bar{n}_0 - \bar{m}_0}$.

Bosonic ladder operators: $[a_i, a_j] = 0$, $[a_i, a_j^\dagger] = \delta_{ij}$

$$\Rightarrow |n_i\rangle = a_i^{\dagger n_i} |0\rangle, a_i |n_i\rangle = n_i |n_i - 1\rangle, a_i^\dagger |n_i\rangle = |n_i + 1\rangle.$$

Similarly for prey, with $[a_i, b_j] = 0 = [a_i, b_j^\dagger]$.

Time-dependent formal state vector:

$$|\Phi(t)\rangle = \sum_{\{n_i\}, \{m_i\}} P(\{n_i\}, \{m_i\}; t) |\{n_i\}, \{m_i\}\rangle$$

$$\Rightarrow \frac{\partial |\Phi(t)\rangle}{\partial t} = -H |\Phi(t)\rangle \quad \text{or} \quad |\Phi(t)\rangle = e^{-Ht} |\Phi(0)\rangle,$$

with $H_{\text{pred}} = -\lambda' \sum_i (a_i^\dagger - b_i^\dagger) a_i^\dagger a_i b_i$.

Expectation values: use projection state $\langle \mathcal{P} | = \langle 0 | \Pi_i e^{a_i b_i}$:

$$\begin{aligned} \langle \mathcal{O}(t) \rangle &= \sum_{\{n_i\}, \{m_i\}} \mathcal{O}(\{n_i\}, \{m_i\}) P(\{n_i\}, \{m_i\}; t) \\ &= \langle \mathcal{P} | \mathcal{O}(\{a_i^\dagger a_i\}, \{b_i^\dagger b_i\}) | \Phi(t) \rangle. \end{aligned}$$

Construct path integral representation with coherent states; continuum action ($\lambda = a_0^d \lambda'$) gives exponential weight:

$$\begin{aligned} S[\hat{a}, a; \hat{b}, b] &= \int d^d x \int dt \left[\hat{a} (\partial_t - D_A \nabla^2) a + \hat{b} (\partial_t - D_B \nabla^2) b \right. \\ &\quad \left. + \mu (\hat{a} - 1) a + \sigma (1 - \hat{b}) \hat{b} b e^{-\rho^{-1} \hat{b} b} + \lambda (\hat{b} - \hat{a}) \hat{a} a b \right] \end{aligned}$$

U.C.T., M.J. Howard, B. Vollmayr-Lee,
J. Phys. A **38** (2005) R79 (review)

Shift $\hat{a} = 1 + \tilde{a}$, $\hat{b} = 1 + \tilde{b}$, and subsequently introduce fluctuating fields $c = a - \langle a \rangle$ and $d = b - \langle b \rangle$:

$$a = \frac{\sigma}{\lambda} \left(1 - \frac{\mu \rho^{-1}}{\lambda} + A_c \right) + c, \quad b = \frac{\mu}{\lambda} (1 + B_c) + d$$

conditions $\langle c \rangle = 0 = \langle d \rangle$ determine counterterms A_c, B_c .
Diagonalize harmonic part of action (assume $D_A = D_B$):

$$c = \frac{1}{\sqrt{2\mu}} \left[\varphi_+ + \varphi_- - \frac{\gamma_0}{i\omega_0} (\varphi_+ - \varphi_-) \right], \quad d = \sqrt{\frac{\mu}{2}} \frac{\varphi_+ - \varphi_-}{i\omega_0}$$

where $\omega_0^2 = \sigma \mu \left(1 - \frac{\mu \rho^{-1}}{\lambda} \right) - \gamma_0^2$, $\gamma_0 = \frac{\sigma \mu \rho^{-1}}{\lambda}$


determine renormalized oscillation frequency, damping, diffusivity in perturbation expansion to one-loop order:

$$\begin{aligned} d = 2 : \quad D_R &= D_0 + \frac{\lambda}{96\pi} \left[1 + 2 \left(\frac{\sigma}{\mu} + \frac{\mu}{\sigma} \right) \right] + \mathcal{O}(\lambda^2) \\ \gamma_R &= \lambda \frac{\omega_0}{D_0} \frac{1}{64} \left[\frac{6}{\pi} \left(\sqrt{\frac{\sigma}{\mu}} - \sqrt{\frac{\mu}{\sigma}} \right) - \left(\frac{\sigma}{\mu} + \frac{\mu}{\sigma} \right) \right] + \mathcal{O}(\lambda^2) \\ \omega_R &= \omega_0 - \lambda \frac{\omega_0}{D_0} \frac{1}{32\pi} \ln \frac{\omega_0}{\gamma_0} \left[1 + \frac{1}{2} \left(\frac{\sigma}{\mu} + \frac{\mu}{\sigma} \right) \right] \\ &\quad + \lambda \frac{\omega_0}{D_0} \frac{3}{32\pi} \left[1 - \frac{\pi}{3} \sqrt{\frac{\sigma}{\mu}} - \frac{1}{4} \left(\frac{\sigma}{\mu} + \frac{\mu}{\sigma} \right) \right] + \mathcal{O}(\lambda^2) \end{aligned}$$


notice symmetry $\mu \leftrightarrow \sigma$ in leading term

U.C.T., J. Phys. Conf. Ser. 319 (2011) 012019;
J. Phys. A **45** (2012) 405002


d = 1:




d = 2:



d = 3:




d = 4:



- $\Delta\gamma_R < 0 \rightarrow$ **instability** against spatial structures
- $\Delta\omega_R < 0 \rightarrow$ drastic frequency reduction; **symmetric** in $\mu \leftrightarrow \sigma$
- $\Delta D_R > 0 \rightarrow$ (diffusive) **spreading accelerated**, fronts faster


Stochastic Lotka-Volterra model in **one** dimension

- no site restriction:
 $\sigma = \mu = \lambda = 0.01$: **diffusion**-dominated




- site occupation restriction:
species **segregation**;
effectively $A + A \rightarrow A$

$$a(t) \sim t^{-1/2} \rightarrow 0$$



- $\sigma = \mu = \lambda = 0.1$: **reaction**-dominated



Stochastic lattice Lotka-Volterra model with spatially varying reaction rates

- 512 x 512 square lattice, up to 1000 particles per site
- reaction probabilities drawn from **Gaussian distribution**, truncated to interval $[0,1]$, **fixed mean**, **different variances**; **fixed** during simulation runs (**quenched** random variables)

Example:

$$\sigma = 0.5,$$

$$\mu = 0.2,$$


$$\lambda = 0.5,$$

$$\Delta\lambda = 0.5$$



initially

$$a(0) = 1,$$

$$b(0) = 1$$




Predator density variation with variance $\Delta\lambda$ (averaged over 50 simulation runs)




- stationary predator **and** prey densities **increase** with $\Delta\lambda$
- amplitude of initial oscillations becomes **larger**
- Fourier peak associated with transient oscillations **broadens**
- relaxation to stationary state **faster**

Spatial correlations and fitness enhancement



- asymptotic population density
- relaxation time, obtained from Fourier peak width
- A/B correlation lengths, from $C_{AA/BB}(r) \sim \exp(-r / l_{corr})$
- A-B typical separation, from zero of $C_{AB}(r)$
- front speed of spreading activity rings into empty region from initially circular prey patch, with predators located in the center






increasing $\Delta\lambda$ leads to more **localized** activity patches, which causes **enhanced** local population fluctuations




*U. Dobramysl, U.C.T.,
Phys. Rev. Lett. **101** (2008) 258102*

Environmental vs. demographic variability

- attach random predation / evasion rates η to lattice sites and *individual* predators / prey; leave $\sigma = 0.5 = \mu$ fixed
- effective predation rate: $\eta' = \frac{1}{2} (\eta_A + \eta_B)$
- offspring rates drawn from Gaussian:
 - centered at *parent* rate (truncated)
 - width $w_\eta \rightarrow$ *mutation* probability
- sharp initial distribution, centered at $\lambda = 0.5$
- lattice: *environmental variability* η with $0 \leq \zeta \leq 1$: $\lambda = \zeta \eta + (1 - \zeta) \eta'$
- extract mean *extinction time* in small systems ($L = 10 \times 10$):
variability enhances robustness




Variable rates attached to individuals; inheritance



w **narrow**: predators/prey evolve to large/small λ ;
no fixation at extremes 1/0 (**n**onlinear dynamics)
 mean-field theory: (semi-)quantitative analysis

w **broad** (uniform):
 predator rate distribution
 stays flat; only prey evolve



Combined spatial environmental and demographic variability:

- characteristic minimum in steady-state population increase at $\zeta \approx 0.3$
 - $\zeta \approx 0 \rightarrow$ A, B optimize, overall neutral
- U. Dobramysl, U.C.T. (2012), arXiv:1206.0973*

Cyclic predation: spatial rock-paper-scissors game






→ **total** particle density ρ **conserved**

$$(a^*, b^*, c^*) = (\sigma, \mu, \lambda) \rho / (\lambda + \sigma + \mu)$$

$$\lambda = 0.2, \sigma = 0.5, \mu = 0.8$$

256 x 256 lattice sites, 500 MCS

without and with site occupation
restrictions: system **well-mixed**



- 1: λ homogeneous, no restrictions
- 2: λ homogeneous, one particle/site
- 3: $0 \leq \lambda \leq 0.4$, no site restrictions
- 4: $0 \leq \lambda \leq 0.4$, one particle/site

→ **negligible disorder effect**;
except: **extreme asymmetry**
 $\lambda \gg \sigma, \mu$: $\rightarrow c^* \approx \rho$,
→ two-species **LV** system

Q. He, M. Mobilia, U.C.T., Phys. Rev. E **82** (2010) 051909;

Eur. Phys. J. B **82** (2011) 97; Q. He, U.C.T., R.K.P. Zia, ibid. **85** (2012) 141

Summary and conclusions

- predator-prey models with **spatial structure** and **stochastic noise**: invalidates Lotka-Volterra mean-field neutral population cycles
- stochastic models yield long-lived erratic **population oscillations**; resonant amplification mechanism for density fluctuations
- lattice site occupation restrictions / **limited resources** induce **predator extinction**; absorbing transition: directed percolation universality class
- spatial stochastic predator-prey systems: complex **spatio-temporal structures**; spreading activity fronts induce **persistent correlations**; stochastic spatial scenario **robust** with respect to model modifications
- fluctuations strongly **renormalize** oscillation properties; fluctuation corrections captured perturbatively through Doi-Peliti **field theory**
- spatial **variability** in the predation rate results in more localized activity patches; population fluctuations in rare favorable regions cause marked **increase** in the population densities / **fitness** of **both** predators and prey
- variable rates attached to **individuals** with **inheritance** and **mutation**: intriguing dynamical **evolution** of rate distributions, **no fixation**
- cyclic **rock-paper-scissors** model: minute effects of disorder, except for extreme **asymmetry** in reaction rates (recovers two-species LV system).

Mapping the Lotka-Volterra reaction kinetics near the predator extinction threshold to directed percolation

Construct Doi–Peliti field theory action, with site occupation restrictions, for the reactions $A \rightarrow \emptyset$ (rate μ), $B \rightarrow B + B$ (rate σ), and $A + B \rightarrow A + A$ (rate λ):

$$S[\hat{a}, a; \hat{b}, b] = \int d^d x \int dt \left[\hat{a} (\partial_t - D_A \nabla^2) a + \mu (\hat{a} - 1) a + \hat{b} (\partial_t - D_B \nabla^2) b + \sigma (1 - \hat{b}) \hat{b} b e^{-\rho^{-1} \hat{b} b} + \lambda (\hat{b} - \hat{a}) \hat{a} a b \right]$$

shift fields $\hat{a} = 1 + \tilde{a}$, $\hat{b} = 1 + \tilde{b}$, expand in ρ^{-1} ($[\rho] = \kappa^d$):

$$S[\tilde{a}, a; \tilde{b}, b] = \int d^d x \int dt \left[\tilde{a} (\partial_t - D_A \nabla^2 + \mu) a + \tilde{b} (\partial_t - D_B \nabla^2 - \sigma) b - \sigma \tilde{b}^2 b + \sigma \rho^{-1} (1 + \tilde{b})^2 \tilde{b} b^2 - \lambda (1 + \tilde{a}) (\tilde{a} - \tilde{b}) a b \right]$$

fluctuating fields $c = b_s - b$, $b_s \approx \rho$, $\langle c \rangle = 0$, $\tilde{c} = -\tilde{b}$:

$$S[\tilde{a}, a; \tilde{c}, c] = \int d^d x \int dt \left[\tilde{a} (\partial_t - D_A \nabla^2 + \mu - \lambda b_s) a + \tilde{c} (\partial_t - D_B \nabla^2 + (2b_s/\rho - 1) \sigma) c + \sigma b_s (2b_s/\rho - 1) \tilde{c}^2 - \sigma \rho^{-1} b_s^2 \tilde{c}^3 - \sigma (4b_s/\rho - 1) \tilde{c}^2 c - \sigma \rho^{-1} (1 + \tilde{c}^2) \tilde{c} c^2 + 2\sigma \rho^{-1} \tilde{c}^2 (c + b_s \tilde{c}) c - \lambda b_s (\tilde{a}^2 + (1 + \tilde{a}) \tilde{c}) a + \lambda (1 + \tilde{a}) (\tilde{a} + \tilde{c}) a c \right]$$

rescale fields $\phi = \sqrt{\sigma} c$, $\tilde{\phi} = \sqrt{\sigma} \tilde{c}$, $\sigma \rightarrow \infty$ ($[\sigma] = \kappa^2$):

$$S_\infty[\tilde{a}, a; \tilde{\phi}, \phi] = \int d^d x \int dt \left[\tilde{a} (\partial_t - D_A \nabla^2 + \mu - \lambda b_s) a - \lambda b_s \tilde{a}^2 a + \tilde{\phi} \phi + b_s \tilde{\phi}^2 \right]$$


add growth-limiting reaction $A + A \rightarrow A$ (rate τ), integrate out fields ϕ and $\tilde{\phi}$, $u = \sqrt{\tau \lambda b_s}$:

$$S_\infty[\tilde{\mathcal{S}}, \mathcal{S}] = \int d^d x \int dt \left[\tilde{\mathcal{S}} (\partial_t + D_A (r_A - \nabla^2)) \mathcal{S} - u \tilde{\mathcal{S}} (\tilde{\mathcal{S}} - \mathcal{S}) \mathcal{S} + \tau \tilde{\mathcal{S}}^2 \mathcal{S}^2 \right]$$

\Rightarrow Reggeon field theory for directed percolation.

Quasi-species mean-field approach

Introduce quasi-species for each rate value: $\lambda_{ij} = \frac{\lambda_i + \lambda_j}{2}$:



f_{ij} denotes the probability for a particle with predation rate λ_i to produce offspring with an assigned rate of λ_j .
Associated mean-field rate equations:


$$\begin{aligned} \dot{a}_i &= -\mu a_i + \sum_{j,k} \lambda_{jk} f_{ki} a_k b_j, \\ \dot{b}_i &= \sigma \sum_k f_{ki} b_k - \sum_j \lambda_{ij} a_j b_i. \end{aligned}$$

General case requires numerical solution; special cases:


- $f_{ij} = \delta_{ij}$, $\lambda_{ij} = \lambda \implies$ standard LV rate equations;
- uniform inheritance distribution $f_{ij} = 1/N \implies$ steady-state solution:

$$a_i = \frac{2\sigma}{N} \sum_j \frac{1}{\lambda_j + \sum_k \lambda_k / N}, \quad b_i = \frac{2\mu}{\lambda_i + \sum_k \lambda_k / N}$$

\implies predator rate distribution uniform,
prey rate distribution inverse linear function.




↓



Quasi-species mean-field approach

Introduce quasi-species for each rate value: $\lambda_{ij} = \frac{\lambda_i + \lambda_j}{2}$:



f_{ij} denotes the probability for a particle with predation rate λ_i to produce offspring with an assigned rate of λ_j .
Associated mean-field rate equations:


$$\begin{aligned} \dot{a}_i &= -\mu a_i + \sum_{j,k} \lambda_{jk} f_{ki} a_k b_j, \\ \dot{b}_i &= \sigma \sum_k f_{ki} b_k - \sum_j \lambda_{ij} a_j b_i. \end{aligned}$$

General case requires numerical solution; special cases:

- $f_{ij} = \delta_{ij}$, $\lambda_{ij} = \lambda \implies$ standard LV rate equations;
- uniform inheritance distribution $f_{ij} = 1/N \implies$ steady-state solution:

$$a_i = \frac{2\sigma}{N} \sum_j \frac{1}{\lambda_j + \sum_k \lambda_k / N}, \quad b_i = \frac{2\mu}{\lambda_i + \sum_k \lambda_k / N}$$

\implies predator rate distribution uniform,
prey rate distribution inverse linear function.



↓

