# Stochastic Compositional Optimization with Compositional Constraints

Shuoguang Yang[1], Wei You[1], Zhe Zhang[2], and Ethan X. Fang[3]

[1]Department of Industrial Engineering and Decision Analytics, The Hong Kong University of Science and Technology

[2]School of Industrial Engineering, Purdue University

[3]Department of Biostatistics and Bioinformatics, Duke University

May 25, 2025

**Abstract**

Stochastic compositional optimization (SCO) has attracted considerable attention due to its broad applicability to important real-world problems. However, existing work on SCO typically assumes that the projection within a solution update is straightforward, which is not the case for problem instances where constraints are in the form of expectations, such as empirical conditional value-at-risk constraints. In this paper, we introduce a novel model that integrates single-level expected value and two-level compositional constraints into the existing SCO framework. Our model has wide applicability to data-driven optimization, fairness optimization, and risk management, including risk-averse optimization and high-moment portfolio selection, and is capable of handling multiple constraints. Additionally, we propose a class of primal-dual algorithms that generate sequences converging to the optimal solution at a rate of $\mathcal{O}(\frac{1}{\sqrt{N}})$ under both single-level and two-level compositional expected-value constraints, where $N$ is the iteration counter, thus establishing benchmarks in expected value constrained SCO. Numerical experiments show the efficiency of our algorithm over real-world applications.

**Keywords:** Stochastic Optimization, Compositional Optimization, Compositional Constraints

## 1 Introduction

Over the past few years, stochastic compositional optimization (SCO) problems have attracted increasing attention. These problems are characterized by their compositional structure, which involves the optimization of functions that are composed of multiple layers of expectations:

$$\min_{x\in\mathcal{X}} \; F(x) = \mathbb{E}_{\xi_1}\Big[f_1(\mathbb{E}_{\xi_2}[f_2(x,\xi_2)],\xi_1)\Big],$$

where $\mathcal{X}\subset\mathbb{R}^{d_x}$ is convex, and $f_2(x,\xi_2)$ and $f_1(y,\xi_1)$ are two functions depending on the random vectors $\xi_2$ and $\xi_1$, respectively. The SCO problems have widespread applications across various

fields, including reinforcement learning (Wang et al. 2017a), risk management (Ahmed et al. 2007, Liu et al. 2022b, Ruszczyński and Shapiro 2006), engineering (Ge et al. 2015, Ghadimi et al. 2020), and machine learning (Chen et al. 2021). Below, we provide a few examples to illustrate these applications.

For example, in reinforcement learning, the policy evaluation problem aims to determine the value function using empirically observed samples. The corresponding Bellman minimization problem can be recast as a two-level SCO problem (Wang et al. 2017b). Another instance is the low-rank matrix estimation problem, which is widely used in engineering applications such as image analysis and recommendation systems. When data arrives sequentially, the online low-rank estimation problem can also be reformulated as a two-level SCO problem (Ghadimi et al. 2020). Additionally, SCO has been applied to Model-Agnostic Meta-Learning (MAML) (Finn et al. 2017, 2019), a powerful approach for training new models from prior related tasks. It was demonstrated by Chen et al. (2021) that MAML falls within the two-level SCO category.

Given the success of SCO in addressing real-world applications, various algorithms have been developed to solve these problems by iteratively updating the solution using projected stochastic gradient descent (SGD) with an approximated gradient (Wang and Liu 2016, Wang et al. 2017b). Despite the rapid advancements in both the theoretical and computational aspects of SCO, most existing studies on constrained SCO assume that the feasible region $\mathcal{X}$ is simple, allowing for efficient projections onto $\mathcal{X}$. Unfortunately, this assumption is not valid in many applications where the constraints are complex, such as nonlinear or high-order polynomials, making projection computationally expensive. The challenge is further exacerbated when the constraints involve expected value functions—whether single-level or compositional—whose explicit forms are not available. These constraints are particularly crucial in practice when incorporating risk-type constraints, such as conditional value-at-risk (CVaR) or high-moment constraints. However, due to the complexity of these constraints, existing SCO approaches are not directly applicable.

In this paper, we aim to solve SCO problems with single-level and two-level compositional expected-value constraints. Specifically, we focus on the following expected value constrained SCO problem that

$$\begin{aligned} &\min_{x\in\mathcal{X}} F(x) = \mathbb{E}_{\xi_1}\Big[f_1(\mathbb{E}_{\xi_2}[f_2(x,\xi_2)],\xi_1)\Big], \\ &\text{s.t. } G^{(j)}(x) \le 0, \text{ for } j = 1,2,\cdots,m. \end{aligned} \tag{1}$$

Here we assume that set $\mathcal{X}$ is convex, and that projection onto it is straightforward. Additionally, we assume that $F(x)$ is convex, and that each $G^{(j)}(x)$ is a convex single-level or two-level compositional function expressed as an expectation, whose explicit form is unknown and must be estimated using observed samples. In Section 2, we consider the case where $G(x)$ is a single-level expected value function and write $G(x) = g(x) = \mathbb{E}_\zeta[g(x,\zeta)]$. In Section 3, we consider the case where $G(x)$ is a two-level compositional function and write $G(x) = \mathbb{E}_{\zeta_1}[g_1(\mathbb{E}_{\zeta_2}[g_2(x,\zeta_2)],\zeta_1)]$.

Our model serves as a powerful tool for incorporating data-driven constraints that arise from large-scale datasets or online scenarios (Yang et al. 2024). Consider a scenario where a decision-maker

has access to a large-scale dataset consisting of $n$ data points $\{w_i\}_{i=1}^n$ and aims to satisfy the data-driven constraint $\frac{1}{n}\sum_{i=1}^n g(x, w_i) \leq 0$ in a decision-making process. When $n$ is large, evaluating the empirical average $\frac{1}{n}\sum_{i=1}^n g(x, w_i)$ and performing projections becomes computationally intractable, even if each function $g(x, w_i)$ is simple. Alternatively, we can reformulate this as a single-level expected value constraint $\mathbb{E}_w[g(x, w)] \leq 0$ where $w$ is a random sample drawn uniformly from the dataset, which is captured by our model (1). Similarly, let $\{w_{1,i}\}_{i=1}^{n_1}$ and $\{w_{2,j}\}_{j=1}^{n_2}$ be two datasets, and let $g_1(z, w_{1,i})$ and $g_2(x, w_{2,j})$ be two functions associated with $w_{1,i}$ and $w_{2,j}$, respectively. The empirical two-level constraint $\frac{1}{n_1}\sum_{i=1}^{n_1} g_1(\frac{1}{n_2}\sum_{j=1}^{n_2} g_2(x, w_{2,j}), w_{1,i}) \leq 0$ can be reformulated as a two-level compositional expected value constraint $\mathbb{E}_{w_1}[g_1(\mathbb{E}_{w_2}[g_2(x, w_2)], w_1)] \leq 0$, where $w_1$ and $w_2$ are random samples from $\{w_{1,i}\}_{i=1}^{n_1}$ and $\{w_{2,j}\}_{j=1}^{n_2}$, respectively. Furthermore, in online scenarios where data arrive sequentially and each data point $w$ induces a loss $g(x, w)$ under decision variable $x$, we can formulate an expected value constraint $\mathbb{E}_w[g(x, w)] \leq 0$.

### 1.1 Motivating Applications

In optimization, most existing approaches treat constraints as penalties on the objective function, with weights assigned to each penalty based on its relative importance. However, in certain cases, hard constraints are necessary to ensure they are strictly satisfied. For example, in finance, regulatory requirements often mandate that the risks of a portfolio do not exceed specific thresholds. In such cases, the commonly used penalty approach may not be suitable, as it could result in solutions that violate these critical hard constraints. In this paper, we propose a model that directly incorporates hard constraints within the big-data regime and introduce an algorithm to solve these problems. Our algorithm can handle multiple hard constraints and guarantees that all are satisfied. Furthermore, our modeling approach is not in conflict with the penalty approach and can be applied when both soft constraints (which can be treated as penalties) and hard constraints (which must be directly modeled) are present. While the penalty approach remains powerful and widely adopted, we believe that our modeling approach offers distinct advantages in addressing the aforementioned scenarios.

We present four motivating applications that requires hard compositional expected-value constraints.

(a) *Conditional value-at-risk.* We first consider the conditional value-at-risk (CVaR) problem (Rockafellar et al. 2000). Let $x \in \mathbb{R}^{d_x}$ be the decision variable and $\ell(x, \zeta)$ be its associated random loss, with $\zeta$ being random variable. We denote by $\mathrm{CVaR}_\alpha(x)$ the $\alpha$-quantile of CVaR. To control the risk of loss such that the $\alpha$-quantile conditional mean of $\ell(x, \zeta)$ is no more than $\gamma$, we can impose a constraint that

$$\Big\{x : \mathrm{CVaR}_\alpha(x) \leq \gamma\Big\} = \Big\{x : \inf_u \Big\{u + \frac{1}{1-\alpha}\mathbb{E}_\zeta\big[(\ell(x, \zeta) - u)_+\big]\Big\} \leq \gamma\Big\},$$

where $u$ is an auxiliary variable. When the distribution of $\zeta$ the above constraint cannot be expressed explicitly and must instead be formulated as an expectation. Additionally, when the loss function $\ell(x, \xi)$ is convex in $x$, the set $\Big\{x : \mathrm{CVaR}_\alpha(x) \leq \gamma\Big\}$ is also convex in $x$.

(b) *Risk-averse mean-deviation constraint.* Risk-averse optimization has attracted significant attention across various communities due to its wide-ranging applications in finance and other industries (Cole et al. 2017, Bruno et al. 2016, Ruszczyński and Shapiro 2006, So et al. 2009). Traditionally, studies in risk-averse optimization have focused on controlling the risk associated with a single utility function. Haneveld and Van Der Vlerk (2006) was one of the first that considered multiple risk levels, and proposed an optimization model that incorporates these risk concerns as constraints. They derived the closed-form of these constraints so that the model can be solved by classical method. However, such an approach is computationally intractable in the big-data regime when the risk-constraints is evaluated on a large dataset.

We consider the big-data regime and propose a model that incorporates multiple risk concerns into the decision-making process using a constrained optimization approach. Specifically, we consider the case where the closed-form of risk constraints are unknown and they have to be written in the form of expectation. Our approach is applicable to instances where constraints can be empirically evaluated, and we don't require the knowledge of constraints' closed form. Letting $x \in \mathbb{R}^{d_x}$ represent the decision variable and $\ell_j(x,\xi)$ for $j = 1, \cdots, K$ be $K$ heterogeneous utility functions, we consider the following set of first-order risk-averse mean-deviation constraints

$$\Big\{x : \mathbb{E}_\xi[\ell_j(x,\xi)] - \mathbb{E}_\xi\Big[(\ell_j(x,\xi) - \mathbb{E}_\xi[\ell_j(x,\xi)])_+\Big] \geq \gamma_j, \text{ for } j = 1, \cdots, K\Big\}.$$

The above set can be reformulated as a region defined by two-level compositional expected-value constraints:

$$\Big\{x : \mathbb{E}_\xi[g_1^{(j)}(\mathbb{E}_\xi\Big[g_2^{(j)}(x,\xi)], \xi)\Big] \leq -\gamma_j, \text{ for } j = 1, 2, \cdots, K\Big\},$$
$$\text{where } g_1^{(j)}\Big((z,x),\xi\Big) = z + (\ell_j(x,\xi) + z)_+ \text{ and } g_2^{(j)}(x,\xi) = (-\ell_j(x,\xi), x).$$

(c) *Portfolio optimization with high-moment constraints.* Since the early 1960s, portfolio managers have been incorporating high-moment considerations into portfolio selection. Let $w \in \mathbb{R}^{d_x}$ represent the random returns of $d_x$ assets, and let $x \in \mathbb{R}^{d_x}$ denote the corresponding portfolio weights. For any even number $p$, the $p$-th order moment of the portfolio's return is given by

$$M_p(x) = \mathbb{E}_w\Big[|w^\top x - \mathbb{E}_w[w^\top x]|^p\Big].$$

High-moment optimization has aroused considerable interest due to its ability to efficiently characterize the distributional properties of portfolios (Harvey et al. 2010). For example, the standardized 4th order moment of a portfolio, known as Kurtosis, is commonly used to capture the tail distribution of portfolios. Portfolios with high Kurtosis often exhibit long tails.

We consider the following portfolio optimization problem with $p$-th order moment constraints

$$\max_x \mathbb{E}_w[w^\top x], \quad \text{s.t. } M_p(x) \leq c_p, \quad \text{for } p = 2, 4, 6, 8, \cdots.$$

Each of the $p$-th order moment constraint can be recast as a two-level compositional expected value constraint

$$\mathbb{E}_w\Big[g_1(\mathbb{E}[g_2(x,w)], w)\Big] \leq c_p,$$
$$\text{where } g_1\Big((z,x),w\Big) = |w^\top x - z|^p \text{ and } g_2(x,w) = (w^\top x, x).$$

(d) *Optimization with group fairness constraints.* Recently, fairness learning has attracted significant attention. In practice, improper use of sensitive attributes, such as gender and race, can lead to discrimination against certain groups, resulting in unfair outcomes. A common approach to address this issue is to introduce fairness constraints. For example, let $w$ be a random feature vector and $z \in 0, 1$ be its associated sensitive attribute. In linear logistic regression, we may enforce the following constraint:

$$\text{Prob}(w^\top x \geq 0 | z = 0) = \text{Prob}(w^\top x \geq 0 | z = 1).$$

This is known as *statistical parity*, which means that the classification decision is independent of the sensitive attribute $z$. Zafar et al. (2019) reformulates the statistical parity constraint as

$$\mathbb{E}[(z - \mathbb{E}[z])w^\top x] = 0,$$

where $\mathbb{E}[\cdot]$ represents the expectation taken over the distribution of $w$ and $z$. In the online setting with streaming data, we typically have no access to the expected value $\mathbb{E}[z]$ of the sensitive variable $z$. To address this challenge, we can introduce a tolerance parameter $c \geq 0$ and impose the following compositional expected-value constraint for fair classification:

$$\mathbb{E}[(z - \mathbb{E}[z])w^\top x] \leq c \text{ and } \mathbb{E}[(z - \mathbb{E}[z])w^\top x] \geq -c.$$

### 1.2 Related Work

**Stochastic compositional optimization.** SCO was first studied by Ermoliev (1988), who analyzed the asymptotic convergence behavior of a two-timescale algorithm employing two sequences of step sizes that preserve different time scales, and established an almost sure convergence guarantee. Wang and Liu (2016) introduced the SCGD algorithm to address two-level SCO, providing the first non-asymptotic convergence result. These convergence rate guarantees were further improved through acceleration techniques (Wang et al. 2017b). Subsequently, Ghadimi et al. (2020) proposed a single-timescale algorithm that achieves an $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate for nonconvex two-level SCO, with various methods being applied to achieve faster convergence rates (Chen et al. 2021, Zhang and Xiao 2021). The extension to multi-level SCO was first explored by Yang et al. (2019), and $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rates for multi-level nonconvex problems were established by Balasubramanian et al. (2022) and Ruszczynski (2021). Liu et al. (2022b) investigated the compositional difference of convex (DC) regime and extensively studied applications in risk management. The convex regime was examined by Zhang and Lan (2024), who established $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rates for both two-level and multi-level SCO using Fenchel conjugates. In all of the above works, the uncertainty only comes from the objective but not the constraints.

**Expected value constrained optimization.** Studies of constrained optimization can be broadly divided into two streams. On one hand, constraints are often treated as penalty terms within the targeted optimization problem to avoid costly projections. Various penalization approaches have been extensively studied (Bertsekas 1997). However, these approaches often require solving

subproblems, which can be computationally expensive and thus limit their applicability. On the other hand, significant efforts have been made to handle constrained optimization problems directly. For deterministic constraints, Nemirovski et al. (2009) proposed a class of gradient-type methods for solving problems with functional constraints. The level-set method, also known as the bundle method, has been explored by Lemaréchal et al. (1995) and Lin et al. (2020). In scenarios where first-order information is unavailable, Lan et al. (2021) developed a class of Frank-Wolfe-based algorithms for solving deterministic optimization problems with complex constraints. Recently, there has been growing interest in addressing problems with stochastic constraints. Lan and Zhou (2020) developed the cooperative stochastic approximation (CSA) algorithm, which updates the solution based on the estimated feasibility of stochastic constraints. Following this, Boob et al. (2023) focused on the smooth scenario and proposed the constraint extrapolation (ConEx) method, a primal-dual algorithm that iteratively updates the primal and dual solutions using accelerated SGD. ConEx achieves a convergence rate of $\mathcal{O}(\frac{1}{\sqrt{N}})$ and sets the benchmark for expected value constrained optimization. Yang et al. (2024) studied expected value constrained problems with minimax objectives, while Yu et al. (2017) investigated the online scenario where an arbitrary objective function is introduced at each online round. For more detailed reviews on constrained optimization, see Boob et al. (2023). It is important to note that all of the aforementioned works focus on single-level stochastic objectives and constraints, and do not address the compositional setting.

### 1.3 Contributions

Our work seeks to incorporate various complex expected-value constraints into the current SCO framework to better address real-world applications. The contributions of this paper are four-fold:

- We introduce a new model that integrates single-level and two-level compositional expected-value constraints into the SCO framework. This model is designed to handle complex expectation constraints that arise in real-world applications involving online or large-scale datasets. We illustrate the applicability of our model with three examples in risk management.
- We address SCO with single-level expected-value constraints by reformulating it as a saddle point optimization problem and proposing a primal-dual algorithm that iteratively updates the primal and dual variables. We demonstrate that the solution path converges to the optimal solution at the optimal rate of $\mathcal{O}(\frac{1}{\sqrt{N}})$. Numerical experiments validate the theoretical convergence rate and showcase the practical effectiveness of our approach.
- Extending our work to SCO with two-level compositional expected-value constraints, we propose a modified primal-dual algorithm to address these challenges. We prove that the solution path achieves the same $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate to the optimal solution.
- We conducted numerical experiments on a stochastic portfolio optimization problem with CVaR constraints. The results suggest that our algorithm is both efficient and stable across problem instances of varying dimensions.

**Notation.** For any vectors $a, b \in \mathbb{R}^n$, we denote by $\langle a, b\rangle = \sum_{i=1}^n a_i b_i$ their inner product and denote by $\max\{a, b\} = (\max\{a_1, b_1\}, \cdots, \max\{a_n, b_n\})^\top \in \mathbb{R}^n$ their component-wise maximum. We denote by $\|a\| = \|a\|_2$ the Euclidean norm for any vector $a \in \mathbb{R}^n$ and denote by $\|A\| = \|A\|_2$ the Euclidean norm for any matrix $A \in \mathbb{R}^{n\times n'}$. For any non-smooth stochastic function $f(x, \omega)$, with some slight abuse of notations, we denote by $f(x) = \mathbb{E}[f(x, \omega)]$, denote by $\partial f(x)$ the collection of its deterministic subgradients, and denote by $\partial f(x, \omega)$ its stochastic subgradient such that $\mathbb{E}_\omega[\partial f(x, \omega)] \in \partial f(x)$.

## 2 Stochastic Compositional Optimization with Single-level Expected-Value Constraints

In this section, we consider the single-level expected value constrained stochastic compositional optimization (EC-SCO) problem

$$\begin{aligned} \min_{x\in\mathcal{X}} \quad & F(x) = f_1 \circ f_2(x) = \mathbb{E}_{\xi_1}\Big[f_1(\mathbb{E}_{\xi_2}[f_2(x, \xi_2)], \xi_1)\Big], \\ \text{s.t.} \quad & G^{(j)}(x) = g^{(j)}(x) = \mathbb{E}_{\zeta_j}[g^j(x, \zeta_j)] \le 0, \text{ for } j = 1, 2, \cdots, m, \end{aligned} \tag{2}$$

where $F$ is *convex*, $\mathcal{X} \subset \mathbb{R}^{d_x}$ is a convex set. Here, $\xi_1$, $\xi_2$, and $\{\zeta_j\}_{j=1}^m$ are random vectors. We assume $\xi_1$, $\xi_2$, and $\zeta = [\zeta_1, \cdots, \zeta_m]$ are independent of each other, and we write $f_1(y, \xi_1), f_2(x, \xi_2)$, $\{g^j(x, \zeta_j)\}_{j=1}^m$ as the sampled function values. With a bit abuse of notation, we write $f_1(y) = \mathbb{E}_{\xi_1}[f_1(y, \xi_1)] : \mathbb{R}^{d_y} \mapsto \mathbb{R}$ as a convex and differentiable function, write $f_2(x) = \mathbb{E}_{\xi_2}[f_2(x, \xi_2)] : \mathbb{R}^{d_x} \mapsto \mathbb{R}^{d_y}$ as a convex function, and write $g^{(j)}(x) = \mathbb{E}_{\zeta_j}[g^{(j)}(x, \zeta_j)] : \mathbb{R}^{d_x} \mapsto \mathbb{R}$ for $j = 1, 2, \cdots, m$ as convex functions. To be specific, we write $f_2(x) = \Big(f_2^{(1)}(x), \cdots, f_2^{(d_y)}(x)\Big)^\top$ and assume each $f_2^{(j)}$ is a convex function. Note that the functions $f_1(y), f_2(x)$, and $g(x)$ are taken in the forms of expectation, whose explicit forms are unknown, and can only be estimated from observed samples. For notational convenience, the functions $f_1, f_2, g$ always refer to their sample-value form when they have two arguments, and always refer to their expectation form when they have one argument.

To facilitate our discussion, we briefly review the Fenchel conjugate here. In particular, for any convex function $h(z) : \mathbb{R}^{d_z} \to \mathbb{R}$, its Fenchel conjugate is $h^*(z) = \max_{u\in\mathbb{R}^{d_z}} \langle u, z\rangle - h(u)$. With a slight abuse of notation, for any function $\tilde{h}(z) : \mathbb{R}^{d_z} \mapsto \mathbb{R}^k$, where each component, $\tilde{h}_j(z)$ for $j = 1, ..., k$, of the function is convex, we let $\tilde{h}_j^*(z) = \max_{w\in\mathbb{R}^{d_z}} \langle w, z\rangle - \tilde{h}_j(w)$, and let $\tilde{h}^*(z) = \Big(\tilde{h}_1^*(z), ..., \tilde{h}_k^*(z)\Big)^\top$. That is, we let the Fenchel conjugate of $\tilde{h}(z)$ be $\tilde{h}^*(z) : \mathbb{R}^{d_z} \to \mathbb{R}^k$, where the $j$-th component of $\tilde{h}^*(z)$, $\tilde{h}_j^*(z)$, is the Fenchel conjugate of $\tilde{h}_j(z)$, the $j$-th component of $\tilde{h}(z)$. For any convex smooth function $f$, we denote by $D_f(x, y) = f(y) - f(x) - \langle \nabla f(x), y - x\rangle$ its associated Bregman's distance for ease of presentation. Given $x \in \mathbb{R}^{n_2}$, we denote by $g(x) = \Big(g^{(1)}(x), \cdots, g^{(m)}(x)\Big)^\top \in \mathbb{R}^m$ the vector of constraint values and denote by $g(x, \zeta) = \Big(g^{(1)}(x, \zeta_1), \cdots, g^{(m)}(x, \zeta_m)\Big)^\top \in \mathbb{R}^m$ the vector of the sample constraint values.

We begin by specifying the stochastic sampling environment and assume access to a blackbox sampling oracle that provides stochastic zeroth- and first-order information for the objective functions

$f_1, f_2$ and constraint function $g$ upon each query. Specifically, we assume the availability of the following *Sampling Oracle* ($\mathcal{SO}$) such that

- Given $x \in \mathcal{X}$, the $\mathcal{SO}$ returns a sample value $f_2(x, \xi_2) \in \mathbb{R}^{d_y}$ and a sample subgradient

$$\partial f_2(x, \xi_2) \in \mathbb{R}^{d_x \times d_y}.$$

- Given $x \in \mathcal{X}$, the $\mathcal{SO}$ returns a sample value $g(x, \zeta) \in \mathbb{R}^m$ and a sample subgradient

$$\partial g(x, \zeta) \in \mathbb{R}^{d_x \times m}.$$

- Given $y \in \mathbb{R}^{n_1}$, the $\mathcal{SO}$ returns a sample value $f_1(y, \xi_1) \in \mathbb{R}$ and a sample gradient

$$\nabla f_1(y, \xi_1) \in \mathbb{R}^{d_y}.$$

Note that in the above $\mathcal{SO}$, we do not impose any smoothness assumption for the inner level function $f_2$ and the constraint function $g$, and we let $\partial f_2(x, \xi_2)$ and $\partial g(x, \zeta)$ be their sample subgradients, respectively. We then introduce the Lagrangian dual function of problem (2) and briefly illustrate the challenges of solving the problem. Specifically, the Lagrangian dual function of problem (2) is

$$\mathcal{L}(x, \lambda) = F(x) + \sum_{j=1}^{m} \lambda_j g^{(j)}(x), \text{ where } \lambda \geq 0. \tag{3}$$

By the convexity of $F$ and $g_j$'s, the above Lagrangian $\mathcal{L}(x, y)$ is convex in $x$ and concave in $\lambda$. We assume that there exists a *saddle point* $(x^*, \lambda^*)$ such that

$$(x^*, \lambda^*) = \operatorname*{argmin}_{x \in \mathcal{X}} \max_{\lambda \in \mathbb{R}^m_+} \mathcal{L}(x, \lambda),$$

and we have that for any $(x, \lambda) \in \mathcal{X} \times \mathbb{R}^m_+$, it holds that

$$\mathcal{L}(x^*, \lambda) \leq \mathcal{L}(x^*, \lambda^*) \leq \mathcal{L}(x, \lambda^*). \tag{4}$$

Note that the existence of such a saddle point is guaranteed under the mild Slater condition (Slater 2014). With a slight abuse of notation, the first-order partial subgradients of $\mathcal{L}$ satisfy

$$\partial_x \mathcal{L}(x, \lambda) = \partial F(x) + \sum_{j=1}^{m} \lambda_j \partial g^{(j)}(x) \text{ and } \nabla_\lambda \mathcal{L}(x, \lambda) = g(x), \quad \forall x \in \mathcal{X}, \lambda \in \mathbb{R}^m_+.$$

**Challenges.** The key difference between expected value constrained SCO and classical minimax optimization (Liu et al. 2022a, Tan et al. 2018, Yan et al. 2019) is the unavailablity of Lipschitz continuity within the Lagrangian dual function $\mathcal{L}(x, \lambda)$ (3). When the classical minimax algorithms are employed to compute the saddle point $(x^*, \lambda^*) \in \operatorname{argmin}_{x \in \mathcal{X}} \max_{\lambda \in \mathbb{R}^m_+} \mathcal{L}(x, \lambda)$, existing works often assume $\mathcal{L}(x, \lambda)$ is Lipschitz continuous in $(x, \lambda)$ for some constants $C_x, C_\lambda > 0$ that

$$\|\mathcal{L}(x_1, \lambda_1) - \mathcal{L}(x_2, \lambda_2)\| \leq C_x \|x_1 - x_2\| + C_\lambda \|\lambda_1 - \lambda_2\|.$$

However, such a Lipschitz continuous property does not naturally hold for expected value constrained optimization because the dual variable is defined on the region $\lambda \in \mathbb{R}^m_+$ of an infinite diameter. As a result, when $F(x)$ and $g(x)$ are Lipschitz continuous such that $\|F(x_1) - F(x_2)\| \leq C_f\|x_1 - x_2\|$ and $\|g(x_1) - g(x_2)\| \leq C_g\|x_1 - x_2\|$, we only have

$$\|\mathcal{L}(x_1, \lambda) - \mathcal{L}(x_2, \lambda)\| \leq (C_f + C_g\|\lambda\|)\|x_1 - x_2\|, \quad \text{for all } \lambda \in \mathbb{R}^m_+, \tag{5}$$

which loses the Lipschitz continuity when $\|\lambda\| \to \infty$. Consequently, existing results on stochastic minimax optimization do not apply to the expected value constrained setting. It requires the development of new algorithms and theories to ensure the boundedness of the generated dual solution $\{\lambda_t\}$ so that $\mathcal{L}(x_t, \lambda_t)$ preserves Lipschitz continuity over the solution path $\{(x_t, \lambda_t)\}$.

In this paper, we aim to develop a class of efficient primal-dual algorithms to solve the expected value constrained optimization, which is challenging from several aspects.

- *Primal update:* At iteration $t$, we first update the primal variable $x_t$ by obtaining an estimator of $\partial_x\mathcal{L}(x_t, \lambda_t)$. The key challenge to solving the EC-SCO problem lies in the unavailability of an unbiased estimator of the subgradient of $F$. Specifically, by employing the chain rule, an unbiased estimator of $\partial F(x)$ is

$$\partial f_2(x, \xi_2)\nabla f_1\Big(\mathbb{E}_{\xi_2}[f_2(x, \xi_2)], \xi_1\Big). \tag{6}$$

  Unfortunately, the absence of knowledge about $\mathbb{E}_{\xi_2}[f_2(x_t, \xi_2)]$ induces a bias when we use the plug-in sample estimator $\nabla f_1(f_2(x_t, \xi_2), \xi_1)$ (Wang et al. 2017a). Without an unbiased estimator of $\partial F(x)$, the primal update is challenging.

- *Dual update:* Letting $\{\lambda_t\}$ be a sequence of generated dual variables, based on (5), we note that $\mathcal{L}(x, \lambda_t)$ is $(C_f + C_g\|\lambda_t\|)$-Lipschitz continuous in $x$ if $\lambda_t$ is bounded. Unfortunately, the boundedness of dual variables $\{\lambda_t\}$ only holds under certain *restrictive* settings, such as vanilla expected value constrained stochastic optimization (Boob et al. 2023, Madavan and Bose 2021). In comparison with the classical single-level stochastic optimization problems, the compositional objective $F(x)$ is much more complicated, and existing results cannot be applied to prove the boundedness of the dual variables.

To address the above-mentioned obstacles, in the next subsection, we develop an efficient algorithm to tackle problem (2). Before presenting this algorithm, we impose some assumptions on the objective functions $f_1$ and $f_2$.

**Assumption 1** *Let $C_{f_1}, C_{f_2}, \sigma_{f_1}, \sigma_{f_2}, L_{f_1}$, and $D_X$ be positive constants. We assume that:*

*(a) The set $\mathcal{X}$ is convex and bounded such that $\sup_{x\in\mathcal{X}} \|x - x^*\| \leq D_X$ where $x^*$ satisfies (4).*

*(b) $f_1, f_2$ are convex, and have equivalent representations by their Fenchel conjugates $f_1^*(\pi_1)$ and $f_2^*(\pi_2)$ that for all $x \in \mathcal{X}, y \in \mathbb{R}^{d_y}$,*

$$f_1(y) = \max_{\pi_1\in\Pi_1} \langle \pi_1, y\rangle - f_1^*(\pi_1), \text{ and } f_2(x) = \max_{\pi_2\in\Pi_2} \langle \pi_2, x\rangle - f_2^*(\pi_2),$$

*where $\Pi_1 \subset \mathbb{R}^{d_y}$ and $\Pi_2 \subset \mathbb{R}^{d_x\times d_y}$ are the domains of $f_1^*$ and $f_2^*$, respectively.*

*(c) $f_1$ is an $L_{f_1}$-smooth function with a Lipschitz continuous gradient*

$$\|\nabla f_1(y_1) - \nabla f_1(y_2)\| \leq L_{f_1}\|y_1 - y_2\|, \ \forall y_1, y_2 \in \mathbb{R}^{d_y}.$$

*(d) The sample value $f_2(x, \xi_2)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\xi_2}[f_2(x, \xi_2)] = f_2(x), \ \textit{and } \mathbb{E}_{\xi_2}[\|f_2(x, \xi_2) - f_2(x)\|^2] = \sigma_{f_2}^2, \ \forall x \in \mathcal{X}.$$

*(e) The sample value $f_1(y, \xi_1)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\xi_1}[f_1(y, \xi_1)] = f_1(y), \ \textit{and } \mathbb{E}_{\xi_1}[(f_1(y, \xi_1) - f_1(y))^2] = \sigma_{f_1}^2, \ \forall y \in \mathbb{R}^{d_y}.$$

*(f) The sample subgradient $\partial f_2(x, \xi_2)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\xi_2}[\partial f_2(x, \xi_2)] \in \partial f_2(x), \ \textit{and } \mathbb{E}_{\xi_2}[\|\partial f_2(x, \xi_2)\|^2] \leq C_{f_2}^2, \forall x \in \mathcal{X}.$$

*(g) The sample gradient $\nabla f_1(y, \xi_1)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\xi_1}[\nabla f_1(y, \xi_1)] = \nabla f_1(y), \ \textit{and } \mathbb{E}_{\xi_1}[\|\nabla f_1(y, \xi_1)\|^2] \leq C_{f_1}^2, \ \forall y \in \mathbb{R}^{d_y}.$$

As stated in (b) above, when $f_1$ and $f_2$ are convex, we have their equivalent representations through their Fenchel conjugates. Also, we recall that for any convex but possibly nonsmooth function $f$, for a dual variable $\pi_y$ associated with a primal variable $y$ such that $\pi_y \in \operatorname{argmax}_{\pi \in \Pi_y} \langle \pi, y \rangle - f^*(\pi)$, we have that $\pi_y$ is a subgradient of $f(y)$. In particular, we have

$$\pi_y \in \operatorname*{argmax}_{\pi \in \Pi_y} \langle \pi, y \rangle - f^*(\pi) \iff \pi_y \in \partial f(y) \iff f(y) = \langle \pi_y, y \rangle - f^*(\pi_y).$$

Further, we note that for any $L_f$-smooth convex function $f$, the Bregman distance function generated by its Fenchel conjugate function is $\frac{1}{L_f}$-strongly convex (see, for example, Theorem 5.26 in Beck (2017)) that

$$\mathcal{D}_{f^*}(\bar{\pi}; \pi) = f^*(\bar{\pi}) - f^*(\pi) - \langle \nabla f^*(\pi), \bar{\pi} - \pi \rangle \geq \tfrac{1}{2L_f}\|\pi - \bar{\pi}\|^2, \quad \forall \pi, \bar{\pi} \in \operatorname{dom}(f^*).$$

Thus the $L_{f_1}$-smoothness assumption of $f_1$ implies that $\mathcal{D}_{f_1^*}$ is $\frac{1}{L_{f_1}}$-strongly convex.

Then, we make the following assumptions for the constraint functions $g^{(j)}$'s.

**Assumption 2** *Let $C_g$ and $\sigma_g$ be positive constants. The constraint functions $g^{(j)}$'s satisfy:*

*(a) For $j = 1, \cdots, m$, each component $g^{(j)}$ is convex such that it admits an equivalent representation using its Fenchel conjugate that*

$$g^{(j)}(x) = \max_{\bar{v}^{(j)} \in V^{(j)}} \langle x, \bar{v}^{(j)} \rangle - [g^{(j)}(v)]^*, \quad \forall x \in \mathcal{X},$$

*where $V^{(j)} \subset \mathbb{R}^{n_2}$ is the domain of $[g^{(j)}]^*$.*

*(b) The sample value $g(x,\zeta)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment such that*

$$\mathbb{E}_\zeta[g(x,\zeta)] = g(x), \text{ and } \mathbb{E}_\zeta[\|g(x,\zeta) - g(x)\|^2] \le \sigma_g^2, \quad \forall x \in \mathcal{X}.$$

*(c) The sample subgradient $\partial g(x,\zeta)$ returned by the $\mathcal{SO}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_\zeta[\partial g(x,\zeta)] \in \partial g(x), \text{ and } \mathbb{E}_\zeta[\|\partial g(x,\zeta)\|^2] \le C_g^2, \quad \forall x \in \mathcal{X}.$$

In this paper, we focus on two scenarios: (i) the inner function $f_2$ is affine; (ii) the inner function $f_2$ is non-affine and the outer function is monotone non-decreasing, which is specified in the next assumption.

**Assumption 3** *For a non-affine function $f_2$, $\nabla f_1(x)$ is always component-wise nonnegative, i.e., $\nabla f_1(y) \ge 0$ for any $y \in \mathbb{R}^{d_y}$. That is, the outer function $f_1$ is monotone non-decreasing if $f_2$ is non-affine.*

For ease of notation, in what follows, we omit the subscripts $\xi_1, \xi_2$, and $\zeta$ within $\mathbb{E}_{\xi_1}[\cdot], \mathbb{E}_{\xi_2}[\cdot]$, and $\mathbb{E}_\zeta[\cdot]$.

### 2.1 A Primal-Dual Algorithm

We now propose our algorithm to tackle problem (2). Our algorithm operates iteratively, alternately updating the primal variable $x$ and dual variable $\lambda$. At the $t$-th iteration, we begin by querying the $\mathcal{SO}$ *twice* at the previous solution $x_t$ to obtain a sample of $f_2(x_t, \xi_{2,t}^1)$ and an independent sample of subgradient $\partial f_2(x_t, \xi_{2,t}^0)$. Then, given the previous estimator $y_t$ for $\mathbb{E}[f_2(x_t,\xi_2)]$ from the previous iteration, we update our estimator. Specifically, we take a weighted average of $y_t$ and $f_2(x_t, \xi_{2,t}^1)$ such that for some $\tau_t \in (0,1)$

$$y_{t+1} = \frac{f_2(x_t, \xi_{2,t}^1) + \tau_t y_t}{1+\tau_t}. \tag{7}$$

Next, given $y_{t+1}$, we query the $\mathcal{SO}$ at $y_{t+1}$ to obtain a sample gradient $\nabla f_1(y_{t+1}, \xi_{1,t}^0)$. In addition, we query the $\mathcal{SO}$ at $x_t$, where $x_t$ is the solution from the previous iteration, for a sample subgradient $\partial g(x_t, \zeta_t^0)$. Then, we update the primal solution $x_t$ by a projected stochastic gradient step such that

$$\begin{aligned} x_{t+1} &= \operatorname*{argmin}_{x\in\mathcal{X}} \left\{ \left\langle \partial f_2(x_t, \xi_{2,t}^0) \nabla f_1(y_{t+1}, \xi_{1,t}^0) + \partial g(x_t, \zeta_t^0)\lambda_t, x \right\rangle + \frac{\eta_t}{2}\|x - x_t\|^2 \right\} \\ &= \mathrm{Proj}_{\mathcal{X}} \left[ x_t - \left\{ \partial f_2(x_t, \xi_{2,t}^0) \nabla f_1(y_{t+1}, \xi_{1,t}^0) + \partial g(x_t, \zeta_t^0)\lambda_t \right\} / \eta_t \right], \end{aligned} \tag{8}$$

where $\eta_t > 0$ is prespecified.

For the dual update part, we query an independent sample of $g(x_t, \zeta_t^1)$, and update the dual variable $\lambda_{t+1}$ by a projected stochastic gradient ascent step that

$$\lambda_{t+1} \in \operatorname*{argmax}_{\lambda\in\mathbb{R}_+^m} \left\{ \left\langle g(x_t, \zeta_t^1), \lambda \right\rangle - \frac{\alpha_t}{2}\|\lambda_t - \lambda\|^2 \right\} = \left[ \lambda_t + g(x_t, \zeta_t^1)/\alpha_t \right]_+, \tag{9}$$

where $\alpha_t > 0$ is prespecified. We summarize the proposed algorithm in Algorithm 1.

**Algorithm 1** Expected Value Constrained Stochastic Compositional Gradient Descent (EC-SCGD)

**Input :** Step-sizes $\{\alpha_t\}$, $\{\eta_t\}$, $\{\tau_t\}$, initial points $x_0 \in \mathcal{X}$, $y_0 = 0$, $\lambda_0 \in \mathbb{R}^m_+$, sampling oracle $\mathcal{SO}$

**for** $t = 0, 1, 2, ..., N-1$ **do**

Query the $\mathcal{SO}$ at $x_t$ twice for the sample values $f_2(x_t, \xi^i_{2,t})$ and sample subgradients $\partial f_2(x_t, \xi^i_{2,t})$, for $i = 0, 1$.

Update

$$y_{t+1} = \frac{f_2(x_t, \xi^1_{2,t}) + \tau_t y_t}{1 + \tau_t}.$$

Query $\mathcal{SO}$ once at $y_{t+1}$ to obtain a sampled gradient $\nabla f_1(y_{t+1}, \xi^0_1)$.

Query the $\mathcal{SO}$ once at $x_t$ to obtain a sampled subgradient $\partial g(x_t, \zeta^0_t)$.

Update the main solution $x_{t+1}$ by

$$x_{t+1} = \mathrm{Proj}_{\mathcal{X}}\left[x_t - \frac{\partial f_2(x_t, \xi^0_{2,t}) \nabla f_1(y_{t+1}, \xi^0_{1,t}) + \partial g(x_t, \zeta^0_t)\lambda_t}{\eta_t}\right].$$

Query the $\mathcal{SO}$ once at $x_t$ to obtain $g(x_t, \zeta^1_t)$, update the dual variable $\lambda_t$ by

$$\lambda_{t+1} = \Big[\lambda_t + g(x_t, \zeta^1_t)/\alpha_t\Big]_+.$$

**end for**

**Output :** $\bar{x}_N = \frac{1}{N}\sum_{t=1}^N x_t$

### 2.2 Stochastic Sequential Dual Interpretation

We provide a stochastic sequential dual (SSD) interpretation (Zhang et al. 2021, Zhang and Lan 2024) of Algorithm 1, which is the key for our later analysis. The SSD interpretation relies on the Fenchel conjugate reformulations for the convex objective and constraints. To build up the SSD framework, we first define the following composite Lagrangian functions for $f_1, f_2$ and constraint functions $g^{(j)}$:

$$\begin{aligned} \mathcal{L}_{f_2}(x, \pi_2) &= \pi_2^\top x - f_2^*(\pi_2), \\ \mathcal{L}_F(x, \pi_{1:2}) &= \mathcal{L}_{f_1}(x, \pi_1, \pi_2) = \pi_1^\top \mathcal{L}_{f_2}(x; \pi_2) - f_1^*(\pi_1), \\ \mathcal{L}_{g^{(j)}}(x, v^{(j)}) &= x^\top v^{(j)} - [g^{(j)}]^*(v^{(j)}), \text{ for } j = 1, \cdots, m. \end{aligned} \tag{10}$$

In what follows, we denote by $v = [v^{(1)}, \cdots, v^{(m)}] \in \mathbb{R}^{d_x \times m}$ the collection of $v^{(j)}$'s, denote by $\mathcal{L}_g(x, v) = [\mathcal{L}_{g^{(1)}}(x, v^{(1)}), \cdots, \mathcal{L}_{g^{(m)}}(x, v^{(m)})]^\top \in \mathbb{R}^m$, and denote by $\mathcal{V} = [V^{(1)}, \cdots, V^{(m)}] \subset \mathbb{R}^{d_x \times m}$ the domain of $g^*$ for ease of presentation. Let $\lambda \in \mathbb{R}^m_+$ be the Lagrangian dual variable associated with the constraints $g^{(j)}$'s. We have the following overall composite Lagrangian function

$$\mathcal{L}(x, \lambda, \pi_{1:2}, v) = \mathcal{L}_F(x, \pi_{1:2}) + \sum_{j=1}^m \lambda_j \mathcal{L}_{g^{(j)}}(x, v^{(j)}), \quad \text{where } \lambda \geq 0.$$

Letting $\Pi = \Pi_1 \times \Pi_2$, problem (2) has an equivalent formulation as the following saddle point problem that

$$\min_{x\in\mathcal{X}}\Big(\max_{\pi_{1:2}\in\Pi}\mathcal{L}_F(x,\pi_{1:2}) + \max_{\lambda\in\mathbb{R}^m_+}\max_{v\in\mathcal{V}}\lambda^\top\mathcal{L}_g(x,v)\Big). \tag{11}$$

The existence of a saddle point to (2) and the first-order optimality condition of $(x^*,\lambda^*)$ ensure that there exist $\pi_2^*\in\partial f_2(x^*)$, $\pi_1^* = \nabla f_1(f_2(x^*))$, and $v^*\in\partial g(x^*)$ such that $(x^*,\lambda^*,\pi_{1:2}^*,v^*)$ constitutes a saddle point to problem (11) (see, for example, Proposition 1 of Zhang and Lan (2024)), i.e., for any $(x,\lambda,\pi_{1:2},v)\in\mathcal{X}\times\mathbb{R}^m_+\times\Pi\times\mathcal{V}$,

$$\mathcal{L}(x,\lambda^*,\pi_{1:2}^*,v^*)\geq\mathcal{L}(x^*,\lambda^*,\pi_{1:2}^*,v^*)\geq\mathcal{L}(x^*,\lambda,\pi_{1:2},v). \tag{12}$$

We then provide interpretations for both primal and dual update steps of Algorithm 1 under the stochastic sequential dual framework. Specifically, for the primal update, given a solution $x_t$, we consider the inner layer $f_2$ and let

$$\pi_{2,t+1}\in\operatorname*{argmax}_{\pi_2\in\Pi_2}\{\pi_2^\top x_t - f_2^*(\pi_2)\}. \tag{13}$$

Here $\pi_{2,t+1}$ is a subgradient of the *deterministic* function $f_2$ at $x_t$ that $\pi_{2,t+1}\in\partial f_2(x_t)$. Correspondingly, we have that

$$\mathcal{L}_{f_2}(x_t,\pi_{2,t+1}) = \pi_{2,t+1}^\top x_t - f_2^*(\pi_{2,t+1}) = f_2(x_t).$$

Under the stochastic setting, we let $\pi_{2,t+1}^0\in\partial f_2(x_t,\xi_{2,t}^0)$ be the *stochastic* dual variable such that

$$\mathbb{E}[\pi_{2,t+1}^0\mid x_t]\in\partial f_2(x_t).$$

Then, recall (7) that $y_{t+1} = (f_2(x_t,\xi_{2,t}^1)+\tau_t y_t)/(1+\tau_t)$. Using $y_{t+1}$ as an estimator for $f_2(x_t)$, we let

$$\pi_{1,t+1}\in\operatorname*{argmax}_{\pi_1\in\Pi_1}\Big\{\pi_1^\top y_{t+1} - f_1^*(\pi_1)\Big\},$$

which equals the gradient of $f_1$ at $y_{t+1}$ that $\pi_{1,t+1} = \nabla f_1(y_{t+1})$. We also define a stochastic dual variable $\pi_{1,t+1}^0 = \nabla f_1(y_{t+1},\xi_{1,t}^0)$ such that

$$\mathbb{E}[\pi_{1,t+1}^0\mid y_{t+1}] = \nabla f_1(y_{t+1}).$$

Meanwhile, for the constraints $g(x)$, we consider its Fenchel conjugate and let

$$\begin{aligned} &v_{t+1}\in\partial g(x_t),\quad v_{t+1}^i\in\partial g(x_t,\xi_t^i),\ \text{for } i=0,1,\\ &g(x_t) = \mathcal{L}_g(x_t,v_{t+1}),\ \text{and } g(x_t,\zeta_t^1) = \mathcal{L}_g(x_t,v_{t+1}^1). \end{aligned} \tag{14}$$

Using the above dual interpretation, the primal update step (8) is equivalent to

$$\begin{aligned} x_{t+1} &\in \operatorname*{argmin}_{x\in\mathcal{X}}\left\{\left\langle\pi_{2,t+1}^0\pi_{1,t+1}^0 + v_{t+1}^0\lambda_t, x\right\rangle + \frac{\eta_t}{2}\|x-x_t\|^2\right\}\\ &= \mathrm{Proj}_{\mathcal{X}}\left[x_t - \frac{\pi_{2,t+1}^0\pi_{1,t+1}^0 + v_{t+1}^0\lambda_t}{\eta_t}\right]. \end{aligned}$$

For the dual update step (9), using the dual interpretation (14), it is equivalent to

$$\lambda_{t+1} \in \underset{\lambda \in \mathbb{R}^m_+}{\operatorname{argmax}} \left\{ \left\langle \mathcal{L}_g\Big(x_t, v^1_{t+1}\Big), \lambda \right\rangle - \frac{\alpha_t}{2}\|\lambda_t - \lambda\|^2 \right\} = \left[\lambda_t + \frac{\mathcal{L}_g\Big(x_t, v^1_{t+1}\Big)}{\alpha_t}\right]_+. \tag{15}$$

The above dual interpretation helps us analyze the structures within the compositional objectives and expectation constraints. In particular, under the above SSD framework, we provide the convergence results of Algorithm 1 in the next subsection.

### 2.3 Convergence Analysis

After presenting Algorithm 1 and its dual interpretation, we now analyze its convergence properties. LetAfter presenting Algorithm 1 and its dual interpretation, we now analyze its convergence properties. Let $\{(x_t, \lambda_t)\}$ be the sequence generated by Algorithm 1, and let $\{(\pi_{1:2,t}, v_t)\}$ represent its corresponding dual sequence. We denote by $\bar{x}_N = \frac{1}{N}\sum_{t=1}^N x_t$. To evaluate the performance of our algorithm, we consider the following metrics: the objective optimality gap $F(\bar{x}_N) - F(x^*)$ and the feasibility residual $\|g(\bar{x}_N)_+\| = \|\max\{g(\bar{x}_N), 0\}\|$. It is clear that for any $x \in \mathcal{X}$ is an optimal solution to problem (2) if and only if $F(x) - F(x^*) = 0$ and $\|g(x)_+\| = 0$. In the remainder of this section, we demonstrate that both of these metrics converge to zero and derive their respective convergence rates.

#### 2.3.1 Convergence of gap function

Letting $z_t = (x_t, \lambda_t, \pi_{1:2,t}, v_t)$ be the solution generated by Algorithm 1 and $z = (x^*, \lambda, \pi_{1:2}, v) \in \mathcal{X} \times \mathbb{R}^m_+ \times \Pi \times \mathcal{V}$ be a general feasible point, to facilitate our analysis, we consider the *gap function*

$$Q(z_t, z) = \mathcal{L}(x_t, \lambda, \pi_{1:2}, v) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,:t}, v_t). \tag{16}$$

In what follows, we provide our first result to characterize the contraction of the above gap function.

**Theorem 1** *Suppose Assumptions 1, 2, and 3 hold and Algorithm 1 generates $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ by setting $\tau_t = t/2$, $\alpha_t = \alpha$, and $\eta_t = \eta$ for $t \le N$. Let $\lambda \in \mathbb{R}^m_+$ be a nonnegative bounded (random) variable such that $\|\lambda\| \le M_\lambda$ uniformly and let $z = (x^*, \lambda, \pi_{1:2}, v)$ be a feasible point, then for any integer $K \le N$, we have*

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}[Q(z_t, z)] + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\
&\le \frac{95KC^2_{f_1}C^2_{f_2}}{2\eta} + \frac{\eta}{2}\|x_0 - x^*\|^2 + \frac{K}{\alpha}D^2_X C^2_g + \sqrt{K}\Big(M_\lambda \sigma_g + L_{f_1}\sigma^2_{f_2} + 3C_{f_1}\sigma_{f_2}\Big) \\
&\quad + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] + \frac{K\sigma^2_g}{\alpha} + \sum_{t=1}^K 5C^2_g\Big(\frac{\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} + \frac{2M^2_\lambda}{\eta} + \frac{\mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta}\Big).
\end{aligned} \tag{17}$$

The above result serves as a crucial building block when analyzing the convergence behavior of Algorithm 1. A key observation is that the dual sequence $\{\lambda_t\}$ appears on both sides of (17),

so that it is not guaranteed to be bounded. In this case, we cannot readily establish the overall convergent property of the gap function $Q(z_t, z)$. In the next two subsections, without imposing any additional assumptions, we show that the dual sequence $\{\lambda_t\}$ is *bounded* by using (17), and further derive the convergence rates of both objective optimality gap and feasibility residual.

### 2.3.2 Boundedness of $\{\lambda_t\}$

To show the boundedness of $\{\lambda_t\}$, we observe that (17) in Theorem 1 preserves a recursive structure. Specifically, in iteration $K$, the dual solution $\lambda_K$ can be bounded by the sum of dual solutions $\{\lambda_t\}_{t=0}^{K-1}$ generated in previous iterations plus an extra constant. Here, we restate a technical result from Lemma 2.8 of Boob et al. (2023) to study the boundedness of such sequences.

**Lemma 1** *Let $\{a_t\}$ be a nonnegative sequence and $m_1, m_2 \geq 0$ be constants such that $a_0 \leq m_1$. Suppose the following relationship holds for all $K \geq 1$.*

$$a_K \leq m_1 + m_2 \sum_{t=0}^{K-1} a_t.$$

*Then we have $a_K \leq m_1(1+m_2)^K$.*

Using the above result, we then show the boundedness of $\{\lambda_t\}$ in the following proposition.

**Proposition 1** *Suppose Assumptions 1, 2, and 3 hold, and Algorithm 1 generates $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ by setting $\tau_t = \frac{t}{2}$, $\alpha_t = \alpha = 2\sqrt{N}$, and $\eta_t = \eta = \frac{15C_g^2\sqrt{N}}{2}$ for all $t \leq N$. Then for any $N \geq 2$ and any integer $K \leq N$, we have*

$$\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq 2Re^2,$$

*where*

$$\begin{aligned} R = &\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + \frac{D_X^2 C_g^2}{2} + 2\|\lambda^*\|^2 + \frac{\sigma_g^2}{2} + L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \|\lambda^*\|\sigma_g \\ &+ \frac{15C_g^2}{4}\|x_0 - x^*\|^2 + \|\lambda_0 - \lambda^*\|^2. \end{aligned} \tag{18}$$

The above result implies that, with properly chosen step-sizes, the dual sequence $\|\lambda_t\|^2$ generated by Algorithm 1 remains bounded in expectation, despite the increased randomness and bias introduced by the compositional setting. Although it is discussed in Section 2 that $\mathcal{L}(x, \lambda)$ generally loses Lipschitz continuity, the above result allows us to conclude that, in expectation, $\mathcal{L}(x, \lambda)$ is $(C_f + C_g(\sqrt{2R}e + \|\lambda^*\|))$-Lipschitz continuous in $x$ over the trajectory $\{(x_t, \lambda_t)\}_{t=1}^N$. This finding serves as a crucial building block for our subsequent convergence analysis.

### 2.3.3 Convergence of value and feasibility metrics

After proving the boundedness of $\{\lambda_t\}$, we derive the convergence rates of the objective optimality gap and the feasibility residual in an ergodic sense.

**Theorem 2** *Suppose that Assumptions 1, 2, and 3 hold, and Algorithm 1 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ *by setting* $\lambda_0 = 0$, $\tau_t = t/2$, $\alpha_t = \alpha = 2\sqrt{N}$, *and* $\eta_t = \eta = \frac{15C_g^2\sqrt{N}}{2}$ *for all* $0 \le t \le N$. *Let* $R$ *be the constant defined in* (18). *Letting* $\bar{x}_N = \sum_{t=1}^N x_t/N$, *for any* $N \ge 2$, *we have*

$$\mathbb{E}[F(\bar{x}_N)] - F(x^*) \le \frac{1}{\sqrt{N}}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \frac{15C_g^2}{4}\|x_0 - x^*\|^2 + \frac{4\|\lambda^*\|^2 + 8Re^2}{3}\Big),$$

*and*

$$\begin{aligned}\mathbb{E}[\|g(\bar{x}_N)_+\|] \le \frac{1}{\sqrt{N}}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \frac{15C_g^2}{4}\|x_0 - x^*\|^2 + \frac{D_X^2 C_g^2}{2} + \frac{\sigma_g^2}{2}\Big) \\ + \frac{(2\|\lambda^*\|^2 + 4Re^2) + 3(\|\lambda^*\| + 1)^2 + (\|\lambda^*\| + 1)\sigma_g}{\sqrt{N}}.\end{aligned}$$

It is important to note that, given any $\lambda_0 \in \mathbb{R}_+^m$, our algorithm still enjoys the $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate for both the objective optimality gap and the feasibility residual. Notably, our approach is sufficiently general to accommodate multiple expected-value constraints. It is also worth noting that the above result matches the optimal $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate for standard convex stochastic optimization, despite the complexity of the compositional objective and expected-value constraints involved. This establishes a new benchmark for stochastic compositional optimization under expected-value constraints. Finally, we emphasize that the primitive $C_g^2$ scales with the number of constraints $m$. Consequently, the hidden constant within $\mathcal{O}(\frac{1}{\sqrt{N}})$ grows *polynomially* in the number of constraints $m$. This highlights the efficiency of our algorithm in handling real-world applications with a large number of expected-value constraints.

# 3 Stochastic Compositional Optimization under Compositional Expected-Value Constraints

In Section 2, we have presented a primal-dual algorithm for expected value constrained stochastic compositional optimization that achieves the optimal rate of convergence. It is natural to ask whether our analytical framework can also handle the compositional expected value constrained applications discussed in Section 1.1. We extend the framework by considering the following two-level compositional expected value constrained stochastic compositional optimization (CoC-SCO) problem:

$$\begin{aligned}&\min_{x\in\mathcal{X}} F(x) = f_1 \circ f_2(x), \\ &\text{s.t.} \quad G(x) = g_1 \circ g_2(x) \le 0,\end{aligned} \tag{19}$$

where $\mathcal{X} \subset \mathbb{R}^{d_x}$ is convex and closed, $f_1(y) = \mathbb{E}_{\xi_1}[f_1(y, \xi_1)] : \mathbb{R}^{d_y} \mapsto \mathbb{R}$ is convex and differentiable, $f_2(x) = \mathbb{E}_{\xi_2}[f_2(x, \xi_2)] : \mathbb{R}^{d_x} \mapsto \mathbb{R}^{d_y}$ is convex, $F(x) : \mathbb{R}^{d_x} \mapsto \mathbb{R}$ is convex, $G(x) : \mathbb{R}^{d_x} \mapsto \mathbb{R}^m$ is convex, $g_1(z) = \mathbb{E}_{\zeta_1}[g_1(z, \zeta_1)] : \mathbb{R}^{d_z} \mapsto \mathbb{R}^m$ is convex and differentiable, and $g_2(x) = \mathbb{E}_{\zeta_2}[g_2(x, \zeta_2)] : \mathbb{R}^{d_x} \mapsto \mathbb{R}^{d_z}$ is convex. In more details, we write $f_2(x) = (f_2^{(1)}(x), \cdots, f_2^{(d_y)}(x))^\top \in$

$\mathbb{R}^{d_y}$, $G(x) = (G^{(1)}(x), \cdots, G^{(m)}(x))^\top \in \mathbb{R}^m$, $g_2(x) = (g_2^{(1)}(x), \cdots, g_2^{(d_z)}(x))^\top \in \mathbb{R}^{d_z}$, and $g_1(z) = (g_1^{(1)}(z), \cdots, g_1^{(m)}(z))^\top \in \mathbb{R}^m$, and assume that $f_2^{(j)}, G^{(j)}, g_2^{(j)}$, and $g_1^{(j)}$ are convex. As in the previous section, we consider that the functions $f_1, f_2, g_1$, and $g_2$ are in the form of expectations and can only be estimated from observed samples. Note that each $G^{(j)}$ represents one compositional constraint and problem (19) allows us to handle $m$ compositional constraints.

We first specify the sampling environment for CoC-SCO. In particular, we assume access to the following *Compositional Sampling Oracle* ($\mathcal{SO}_{\text{co}}$) such that

- Given $x \in \mathcal{X}$, the $\mathcal{SO}_{\text{co}}$ returns a sample value $f_2(x, \xi_2) \in \mathbb{R}^{d_y}$ and a sample subgradient
$$\partial f_2(x, \xi_2) \in \mathbb{R}^{d_x \times d_y}.$$
- Given $y \in \mathbb{R}^{d_y}$, the $\mathcal{SO}_{\text{co}}$ returns a sample value $f_1(y, \xi_1) \in \mathbb{R}$ and a sample gradient
$$\nabla f_1(y, \xi_1) \in \mathbb{R}^{d_y}.$$
- Given $x \in \mathcal{X}$, the $\mathcal{SO}_{\text{co}}$ returns a sample value $g_2(z, \zeta_2) \in \mathbb{R}^{d_z}$ and a sample subgradient
$$\partial g_2(x, \zeta_2) \in \mathbb{R}^{d_x \times d_z}.$$
- Given $z \in \mathbb{R}^{d_z}$, the $\mathcal{SO}_{\text{co}}$ returns a sample value $g_1(z, \zeta_1) \in \mathbb{R}^m$ and a sample gradient
$$\nabla g_1(z, \zeta_1) \in \mathbb{R}^{d_z \times m}.$$

Here we do not impose any smoothness assumptions on the inner-level functions $f_2$ and $g_2$. For the compositional expected value constrained problem (19), with a slight abuse of notation, we write its Lagrangian dual function as

$$\mathcal{L}(x, \lambda) = F(x) + \langle G(x), \lambda \rangle, \text{ where } \lambda \in \mathbb{R}^m_+.$$

By the convexity of $F$ and $G^{(j)}$'s, the above Lagrangian $\mathcal{L}(x, y)$ is convex in $x$ and concave in $\lambda$. We assume the Slater condition holds such that there exists a *saddle point* $(x^*, \lambda^*)$ satisfying

$$\mathcal{L}(x^*, \lambda) \le \mathcal{L}(x^*, \lambda^*) \le \mathcal{L}(x, \lambda^*), \ \forall (x, \lambda) \in \mathcal{X} \times \mathbb{R}^m_+.$$

Letting $\partial F(x) \in \mathbb{R}^{d_x}$ and $\partial G(x) \in \mathbb{R}^{d_x \times m}$ be the subgradients of $F(x)$ and $G(x)$, respectively, the first-order partial subgradients of $\mathcal{L}$ satisfy

$$\partial_x \mathcal{L}(x, \lambda) = \partial F(x) + \partial G(x)\lambda \text{ and } \nabla_\lambda \mathcal{L}(x, \lambda) = G(x), \quad \forall x \in \mathcal{X}, \lambda \in \mathbb{R}^m_+.$$

Compared with EC-SCO discussed in Section 2, the compositional expected value constrained problem is more challenging, because of the lack of unbiased estimators of $\partial G(x)$ and $G(x)$. Specifically, the unbiased estimators of $\partial G(x)$ and $G(x)$ are

$$\partial g_2(x, \zeta_2) \nabla g_1(\mathbb{E}_{\zeta_2}[g_2(x, \zeta_2)], \zeta_1) \text{ and } g_1(\mathbb{E}_{\zeta_2}[g_2(x, \zeta_2)], \zeta_1),$$

respectively. Lacking the knowledge of the expected value $\mathbb{E}_{\zeta_2}[g_2(x,\zeta_2)]$, the plug-in estimators, $\partial g_2(x,\zeta_2)\nabla g_1(g_2(x,\zeta_2),\zeta_1)$ and $g_1(g_2(x,\zeta_2),\zeta_1)$, induce biases in both primal and dual update steps. Such biases would corrupt both our primal and dual updates. For the primal update step, in addition to the bias in estimating $\partial F(x)$ discussed earlier, the biased estimator of $\partial G(x)$ induces further biases to our estimation of $\partial_x \mathcal{L}(x,\lambda)$. Furthermore, for the dual update step, the bias induced by estimating $G(x)$ would compound with the randomness within updating $\lambda_t$. Consequently, it is more challenging to analyze the boundedness of the dual sequence $\{\lambda_t\}$ and the convergence behavior of the solution sequence $\{x_t\}$ for problem (19).

To facilitate our discussion, in addition to Assumptions 1 and 3 for the objective functions $f_1$ and $f_2$, we impose the following convexity, smoothness, and boundedness assumptions for the constraint functions $g_1$ and $g_2$.

**Assumption 4** *Let $C_{g_1}, C_{g_2}, \sigma_{g_1}, \sigma_{g_2}$ and $L_{g_1}$ be positive scalars. The constraint functions $g_1$ and $g_2$ satisfy:*

*(a) $g_1, g_2$ are convex such that they could be reformulated using the following Fenchel conjugates*

$$g_1(z) = \max_{v_1 \in \mathcal{V}_1} v_1^\top z - g_1^*(v_1), \text{ and } g_2(x) = \max_{v_2 \in \mathcal{V}_2} v_2^\top x - g_2^*(v_2), \quad \forall\, x \in \mathcal{X},\, z \in \mathbb{R}^{d_z},$$

*where $\mathcal{V}_1 \subset \mathbb{R}^{d_z \times m}$ and $\mathcal{V}_2 \subset \mathbb{R}^{d_x \times d_z}$ are the domains of $g_1^*$ and $g_2^*$, respectively.*

*(b) The sample value $g_1(z,\zeta_1)$ returned by the $\mathcal{SO}_{co}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\zeta_1}[g_1(z,\zeta_1)] = g_1(z) \text{ and } \mathbb{E}_{\zeta_1}[(g_1(z,\zeta_1) - g_1(z))^2] \le \sigma_{g_1}^2, \ \forall\, z \in \mathbb{R}^{d_z}.$$

*(c) The sample value $g_2(x,\zeta_2)$ returned by the $\mathcal{SO}_{co}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\zeta_2}[g_2(x,\zeta_2)] = g_2(x) \text{ and } \mathbb{E}_{\zeta_2}[\|g_2(x,\zeta_1) - g_2(x)\|^2] \le \sigma_{g_2}^2, \ \forall\, x \in \mathcal{X}.$$

*(d) The sample subgradient $\partial g_2(x,\zeta_2)$ returned by the $\mathcal{SO}_{co}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\zeta_2}[\partial g_2(x,\zeta_2)] \in \partial g_2(x) \text{ and } \mathbb{E}_{\zeta_2}[\|\partial g_2(x,\zeta_2)\|^2] \le C_{g_2}^2, \ \forall\, x \in \mathcal{X}.$$

*(e) The sample gradient $\nabla g_1(y,\zeta_1)$ returned by the $\mathcal{SO}_{co}$ is unbiased and has a bounded second moment*

$$\mathbb{E}_{\zeta_1}[\nabla g_1(z,\zeta_1)] = \nabla g_1(z) \text{ and } \mathbb{E}_{\zeta_1}[\|\nabla g_1(z,\zeta_1)\|^2] \le C_{g_1}^2, \ \forall\, z \in \mathbb{R}^{d_z}.$$

*(f) $g_1$ is an $L_{g_1}$-smooth function with a Lipschitz continuous gradient such that*

$$\|\nabla g_1(z_1) - \nabla g_1(z_2)\| \le L_{g_1}\|z_1 - z_2\|, \quad \forall z_1, z_2 \in \mathbb{R}^{d_z}.$$

*(g) For each constraint $G^{(j)}$, its outer level function is monotone non-decreasing if its inner-level function is non-affine.*

Note that (g) above is essentially a generalization of Assumption 3 to handle the mixture of affine and non-affine inner-level compositional constraints. Specifically, we allow some constraints $G^{(j)}$ to have affine inner-level functions, while allowing others to have non-affine inner-level and monotone non-decreasing outer-level functions. Also, we only assume smoothness for the outer-level constraint $g_1$ while allowing the inner-level constraint $g_2$ to be non-smooth. We omit the subscripts $\xi_1, \xi_2, \zeta_1$, and $\zeta_2$ within $\mathbb{E}_{\xi_1}[\cdot], \mathbb{E}_{\xi_2}[\cdot], \mathbb{E}_{\zeta_1}[\cdot]$, and $\mathbb{E}_{\zeta_2}[\cdot]$ for notational convenience.

The above assumptions are mild and our framework can handle the compositional expected value constrained risk management applications provided in Section 1.1 (b) and (c). Specifically, in the risk-averse mean-deviation application (b), when each utility function $\ell_j(x, \xi)$ is concave and monotone nondecreasing in $x$, we observe that $g_2^{(j)}(x, \xi)$ is convex in $x$, and $g_1^{(j)}\Big((z, x), \xi\Big)$ is convex and monotone nondecreasing in $(z, x)$. Thus, this application can be handled after smoothing the $(\cdot)_+$ operator within the outer-layer function. For the high-moment portfolio selection application (c), given any random return $w$, $g_2(x, w)$ is an affine function while $g_1\Big((z, x), w\Big)$ is convex in $(z, x)$. Consequently, by taking expectation over the random return $w$, $\mathbb{E}_w[g_2(x, w)]$ is affine and $\mathbb{E}_w[g_1\Big((z, x), w\Big)]$ is convex in $(z, x)$.

### 3.1 Algorithm

Now we propose our algorithm to solve problem (19). In iteration $t$, we first construct an estimator $y_{t+1}$ for $\mathbb{E}_{\xi_2}[f_2(x_t, \xi_2)]$, or the limit point of $\{\mathbb{E}_{\xi_2}[f_2(x_t, \xi_2)]\}_t$ to be precise, by letting $y_{t+1} = \frac{f_2(x_t, \xi_{2,t}^1) + \tau_t y_t}{1+\tau_t}$ for some $\tau_t > 0$. We query $\mathcal{SO}_{\rm co}$ for a sample gradient $\nabla f_1(y_{t+1}, \xi_{1,t}^0)$ and a sample subgradient $\partial f_2(x_t, \xi_{2,t}^0)$. Letting $z_t$ be an estimator of $g_2(x_{t-1})$, we query $\mathcal{SO}_{\rm co}$ to obtain $g_2(x_t, \zeta_{2,t}^1)$, and update

$$z_{t+1} = \frac{g_2(x_t, \zeta_{2,t}^1) + \rho_t z_t}{1 + \rho_t},$$

where $\rho_t > 0$ is prespecified. Next, we query $\mathcal{SO}_{\rm co}$ at $x_t$ to obtain a sample subgradient $\partial g_2(x_t, \zeta_{2,t}^0)$ and query $\mathcal{SO}_{\rm co}$ at $z_{t+1}$ to obtain another sample gradient $\nabla g_1(z_{t+1}, \zeta_{1,t}^0)$. We then employ $\partial f_2(x_t, \xi_{2,t}^0)\nabla f_1(y_{t+1}, \xi_{1,t}^0)$ as an estimator of $\partial F(x_t)$, use $\partial g_2(x_t, \zeta_{2,t}^0)\nabla g_1(z_{t+1}, \zeta_{1,t}^0)$ as an estimator of $\partial G(x_t)$, and update the primal solution $x_t$ by a projected stochastic gradient step that

$$x_{t+1} = \mathrm{Proj}_{\mathcal{X}}\left[x_t - \frac{\partial f_2(x_t, \xi_{2,t}^0)\nabla f_1(y_{t+1}, \xi_{1,t}^0) + \partial g_2(x_t, \zeta_{2,t}^0)\nabla g_1(z_{t+1}, \zeta_{1,t}^0)\lambda_t}{\eta_t}\right], \tag{20}$$

where $\eta_t > 0$ is prespecified. Finally, we consider the dual variable $\lambda \in \mathbb{R}_+^m$, query $\mathcal{SO}_{\rm co}$ to obtain $g_1(z_{t+1}, \zeta_{1,t}^1)$, $\nabla g_1(z_{t+1}, \zeta_{1,t}^1)$, and $g_2(x_t, \zeta_{2,t}^2)$, and set

$$H_{t+1} = g_1(z_{t+1}, \zeta_{1,t}^1) - \nabla g_1(z_{t+1}, \zeta_{1,t}^1)z_{t+1} + \nabla g_1(z_{t+1}, \zeta_{1,t}^1)g_2(x_t, \zeta_{2,t}^2) \tag{21}$$

as an estimator of $G(x)$. We update $\lambda_{t+1}$ by a projected stochastic gradient step that

$$\lambda_{t+1} \in \operatorname*{argmax}_{\lambda \in \mathbb{R}_+^m}\left\{\lambda^\top H_{t+1} - \frac{\alpha_t}{2}\|\lambda_t - \lambda\|^2\right\} = \Big[\lambda_t + H_{t+1}/\alpha_t\Big]_+, \tag{22}$$

for some $\alpha_t > 0$. We summarize the details of the above process in Algorithm 2. We point out that given $x_t, \lambda_t, z_t$, and $y_t$, the subsequent updates $x_{t+1}, \lambda_{t+1}$ are independent, because we employ independent stochastic samples within our update scheme.

**Algorithm 2** Compositional-Constrained Stochastic Compositional Gradient Descent (CC-SCGD)

**Input:** Step-sizes $\{\alpha_k\}$, $\{\eta_k\}$, $\{\tau_k\}$, $\{\rho_k\}$, initial points $x_0 \in \mathcal{X}$, $y_0 = 0$, $z_0 = 0$, $\lambda_0 \in \mathbb{R}^m_+$, sampling oracle $\mathcal{SO}_{\text{co}}$.

**for** $t = 1, 2, ..., N$ **do**

Query the $\mathcal{SO}_{\text{co}}$ at $x_t$ for the sample values $f_2(x_t, \xi^1_{2,t}), g_2(x_t, \zeta^1_{2,t}), g_2(x_t, \zeta^2_{2,t})$ and sample subgradients $\partial f_2(x_t, \xi^0_{2,t}), \partial g_2(x_t, \zeta^0_{2,t})$.

Update

$$y_{t+1} = \frac{f_2(x_t, \xi^1_{2,t}) + \tau_t y_t}{1 + \tau_t} \text{ and } z_{t+1} = \frac{g_2(x_t, \zeta^1_{2,t}) + \rho_t z_k}{1 + \rho_t}.$$

Query $\mathcal{SO}_{\text{co}}$ at $y_{t+1}$ to obtain $\nabla f_1(y_{t+1}, \xi^0_1)$ and query $\mathcal{SO}_{\text{co}}$ at $z_{t+1}$ to obtain $\nabla g_1(z_{t+1}, \zeta^0_1)$.

Update the main solution $x_{t+1}$ by

$$x_{t+1} = \text{Proj}_{\mathcal{X}}\left[x_t - \frac{\partial f_2(x_t, \xi^0_{2,t}) \nabla f_1(y_{t+1}, \xi^0_{1,t}) + \partial g_2(x_t, \zeta^0_{2,t}) \nabla g_1(z_{t+1}, \zeta^0_{1,t}) \lambda_t}{\eta_t}\right].$$

Query $\mathcal{SO}_{\text{co}}$ at $y_{t+1}$ to obtain $g_1(z_{t+1}, \zeta^1_{1,t})$ and $\nabla g_1(z_{t+1}, \zeta^1_{1,t})$. Set $H_{t+1}$ by

$$H_{t+1} = g_1(z_{t+1}, \zeta^1_{1,t}) - \nabla g_1(z_{t+1}, \zeta^1_{1,t}) z_{t+1} + \nabla g_1(z_{t+1}, \zeta^1_{1,t}) g_2(x_t, \zeta^2_{2,t}).$$

Update the dual variable $\lambda_{t+1}$ by

$$\lambda_{t+1} = \Big[\lambda_t + H_{t+1}/\alpha_t\Big]_+.$$

**end for**

**Output :** $\bar{x}_N = \frac{1}{N}\sum_{t=1}^N x_t$.

### 3.2 Stochastic Sequential Dual Interpretation

After introducing Algorithm 2, we now present its interpretation under the SSD framework to facilitate our analysis. We first define the following composite Lagrangian functions for $f_1, f_2$:

$$\mathcal{L}_{f_2}(x, \pi_2) = \pi_2^\top x - f_2^*(\pi_2), \ \mathcal{L}_F(x, \pi_{1:2}) = \pi_1^\top \mathcal{L}_{f_2}(x, \pi_2) - f_1^*(\pi_1),$$

and define the composite Lagrangians for constraint functions $g_1, g_2$ that

$$\mathcal{L}_{g_2}(x, v_2) = v_2^\top x - g_2^*(v_2),$$
$$\mathcal{L}_G(x, v_{1:2}) = \mathcal{L}_{g_1}(x, v_1, v_2) = v_1^\top \mathcal{L}_{g_2}(x, v_2) - g_1^*(v_1).$$

We then have the overall composite Lagrangian function

$$\mathcal{L}(x, \lambda, \pi_{1:2}, v_{1:2}) = \mathcal{L}_F(x, \pi_{1:2}) + \lambda^\top \mathcal{L}_G(x, v_{1:2}).$$

Letting $\Pi = \Pi_1 \times \Pi_2$ and $\mathcal{V} = \mathcal{V}_1 \times \mathcal{V}_2$, problem (19) can be reformulated as a saddle point problem

$$\min_{x \in \mathcal{X}} \Big( \max_{\pi_{1:2} \in \Pi} \mathcal{L}_F(x, \pi_{1:2}) + \max_{\lambda \in \mathbb{R}^m_+} \max_{v_{1:2} \in \mathcal{V}} \lambda^\top \mathcal{L}_G(x, v_{1:2}) \Big). \tag{23}$$

There exist $\pi_2^* \in \partial f_2(x^*), \pi_1^* = \nabla f_1(f_2(x^*)), v_2^* \in \partial g_2(x^*)$, and $v_1^* = \nabla g_1(g_2(x^*))$ such that $(x^*, \lambda^*, \pi_{1:2}^*, v_{1:2}^*)$ serves as a saddle point to (23). Specifically, for any feasible $(x, \lambda, \pi_{1:2}, v_{1:2}) \in \mathcal{X} \times \mathbb{R}_+^m \times \Pi \times \mathcal{V}$,

$$\mathcal{L}(x, \lambda^*, \pi_{1:2}^*, v_{1:2}^*) \geq \mathcal{L}(x^*, \lambda^*, \pi_{1:2}^*, v_{1:2}^*) \geq \mathcal{L}(x^*, \lambda, \pi_{1:2}, v_{1:2}). \tag{24}$$

For any solution $x_t$, we define $\pi_{1,t}, \pi_{1,t+1}^0, \pi_{2,t}, \pi_{2,t+1}^0$ as

$$\pi_{2,t+1} \in \operatorname*{argmax}_{\pi_2 \in \Pi_2} \{\pi_2^\top x_t - f_2^*(\pi_2)\}, \;\; \pi_{2,t+1}^0 \in \partial f_2(x_t, \xi_{2,t}^0),$$
$$\pi_{1,t+1} \in \operatorname*{argmax}_{\pi_1 \in \Pi_1} \{\pi_1^\top y_{t+1} - f_1^*(\pi_1)\}, \text{ and } \pi_{1,t+1}^0 = \nabla f_1(y_{t+1}, \xi_{1,t}^0).$$

Consequently, we have

$$\mathcal{L}_{f_2}(x_t, \pi_{2,t+1}) = f_2(x_t), \;\; \mathbb{E}[\pi_{2,t+1}^0 \mid x_t] \in \partial f_2(x_t), \text{ and } \mathbb{E}[\pi_{1,t+1}^0 \mid y_{t+1}] = \nabla f_1(y_{t+1}).$$

Then for the constraints $g_1, g_2$, we define

$$v_{2,t+1} \in \operatorname*{argmax}_{v_2 \in \mathcal{V}_2} \{v_2^\top x_t - g_2^*(v_2)\}, \tag{25}$$

and observe that $v_{2,t+1}$ is the subgradient of $g_2$ at $x_t$. That is, $v_{2,t+1} \in \partial g_2(x_t)$. Correspondingly, we have

$$\mathcal{L}_{g_2}(x_t, v_{2,t+1}) = v_{2,t+1}^\top x_t - g_2^*(v_{2,t+1}) = g_2(x_t).$$

Meanwhile, we let the stochastic dual variable be $v_{2,t+1}^0 \in \partial g_2(x_t, \zeta_{2,t}^0)$, and observe that

$$\mathbb{E}[v_{2,t+1}^0 \mid x_t] \in \partial g_2(x_t).$$

Then, recall that $z_{t+1} = \frac{g_2(x_t, \zeta_{2,t}^1) + \rho_t z_t}{1+\rho_t}$, taking $z_{t+1}$ as an estimator for $g_2(x_t)$, we have that

$$v_{1,t+1} \in \operatorname*{argmax}_{v_1 \in \mathcal{V}_1} \Big\{ v_1^\top z_{t+1} - g_1^*(v_1) \Big\}$$

is the gradient of $g_1$ at $z_{t+1}$. In particular, we have

$$v_{1,t+1} = \nabla g_1(z_{t+1}).$$

We also define a stochastic dual variable $v_{1,t+1}^0 = \nabla g_1(z_{t+1}, \zeta_{1,t}^0)$ such that

$$\mathbb{E}[v_{1,t+1}^0 \mid z_{t+1}] = v_{1,t+1}.$$

After understanding the stochastic gradients under the stochastic sequential dual framework, the primal update step (20) can be viewed as

$$x_{t+1} = \operatorname*{argmin}_{x \in \mathcal{X}} \left\{ \left\langle \pi_{2,t+1}^0 \pi_{1,t+1}^0 + v_{2,t+1}^0 v_{1,t+1}^0 \lambda_t, x \right\rangle + \frac{\eta_t}{2} \|x - x_t\|^2 \right\}. \tag{26}$$

For the dual update step (22), by the following lemma, we observe that $H_{t+1}$ is an unbiased estimator of $\mathcal{L}_G(x_t, v_{1:2,t+1})$ with finite variance. This implies that (22) is equivalent to

$$\lambda_{t+1} \in \operatorname*{argmax}_{\lambda \in \mathbb{R}_+^m} \left\{ \lambda^\top H_{t+1} - \frac{\alpha_t}{2} \|\lambda_t - \lambda\|^2 \right\}, \quad \text{where} \quad \mathbb{E}[H_{t+1} | x_t, v_{1,t+1}, v_{2,t+1}] = \mathcal{L}_G(x_t, v_{1:2,t+1}).$$

**Lemma 2** *Suppose Assumptions 1, 3, and 4 hold, and Algorithm 2 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\}_{t=1}^N$. *We have*

*(a)* $H_t$ *is an unbiased estimator of* $\mathcal{L}_G(x_{t-1}, v_{1:2,t})$ *such that*

$$\mathbb{E}\Big[H_t \mid x_{t-1}, v_{1,t}, v_{2,t}\Big] = \mathcal{L}_G(x_{t-1}, v_{1:2,t}).$$

*(b) The variance of* $H_t$ *is upper bounded by a constant* $\sigma_H > 0$ *such that*

$$\mathrm{Var}\Big(H_t \mid x_{t-1}, v_{1,t}, v_{2,t}\Big) \le \sigma_H^2. \tag{27}$$

*Proof:* (a) By recalling the definition of $H_t$ (21) and noting that $g_1(z_t, \zeta_{1,t-1}^1)$, $\nabla g_1(z_t, \zeta_{1,t-1}^1)$, and $g_2(x_{t-1}, \zeta_{2,t-1}^2)$ are independently sampled, we have

$$\begin{aligned}
\mathbb{E}\Big[H_t \mid x_{t-1}, v_{1,t}, v_{2,t}\Big] &= \mathbb{E}\Big[g_1(z_t, \zeta_{1,t-1}^1) - [v_{1,t}^1]^\top z_t + [v_{1,t}^1]^\top g_2(x_{t-1}, \zeta_{2,t-1}^2) \;\Big|\; x_{t-1}, v_{1,t}, v_{2,t}\Big] \\
&= g_1(z_t) + v_{1,t}^\top g_2(x_{t-1}) - v_{1,t}^\top z_t = v_{1,t}^\top \mathcal{L}_{g_2}(x_{t-1}, v_{2,t}) - g_1^*(v_{1,t}) \\
&= \mathcal{L}_G(x_{t-1}, v_{1:2,t}),
\end{aligned}$$

where the third equality holds since $v_{1,t} = \nabla g_1(z_t)$ and $v_{2,t} \in \partial g_2(x_{t-1})$ such that $g_1(z_t) = v_{1,t}^\top z_t - g_1^*(v_{1,t})$ and $g_2(x_{t-1}) = \mathcal{L}_{g_2}(x_{t-1}, v_{2,t})$. This proves part (a).

(b) Next, given $x_{t-1}, v_{1,t}, v_{2,t}$, by the definition of $H_t$ (21) and the fact that $g_1(z_t, \zeta_{1,t-1}^1), v_{1,t}^1$, and $g_2(x_{t-1}, \zeta_{2,t-1}^2)$ are independently sampled, we have that each component of $H_t$ preserves a bounded second moment. Therefore, $H_t$ has finite variance, completing the proof. ∎

### 3.3 Convergence Analysis

We then analyze the convergence behavior of Algorithm 2. With a slight abuse notation, we denote by $(x_t, \lambda_t, \pi_{1:2,t}, v_{1:2,t})$ the solution generated by Algorithm 2 at the $t$-th iteration and denote by $(x^*, \lambda, \pi_{1:2}, v_{1:2}) \in \mathcal{X} \times \mathbb{R}_+^m \times \Pi \times \mathcal{V}$ a general feasible point. We start our analysis by decomposing the composite Lagrangian difference:

$$\begin{aligned}
&\mathcal{L}(x_t, \lambda, \pi, v) - \mathcal{L}(x^*, \lambda_t, \pi_t, v_t) \\
&= \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda^\top \Big(\mathcal{L}_G(x_t, v_{1:2}) - \mathcal{L}_G(x_t, v_{1:2,t})\Big) \\
&\quad + \lambda_t^\top \Big(\mathcal{L}_G(x_t, v_{1:2,t}) - \mathcal{L}_G(x^*, v_{1:2,t})\Big) + (\lambda - \lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t}) \\
&= \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t}) + \lambda^\top \Big(\mathcal{L}_G(x_t, v_{1:2}) - \mathcal{L}_G(x_t, v_{1:2,t})\Big) \\
&\quad + (\lambda - \lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t}) + \mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) \\
&\quad + \lambda_t^\top \Big(\mathcal{L}_G(x_t, v_{1:2,t}) - \mathcal{L}_G(x^*, v_{1:2,t})\Big).
\end{aligned} \tag{28}$$

By Lemma 2 that $H_t$ is an unbiased estimator for $\mathcal{L}_G(x_{t-1}, v_{1:2,t})$ with finite variance, we first bound the term $(\lambda - \lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t})$ in expectation in the next lemma.

**Lemma 3** *Suppose Assumptions 1, 3, and 4 hold. Suppose Algorithm 2 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\}_{t=1}^N$ *by setting* $\tau_t = \rho_t = t/2$, $\alpha_t = \alpha > 0$, *and* $\eta_t = \eta > 0$ *for* $t = 1, 2, \cdots, N$. *Let* $\lambda \in \mathbb{R}_+^m$ *be a bounded random dual variable such that* $\|\lambda\| \leq M_\lambda$ *uniformly, then there exists a constant* $\sigma_H > 0$ *such that for any* $K \leq N$,

$$\begin{aligned}
&\mathbb{E}\Big[\sum_{t=1}^K (\lambda - \lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t})\Big] + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\
&\quad \leq \sum_{t=1}^K \Big(\frac{\sigma_H^2}{\alpha} - \frac{\alpha \mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{5C_{g_1}^2 C_{g_2}^2 \mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10}\Big) \\
&\qquad + \sqrt{K} M_\lambda \sigma_H + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2].
\end{aligned}$$

Next, we show the contraction of the composite Lagrangian difference. Note that we first assume that the dual variable $\lambda$ is bounded, and we justify this assumption in Proposition 2.

**Theorem 3** *Suppose Assumptions 1, 3, and 4 hold, and Algorithm 2 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\}_{t=1}^N$ *by setting* $\tau_t = \rho_t = t/2$, $\eta_t = \eta$ *and* $\alpha_t = \alpha$ *for* $1 \leq t \leq N$. *Letting* $\lambda \in \mathbb{R}_+^m$ *be a bounded random variable such that* $\|\lambda\| \leq M_\lambda$, *then for any integer* $K \leq N$ *and any feasible point* $(x^*, \lambda, \pi_{1:2}, v_{1:2})$, *we have*

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\Big(\mathcal{L}(x_t, \lambda, \pi_{1:2}, v_{1:2}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\Big) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\
&\quad \leq \frac{40KC_{g_1}^2 C_{g_2}^2 M_\lambda^2}{\eta} + 2\sqrt{K} L_{g_1} \sigma_{g_2}^2 M_\lambda + 3\sqrt{K} C_{g_1} \sigma_{g_2} M_\lambda + \frac{\eta}{2}\|x_0 - x^*\|^2 \\
&\qquad + \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \frac{K}{\alpha}(D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] + \sqrt{K} M_\lambda \sigma_H \\
&\qquad + 2\sqrt{K} L_{f_1} \sigma_{f_2}^2 + 3\sqrt{K} C_{f_1} \sigma_{f_2} + \sum_{t=1}^K \frac{5C_{g_1}^2 C_{g_2}^2}{2\eta}\Big(\mathbb{E}[\|\lambda - \lambda_t\|^2] + \mathbb{E}[\|\lambda_{t-1}\|^2]\Big),
\end{aligned}$$

*where* $\sigma_H > 0$ *is provided in* (27).

In the rest of this section, Proposition 3.1 demonstrates the boundedness of the sequence $\{\lambda_t\}$, and while it references Theorem 3.1, the theorem itself does not assume the boundedness of $\{\lambda_t\}$. In Theorem 3.1, we construct a random variable $\lambda$ without assuming that either $\{\lambda_t\}$ or $\{\|\lambda_t - \lambda\|\}$ is bounded.

We now show that the dual variables $\{\lambda_t\}$ are bounded in the next proposition.

**Proposition 2** *Suppose Assumptions 1, 3, and 4 hold. Let* $\{\lambda_t\}_{t=1}^N$ *be the sequence of dual variables generated by Algorithm 2 with* $\tau_t = \rho_t = t/2$, $\alpha_t = 2\sqrt{N}$, *and* $\eta_t = \frac{15C_{g_1}^2 C_{g_2}^2 \sqrt{N}}{2}$ *for all* $t \leq N$. *Then for any* $N \geq 2$ *and any integer* $K \leq N$, *we have*

$$\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq 2Re^2,$$

*where* $\sigma_H > 0$ *is provided in* (27),

$$\begin{aligned}
Q &= 2L_{g_1}\sigma_{g_2}^2\|\lambda^*\| + 3C_{g_1}\sigma_{g_2}\|\lambda^*\| + \|\lambda^*\|\sigma_H + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}, \\
\text{and } R &= \frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2}^2} + 6\|\lambda^*\| + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2} + \frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \|\lambda_0 - \lambda^*\|^2 + Q.
\end{aligned} \tag{29}$$

In the above, we show that the dual sequence $\{\lambda_t\}$ generated by Algorithm 2 is also bounded in expectation, serving as a building block for analyzing the convergence behavior of the solution path $\{x_t\}$.

Letting $\bar{x}_N = \frac{1}{N}\sum_{t=1}^{N} x_t$, by the boundedness property, we derive the convergence rates for the objective optimality gap $F(\bar{x}_N) - F(x^*)$ and feasibility residual $G(\bar{x}_N)_+ = \max\{G(x), 0\}$ in the next theorem.

**Theorem 4** *Suppose Assumptions 1, 3, and 4 hold, and Algorithm 2 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\}_{t=0}^{N}$ *by setting* $\lambda_0 = 0$, $\tau_t = \rho_t = t/2$, $\alpha_t = 2\sqrt{N}$, *and* $\eta_t = \frac{15C_{g_1}^2 C_{g_2}^2 \sqrt{N}}{2}$ *for* $t = 1, \cdots, N$, *and* $N \geq 2$. *Letting* $\bar{x}_N = \frac{1}{N}\sum_{t=1}^{N} x_t$, *we have*

$$\begin{aligned}
&\mathbb{E}[F(\bar{x}_N) - F(x^*)] \\
&\leq \frac{1}{\sqrt{N}}\Big(\frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2^2}^2} + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2} + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}\Big) \\
&\quad + \frac{4}{3\sqrt{N}}\Big(\|\lambda^*\|^2 + 2Re^2\Big),
\end{aligned}$$

*and*

$$\begin{aligned}
&\mathbb{E}[\|G(\bar{x}_N)_+\|_2] \\
&\leq \frac{1}{\sqrt{N}}\Big(\frac{16}{3}\|\tilde{\lambda}\|^2 + 2L_{g_1}\sigma_{g_2}^2\|\tilde{\lambda}\| + 3C_{g_1}\sigma_{g_2}\|\tilde{\lambda}\| + \frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2}^2}\Big) \\
&\quad + \frac{1}{\sqrt{N}}\Big(\frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2} + \|\tilde{\lambda}\|^2 + \sigma_g\|\tilde{\lambda}\| + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}\Big) \\
&\quad + \frac{1}{\sqrt{N}}\Big(\|\tilde{\lambda}\| + \|\lambda^*\|^2 + 2Re^2\Big),
\end{aligned}$$

*where* $\tilde{\lambda} = (\|\lambda^*\| + 1)\frac{G(\bar{x}_N)_+}{\|G(\bar{x}_N)_+\|_2}$ *is bounded such that* $\|\tilde{\lambda}\| = \|\lambda^*\| + 1$ *and* $\|\tilde{\lambda}\|^2 = (\|\lambda^*\| + 1)^2$, *and* $\sigma_H, R > 0$ *are constants defined in* (27) *and* (29), *respectively.*

The above result indicates that Algorithm 2 achieves the optimal $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate for more complex compositional expected value constrained problems. Similar to Theorem 2, the hidden constant within the above $\mathcal{O}(\frac{1}{\sqrt{N}})$ convergence rate still grows polynomially with the number of constraints $m$, underscoring the efficiency of our algorithm in managing real-world applications that involve a large number of compositional constraints.

# 4 Numerical experiments

In this section, we investigate the empirical performance of the proposed algorithm through application to a portfolio optimization problem involving CVaR constraints. This scenario corresponds to the single-level expected value constrained problem introduced in Section 1.1.

Let $w \in \mathbb{R}^{d_x}$ denote random returns for $d_x$ assets, and let $x$ represent corresponding portfolio weights. We focus exclusively on long positions, constraining $x$ to the $(d_x - 1)$-dimensional probability

simplex $\Delta = \{x \in \mathbb{R}^d_+ : \sum_{i=1}^{d_x} x_i = 1\}$. We formulate the portfolio optimization problem with multiple CVaR constraints as follows:

$$\min_{x\in\Delta} F(x), \quad \text{s.t. } \mathrm{CVaR}_{\delta_i}(x) \le \gamma_i, \quad \forall i = 1, \dots, m, \tag{30}$$

where $\delta_i$ are CVaR constraint levels, $\gamma_i$ are their corresponding upper bounds, the objective function is given by $F(x) = \mathbb{E}_w[-w^\top x] + c\mathbb{E}_w\left[(w^\top x - \mathbb{E}[w^\top x])^4\right]$, and the CVaR of the portfolio's negative returns is defined as $\mathrm{CVaR}_{\delta_i}(x) = \mathrm{CVaR}_{\delta_i}(-w^\top x)$.

We now reformulate problem (30) into the standard structure from Section 2. Define the following:

$$\begin{aligned} f_1(y, w) = f_1(x, z, w) &= -z + \xi(w^\top x - z)^4 \in \mathbb{R}^1, \\ f_2(x, \boldsymbol{u}, w) &= [x, w^\top x] \in \mathbb{R}^{d+1}, \\ g_i(x, u_i, w) &= u_i + \frac{1}{1-\delta_i}[-w^\top x - u_i]_+ - \gamma_i \in \mathbb{R}^1, \end{aligned}$$

where $z \in \mathbb{R}$, $\boldsymbol{u} \in \mathbb{R}^m$, and $y = (x, z)$. Let the expected functions be:

$$f_1(x, z) = \mathbb{E}_w[f_1(x, z, w)], \quad f_2(x, \boldsymbol{u}) = \mathbb{E}_w[f_2(x, \boldsymbol{u}, w)], \quad g_i(x, u_i) = \mathbb{E}_w[g_i(x, u_i, w)].$$

Then, problem (30) takes the form:

$$\min_{x\in\Delta, u\in\mathbb{R}} f_1 \circ f_2(x, u), \quad \text{s.t. } g_i(x, u_i) \le 0, \quad \forall i = 1, \dots, m.$$

**Single CVaR constraint.** First, we analyze the single CVaR constraint scenario. Consider $w$ as a multivariate normal random vector with mean $\mu$ and covariance $\Sigma$. Random instances are generated with $\mu \sim \mathrm{Unif}[-1, 1]^{d_x}$, using two covariance structures: independent case ($\Sigma = I_{d_x}$) and positively correlated Toeplitz matrix with entries $\Sigma_{ij} = 0.5^{|i-j|}$. Parameters are set as follows: risk-aversion parameter $c = 0.5$, CVaR level $\delta = 0.95$, and upper bound $\gamma = 0.6\mathrm{CVaR}_0 + 0.4\mathrm{CVaR}_{\min}$, where $\mathrm{CVaR}_0$ corresponds to the optimal portfolio without CVaR constraint and $\mathrm{CVaR}_{\min}$ is the minimum achievable CVaR. Algorithm 1 is implemented with step-sizes $\alpha_t = \max\{20d_x, 0.02d_x\sqrt{t}\}$, $\eta_t = 300\sqrt{t}$, $\tau_t = 0.02t$. We consider dimensions $d_x = 10, 1000$ for $10^7$ iterations with 10 repetitions each. Results depicted in Figure 1 show the trajectories of optimality gaps $F(x^*) - F(x_t)$ and feasibility residuals $(\mathrm{CVaR}_\delta(x_t) - \gamma)_+$, averaged over 10 runs, with 95% confidence intervals shaded. Observations confirm theoretical convergence rates from Theorem 2, showing slope close to $-1/2$, and feasibility residual quickly approaching zero.

**Multiple CVaR constraints.** We extend the analysis to multiple CVaR constraints, maintaining identical settings for noise distributions and objective function. The upper bounds are set as $\gamma_i = 0.6\mathrm{CVaR}_{0_i} + 0.4\mathrm{CVaR}_{\min_i}$, with CVaR levels $\delta_i = 0.01, 0.02, 0.05, 0.1, 0.2$, where $\mathrm{CVaR}_{0_i}$ is the CVaR at level $\delta_i$ of the optimal portfolio without the CVaR constraints, and $\mathrm{CVaR}_{\min_i}$ is the smallest achievable CVaR at level $\delta_i$. Algorithm 1 parameters remain unchanged, with dimension $d_x = 10$ and $10^7$ iterations over 10 repetitions. Results presented in Figure 2 show the trajectories of





Figure 1: Empirical convergence of EC-SCGD algorithm for single CVaR constraint. Optimality gap and feasibility residual averaged over 10 runs with shaded 95% confidence intervals.

optimality gaps $F(x^*) - F(x_t)$ and feasibility residuals $\|(\mathrm{CVaR}_\delta(x_t) - \gamma)_+\|_2$, averaged over 10 runs, with 95% confidence intervals shaded. We observe that the slope of the trajectory of the optimality gap closely matches $-1/2$, confirming the theoretical convergence rate established in Theorem 2. Additionally, the feasibility residual converges quickly to zero.





Figure 2: Empirical convergence of EC-SCGD for multiple CVaR constraints. Optimality gap and feasibility residual averaged over 10 runs, with shaded 95% confidence intervals.

# 5 Conclusion

In this paper, we develop a novel model for stochastic compositional optimization (SCO) with single-level and two-level compositional expected-value constraints. Our formulation is handles multiple expected-value constraints and has broad applications in data-driven optimization and risk management. We propose a class of primal-dual algorithms, which we interpret within the stochastic sequential dual framework. For convex objectives and constraints, we demonstrate that our algorithms achieve the optimal convergence rate of $\mathcal{O}(\frac{1}{\sqrt{N}})$ in both single-level and two-level compositional expected value constrained scenarios. These results match the optimal convergence rate for standard convex stochastic optimization and set new benchmarks.

**Acknowledgement.** Shuoguang Yang received support from the Hong Kong Research Grants Council [ECS Grant 26209422]. Wei You received support from the Hong Kong Research Grants Council [ECS Grant 26212320 and GRF Grant 16212823].

# References


Shabbir Ahmed, Ulaş Çakmak, and Alexander Shapiro. Coherent risk measures in inventory problems. *European Journal of Operational Research*, 182(1):226–238, 2007.

Krishnakumar Balasubramanian, Saeed Ghadimi, and Anthony Nguyen. Stochastic multilevel composition optimization algorithms with level-independent convergence rates. *SIAM Journal on Optimization*, 32 (2):519–544, 2022.

Amir Beck. *First-order methods in optimization*. SIAM, 2017.

Dimitri P Bertsekas. Nonlinear programming. *Journal of the Operational Research Society*, 48(3):334–334, 1997.

Digvijay Boob, Qi Deng, and Guanghui Lan. Stochastic first-order methods for convex and nonconvex functional constrained optimization. *Mathematical Programming*, 197(1):215–279, 2023.

Sergio Bruno, Shabbir Ahmed, Alexander Shapiro, and Alexandre Street. Risk neutral and risk averse approaches to multistage renewable investment planning under uncertainty. *European Journal of Operational Research*, 250(3):979–989, 2016.

Tianyi Chen, Yuejiao Sun, and Wotao Yin. Solving stochastic compositional optimization is nearly as easy as solving stochastic optimization. *IEEE Transactions on Signal Processing*, 69:4937–4948, 2021.

Shawn Cole, Xavier Giné, and James Vickery. How does risk management influence production decisions? evidence from a field experiment. *The Review of Financial Studies*, 30(6):1935–1970, 2017.

Y. Ermoliev. Stochastic quasigradient methods. In Y. Ermoliev and R. J.-B. Wets, editors, *Numerical Techniques for Stochastic Optimization*, pages 141–186. Springer-Verlag, New York, 1988.

Chelsea Finn, Pieter Abbeel, and Sergey Levine. Model-agnostic meta-learning for fast adaptation of deep networks. In *International conference on machine learning*, pages 1126–1135. PMLR, 2017.

Chelsea Finn, Aravind Rajeswaran, Sham Kakade, and Sergey Levine. Online meta-learning. In Kamalika Chaudhuri and Ruslan Salakhutdinov, editors, *Proceedings of the 36th International Conference on Machine Learning*, volume 97 of *Proceedings of Machine Learning Research*, pages 1920–1930. PMLR, 09–15 Jun 2019.

Rong Ge, Furong Huang, Chi Jin, and Yang Yuan. Escaping from saddle points—online stochastic gradient for tensor decomposition. In Peter Grünwald, Elad Hazan, and Satyen Kale, editors, *Proceedings of The 28th Conference on Learning Theory*, volume 40 of *Proceedings of Machine Learning Research*, pages 797–842, Paris, France, 03–06 Jul 2015. PMLR.

Saeed Ghadimi, Andrzej Ruszczynski, and Mengdi Wang. A single timescale stochastic approximation method for nested stochastic optimization. *SIAM Journal on Optimization*, 30(1):960–979, 2020.

Willem K Klein Haneveld and Maarten H Van Der Vlerk. Integrated chance constraints: reduced forms and an algorithm. *Computational Management Science*, 3:245–269, 2006.

Campbell R Harvey, John C Liechty, Merrill W Liechty, and Peter Müller. Portfolio selection with higher moments. *Quantitative Finance*, 10(5):469–485, 2010.

Guanghui Lan. *First-order and Stochastic Optimization Methods for Machine Learning*. Springer Nature, 2020.

Guanghui Lan and Zhiqiang Zhou. Algorithms for stochastic optimization with function or expectation constraints. *Computational Optimization and Applications*, 76(2):461–498, 2020.

Guanghui Lan, Edwin Romeijn, and Zhiqiang Zhou. Conditional gradient methods for convex optimization with general affine and nonlinear constraints. *SIAM Journal on Optimization*, 31(3):2307–2339, 2021.

Claude Lemaréchal, Arkadii Nemirovskii, and Yurii Nesterov. New variants of bundle methods. *Mathematical Programming*, 69(1):111–147, 1995.

Qihang Lin, Selvaprabu Nadarajah, Negar Soheili, and Tianbao Yang. A data efficient and feasible level set method for stochastic convex optimization with expectation constraints. *Journal of Machine Learning Research*, 21(1), January 2020. ISSN 1532-4435.

Huikang Liu, Xiaolu Wang, and Anthony Man-Cho So. Adaptive coordinate sampling for stochastic primal–dual optimization. *International Transactions in Operational Research*, 29(1):24–47, 2022a.

Junyi Liu, Ying Cui, and Jong-Shi Pang. Solving nonsmooth and nonconvex compound stochastic programs with applications to risk measure minimization. *Mathematics of Operations Research*, 47(4):3051–3083, 2022b.

Avinash N. Madavan and Subhonmesh Bose. A stochastic primal-dual method for optimization with conditional value at risk constraints. *Journal of Optimization Theory and Applications*, 190(2):428–460, August 2021.

Arkadi Nemirovski, Anatoli Juditsky, Guanghui Lan, and Alexander Shapiro. Robust stochastic approximation approach to stochastic programming. *SIAM Journal on Optimization*, 19(4):1574–1609, 2009.

R Tyrrell Rockafellar, Stanislav Uryasev, et al. Optimization of conditional value-at-risk. *Journal of Risk*, 2: 21–42, 2000.

A. Ruszczyński and A. Shapiro. Optimization of convex risk functions. *Mathematics of Operations Research*, 31(3):433–452, 2006.

Andrzej Ruszczynski. A stochastic subgradient method for nonsmooth nonconvex multilevel composition optimization. *SIAM Journal on Control and Optimization*, 59(3):2301–2320, 2021.

Morton Slater. Lagrange multipliers revisited. In *Traces and Emergence of Nonlinear Programming*, pages 293–306. Springer, 2014.

Anthony Man-Cho So, Jiawei Zhang, and Yinyu Ye. Stochastic combinatorial optimization with controllable risk aversion level. *Mathematics of Operations Research*, 34(3):522–537, 2009.

Conghui Tan, Tong Zhang, Shiqian Ma, and Ji Liu. Stochastic primal-dual method for empirical risk minimization with O(1) per-iteration complexity. *Advances in Neural Information Processing Systems*, 31, 2018.

M. Wang and J. Liu. A stochastic compositional gradient method using markov samples. In *Proceedings of the 2016 Winter Simulation Conference*, pages 702–713. IEEE Press, 2016.

M. Wang, E. X. Fang, and H. Liu. Stochastic compositional gradient descent: Algorithms for minimizing compositions of expected-value functions. *Mathematical Programming*, 161(1-2):419–449, 2017a.

M. Wang, J. Liu, and E. X. Fang. Accelerating stochastic composition optimization. *The Journal of Machine Learning Research*, 18(1):3721–3743, 2017b.

Yan Yan, Yi Xu, Qihang Lin, Lijun Zhang, and Tianbao Yang. Stochastic primal-dual algorithms with faster convergence than $o(1/\sqrt{T})$ for problems without bilinear structure. *arXiv preprint arXiv:1904.10112*, 2019.

S. Yang, M. Wang, and E. X. Fang. Multilevel stochastic gradient methods for nested composition optimization. *SIAM Journal on Optimization*, 29(1):616–659, 2019.

Shuoguang Yang, Xudong Li, and Guanghui Lan. Data-driven minimax optimization with expectation constraints. *Operations Research*, 2024.

Hao Yu, Michael Neely, and Xiaohan Wei. Online convex optimization with stochastic constraints. *Advances in Neural Information Processing Systems*, 30, 2017.

Muhammad Bilal Zafar, Isabel Valera, Manuel Gomez-Rodriguez, and Krishna P Gummadi. Fairness constraints: A flexible approach for fair classification. *The Journal of Machine Learning Research*, 20 (1):2737–2778, 2019.

Junyu Zhang and Lin Xiao. Multilevel composite stochastic optimization via nested variance reduction. *SIAM Journal on Optimization*, 31(2):1131–1157, 2021.

Zhe Zhang and Guanghui Lan. Optimal methods for convex nested stochastic composite optimization. *Mathematical Programming*, pages 1–48, 2024.

Zhe Zhang, Shabbir Ahmed, and Guanghui Lan. Efficient algorithms for distributionally robust stochastic optimization with discrete scenario support. *SIAM Journal on Optimization*, 31(3):1690–1721, 2021.

# Appendix

## A Proof of Results in Section 2

Letting Assumptions 1, 2, and 3 hold and letting $(x^*, \lambda, \pi_{1:2}, v) \in \mathcal{X} \times \mathbb{R}^m_+ \times \Pi \times \mathcal{V}$ be a general feasible point, we first introduce two technical lemmas to facilitate our analysis.

**Lemma 4 (Lemma 3.8 of Lan 2020)** *Assume function $g$ is $\mu$-strongly convex with respect to some Bergman distance $V$, i.e., $g(y) - g(\bar{y}) - g'(\bar{y})^\top(y - \bar{y}) \geq \mu V(y, \bar{y})$. If $\hat{y} \in \operatorname{argmin}_{y \in Y}\{\pi^\top y + g(y) + \tau V(y, \bar{y})\}$, then*

$$(\hat{y} - y)^\top \pi + g(\hat{y}) - g(y) \leq \tau V(y, \bar{y}) - (\tau + \mu) V(y, \hat{y}) - \tau V(\hat{y}, \bar{y}).$$

### A.1 Proof of Theorem 1

*Proof:* Recall that $\mathcal{L}(x, \pi_{1:2}, v, \lambda) = \mathcal{L}_F(x, \pi_{1:2}) + \lambda^\top \mathcal{L}_g(x, v)$ for any $(x, \lambda, \pi_{1:2}, v) \in \mathcal{X} \times \mathbb{R}^m_+ \times \Pi \times \mathcal{V}$, we consider $Q(z_t, z)$ and decompose it as

$$\begin{aligned}
Q(z_t, z) &= \mathcal{L}(x_t, \lambda, \pi_{1:2}, v) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t) \\
&= \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda^\top \mathcal{L}_g(x_t, v) - \lambda_t^\top \mathcal{L}_g(x^*, v_t) \\
&= \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda^\top \Big( \mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t) \Big) \\
&\quad + \lambda_t^\top \Big( \mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t) \Big) + (\lambda - \lambda_t)^\top \mathcal{L}_g(x_t, v_t) \\
&= \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t}) + \mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) \\
&\quad + \lambda^\top \Big( \mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t) \Big) + \lambda_t^\top \Big( \mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t) \Big) + (\lambda - \lambda_t)^\top \mathcal{L}_g(x_t, v_t).
\end{aligned} \tag{31}$$

We then provide bounds for the terms above in expectation. First, by Lemma 6 in Appendix Section A.2, we have

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\Big[ \mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t}) \Big] \\
&\quad \leq \frac{40 K C_{f_1}^2 C_{f_2}^2}{\eta_0} + \sum_{t=1}^K \frac{\eta_{t-1}}{10} \mathbb{E}[\|x_t - x_{t-1}\|^2] \quad + \sqrt{K} L_{f_1} \sigma_{f_2}^2 + 3\sqrt{K} C_{f_1} \sigma_{f_2}.
\end{aligned} \tag{32}$$

Secondly, by Lemma 7 in Appendix Section A.3, we obtain that

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\Big[ \mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big( \mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t) \Big) \Big] \\
&\quad \leq \frac{\eta}{2} \|x_0 - x^*\|^2 + \sum_{t=1}^K \Big( \frac{5 C_{f_1}^2 C_{f_2}^2}{2\eta} - \frac{3\eta}{10} \mathbb{E}[\|x_{t-1} - x_t\|^2] \Big) \\
&\qquad + \sum_{t=1}^K \Big( \frac{\alpha}{4} \mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \frac{D_X^2 C_g^2}{\alpha} + \frac{5 \mathbb{E}[\|\lambda_{t-1}\|^2] C_g^2}{2\eta} \Big).
\end{aligned} \tag{33}$$

Thirdly, by Lemma 8 in Appendix Section A.4 and the condition that $\|\lambda\| \leq M_\lambda$, we obtain

$$\mathbb{E}\Big[\sum_{t=1}^{K} \lambda^\top \Big(\mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t)\Big)\Big] \leq \sum_{t=1}^{K} \frac{10C_g^2 M_\lambda^2}{\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10}. \tag{34}$$

Finally, by Lemma 9 in Appendix Section A.5, we obtain that

$$\begin{aligned}
&\mathbb{E}\Big[\sum_{t=1}^{K} (\lambda - \lambda_t)^\top \mathcal{L}_g(x_t, v_t)\Big] + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\
&\quad \leq \sqrt{K} M_\lambda \sigma_g + \sum_{t=1}^{K} \Big(\frac{5C_g^2 \mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta} + \frac{\eta}{10}\mathbb{E}[\|x_{t-1} - x_t\|^2] - \frac{\alpha}{4}\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]\Big) \\
&\qquad + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] + \frac{K\sigma_g^2}{\alpha}.
\end{aligned} \tag{35}$$

Summing (31) over $t = 1, 2, \cdots, K$, taking expectations, and combining (32), (33), (34), and (35), we conclude that

$$\begin{aligned}
&\sum_{t=1}^{K} \mathbb{E}\Big(\mathcal{L}(x_t, \lambda, \pi_{1:2}, v) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\
&\leq \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \sqrt{K} L_{f_1} \sigma_{f_2}^2 + 3\sqrt{K} C_{f_1} \sigma_{f_2} + \frac{\eta}{2}\|x_0 - x^*\|^2 + \frac{K}{\alpha} D_X^2 C_g^2 + \sqrt{K} M_\lambda \sigma_g \\
&\quad + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] + \frac{K\sigma_g^2}{\alpha} + \sum_{t=1}^{K} \Big(\frac{5C_g^2 \mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} + \frac{10C_g^2 M_\lambda^2}{\eta} + \frac{5C_g^2 \mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta}\Big),
\end{aligned}$$

which completes the proof. ∎

**Lemma 5 (Lemma 2 of Zhang and Lan 2024)** *Let $\mathcal{F}_t$ be a filtration and $\{\delta_t\}_{t=1}^N$ be a martingale noise sequence such that $\delta_j \in \mathcal{F}_{t+1}$ for $j = 1, 2, \cdots, t$, $\mathbb{E}[\delta_t | \mathcal{F}_t] = 0$, and $\mathbb{E}[\|\delta_t\|^2 | \mathcal{F}_t] \leq \sigma^2$. For any random variable $\pi \in \Pi$ correlated with $\{\delta_t\}_{t=1}^N$, suppose it is bounded such that $\|\pi\| \leq M_\pi$ uniformly, then*

$$\mathbb{E}[\sum_{t=1}^{N} \hat{\pi}^\top \delta_t] \leq \sqrt{N} M_\pi \sigma.$$

## A.2 Lemma 6 and Its Proof

**Lemma 6** *Suppose Assumptions 1, 2 and 3 hold, and Algorithm 1 generates $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ by setting $\tau_t = t/2$ and $\eta_t = \eta$ for $t = 0, 1, \cdots, N$. Then for any integer $K \leq N$, we have*

$$\begin{aligned}
&\sum_{t=1}^{K} \mathbb{E}\Big[\mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t})\Big] \\
&\quad \leq \sum_{t=1}^{K} \frac{\eta_{t-1}}{10}\mathbb{E}[\|x_t - x_{t-1}\|^2] + \frac{40KC_{f_1}^2 C_{f_2}^2}{\eta} + 2\sqrt{K} L_{f_1} \sigma_{f_2}^2 + 3\sqrt{K} C_{f_1} \sigma_{f_2}.
\end{aligned}$$

*Proof:* This result can be derived by combining Propositions 5 and 6 of Zhang and Lan (2024). Here we provide the proof sketch. By decomposing $\mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t})$, we have

$$\begin{aligned}
&\mathcal{L}_F(x_t, \pi_{1:2}) - \mathcal{L}_F(x_t, \pi_{1:2,t}) \\
&= \mathcal{L}_{f_1}(x_t, \pi_1, \pi_2) - \mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t}) + \mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t}) - \mathcal{L}_{f_1}(x_t, \pi_{1,t}, \pi_{2,t}).
\end{aligned}$$

We provide bound for each term after the above decomposition. First, we consider $\mathcal{L}_{f_1}(x_t, \pi_1, \pi_2) - \mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t})$ and obtain

$$\begin{aligned}
\sum_{t=1}^K \mathbb{E}\Big(\mathcal{L}_{f_1}(x_t, \pi_1, \pi_2) - \mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t})\Big) &= \sum_{t=1}^K \mathbb{E}[\Big\langle \mathcal{L}_{f_2}(x_t, \pi_2) - \mathcal{L}_{f_2}(x_t, \pi_{2,t}), \pi_1 \Big\rangle] \\
&= \sum_{t=1}^K \mathbb{E}\Big[ \Big\langle (\pi_2 - \pi_{2,t})^\top x_t - f_2^*(\pi_2) + f_2^*(\pi_{2,t}), \pi_1 \Big\rangle \Big] \\
&= \sum_{t=1}^K \mathbb{E}\Big[ \Big\langle (\pi_2 - \pi_{2,t})^\top x_{t-1} - f_2^*(\pi_2) + f_2^*(\pi_{2,t}), \pi_1 \Big\rangle \Big] \\
&\quad + \sum_{t=1}^K \mathbb{E}\Big[ \Big\langle (\pi_2 - \pi_{2,t})^\top (x_t - x_{t-1}), \pi_1 \Big\rangle \Big],
\end{aligned}$$

By recalling (13) that $\pi_{2,t} \in \operatorname{argmax}_{\pi_2 \in \Pi_2}\{\langle \pi_2, x_{t-1}\rangle - f_2^*(\pi_2)\}$ and using Assumptions 3 that $\pi_1 \geq 0$ for non-affine $f_2$, we obtain for both affine ($\pi_2$ is a constant) and non-affine $f_2$,

$$\mathbb{E}\Big[ \Big\langle (\pi_2 - \pi_{2,t})^\top x_{t-1} - f_2^*(\pi_2) + f_2^*(\pi_{2,t}), \pi_1 \Big\rangle \Big] \leq 0,$$

which further implies that

$$\begin{aligned}
\sum_{t=1}^K \mathbb{E}\Big(\mathcal{L}_{f_1}(x_t, \pi_1, \pi_2) - \mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t})\Big) &\leq \sum_{t=1}^K \mathbb{E}[\langle (\pi_2 - \pi_{2,t})\pi_1, x_t - x_{t-1}\rangle] \\
&\leq \sum_{t=1}^K \Big[ \frac{5\mathbb{E}[\|(\pi_2 - \pi_{2,t})\pi_1\|^2]}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2] \Big] \\
&\leq \sum_{t=1}^K \Big[ \frac{20 C_{f_1}^2 C_{f_2}^2}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2] \Big].
\end{aligned} \tag{36}$$

The next part is similar to Proposition 6 of Zhang and Lan (2024). Consider the following decomposition:

$$\begin{aligned}
\mathbb{E}[\mathcal{L}_{f_1}(x_t, \pi_1, \pi_{2,t}) - \mathcal{L}_{f_1}(x_t, \pi_{1,t}, \pi_{2,t})] &= \mathbb{E}[\langle \pi_1 - \pi_{1,t}, f_2(x_t)\rangle - \{f_1^*(\pi_1) - f_1^*(\pi_{1,t})\}] \\
&= \mathbb{E}[\Big\langle \pi_1 - \pi_{1,t}, f_2(x_{t-1}, \xi_{2,t}^1)\Big\rangle - \{f_1^*(\pi_1) - f_1^*(\pi_{1,t})\}] \\
&\quad + \mathbb{E}[\Big\langle \pi_{1,t} - \pi_{1,t-1}, f_2(x_{t-1}) - f_2(x_{t-1}, \xi_{2,t}^1)\Big\rangle] + \mathbb{E}[\langle \pi_1 - \pi_{1,t}, f_2(x_t) - f_2(x_{t-1})\rangle] \\
&\quad + \mathbb{E}[\Big\langle \pi_1, f_2(x_{t-1}) - f_2(x_{t-1}, \xi_{2,t}^t)\Big\rangle] + \underbrace{\mathbb{E}[\Big\langle \pi_{1,t-1}, f_2(x_{t-1}) - f_2(x_{t-1}, \xi_{2,t}^1)\Big\rangle]}_{=0}.
\end{aligned} \tag{37}$$

Among these terms, the update rule of $\pi_{1,t}$, the strong convexity of $f_1^*$ with respect to $D_{f_1^*}$, and the choice of $\tau_t$ imply

$$\begin{aligned}
&\mathbb{E}[\sum_{t=1}^{K}\Big\langle \pi_1 - \pi_{1,t}, f_2(x_{t-1},\xi_{2,t}^1)\Big\rangle - \{f_1^*(\pi_1) - f_1^*(\pi_{1,t})\}] \\
&\quad\le \mathbb{E}[\sum_{t=1}^{K}\tau_t D_{f_1^*}(\pi_1;\pi_{1,t-1}) - (\tau_t+1)D_{f_1^*}(\pi_1;\pi_{1,t-1}) - \tau_t D_{f_1^*}(\pi_{1,t};\pi_{1,t-1})\}] \\
&\quad\le -\mathbb{E}[\sum_{t=1}^{K}\tau_t D_{f_1^*}(\pi_{1,t};\pi_{1,t-1})\}].
\end{aligned}$$

The $1/L_{f_1}$-strong convexity of $D_{f_1^*}$, the choice of $\tau_t$ and the Cauchy-Schwartz inequality imply

$$\begin{aligned}
&\mathbb{E}\sum_{t=1}^{K}\Big[(\pi_{1,t}-\pi_{1,t-1})(f_2(x_{t-1},\xi_{2,t}^1) - f_2(x_{t-1})) - \tau_t \mathcal{D}_{f_1^*}(\pi_{1,t};\pi_{1,t-1})\Big] \\
&\quad\le \sum_{t=2}^{K}\frac{L_{f_1}\sigma_{f_2}^2}{\tau_t} + \sqrt{\mathbb{E}[\|\pi_{1,1}-\pi_{1,0}\|^2]}\sqrt{\mathbb{E}[\Big\|f_2(x_0,\xi_{2,0}^1) - f_2(x_0)\Big\|^2]} \\
&\quad\le 2L_{f_1}\sigma_{f_2}^2\log K + 2C_{f_1}\sigma_{f_2} \\
&\quad\le 2\sqrt{K}L_{f_1}\sigma_{f_2}^2 + 2\sqrt{K}C_{f_1}\sigma_{f_2}.
\end{aligned}$$

Lemma 2 of Zhang and Lan (2024) implies that

$$\mathbb{E}\left[\sum_{t=1}^{K}\Big\langle \pi_1, f_2(x_{t-1}) - f_2(x_{t-1},\xi_{2,t}^t)\Big\rangle\right] \le \sqrt{K}C_{f_1}\sigma_{f_2}.$$

Similar to (36), we also have

$$\mathbb{E}[\sum_{t=1}^{K}\langle \pi_1 - \pi_{1,t}, f_2(x_t) - f_2(x_{t-1})\rangle] \le \sum_{t=1}^{K}\left[\frac{20C_{f_1}^2C_{f_2}^2}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2]\right].$$

Combining them with the decomposition in (37), we obtain

$$\begin{aligned}
&\sum_{t=1}^{K}\mathbb{E}\Big(\mathcal{L}_{f_1}(x_t,\pi_1,\pi_{2,t}) - \mathcal{L}_{f_1}(x_t,\pi_{1,t},\pi_{2,t})\Big) \\
&\quad\le \sum_{t=1}^{K}\left[\frac{20C_{f_1}^2C_{f_2}^2}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2]\right] + 2\sqrt{K}L_{f_1}\sigma_{f_2}^2 + 3\sqrt{K}C_{f_1}\sigma_{f_2}.
\end{aligned}$$

The desired result then follows from adding to preceding inequality to (36). ∎

### A.3 Lemma 7 and Its Proof

**Lemma 7** *Suppose Assumptions 1, 2, and 3 hold, and Algorithm 1 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ *by setting* $\tau_t = t/2$, $\alpha_t = \alpha$, *and* $\eta_t = \eta$ *for all* $0 \le t \le N$. *We have*

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\Big[\mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big(\mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t)\Big)\Big] \\
&\quad \le \frac{\eta}{2}\mathbb{E}[\|x_{t-1} - x^*\|^2] - \sum_{t=1}^K \frac{3\eta}{10}\mathbb{E}\Big[\|x_{t-1} - x_t\|^2\Big] + \sum_{t=1}^K \frac{5C_{f_1}^2 C_{f_2}^2}{2\eta} \\
&\qquad + \sum_{t=1}^K \frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \sum_{t=1}^K \frac{1}{\alpha} D_X^2 C_g^2 + \sum_{t=1}^K \frac{5\mathbb{E}[\|\lambda_{t-1}\|^2]C_g^2}{2\eta}].
\end{aligned}$$

*Proof.* Recall Algorithm 1, for $t = 1, 2, \cdots, K$, we note that $\pi_{2,t}\pi_{1,t}, v_t\lambda_t \in \mathbb{R}^{d_x}$ and obtain

$$\begin{aligned}
&\mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big(\mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t)\Big) \\
&\quad = \langle \pi_{2,t}\pi_{1,t}, x_t - x^* \rangle + \langle v_t\lambda_t, x_t - x^* \rangle \\
&\quad = \Big\langle \pi_{2,t}^0\pi_{1,t}^0, x_t - x^* \Big\rangle + \Big\langle v_t^0\lambda_{t-1}, x_t - x^* \Big\rangle \\
&\qquad + \Big\langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0, x_t - x^* \Big\rangle + \Big\langle v_t\lambda_t - v_t^0\lambda_{t-1}, x_t - x^* \Big\rangle,
\end{aligned}$$

and

$$x_t = \operatorname*{argmin}_{x\in\mathcal{X}} \Big\{ \Big\langle \pi_{2,t}^0\pi_{1,t}^0 + v_t^0\lambda_{t-1}, x \Big\rangle + \frac{\eta_{t-1}}{2}\|x_{t-1} - x\|^2 \Big\}.$$

By applying Lemma 4 to the above update rule, since $x^* \in \mathcal{X}$, we have

$$\Big\langle \pi_{2,t}^0\pi_{1,t}^0 + v_t^0\lambda_{t-1}, x_t - x^* \Big\rangle \le \frac{\eta}{2}\|x_{t-1} - x^*\|^2 - \frac{\eta}{2}\|x_{t-1} - x_t\|^2 - \frac{\eta}{2}\|x_t - x^*\|^2,$$

which further yields

$$\begin{aligned}
&\mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big(\mathcal{L}_g(x_t, v_t) - \mathcal{L}_g(x^*, v_t)\Big) \\
&\quad \le \frac{\eta}{2}\|x_{t-1} - x^*\|^2 - \frac{\eta}{2}\|x_{t-1} - x_t\|^2 - \frac{\eta}{2}\|x_t - x^*\| \\
&\qquad + \underbrace{\Big\langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0, x_t - x^* \Big\rangle}_{\Delta_0^t} + \underbrace{\Big\langle v_t\lambda_t - v_t^0\lambda_{t-1}, x_t - x^* \Big\rangle}_{\Lambda_0^t}.
\end{aligned} \tag{38}$$

Then, we decompose $\Delta_0^t$ by

$$\Delta_0^t = \Big(\pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0\Big)(x_{t-1} - x^*) + \Big(\pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0\Big)(x_t - x_{t-1}).$$

We note that given $x_{t-1}, y_t$,

$$\mathbb{E}\Big[\pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0 \Big| x_{t-1}, y_t\Big] = 0,$$

and

$$\begin{aligned}
&\mathbb{E}\Big[\|\pi_{2,t}\pi_{1,t} - \pi_{2,t}^0\pi_{1,t}^0\|^2 | x_{t-1}, y_t\Big] \\
&\quad \le \mathbb{E}\Big[\|\pi_{1,t} - \pi_{1,t}^0\|^2 \Big| x_{t-1}, y_t\Big]\mathbb{E}\Big[\|\pi_{2,t}^0\|^2 \Big| x_{t-1}\Big] + \mathbb{E}\Big[\|(\pi_{2,t}^0 - \pi_{2,t})\pi_{1,t}\|^2 | x_{t-1}, y_t\Big] \\
&\quad \le 2C_{f_1}^2 C_{f_2}^2.
\end{aligned}$$

By the independence between $(x_{t-1}-x^*)$ and the mean-zero term $(\pi_{2,t}\pi_{1,t}-\pi^0_{2,t}\pi^0_{1,t})$, we have $\mathbb{E}\Big[\Big(\pi_{2,t}\pi_{1,t}-\pi^0_{2,t}\pi^0_{1,t}\Big)^\top(x_{t-1}-x^*)\Big]=0$. Moreover, to handle the possible correlation between $(x_t-x_{t-1})$ and $(\pi_{2,t}\pi_{1,t}-\pi^0_{2,t}\pi^0_{1,t})$, we utilize the fact that $ab\le\frac{5a^2}{2}+\frac{b^2}{10}$ for all $a,b\in\mathbb{R}$, and obtain

$$\begin{aligned}\mathbb{E}[\sum_{t=1}^K\Delta_0^t]&=\mathbb{E}\Big[\sum_{t=1}^K\Big\langle\pi_{2,t}\pi_{1,t}-\pi^0_{2,t}\pi^0_{1,t},x_t-x_{t-1}\Big\rangle\Big]\\&\le\sum_{t=1}^K\Big(\frac{5C_{f_1}^2C_{f_2}^2}{2\eta}+\frac{\eta\mathbb{E}[\|x_t-x_{t-1}\|^2]}{10}\Big).\end{aligned}\tag{39}$$

We then decompose $\Lambda_0^t$ as

$$\Lambda_0^t=\underbrace{\langle v_t(\lambda_t-\lambda_{t-1}),x_t-x^*\rangle}_{\Lambda_1^t}+\underbrace{\Big\langle(v_t-v_t^0)\lambda_{t-1},x_t-x^*\Big\rangle}_{\Lambda_2^t}.$$

By the fact that $2ab\le a^2+b^2$ and Assumptions 1 and 2 that $\|x_t-x^*\|\le D_X$ and $\|v_t\|\le C_g$, for any $\alpha>0$, we have

$$\begin{aligned}\mathbb{E}[\Lambda_1^t]&\le\frac{\alpha}{4}\mathbb{E}[\|\lambda_t-\lambda_{t-1}\|^2]+\frac{1}{\alpha}\mathbb{E}[\|v_t^\top(x_t-x^*)\|^2]\\&\le\frac{\alpha}{4}\mathbb{E}[\|\lambda_t-\lambda_{t-1}\|^2]+\frac{1}{\alpha}D_X^2C_g^2.\end{aligned}\tag{40}$$

Meanwhile, since $\mathbb{E}[v_t-v_t^0]=0$ and $\mathbb{E}[\|v_t-v_t^0\|^2]\le C_g^2$, we obtain

$$\begin{aligned}\mathbb{E}[\sum_{t=1}^K\Lambda_2^t]&=\mathbb{E}\Big[\sum_{t=1}^K\Big\langle(v_t-v_t^0)\lambda_{t-1},x_{t-1}-x^*\Big\rangle\Big]+\mathbb{E}\Big[\sum_{t=1}^K\Big\langle(v_t-v_t^0)\lambda_{t-1},x_t-x_{t-1}\Big\rangle\Big]\\&=\sum_{t=1}^K\mathbb{E}\Big[\Big\langle(v_t-v_t^0)\lambda_{t-1},x_t-x_{t-1}\Big\rangle\Big]\\&\le\sum_{t=1}^K\mathbb{E}\Big[\frac{5\|(v_t-v_t^0)\lambda_{t-1}\|^2}{2\eta}\Big]+\sum_{t=1}^K\frac{\eta\mathbb{E}[\|x_t-x_{t-1}\|^2]}{10}\\&\le\sum_{t=1}^K\frac{5C_g^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta}]+\sum_{t=1}^K\frac{\eta\mathbb{E}[\|x_t-x_{t-1}\|^2]}{10},\end{aligned}\tag{41}$$

where the second equality holds by the independence between $(v_t-v_t^0)\lambda_{t-1}$ and $(x_{t-1}-x^*)$, and the second last inequality holds due to the fact that $ab\le\frac{5a^2}{2}+\frac{b^2}{10}$. Taking expectations on both sides of (38), summing over $t=1,\cdots,N$, and substituting (39), (40), and (41) into it, we obtain

$$\begin{aligned}&\sum_{t=1}^K\mathbb{E}\Big[\mathcal{L}_F(x_t,\pi_{1:2,t})-\mathcal{L}_F(x^*,\pi_{1:2,t})+\lambda_t^\top\Big(\mathcal{L}_g(x_t,v_t)-\mathcal{L}_g(x^*,v_t)\Big)\Big]\\&\le\sum_{t=1}^K\mathbb{E}\Big[\frac{\eta}{2}\|x_t-x^*\|^2-\frac{\eta}{2}\|x_t-x_{t+1}\|^2-\frac{\eta}{2}\|x_{t+1}-x^*\|+\Delta_0^t+\Lambda_1^t+\Lambda_2^t\Big]\\&\le\frac{\eta}{2}\mathbb{E}[\|x_{t-1}-x^*\|^2]-\sum_{t=1}^K\frac{3\eta}{10}\mathbb{E}\Big[\|x_{t-1}-x_t\|^2\Big]+\sum_{t=1}^K\frac{5C_{f_1}^2C_{f_2}^2}{2\eta_{t-1}}\\&\quad+\sum_{t=1}^K\frac{\alpha}{4}\mathbb{E}[\|\lambda_t-\lambda_{t-1}\|^2]+\sum_{t=1}^K\frac{D_X^2C_g^2}{\alpha}+\sum_{t=1}^K\frac{5\mathbb{E}[\|\lambda_{t-1}\|^2]C_g^2}{2\eta}],\end{aligned}$$

where the last inequality holds by the fact that $\eta_t = \eta$ for $t = 1, 2, \cdots, N$. This completes the proof. ∎

### A.4 Lemma 8 and Its Proof

**Lemma 8** *Suppose Assumptions 1, 2 and 3 hold, and Algorithm 1 generates* $\{(x_t, \lambda_t, \pi_t, v_t)\}_{t=1}^N$ *by setting* $\tau_t = t/2$, $\alpha_t = \alpha$, *and* $\eta_t = \eta$ *for all* $0 \le t \le N$. *Let* $\lambda \in \mathbb{R}_+^m$ *be a nonnegative random variable whose norm is bounded such that* $\|\lambda\| \le M_\lambda$ *uniformly. Then for any* $K \le N$, *we have*

$$\mathbb{E}\Big[\sum_{t=1}^K \lambda^\top \Big(\mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t)\Big)\Big] \le \frac{10 M_\lambda^2 C_g^2}{\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10}.$$

*Proof.* Recall that the update rule is that

$$v_t \in \operatorname*{argmax}_{v \in \mathcal{V}} \Big\{ \langle v, x_{t-1} \rangle - g^*(v) \Big\}.$$

By the fact that $\lambda \ge 0$, we have

$$\begin{aligned}
&\lambda^\top \Big(\mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t)\Big) \\
&= \lambda^\top \Big((v - v_t)^\top x_{t-1} - (g^*(v) - g^*(v_t)) + (v - v_t)^\top (x_t - x_{t-1})\Big) \\
&\le \lambda^\top \Big((v - v_t)^\top (x_t - x_{t-1})\Big).
\end{aligned}$$

By taking expectations on both sides of the above inequality, we have

$$\begin{aligned}
\mathbb{E}\left[\lambda^\top \Big(\mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t)\Big)\right] &\le \mathbb{E}\Big[\frac{5\|(v - v_t)\lambda\|^2}{2\eta} + \frac{\eta \|x_t - x_{t-1}\|^2}{10}\Big] \\
&\le \frac{5 M_\lambda^2 \mathbb{E}[\|v - v_t\|^2]}{2\eta_{t-1}} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10} \le \frac{10 C_g^2 M_\lambda^2}{\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10},
\end{aligned}$$

where the first inequality uses the fact that $ab \le \frac{5a^2}{2} + \frac{b^2}{10}$, the second inequality holds by the condition that $\|\lambda\| \le M_\lambda$, and the third inequality holds by Assumption 2 that $\|v - v_t\|^2 \le 2\|v\|^2 + 2\|v_t\|^2 \le 4C_g^2$. Summing the above inequality over $t = 1, \cdots, K$, we obtain

$$\mathbb{E}\Big[\sum_{t=1}^K \lambda^\top \Big(\mathcal{L}_g(x_t, v) - \mathcal{L}_g(x_t, v_t)\Big)\Big] \le \frac{10 C_g^2 M_\lambda^2}{\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10},$$

which completes the proof. ∎

### A.5 Lemma 9 and Its Proof

**Lemma 9** *Suppose Assumptions 1, 2, and 3 hold, and Algorithm 1 generates* $\{(x_t, \lambda_t, \pi_{1:2,t}, v_t)\}_{t=1}^N$ *by setting* $\tau_t = t/2$, $\alpha_t = \alpha$, *and* $\eta_t = \eta$ *for all* $0 \le t \le N$. *Let* $\lambda \in \mathbb{R}_+^m$ *be a bounded nonnegative*

*random variable such that* $\|\lambda\| \leq M_\lambda$. *For any integer* $K \leq N$, *we have*

$$\begin{aligned}
&\mathbb{E}\Big[\sum_{t=1}^{K}(\lambda-\lambda_t)^\top \mathcal{L}_g(x_t,v_t)\Big] + \frac{\alpha_{K-1}}{2}\mathbb{E}[\|\lambda_K-\lambda\|^2] \\
&\quad \leq \sqrt{K}M_\lambda\sigma_g + \sum_{t=1}^{K}\left(\frac{5C_g^2\mathbb{E}[\|\lambda-\lambda_t\|^2]}{2\eta} + \frac{\eta}{10}\mathbb{E}[\|x_{t-1}-x_t\|^2] - \frac{\alpha}{4}\mathbb{E}[\|\lambda_{t-1}-\lambda_t\|^2]\right) \\
&\qquad + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0-\lambda\|^2] + \frac{K\sigma_g^2}{\alpha}.
\end{aligned}$$

*Proof:* By decomposing $(\lambda-\lambda_t)^\top \mathcal{L}_g(x_t,v_t)$, we have,

$$\begin{aligned}
&(\lambda-\lambda_t)^\top \mathcal{L}_g(x_t,v_t) \\
&\quad = (\lambda-\lambda_t)^\top\Big(\mathcal{L}_g(x_t,v_t)-\mathcal{L}_g(x_{t-1},v_t)\Big) + (\lambda-\lambda_t)^\top[\mathcal{L}_g(x_{t-1},v_t)-g(x_{t-1},\zeta_{t-1}^1)] \\
&\qquad + (\lambda-\lambda_t)g(x_{t-1},\zeta_{t-1}^1) \\
&= \underbrace{\langle v_t(\lambda-\lambda_t), x_{t-1}-x_t\rangle}_{T_{1,t}} + \underbrace{(\lambda-\lambda_t)^\top[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)]}_{T_{2,t}} \\
&\qquad + \underbrace{(\lambda-\lambda_t)^\top g(x_{t-1},\zeta_{t-1}^1)}_{T_{3,t}}.
\end{aligned} \tag{42}$$

We provide bounds for $T_{1,t}, T_{2,t}$, and $T_{3,t}$ in our analysis.

First, for $T_{1,t}$, by using the fact that $\langle a,b\rangle \leq \frac{5\|a\|^2}{2} + \frac{\|b\|^2}{10}$ and Assumption 2 that $\|v_t\|^2 \leq C_g^2$, we obtain

$$\begin{aligned}
\mathbb{E}[\langle v_t(\lambda-\lambda_t), x_{t-1}-x_t\rangle] &\leq \frac{5\mathbb{E}[\|v_t(\lambda-\lambda_t)\|^2]}{2\eta} + \frac{\eta\mathbb{E}[\|x_{t-1}-x_t\|^2]}{10} \\
&\leq \frac{5C_g^2\mathbb{E}[\|\lambda-\lambda_t\|^2]}{2\eta} + \frac{\eta\mathbb{E}[\|x_{t-1}-x_t\|^2]}{10}.
\end{aligned} \tag{43}$$

Second, consider $T_{2,t}$, by the independence between $\lambda_{t-1}$ and the fact that $\mathbb{E}[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)] = 0$, we have

$$\mathbb{E}\Big[\lambda_{t-1}^\top(g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1))\Big] = 0,$$

which further implies that

$$\begin{aligned}
\mathbb{E}[T_{2,t}] &= \mathbb{E}\Big[(\lambda-\lambda_{t-1})^\top[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)]\Big] \\
&\quad + \mathbb{E}\Big[(\lambda_{t-1}-\lambda_t)^\top[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)]\Big] \\
&= \mathbb{E}\Big[\lambda^\top[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)]\Big] + \mathbb{E}\Big[(\lambda_{t-1}-\lambda_t)^\top[g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)]\Big].
\end{aligned} \tag{44}$$

Recall that $\lambda$ is random but bounded such that $\|\lambda\| \leq M_\lambda$, by Lemma 5 and Assumption 2 that $\mathbb{E}[\|g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)\|^2] \leq \sigma_g^2$, we obtain

$$\mathbb{E}\Big[\sum_{t=1}^{K}\lambda^\top\Big(g(x_{t-1})-g(x_{t-1},\zeta_{t-1}^1)\Big)\Big] \leq \sqrt{K}M_\lambda\sigma_g. \tag{45}$$

Meanwhile, by the fact that $\langle a, b\rangle \le \frac{\alpha\|a\|^2}{4} + \frac{\|b\|^2}{\alpha}$ for $\alpha > 0$, we bound the second term within the right side of (44) by

$$\begin{aligned}&\mathbb{E}\Big[(\lambda_{t-1} - \lambda_t)^\top [g(x_{t-1}) - g(x_{t-1}, \zeta_{t-1}^1)]\Big] \\ &\le \frac{\alpha\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{\mathbb{E}[\|g(x_{t-1}) - g(x_{t-1}, \zeta_{t-1}^1)\|^2]}{\alpha} \le \frac{\alpha\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{\sigma_g^2}{\alpha}.\end{aligned} \tag{46}$$

Summing (44) over $t = 1, \cdots, K$ and applying (45) and (46), we obtain

$$\sum_{t=1}^K \mathbb{E}[T_{2,t}] \le \sqrt{K} M_\lambda \sigma_g + \sum_{t=1}^K \Big(\frac{\alpha\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{\sigma_g^2}{\alpha}\Big). \tag{47}$$

Third, consider $T_{3,t}$, by recalling (15) that

$$\lambda_t \in \operatorname*{argmax}_{\lambda \in \mathbb{R}_+^m} \Big\{ \Big\langle g(x_{t-1}, \zeta_{t-1}^1), \lambda \Big\rangle - \frac{\alpha}{2}\|\lambda_t - \lambda\|^2 \Big\},$$

and by Lemma 4, we have

$$\begin{aligned}T_{3,t} &= (\lambda - \lambda_t)^\top g(x_{t-1}, \zeta_{t-1}^1) = -(\lambda_t - \lambda)^\top g(x_{t-1}, \zeta_{t-1}^1) \\ &\le \frac{\alpha}{2}\|\lambda_{t-1} - \lambda\|^2 - \frac{\alpha}{2}\|\lambda_{t-1} - \lambda_t\|^2 - \frac{\alpha}{2}\|\lambda_t - \lambda\|^2.\end{aligned} \tag{48}$$

Finally, summing (42) over $t = 1, \cdots, N$, taking expectations on both sides, and plugging in (43), (47), and (48), we conclude that

$$\begin{aligned}&\mathbb{E}\Big[\sum_{t=1}^K (\lambda - \lambda_t)^\top \mathcal{L}_g(x_t, v_t)\Big] + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] \\ &\le \sqrt{K} M_\lambda \sigma_g + \sum_{t=1}^K \Big(\frac{5C_g^2\mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta} + \frac{\eta}{10}\mathbb{E}[\|x_{t-1} - x_t\|^2] - \frac{\alpha}{4}\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]\Big) \\ &\quad + \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] + \frac{K\sigma_g^2}{\alpha},\end{aligned}$$

which completes the proof. ∎

### A.6 Proof of Proposition 1

By applying Theorem 1 with $\lambda = \lambda^*$, $\pi_{1:2} = \pi_{1:2}^*$, and $v = v^*$ defined in (12), we set $M_{\lambda^*} = \|\lambda^*\|$ and obtain

$$\begin{aligned}&\sum_{t=1}^K \mathbb{E}\Big(\mathcal{L}(x_t, \lambda^*, \pi_{1:2}^*, v^*) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\ &\le \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \sqrt{K} L_{f_1} \sigma_{f_2}^2 + 3\sqrt{K} C_{f_1}\sigma_{f_2} + \frac{\eta}{2}\|x_0 - x^*\|^2 \\ &\quad + \frac{KD_X^2 C_g^2}{\alpha} + \frac{10KC_g^2\|\lambda^*\|^2}{\eta} + \sum_{t=1}^K \frac{5C_g^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} + \sqrt{K}\|\lambda^*\|\sigma_g \\ &\quad + \frac{\alpha}{2}\|\lambda_0 - \lambda^*\|^2 + \frac{K\sigma_g^2}{\alpha} + \sum_{t=1}^K \frac{5C_g^2\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{2\eta}.\end{aligned}$$

By using the fact that $\|\lambda_t\|^2 \leq 2\|\lambda_t - \lambda^*\|^2 + 2\|\lambda^*\|^2$, we further express the above inequality as

$$\begin{aligned}
&\sum_{t=1}^{K} \mathbb{E}\Big(\mathcal{L}(x_t, \lambda^*, \pi^*_{1:2}, v^*) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\
&\quad \leq \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \sqrt{K}L_{f_1}\sigma_{f_2}^2 + 3\sqrt{K}C_{f_1}\sigma_{f_2} + \frac{\eta}{2}\|x_0 - x^*\|^2 \\
&\qquad + \frac{KD_X^2 C_g^2}{\alpha} + \sum_{t=1}^{K} \frac{15C_g^2\|\lambda^*\|^2}{\eta} + \sum_{t=1}^{K} \frac{5C_g^2\mathbb{E}[\|\lambda_{t-1} - \lambda^*\|^2]}{\eta} + \sqrt{K}\|\lambda^*\|\sigma_g \\
&\qquad + \frac{\alpha}{2}\|\lambda_0 - \lambda^*\|^2 + \frac{K\sigma_g^2}{\alpha} + \sum_{t=1}^{K} \frac{5C_g^2\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{2\eta}.
\end{aligned}$$

By using the min-max relationship (12) that

$$\mathcal{L}(x_t, \lambda^*, \pi^*_{1:2}, v^*) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1,t}) \geq 0,$$

dividing both sides of the above inequality by $\frac{\alpha}{2}$, and setting $\alpha = 2\sqrt{N}$ and $\eta = \frac{15C_g^2\sqrt{N}}{2}$, we obtain

$$\begin{aligned}
\mathbb{E}[\|\lambda_K - \lambda^*\|^2] &\leq \frac{2}{\alpha}\sum_{t=1}^{K} \mathbb{E}\Big(\mathcal{L}(x_t, \lambda^*, \pi^*_{1:2}, v^*) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) + \mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\
&\leq R_K + \frac{\mathbb{E}[\|\lambda^* - \lambda_K\|^2]}{3N} + \sum_{t=0}^{K-1} \frac{\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{N},
\end{aligned}$$

where

$$\begin{aligned}
R_K = \frac{K}{N}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + \frac{D_X^2 C_g^2}{2} + 2\|\lambda^*\|^2 + \frac{\sigma_g^2}{2}\Big) + \frac{15C_g^2}{4}\|x_0 - x^*\|^2 + \|\lambda_0 - \lambda^*\|^2 \\
+ \frac{\sqrt{K}}{\sqrt{N}}\Big(L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \|\lambda^*\|\sigma_g\Big).
\end{aligned}$$

Next, letting $R$ be the constant defined in (18), we note that $R_K \leq R$, and the above inequality further implies that

$$\Big(1 - \frac{1}{3N}\Big)\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq R + \sum_{t=0}^{K-1} \frac{\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{N}.$$

Dividing both sides of the above inequality by $(1 - \frac{1}{3N})$, using the fact that $(1 - \frac{1}{3N}) \geq 1/2$ for $N \geq 2$, and applying Lemma 1, we conclude that

$$\begin{aligned}
\mathbb{E}[\|\lambda_K - \lambda^*\|^2] &\leq \Big(1 - \frac{1}{3N}\Big)^{-1}\Big(R + \sum_{t=0}^{K-1} \frac{\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{N}\Big) \\
&\leq 2R + 2\sum_{t=0}^{K-1} \frac{\mathbb{E}[\|\lambda^* - \lambda_t\|^2]}{N} \leq 2R\Big(1 + \frac{2}{N}\Big)^K \leq 2Re^2,
\end{aligned}$$

for all $1 \leq K \leq N$. This completes the proof. ∎

## A.7 Proof of Theorem 2

*Proof:* In our analysis, we set $\lambda_0 = 0$ for ease of presentation. Recall that $\bar{x}_N = \frac{1}{N}\sum_{t=1}^N x_t$, for any fixed $(\lambda, \pi_{1:2}, v)$, by the convexity of $\mathcal{L}(x, \lambda, \pi_{1:2}, v)$ with respect to $x$, we have

$$\frac{1}{N}\sum_{t=1}^N \mathcal{L}(x_t, \lambda, \pi_{1:2}, v) \geq \mathcal{L}(\bar{x}_N, \lambda, \pi_{1:2}, v). \tag{49}$$

Meanwhile, we denote by

$$\hat{\pi}_2 \in \partial f_2(\bar{x}_N),\ \hat{\pi}_1 = \nabla f_1\Big(f_2(\bar{x}_N)\Big),\ \text{and}\ \ \hat{v} \in \partial g(\bar{x}_N), \tag{50}$$

the dual variables associated with $\bar{x}_N$. By the definition of composite Lagrangian (10), we have

$$F(\bar{x}_N) = \mathcal{L}_F(\bar{x}_N, \hat{\pi}_{1:2}) = \mathcal{L}(\bar{x}_N, 0, \hat{\pi}_{1:2}, \hat{v}). \tag{51}$$

First, we derive the convergence rate of the objective optimality gap $F(\bar{x}_N) - F(x^*)$. Let $\bar{\lambda}_N = \frac{1}{N}\sum_{t=1}^N \lambda_t$, $\bar{\pi}_{1:2,N} = \frac{1}{N}\sum_{t=1}^N \pi_{1:2,t}$, and $\bar{v}_N = \frac{1}{N}\sum_{t=1}^N v_t$. By setting $\lambda = 0$, $\pi_{1:2} = \hat{\pi}_{1:2}$, and $v = \hat{v}$ within (50), and using (12) that $\mathcal{L}(x^*, \lambda^*, \pi^*_{1:2}, v^*) \geq \mathcal{L}(x^*, \lambda, \pi_{1:2}, v)$ for any $(\lambda, \pi_{1:2}, v) \in \mathbb{R}^m_+ \times \Pi \times \mathcal{V}$, we have

$$\begin{aligned}
&\frac{1}{N}\sum_{t=1}^N \Big(\mathcal{L}(x_t, 0, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) \\
&\qquad \geq \mathcal{L}(\bar{x}_N, 0, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda^*, \pi^*_{1:2}, v^*) = F(\bar{x}_N) - F(x^*),
\end{aligned} \tag{52}$$

where the last equality holds by (51). Therefore, by (52), setting the feasible point as $(x^*, \lambda, \pi_{1:2}, v) = (x^*, 0, \hat{\pi}_{1:2}, \hat{v})$, and applying Theorem 1 with the fact $\|\lambda\| = 0$ when $\lambda = 0$, we have

$$\begin{aligned}
\mathbb{E}[F(\bar{x}_N)] - F(x^*) &\leq \frac{1}{N}\sum_{t=1}^N \mathbb{E}\Big(\mathcal{L}(x_t, 0, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t)\Big) \\
&\leq \frac{1}{\sqrt{N}}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \frac{15C_g^2}{4}\|x_0 - x^*\|^2\Big) \\
&\quad + \frac{1}{N}\sum_{t=1}^N \Big(\frac{\mathbb{E}[\|\lambda_{t-1}\|^2]}{3\sqrt{N}} + \frac{\mathbb{E}[\|\lambda_t\|^2]}{3\sqrt{N}}\Big) \\
&\leq \frac{1}{\sqrt{N}}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2} + \frac{15C_g^2}{4}\|x_0 - x^*\|^2\Big) + \frac{(4\|\lambda^*\|^2 + 8Re^2)}{3\sqrt{N}},
\end{aligned}$$

where the last inequality comes from the bound of $\|\lambda_t\|^2$ provided by Proposition 1 that

$$\mathbb{E}[\|\lambda_t\|^2] \leq 2\|\lambda^*\|^2 + 2\mathbb{E}[\|\lambda_t - \lambda^*\|^2] \leq 2\|\lambda^*\|^2 + 4Re^2,\ \text{for all}\ t = 1, 2, \cdots, N. \tag{53}$$

This establishes the convergence rate for the objective optimality gap $F(\bar{x}_N) - F(x^*)$.

Second, we consider the feasibility residual $\|g(\bar{x}_N)_+\|_2$. Recall that $\hat{\pi}_{1:2}$ and $\hat{v}$ are the dual variables associated with $\bar{x}_N$ such that $\hat{\pi}_{1:2} \in \operatorname{argmax}_{\pi_{1,2}\in\Pi} \mathcal{L}_F(\bar{x}_N, \pi_{1:2})$ and $\hat{v} \in \operatorname{argmax}_{v\in\mathcal{V}} \mathcal{L}_g(\bar{x}_N, v)$, we have

$$F(\bar{x}_N) = \mathcal{L}_F(\bar{x}_N, \hat{\pi}_{1:2}) \geq \mathcal{L}_F(\bar{x}_N, \pi^*_{1:2}) \text{ and } g(\bar{x}_N) = \mathcal{L}_g(\bar{x}_N, \hat{v}) \geq \mathcal{L}_g(\bar{x}_N, v^*).$$

Since $\lambda^* \geq 0$, by the above inequalities and the min-max relationship (12), we can see

$$\begin{aligned}
&\mathcal{L}(\bar{x}_N, \lambda^*, \hat{\pi}_{1:2}, \hat{v}) = \mathcal{L}_F(\bar{x}_N, \hat{\pi}) + \langle \lambda^*, \mathcal{L}_g(\bar{x}_N, \hat{v}) \rangle \\
&\quad \geq \mathcal{L}_F(\bar{x}_N, \pi^*_{1:2}) + \langle \lambda^*, \mathcal{L}_g(\bar{x}_N, v^*) \rangle = \mathcal{L}(\bar{x}_N, \lambda^*, \pi^*_{1:2}, v^*) \geq \mathcal{L}(x^*, \lambda^*, \pi^*_{1:2}, v^*),
\end{aligned}$$

which further implies

$$F(\bar{x}_N) + \langle \lambda^*, g(\bar{x}_N) \rangle - F(x^*) \geq 0.$$

Meanwhile, due to the facts that $\lambda^* \geq 0$ and $g(\bar{x}_N) \leq g(\bar{x}_N)_+$, we have $\langle \lambda^*, g(\bar{x}_N) \rangle \leq \langle \lambda^*, g(\bar{x}_N)_+ \rangle$, hence,

$$F(\bar{x}_N) + \|\lambda^*\| \|g(\bar{x}_N)_+\| - F(x^*) \geq F(\bar{x}_N) + \langle \lambda^*, g(\bar{x}_N) \rangle - F(x^*) \geq 0. \tag{54}$$

Let $\tilde{\lambda} = (\|\lambda^*\|_2 + 1) g(\bar{x}_N)_+ / \|g(\bar{x}_N)_+\|$. Consider another feasible point $(x^*, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v})$, by (49) and the facts that $g(\bar{x}_N)^\top g(\bar{x}_N)_+ = \|g(\bar{x}_N)_+\|^2$ and $\mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t) \leq \mathcal{L}(x^*, \lambda^*, \pi^*_{1:2}, v^*)$, we have $\langle g(\bar{x}_N), \tilde{\lambda} \rangle = (\|\lambda^*\|_2 + 1) \|g(\bar{x}_N)_+\|$, which further yields that

$$\begin{aligned}
&\frac{1}{N} \sum_{t=1}^{N} \Big( \mathcal{L}(x_t, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t) \Big) \geq \mathcal{L}(\bar{x}_N, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda^*, \pi^*_{1:2}, v^*) \\
&\quad = F(\bar{x}_N) + \langle \tilde{\lambda}, g(\bar{x}_N) \rangle - F(x^*) = F(\bar{x}_N) + (\|\lambda^*\|_2 + 1) \|g(\bar{x}_N)_+\| - F(x^*).
\end{aligned}$$

By rearranging the terms in the inequality above and applying (54), we obtain

$$\begin{aligned}
\|g(\bar{x}_N)_+\| &\leq \frac{1}{N} \sum_{t=1}^{N} \Big( \mathcal{L}(x_t, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}) - (\mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t) \Big) \\
&\quad - \Big( F(\bar{x}_N) + \|\lambda^*\| \|g(\bar{x}_N)_+\| - F(x^*) \Big) \\
&\leq \frac{1}{N} \sum_{t=1}^{N} \Big( \mathcal{L}(x_t, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_t) \Big).
\end{aligned}$$

Next, we note that $\|\tilde{\lambda}\| = \|\lambda^*\| + 1$. By the inequality above, considering the feasible point $(x^*, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v})$, and applying Theorem 1 with $\lambda_0 = 0$, $M_{\tilde{\lambda}} = \|\tilde{\lambda}\| = \|\lambda^*\| + 1$, $\alpha = 2\sqrt{N}$, and $\eta = \frac{15 C_g^2 \sqrt{N}}{2}$, we have

$$\begin{aligned}
&\mathbb{E}[\|g(\bar{x}_N)_+\|] \\
&\leq \frac{95 C_{f_1}^2 C_{f_2}^2}{2\eta} + \frac{L_{f_1} \sigma_{f_2}^2}{\sqrt{N}} + \frac{3 C_{f_1} \sigma_{f_2}}{\sqrt{N}} + \frac{\eta}{2N} \|x_0 - x^*\|^2 + \frac{D_X^2 C_g^2}{\alpha} + \frac{\|\tilde{\lambda}\| \sigma_g}{\sqrt{N}} \\
&\quad + \frac{1}{N} \Big[ \frac{\alpha}{2} \|\tilde{\lambda}\|^2 + \frac{N \sigma_g^2}{\alpha} + \sum_{t=1}^{N} \Big( \frac{5 C_g^2 \mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} + \frac{10 C_g^2 \mathbb{E}[\|\tilde{\lambda}\|^2]}{\eta} + \frac{5 C_g^2 \mathbb{E}[\|\tilde{\lambda} - \lambda_t\|^2]}{2\eta} \Big) \Big] \\
&\leq \frac{1}{\sqrt{N}} \Big( \frac{7 C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1} \sigma_{f_2}^2 + 3 C_{f_1} \sigma_{f_2} + \frac{15}{4} \|x_0 - x^*\|^2 + \frac{D_X^2 C_g^2}{2} + \|\tilde{\lambda}\| \sigma_g + \|\tilde{\lambda}\|^2 \Big) \\
&\quad + \frac{\sigma_g^2}{2\sqrt{N}} + \frac{1}{N} \sum_{t=1}^{N} \Big( \frac{\mathbb{E}[\|\lambda_{t-1}\|^2]}{3\sqrt{N}} + \frac{4\|\tilde{\lambda}\|^2}{3\sqrt{N}} + \frac{2(\|\tilde{\lambda}\|^2 + \mathbb{E}[\|\lambda_t\|^2])}{3\sqrt{N}} \Big) \\
&\leq \frac{1}{\sqrt{N}} \Big( \frac{7 C_{f_1}^2 C_{f_2}^2}{C_g^2} + L_{f_1} \sigma_{f_2}^2 + 3 C_{f_1} \sigma_{f_2} + \frac{15 C_g^2}{4} \|x_0 - x^*\|^2 + \frac{D_X^2 C_g^2}{2} + \frac{\sigma_g^2}{2} \Big) \\
&\quad + \frac{1}{\sqrt{N}} \Big( (2\|\lambda^*\|^2 + 4Re^2) + 3\|\tilde{\lambda}\|^2 + \|\tilde{\lambda}\| \sigma_g \Big),
\end{aligned}$$

where the second inequality holds because $\|\tilde{\lambda}-\lambda_t\|^2 \le 2\|\tilde{\lambda}\|^2+2\|\lambda_t\|^2$, and the last inequality holds by the bound of $\|\lambda_t\|^2$ in (53). The desired result can be acquired by substituting $\|\tilde{\lambda}\| = \|\lambda^*\|_2+1$ and $\|\tilde{\lambda}\|^2 = (\|\lambda^*\|_2+1)^2$ into the above inequality. This completes the proof. ∎

# B Proof of Results in Section 3

## B.1 Proof of Lemma 3

First, we decompose $(\lambda-\lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t})$ that

$$\begin{aligned}
&(\lambda-\lambda_t)^\top \mathcal{L}_G(x_t, v_{1:2,t}) \\
&\quad= (\lambda-\lambda_t)^\top \Big(\mathcal{L}_G(x_t, v_{1:2,t}) - \mathcal{L}_G(x_{t-1}, v_{1:2,t})\Big) + (\lambda-\lambda_t)^\top \mathcal{L}_G(x_{t-1}, v_{1:2,t}) \\
&\quad= \underbrace{\langle v_{2,t}v_{1,t}(\lambda-\lambda_t), x_t - x_{t-1}\rangle}_{T_{1,t}} + \underbrace{(\lambda-\lambda_t)^\top[\mathcal{L}_G(x_{t-1}, v_{1:2,t}) - H_t]}_{T_{2,t}} + \underbrace{(\lambda-\lambda_t)^\top H_t}_{T_{3,t}}.
\end{aligned} \tag{55}$$

We now provide bounds for the three terms $T_{1,t}, T_{2,t}$, and $T_{3,t}$. First, for $T_{1,t}$, by the fact that $\langle a, b\rangle \le \frac{5\|a\|^2}{2\eta} + \frac{\eta\|b\|^2}{10}$ for any vectors $a, b$, we have

$$\begin{aligned}
\sum_{t=1}^K \mathbb{E}[T_{1,t}] &= \sum_{t=1}^K \mathbb{E}[(x_t - x_{t-1})^\top v_{2,t}v_{1,t}(\lambda-\lambda_t)] \\
&\le \sum_{t=1}^K \frac{5\mathbb{E}[\|v_{2,t}v_{1,t}(\lambda-\lambda_t)\|^2]}{2\eta} + \sum_{t=1}^K \frac{\eta\mathbb{E}[\|x_t - x_{t-1}\|^2]}{10} \\
&\le \sum_{t=1}^K \frac{5C_{g_1}^2 C_{g_2}^2 \mathbb{E}[\|\lambda-\lambda_t\|^2]}{2\eta} + \sum_{t=1}^K \frac{\eta\mathbb{E}[\|x_t - x_{t-1}\|^2]}{10},
\end{aligned} \tag{56}$$

where the last inequality holds by Assumption 4 (d)-(e) that $\|v_{1,t}\|^2 \le C_{g_1}^2$ and $\|v_{2,t}\|^2 \le C_{g_2}^2$. Second, we consider $T_{2,t}$ and denote by $\tilde{\Delta}_{2,t} = \mathcal{L}_G(x_{t-1}, v_{1:2,t}) - H_t$. Given $x_{t-1}$, we observe from Algorithm 2 that $H_t$ is conditionally independent of $\lambda_{t-1}$. Together with Lemma 2 (a) that $H_t$ is also an unbiased estimator for $\mathcal{L}_G(x_{t-1}, v_{1:2,t})$, we have $\mathbb{E}[\lambda_{t-1}^\top \tilde{\Delta}_{2,t}] = 0$, which further implies that

$$\begin{aligned}
\mathbb{E}[T_{2,t}] &= \mathbb{E}[\lambda^\top \tilde{\Delta}_{2,t}] - \mathbb{E}[\tilde{\Delta}_{2,t}^\top \lambda_{t-1}] + \mathbb{E}[(\lambda_{t-1}-\lambda_t)^\top \tilde{\Delta}_{2,t}] \\
&= \mathbb{E}[\lambda^\top \tilde{\Delta}_{2,t}] + \mathbb{E}[(\lambda_{t-1}-\lambda_t)^\top \tilde{\Delta}_{2,t}].
\end{aligned}$$

By setting $M_\lambda = \|\lambda\|$ and recalling Lemma 2 (b) that $\mathrm{Var}(\tilde{\Delta}_{2,t}) \le \sigma_H^2$, we further have

$$\begin{aligned}
\sum_{t=1}^K \mathbb{E}[T_{2,t}] &= \sum_{t=1}^K \mathbb{E}\Big[\lambda^\top \tilde{\Delta}_{2,t}\Big] + \sum_{t=1}^K \mathbb{E}\Big[(\lambda_{t-1}-\lambda_t)^\top \tilde{\Delta}_{2,t}\Big] \\
&\le \sqrt{K} M_\lambda \sigma_H + \sum_{t=1}^K \mathbb{E}\Big[(\lambda_{t-1}-\lambda_t)^\top \tilde{\Delta}_{2,t}\Big] \\
&\le \sqrt{K} M_\lambda \sigma_H + \sum_{t=1}^K \Big(\frac{\alpha\mathbb{E}[\|\lambda_{t-1}-\lambda_t\|^2]}{4} + \frac{\sigma_H^2}{\alpha}\Big),
\end{aligned} \tag{57}$$

where the first inequality holds by Lemma 5 in Appendix Section A, and the last inequality holds by the fact $\langle a, b\rangle \leq \frac{\|a\|^2}{4} + \|b\|^2$ for any vectors $a, b$.

Finally, for $T_{3,t}$, by applying the three-point Lemma 4 to the update rule that

$$\lambda_t \in \operatorname*{argmax}_{\lambda \in \mathbb{R}_+^m} \left\{ \lambda^\top H_t - \frac{\alpha}{2}\|\lambda_{t-1} - \lambda\|^2 \right\},$$

we have that for any $\lambda \in \mathbb{R}_+^m$,

$$(\lambda - \lambda_t)^\top H_t = -(\lambda_t - \lambda)^\top H_t \leq \frac{\alpha}{2}\|\lambda_{t-1} - \lambda\|^2 - \frac{\alpha}{2}\|\lambda_{t-1} - \lambda_t\|^2 - \frac{\alpha}{2}\|\lambda_t - \lambda\|^2,$$

which implies

$$\mathbb{E}[T_{3,t}] \leq \sum_{t=1}^{K} \left( \frac{\alpha}{2}\|\lambda_{t-1} - \lambda\|^2 - \frac{\alpha}{2}\|\lambda_{t-1} - \lambda_t\|^2 - \frac{\alpha}{2}\|\lambda_t - \lambda\|^2 \right). \tag{58}$$

Summing (55) over $t = 1, \cdots, K$, taking expectations, and plugging (56), (57), and (58) in, we obtain that for all $\lambda \in \mathbb{R}_+^m$,

$$\begin{aligned}
&\mathbb{E}\Big[ \sum_{t=1}^{K} (\lambda - \lambda_t)^\top \mathcal{L}_G(x_t, v_{1,t}, v_{2,t}) \Big] \\
&\leq \frac{\alpha}{2}\mathbb{E}[\|\lambda_0 - \lambda\|^2] - \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] - \sum_{t=1}^{K} \frac{\alpha}{2}\mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2] + \sqrt{K} M_\lambda \sigma_H \\
&\quad + \sum_{t=1}^{K} \left( \frac{\alpha \mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{\sigma_H^2}{\alpha} + \frac{5 C_{g_1}^2 C_{g_2}^2 \mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10} \right) \\
&\leq \frac{\alpha}{2}\|\lambda_0 - \lambda\|^2 - \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda\|^2] + \sqrt{K} M_\lambda \sigma_H \\
&\quad + \sum_{t=1}^{K} \left( \frac{\sigma_H^2}{\alpha} - \frac{\alpha \mathbb{E}[\|\lambda_{t-1} - \lambda_t\|^2]}{4} + \frac{5 C_{g_1}^2 C_{g_2}^2 \mathbb{E}[\|\lambda - \lambda_t\|^2]}{2\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10} \right),
\end{aligned}$$

which completes the proof. ∎

## B.2 Proof of Theorem 3

We provide bound on each term after decomposing (28). Consider any integer $K$ such that $K \leq N$. First, by Lemma 10 in Appendix Section B, we obtain

$$\begin{aligned}
&\sum_{t=1}^{K} \mathbb{E}\left[ \lambda^\top \Big( \mathcal{L}_G(x_t, v_{1:2}) - \mathcal{L}_G(x_t, v_{1:2,t}) \Big) \right] \\
&\quad \leq \sum_{t=1}^{K} \frac{\eta}{10} \mathbb{E}\Big[ \|x_t - x_{t-1}\|^2 \Big] + \frac{40 K C_{g_1}^2 C_{g_2}^2 M_\lambda^2}{\eta} + 2\sqrt{K} L_{g_1} \sigma_{g_2}^2 M_\lambda + 3\sqrt{K} C_{g_1} \sigma_{g_2} M_\lambda.
\end{aligned} \tag{59}$$

Second, by Lemma 11 in Appendix Section B, we have

$$\begin{aligned}
&\sum_{t=1}^{K}\mathbb{E}\Big[\mathcal{L}_F(x_t,\pi_{1:2,t})-\mathcal{L}_F(x^*,\pi_{1:2,t})+\lambda_t^\top\Big(\mathcal{L}_G(x_t,v_{1:2,t})-\mathcal{L}_G(x^*,v_{1:2,t})\Big)\Big]\\
&\quad\le\frac{\eta}{2}\|x_0-x^*\|^2-\sum_{t=1}^{K}\frac{3\eta}{10}\mathbb{E}\Big[\|x_{t-1}-x_t\|^2\Big]+\sum_{t=1}^{K}\frac{5C_{f_1}^2C_{f_2}^2}{2\eta}\\
&\qquad+\sum_{t=1}^{K}\frac{\alpha}{4}\mathbb{E}\Big[\|\lambda_t-\lambda_{t-1}\|^2\Big]+\sum_{t=1}^{K}\frac{D_X^2C_{g_1}^2C_{g_2}^2}{\alpha}+\sum_{t=1}^{K}\frac{5C_{g_1}^2C_{g_2}^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta}.
\end{aligned}\tag{60}$$

Meanwhile, Lemma 6 in Appendix Section A.2 implies that

$$\begin{aligned}
&\sum_{t=1}^{K}\mathbb{E}\Big[\mathcal{L}_F(x_t,\pi_{1:2})-\mathcal{L}_F(x_t,\pi_{1:2,t})\Big]\\
&\quad\le\sum_{t=1}^{K}\frac{\eta}{10}\mathbb{E}\Big[\|x_t-x_{t-1}\|^2\Big]+\frac{40KC_{f_1}^2C_{f_2}^2}{\eta}+2\sqrt{K}L_{f_1}\sigma_{f_2}^2+3\sqrt{K}C_{f_1}\sigma_{f_2}.
\end{aligned}\tag{61}$$

Finally, by Lemma 3 and the assumption that $\|\lambda\|\le M_\lambda$, we have

$$\begin{aligned}
&\mathbb{E}\Big[\sum_{t=1}^{K}(\lambda-\lambda_t)^\top\mathcal{L}_G(x_t,v_{1:2,t})\Big]+\frac{\alpha}{2}\mathbb{E}\Big[\|\lambda_K-\lambda\|^2\Big]\\
&\quad\le\sum_{t=1}^{K}\Big(\frac{\sigma_H^2}{\alpha}-\frac{\alpha\mathbb{E}[\|\lambda_{t-1}-\lambda_t\|^2]}{4}+\frac{5C_{g_1}^2C_{g_2}^2\mathbb{E}[\|\lambda-\lambda_t\|^2]}{2\eta}+\frac{\eta\mathbb{E}[\|x_t-x_{t-1}\|^2]}{10}\Big)\\
&\qquad+\frac{\alpha}{2}\mathbb{E}\Big[\|\lambda_0-\lambda\|^2\Big]+\sqrt{K}M_\lambda\sigma_H.
\end{aligned}\tag{62}$$

By substituting (59), (60), (61), and (62) into (28), and omitting some algebras, we obtain

$$\begin{aligned}
&\sum_{t=1}^{K}\mathbb{E}\Big(\mathcal{L}(x_t,\lambda,\pi_{1:2},v_{1:2})-\mathcal{L}(x^*,\lambda_t,\pi_{1:2,t},v_{1:2,t})\Big)+\frac{\alpha}{2}\mathbb{E}[\|\lambda_K-\lambda\|^2]\\
&\quad\le\frac{40KC_{g_1}^2C_{g_2}^2M_\lambda^2}{\eta}+2\sqrt{K}L_{g_1}\sigma_{g_2}^2M_\lambda+3\sqrt{K}C_{g_1}\sigma_{g_2}M_\lambda+\frac{\eta}{2}\|x_0-x^*\|^2\\
&\qquad+\frac{95KC_{f_1}^2C_{f_2}^2}{2\eta}+\frac{K}{\alpha}(D_X^2C_{g_1}^2C_{g_2}^2+\sigma_H^2)+\frac{\alpha}{2}\mathbb{E}[\|\lambda_0-\lambda\|^2]+\sqrt{K}M_\lambda\sigma_H\\
&\qquad+2\sqrt{K}L_{f_1}\sigma_{f_2}^2+3\sqrt{K}C_{f_1}\sigma_{f_2}+\sum_{t=1}^{K}\frac{5C_{g_1}^2C_{g_2}^2}{2\eta}\Big(\mathbb{E}[\|\lambda-\lambda_t\|^2]+\mathbb{E}[\|\lambda_{t-1}\|^2]\Big).
\end{aligned}$$

This completes the proof. ∎

**Lemma 10** *Suppose Assumptions 1, 3, and 4 hold, and Algorithm 2 generates* $\{(x_t,\lambda_t,\pi_{1:2,t},v_{1:2,t})\}_{t=1}^N$ *by setting* $\tau_t=\rho_t=t/2$, $\eta_t=\eta$, *and* $\alpha_t=\alpha$ *for* $t=1,2,\cdots,N$. *Letting* $\lambda\in\mathbb{R}_+^m$ *be a bounded random variable satisfying* $\|\lambda\|\le M_\lambda$ *uniformly, then for any* $K\le N$, *we have*

$$\begin{aligned}
&\sum_{t=1}^{K}\mathbb{E}\Big[\lambda^\top\Big(\mathcal{L}_G(x_t,v_{1:2})-\mathcal{L}_G(x_t,v_{1:2,t})\Big)\Big]\\
&\quad\le\sum_{t=1}^{K}\frac{\eta}{10}\mathbb{E}[\|x_t-x_{t-1}\|^2]+\frac{60KC_{g_1}^2C_{g_2}^2M_\lambda^2}{\eta}+2\sqrt{K}L_{g_1}M_\lambda\sigma_{g_2}^2+4\sqrt{K}M_\lambda C_{g_1}\sigma_{g_2}.
\end{aligned}$$

*Proof:* This result is similar to Lemma 6 with an additional nonnegative (random) variable $\lambda$. We start with the following decomposition that

$$\begin{aligned}
&\lambda^\top(\mathcal{L}_G(x_t, v_{1:2}) - \mathcal{L}_G(x_t, v_{1:2,t})) \\
&\quad = \lambda^\top(\mathcal{L}_{g_1}(x_t, v_1, v_2) - \mathcal{L}_{g_1}(x_t, v_1, v_{2,t})) + \lambda^\top(\mathcal{L}_{g_1}(x_t, v_1, v_{2,t}) - \mathcal{L}_{g_1}(x_t, v_{1,t}, v_{2,t})).
\end{aligned}$$

First, we recall (25) that $v_{2,t} \in \operatorname{argmax}_{v_2 \in \mathcal{V}_2}\{v_2^\top x_{t-1} - g_2^*(v_2)\}$. By Assumption 4 (g) that for each compositional constraint $G^{(j)} = g_1^{(j)} \circ g_2^{(j)}(x)$, the outer-level function $g_1^{(j)}$ is monotone nondecreasing for non-affine inner-level function $g_2^{(j)}$, we obtain

$$\mathbb{E}\bigg[\Big\langle (v_2 - v_{2,t})^\top x_{t-1} - g_2^*(v_2) + g_2^*(v_{2,t}), v_1 \Big\rangle\bigg] \le 0.$$

Because $\lambda \ge 0$, by following (36) in Lemma 6, we have

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\bigg[\lambda^\top\Big(\mathcal{L}_{g_1}(x_t, v_1, v_2) - \mathcal{L}_{g_1}(x_t, v_1, v_{2,t})\Big)\bigg] \le \sum_{t=1}^K \mathbb{E}[\langle (v_2 - v_{2,t})v_1\lambda, x_t - x_{t-1}\rangle] \\
&\quad \le \sum_{t=1}^K \bigg[\frac{10\mathbb{E}[\|(v_2 - v_{2,t})v_1\lambda\|^2]}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2]\bigg] \\
&\quad \le \sum_{t=1}^K \bigg[\frac{40C_{g_1}^2 C_{g_2}^2 M_\lambda^2}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2]\bigg],
\end{aligned}$$

where the last inequality uses the fact that $\|\lambda\| \le M_\lambda$. Second, by Proposition 15 in Zhang and Lan (2024), the $L_{g_1}$ smoothness of $g_1$ implies $\lambda^\top D_{g_1^*}(v_1; \bar{v}_1) \ge \frac{\|\lambda^\top(v_1 - \bar{v}_1)\|^2}{2\|\lambda\|L_{g_1}} \ge \frac{\|\lambda^\top(v_1 - \bar{v}_1)\|^2}{2M_\lambda L_{g_1}}$, which further yields

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\left[\lambda^\top(v_{1,t} - v_{1,t-1})^\top(g_2(x_{t-1}, \zeta_{2,t-1}^1) - g_2(x_{t-1})) - \rho_t \lambda^\top D_{g_1^*}(v_{1,t}; v_{1,t-1})\right] \\
&\quad \le \sum_{t=2}^K \frac{M_\lambda L_{g_1}\sigma_{g_2}^2}{\rho_t} + \sqrt{\mathbb{E}[\|\lambda\|^2\|v_{1,1} - v_{1,0}\|^2]}\sqrt{\mathbb{E}\Big[\Big\|g_2(x_0, \zeta_{2,0}^1) - g_2(x_0)\Big\|^2\Big]} \\
&\quad \le 2L_{g_1}M_\lambda\sigma_{g_2}^2 \log K + 2M_\lambda C_{g_1}\sigma_{g_2} \\
&\quad \le 2\sqrt{K}L_{g_1}M_\lambda\sigma_{g_2}^2 + 2\sqrt{K}M_\lambda C_{g_1}\sigma_{g_2}.
\end{aligned}$$

By substituting the above bound into Proposition 6 of Zhang and Lan (2024), we further obtain that

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\bigg[\lambda^\top\Big(\mathcal{L}_{g_1}(x_t, v_1, v_{2,t}) - \mathcal{L}_{g_1}(x_t, v_{1,t}, v_{2,t})\Big)\bigg] \\
&\quad \le \sum_{t=1}^K \bigg[\frac{20C_{g_1}^2 C_{g_2}^2 M_\lambda^2}{\eta} + \frac{\eta}{20}\mathbb{E}[\|x_t - x_{t-1}\|^2]\bigg] + 2\sqrt{K}L_{g_1}M_\lambda\sigma_{g_2}^2 + 4\sqrt{K}M_\lambda C_{g_1}\sigma_{g_2}.
\end{aligned}$$

The desired result then follows by combining the preceding inequalities. ∎

**Lemma 11** *Suppose Assumptions 1, 3, and 4 hold, and Algorithm 2 generates* $\{(x_t, \lambda_t, \pi_t, v_t)\}_{t=1}^N$ *by setting* $\rho_t = \tau_t = \frac{t}{2}$, $\eta_t = \eta$, *and* $\alpha_t = \alpha$ *for* $t \le N$. *Then for any integer* $K \le N$, *we have*

$$\begin{aligned}
&\sum_{t=1}^K \mathbb{E}\Big[\mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big(\mathcal{L}_G(x_t, v_{1:2,t}) - \mathcal{L}_G(x^*, v_{1:2,t})\Big)\Big] \\
&\quad \le \frac{\eta}{2}\|x_0 - x^*\|^2 - \sum_{t=1}^K \frac{3\eta_{t-1}}{10}\mathbb{E}\Big[\|x_{t-1} - x_t\|^2\Big] + \sum_{t=1}^K \frac{5C_{f_1}^2 C_{f_2}^2}{2\eta} \\
&\qquad + \sum_{t=1}^K \frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \sum_{t=1}^K \frac{D_X^2 C_{g_1}^2 C_{g_2}^2}{\alpha} + \sum_{t=1}^K \frac{5C_{g_1}^2 C_{g_2}^2 \mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta}.
\end{aligned}$$

*Proof:* Recall the update rule for $x_t$ (26) that

$$x_t = \operatorname*{argmin}_{x \in \mathcal{X}} \Big\{\Big\langle \pi_{2,t}^0 \pi_{1,t}^0 + v_{2,t}^0 v_{1,t}^0 \lambda_{t-1}, x\Big\rangle + \frac{\eta}{2}\|x_{t-1} - x\|^2\Big\}.$$

By following analogous analysis in Lemma 7, and applying the three-point Lemma 4 to the above update rule, we have

$$\begin{aligned}
&\mathcal{L}_F(x_t, \pi_{1:2,t}) - \mathcal{L}_F(x^*, \pi_{1:2,t}) + \lambda_t^\top \Big(\mathcal{L}_G(x_t, v_{1:2,t}) - \mathcal{L}_G(x^*, v_{1:2,t})\Big) \\
&\quad = (x_t - x^*)^\top \pi_{2,t}\pi_{1,t} + (x_t - x^*)^\top v_{2,t} v_{1,t} \lambda_t \\
&\quad \le \frac{\eta}{2}\|x_{t-1} - x^*\|^2 - \frac{\eta}{2}\|x_{t-1} - x_t\|^2 - \frac{\eta}{2}\|x_t - x^*\| \\
&\qquad + \langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0 \pi_{1,t}^0, x_t - x^*\rangle + \tilde{\Lambda}_{1,t} + \tilde{\Lambda}_{2,t},
\end{aligned} \tag{63}$$

where

$$\tilde{\Lambda}_{1,t} = (x_t - x^*)^\top v_{2,t} v_{1,t}(\lambda_t - \lambda_{t-1}) \text{ and } \tilde{\Lambda}_{2,t} = (x_t - x^*)^\top \Big(v_{2,t} v_{1,t} - v_{2,t}^0 v_{1,t}^0\Big)\lambda_{t-1}.$$

We note that a bound for $\Big\langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0 \pi_{1,t}^0, x_t - x^*\Big\rangle$ has been provided in (39). Specifically, we have

$$\begin{aligned}
\mathbb{E}\Big[\sum_{t=1}^K \Big\langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0 \pi_{1,t}^0, x_t - x^*\Big\rangle\Big] &= \mathbb{E}\Big[\sum_{t=1}^K \Big\langle \pi_{2,t}\pi_{1,t} - \pi_{2,t}^0 \pi_{1,t}^0, x_t - x_{t-1}\Big\rangle\Big] \\
&\le \sum_{t=1}^K \Big(\frac{5C_{f_1}^2 C_{f_2}^2}{2\eta} + \frac{\eta \mathbb{E}[\|x_t - x_{t-1}\|^2]}{10}\Big).
\end{aligned} \tag{64}$$

To bound $\tilde{\Lambda}_{1,t}$, by following the analysis of (40), we have

$$\begin{aligned}
\mathbb{E}[\tilde{\Lambda}_{1,t}] &\le \frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \frac{1}{\alpha}\mathbb{E}[\|(x_t - x^*)^\top v_{2,t} v_{1,t}\|^2] \\
&\le \frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2}{\alpha}.
\end{aligned} \tag{65}$$

Meanwhile, we observe from Algorithm 2 that $\mathbb{E}[x_{t-1}^\top (v_{2,t}^0 v_{1,t}^0 - v_{2,t} v_{1,t})\lambda_{t-1}] = 0$. By following the

analysis of (41), we have

$$\begin{aligned}\mathbb{E}[\sum_{t=1}^{K}\tilde{\Lambda}_{2,t}] &= \sum_{t=1}^{K}\mathbb{E}[(x_t - x_{t-1})^\top (v_{2,t}v_{1,t} - v_{2,t}^0 v_{1,t}^0)\lambda_{t-1}] \\ &\leq \sum_{t=1}^{K}\mathbb{E}[\frac{5\|(v_{2,t}^0 v_{1,t}^0 - v_{2,t}v_{1,t})\lambda_{t-1}\|^2}{2\eta}] + \sum_{t=1}^{K}\frac{\eta\mathbb{E}[\|x_t - x_{t-1}\|^2]}{10} \\ &\leq \sum_{t=1}^{K}\frac{5C_{g_1}^2 C_{g_2}^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} + \sum_{t=1}^{K}\frac{\eta\mathbb{E}[\|x_t - x_{t-1}\|^2]}{10},\end{aligned} \tag{66}$$

where the last inequality uses the fact that $\lambda_{t-1}$ and $v_{2,t}^0 v_{1,t}^0 - v_{2,t}v_{1,t}$ are independent. Summing (63) over $t = 1, 2, \cdots, K$, and combining (64), (65), and (66), we obtain

$$\begin{aligned}&\sum_{t=1}^{K}\mathbb{E}\Big[\mathcal{L}_F(x_t, \pi_{1,t}, \pi_{2,t}) - \mathcal{L}_F(x^*, \pi_{1,t}, \pi_{2,t}) + \lambda_t^\top\Big(\mathcal{L}_G(x_t, v_t) - \mathcal{L}_G(x^*, v_t)\Big)\Big] \\ &\leq \frac{\eta}{2}\|x_0 - x^*\|^2 + \sum_{t=1}^{K}\Big(\frac{\eta}{10} + \frac{\eta}{10} - \frac{\eta}{2}\Big)\mathbb{E}\Big[\|x_{t-1} - x_t\|^2\Big] + \sum_{t=1}^{K}\frac{5C_{f_1}^2 C_{f_2}^2}{2\eta} \\ &\quad + \sum_{t=1}^{K}\frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \sum_{t=1}^{K}\frac{D_X^2 C_{g_1}^2 C_{g_2}^2}{\alpha} + \sum_{t=1}^{K}\frac{5C_{g_1}^2 C_{g_2}^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta} \\ &= \frac{\eta}{2}\|x_0 - x^*\|^2 - \sum_{t=1}^{K}\frac{3\eta}{10}\mathbb{E}\Big[\|x_{t-1} - x_t\|^2\Big] + \sum_{t=1}^{K}\frac{5C_{f_1}^2 C_{f_2}^2}{2\eta} \\ &\quad + \sum_{t=1}^{K}\frac{\alpha}{4}\mathbb{E}[\|\lambda_t - \lambda_{t-1}\|^2] + \sum_{t=1}^{K}\frac{D_X^2 C_{g_1}^2 C_{g_2}^2}{\alpha} + \sum_{t=1}^{K}\frac{5C_{g_1}^2 C_{g_2}^2\mathbb{E}[\|\lambda_{t-1}\|^2]}{2\eta}.\end{aligned}$$

This completes the proof. ∎

### B.3 Proof of Proposition 2

For the feasible point $(x^*, \lambda, \pi_{1:2}, v_{1:2})$, we set $\pi_{1:2} = \pi_{1:2}^*$, $v_{1:2} = v_{1:2}^*$, and $\lambda = \lambda^*$ as defined in (24). By applying Theorem 3 with $M_{\lambda^*} = \|\lambda^*\|$, we obtain

$$\begin{aligned}&\sum_{t=1}^{K}\mathbb{E}\Big(\mathcal{L}(x_t, \lambda^*, \pi_{1:2}^*, v_{1:2}^*) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\Big) + \frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\ &\leq \frac{40KC_{g_1}^2 C_{g_2}^2\|\lambda^*\|}{\eta} + 2\sqrt{K}L_{g_1}\sigma_{g_2}^2\|\lambda^*\| + 3\sqrt{K}C_{g_1}\sigma_{g_2}\|\lambda^*\| + \frac{\eta}{2}\|x_0 - x^*\|^2 \\ &\quad + \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \frac{K}{\alpha}(D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2) + \frac{\alpha}{2}\|\lambda_0 - \lambda^*\|^2 + \sqrt{K}\|\lambda^*\|\sigma_H \\ &\quad + 2\sqrt{K}L_{f_1}\sigma_{f_2}^2 + 3\sqrt{K}C_{f_1}\sigma_{f_2} + \sum_{t=1}^{K}\frac{5C_{g_1}^2 C_{g_2}^2}{2\eta}\Big(\mathbb{E}[\|\lambda^* - \lambda_t\|^2] + \mathbb{E}[\|\lambda_{t-1}\|^2]\Big).\end{aligned}$$

We note that $\mathcal{L}(x_t, \lambda^*, \pi^*_{1:2}, v^*_{1:2}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1:2,t}) \geq 0$ by the definition of a saddle point (24). Thus, we have

$$\begin{aligned}
&\frac{\alpha}{2}\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\
&\leq \frac{40KC_{g_1}^2 C_{g_2}^2 \|\lambda^*\|}{\eta} + \frac{\eta}{2}\|x_0 - x^*\|^2 + \frac{95KC_{f_1}^2 C_{f_2}^2}{2\eta} + \frac{K}{\alpha}(D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2) \\
&\quad + \frac{\alpha}{2}\|\lambda_0 - \lambda^*\|^2 + \sum_{t=1}^{K} \frac{5C_{g_1}^2 C_{g_2}^2}{2\eta}\Big(\mathbb{E}[\|\lambda^* - \lambda_t\|^2] + \mathbb{E}[\|\lambda_{t-1}\|^2]\Big) + \sqrt{K}Q,
\end{aligned}$$

where $Q = 2L_{g_1}\sigma_{g_2}^2\|\lambda^*\| + 3C_{g_1}\sigma_{g_2}\|\lambda^*\| + \|\lambda^*\|\sigma_H + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}$ is defined in (29). By using the fact that $\|\lambda_t\|^2 \leq 2\|\lambda_t - \lambda^*\|^2 + 2\|\lambda^*\|^2$, setting $\alpha = 2\sqrt{N}$ and $\eta = \frac{15C_{g_1}^2 C_{g_2}^2\sqrt{N}}{2}$ , and dividing both sides of the above inequality by $\sqrt{N}$, we obtain

$$\begin{aligned}
&\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \\
&\leq \frac{16K\|\lambda^*\|}{3N} + \frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \frac{19KC_{f_1}^2 C_{f_2}^2}{3C_{g_1}^2 C_{g_2}^2 N} + \frac{K}{2N}(D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2) \\
&\quad + \|\lambda_0 - \lambda^*\|^2 + \frac{1}{3N}\sum_{t=1}^{K}\Big(\mathbb{E}[\|\lambda^* - \lambda_t\|^2] + 2\mathbb{E}[\|\lambda_{t-1} - \lambda^*\|^2] + 2\|\lambda^*\|^2\Big) + \frac{\sqrt{K}Q}{\sqrt{N}} \\
&= R_K + \frac{1}{3N}\mathbb{E}[\|\lambda^* - \lambda_K\|^2] + \frac{1}{N}\sum_{t=1}^{K}\mathbb{E}[\|\lambda_{t-1} - \lambda^*\|^2],
\end{aligned}$$

where

$$R_K = \frac{K}{N}\Big(6\|\lambda^*\| + \frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2}^2} + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2}\Big) + \frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \|\lambda_0 - \lambda^*\|^2 + \frac{\sqrt{K}Q}{\sqrt{N}}.$$

By noting that $R_K \leq R$ defined in (29), the above inequality further implies that

$$\Big(1 - \frac{1}{3N}\Big)\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq R + \sum_{t=1}^{K}\frac{\mathbb{E}[\|\lambda_{t-1} - \lambda^*\|^2]}{N}.$$

Assuming $N \geq 2$, for all $K \leq N$, we have

$$\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq 2R + \sum_{t=0}^{K-1}\frac{2}{N}\cdot\mathbb{E}[\|\lambda_t - \lambda^*\|^2].$$

Finally, by applying Lemma 1, we recursively bound $\mathbb{E}[\|\lambda_K - \lambda^*\|^2]$ by

$$\mathbb{E}[\|\lambda_K - \lambda^*\|^2] \leq 2Re^2$$

for all $K \leq N$, which completes the proof. ∎

## B.4 Proof of Theorem 4

We set $\lambda_0 = 0$ throughout our analysis. We first consider the objective optimality gap $F(\bar{x}_N) - F(x^*)$. We denote by

$$\hat{\pi}_2 \in \partial f_2(\bar{x}_N),\ \hat{\pi}_1 = \nabla f_1\Big(f_2(\bar{x}_N)\Big),\ \hat{v}_2 \in \partial g_2(\bar{x}_N),\ \text{and}\ \ \hat{v}_1 = \nabla g_1(g_2(\bar{x}_N)).$$

Setting the feasible point as $(x^*, 0, \hat{\pi}_{1:2}, \hat{v}_{1:2})$ and following Theorem 2 with $M_\lambda = 0$ for $\lambda = 0$, $\alpha_t = 2\sqrt{N}$, and $\eta_t = \frac{15C_{g_1}^2 C_{g_2}^2 \sqrt{N}}{2}$, we obtain that

$$\begin{aligned}
\mathbb{E}[F(\bar{x}_N)] - F(x^*) &\le \frac{1}{N}\sum_{t=1}^{N} \mathbb{E}\Big(\mathcal{L}(x_t, 0, \hat{\pi}_{1:2}, \hat{v}_{1:2}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\Big) \\
&\le \frac{1}{\sqrt{N}}\Big(\frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2 + \frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2^2}} + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2} + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}\Big) \\
&\quad + \frac{1}{N}\sum_{t=1}^{N}\frac{1}{3\sqrt{N}}\Big(\mathbb{E}[\|\lambda_t\|^2] + \mathbb{E}[\|\lambda_{t-1}\|^2]\Big).
\end{aligned}$$

We then derive the convergence rate of the optimality gap by the bound of $\|\lambda_t\|^2$ in Proposition 2 that

$$\mathbb{E}[\|\lambda_t\|^2] \le 2\|\lambda^*\|^2 + 2\mathbb{E}[\|\lambda_t - \lambda^*\|^2] \le 2\|\lambda^*\|^2 + 2Re^2,\ \text{for all}\ t = 1, 2, \cdots, N.$$

Second, we consider the feasibility residual $G(\bar{x}_N)_+$. By adopting the feasible point $(x^*, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}_{1:2})$ where $\tilde{\lambda} = (\lambda^* + 1)\frac{G(\bar{x}_N)_+}{\|G(\bar{x}_N)_+\|_2}$ and following Theorem 2 with $M_{\tilde{\lambda}} = \|\tilde{\lambda}\|$, $\alpha_t = 2\sqrt{N}$, and $\eta_t = \frac{15C_{g_1}^2 C_{g_2}^2 \sqrt{N}}{2}$, we obtain

$$\begin{aligned}
\mathbb{E}[\|G(\bar{x}_N)_+\|_2] &\le \frac{1}{N}\sum_{t=1}^{N} \mathbb{E}\Big(\mathcal{L}(x_t, \tilde{\lambda}, \hat{\pi}_{1:2}, \hat{v}_{1:2}) - \mathcal{L}(x^*, \lambda_t, \pi_{1:2,t}, v_{1:2,t})\Big) \\
&\le \frac{1}{\sqrt{N}}\Big(\frac{16\|\tilde{\lambda}\|^2}{3} + 2L_{g_1}\sigma_{g_2}^2\|\tilde{\lambda}\| + 3\sqrt{K}C_{g_1}\sigma_{g_2}\|\tilde{\lambda}\| + \frac{15C_{g_1}^2 C_{g_2}^2}{4}\|x_0 - x^*\|^2\Big) \\
&\quad + \frac{1}{\sqrt{N}}\Big(\frac{7C_{f_1}^2 C_{f_2}^2}{C_{g_1}^2 C_{g_2^2}} + \frac{D_X^2 C_{g_1}^2 C_{g_2}^2 + \sigma_H^2}{2} + \|\tilde{\lambda}\|^2 + \sigma_H\|\tilde{\lambda}\| + 2L_{f_1}\sigma_{f_2}^2 + 3C_{f_1}\sigma_{f_2}\Big) \\
&\quad + \frac{1}{N}\sum_{t=1}^{N}\frac{1}{3\sqrt{N}}\Big(\mathbb{E}[\|\tilde{\lambda} - \lambda_t\|^2] + \mathbb{E}[\|\lambda_{t-1}\|^2]\Big).
\end{aligned}$$

We obtain the convergence rate of feasibility residual $\|G(x)_+\|_2$ by using the facts that

$$\|\lambda_t\|^2 \le 2\|\lambda^*\|^2 + 2\|\lambda_t - \lambda^*\|^2 \text{ and } \|\tilde{\lambda} - \lambda_t\|^2 \le 3\|\tilde{\lambda}\|^2 + 3\|\lambda^*\|^2 + 3\|\lambda_t - \lambda^*\|^2,$$

and applying $\mathbb{E}[\|\lambda_t - \lambda^*\|^2] \le 2Re^2$ provided by Proposition 2. This completes the proof. ∎